

Reduction of cardiovascular event rate: different effects of cardiac rehabilitation in CABG and PCI patients

Non Peer-reviewed author version

HANSEN, Dominique; DENDALE, Paul; Leenders, M.; BERGER, Jan; Raskin, A.; Vaes, J. & Meeusen, R. (2009) Reduction of cardiovascular event rate: different effects of cardiac rehabilitation in CABG and PCI patients. In: ACTA CARDIOLOGICA, 64(5). p. 639-644.

DOI: 10.2143/AC.64.5.2042694

Handle: <http://hdl.handle.net/1942/10316>

08-3420 OAreview

Original article

Reduction of cardiovascular event rate: different effects of cardiac rehabilitation in CABG and PCI patients

Dominique HANSEN^{1,2}, PhD; Paul DENDALE^{1,3}, MD, PhD; Marika LEENDERS¹, MSc; Jan BERGER¹, BSc; Anita RASKIN⁴, RN; Johan VAES¹, MD; Romain MEEUSEN², PhD

¹Rehabilitation and Health Centre, Virga Jesse Hospital, Hasselt; ²Vrije Universiteit Brussel (VUB), Faculty LK, Department of Human Physiology and Sportsmedicine, Brussels; ³Hasselt University, Faculty of Medicine, Hasselt; ⁴Department of Cardiology, Virga Jesse Hospital, Hasselt, Belgium.

Running title: [Rehabilitation in PCI and CABG patients](#)

Address for correspondence:

Prof. Dr. Romain Meeusen, [PhD](#),

Vrije Universiteit Brussel (VUB),

Department of Human Physiology and Sportsmedicine, Faculty LK,

Pleinlaan 2, 1050 Brussels,

Belgium.

E-mail: rmeeusen@vub.ac.be

Received 29 August 2008; revision accepted for publication 9 March 2009.

Abstract

Objective

In coronary artery disease, the implementation of a cardiac rehabilitation (CR) programme favourably affects cardiovascular prognosis. However, it remains uncertain whether patients benefit to a similar extent from CR after coronary artery bypass graft surgery (CABG) or percutaneous coronary intervention (PCI). In this study, we have assessed whether CR is equally effective for suppressing the two-year cardiovascular event incidence after CABG or PCI.

Methods and results

194 PCI and 149 CABG patients participated in a three-month CR programme, while 245 PCI and 89 CABG patients received standard care. After the completion of CR during a two-year follow-up, data on cardiovascular risk factors, medication and cardiovascular events (repeat coronary revascularisation, acute myocardial infarction, and death) were collected from hospital files. Both CABG and PCI patients included into CR showed a significantly lower mortality, as compared to control patients (0.6% vs. 4.2%, $P < 0.05$). However, total cardiovascular disease incidence was significantly lower as a result of CR in CABG patients (4.7% vs. 14.0%, $P < 0.05$), but not in PCI patients (19.1% vs. 22.4%, $P > 0.05$).

Conclusion

When following a similar 3-month cardiac rehabilitation programme, the reduction of cardiovascular disease incidence during 2-years of follow-up is different between PCI and CABG patients.

Keywords: PCI – CABG – rehabilitation – mortality – cardiovascular disease.

Introduction

Cardiac rehabilitation lowers cardiovascular mortality and morbidity in patients who underwent coronary artery bypass graft surgery (CABG) or percutaneous coronary intervention (PCI)¹. Data indicate that especially aerobic exercise, a component of cardiac rehabilitation, is an important treatment modality for reducing cardiovascular mortality in these patients. Previous research reported an inverse relationship between peak whole-body oxygen uptake capacity (VO_{2peak}), a surrogate for aerobic exercise capacity, and mortality². In addition, improved VO_{2peak} as a result of exercise training is significantly related to reduced mortality during follow-up^{3,4}. Besides increasing VO_{2peak} , exercise training also improves traditional cardiovascular disease risk factors, blood rheology and endothelial function. It lowers low-degree inflammation, reduces coronary atherosclerosis, and facilitates coronary collateralisation⁵⁻⁹.

However, there seem to be significant differences in cardiovascular disease incidence between CABG and PCI patients, unrelated to exercise interventions. Due to the high need for repeat coronary revascularisations in PCI patients, CABG patients show a lower total cardiovascular event incidence¹⁰⁻¹³. In addition to these findings, important differences between CABG and PCI are apparent at entry of rehabilitation. As a result of median sternotomy and a longer hospitalisation period^{14,15}, CABG patients often present with a reduced pulmonary function, skeletal muscle mass loss, and a lower VO_{2peak} .

Therefore, as a result of cardiac rehabilitation, the cardiovascular disease incidence reduction might differ between CABG and PCI patients. Even though CABG and PCI patients have been investigated separately in a limited number of studies¹⁶⁻²¹, for an identical rehabilitation programme, no single study has compared the cardiovascular disease incidence reduction between those two groups.

In this study, we have investigated during two years the effects of a three-month cardiac rehabilitation programme on the cardiovascular event incidence in CABG and PCI patients.

Methods

Six hundred and seventy-seven coronary artery disease patients presented at our coronary revascularisation unit with a critical stenosis and/or acute myocardial infarction. From this group, 439 patients were revascularised with a percutaneous coronary intervention (PCI) and 238 patients with coronary artery bypass graft surgery (CABG). CABG was preferred in the case of three-vessel disease and/or reduced left ventricular contractility, diabetes mellitus, and/or the inability to achieve revascularisation by PCI. After hospital discharge, 194 PCI patients and 149 CABG patients were referred to a cardiac rehabilitation programme in the same hospital. 245 PCI and 89 CABG patients were referred from another hospital and offered standard care as there were no cardiac rehabilitation facilities. As a result, we have compared patients with and without the opportunity to rehabilitate in a multidisciplinary centre. The control and rehabilitation groups were closely matched for baseline demographic characteristics and cardiovascular disease risk factors. Follow-up included at least two clinical examinations in the first six months after revascularisation and a yearly follow-up visit afterwards. For each patient, a follow-up period of two years was analysed. In case of recurrent myocardial ischaemia, the cardiologist decided if a change of therapy or a new intervention was indicated.

No patient had a CABG or PCI before this study. Patients were excluded if CABG or PCI were not successful, or after significant complications (life-threatening haemorrhage, need for re-intervention during hospitalisation, acute renal failure). Patients with a cardiovascular event within two weeks post-PCI or four weeks post-CABG were excluded from the study. Patients with symptomatic or important co-morbidity were excluded.

This study was approved by the hospital's ethical committee.

Outcome measures

All first-count cardiovascular events (acute myocardial infarction, need for repeat coronary revascularisation, death) were collected from hospital files and from general practitioners.

Follow-up started after completion of the rehabilitation programme. Also the relationship between baseline cardiovascular risk factors and cardiovascular incidents was analysed.

Rehabilitation programme

The multidisciplinary ambulatory rehabilitation programme was started one or two weeks after hospital discharge, for a total duration of three months. It included at least 24 hospital-based exercise sessions: each week three one-hour aerobic exercise training sessions (treadmill, cycling and arm cranking). The patients trained at their ventilatory threshold level, tested by maximal ergospirometry (V-slope method).

All patients received psychological and dietary counselling. If necessary, they were advised to participate in a smoking cessation programme.

Both control and rehabilitation programme patients were advised to perform endurance exercise at home for at least 30 minutes each session, at least three days per week (from hospital discharge in the control patients, and after completion of the rehabilitation programme in the intervention patients).

Data analysis

Statistical analysis of patient characteristics, cardiovascular disease risk factors, and medication use in the 4 subgroups was performed with the SPSS-programme (version 14.0, Microsoft Windows): (i) a chi-square test for nominal data; (ii) analysis of variance (ANOVA) for ordinal data. A chi-square test compared the cardiovascular event rate between rehabilitation and control groups. In addition, odds ratios (OR) were calculated for the

rehabilitation and control groups. Logistic regression tested the relationship between the total cardiovascular event rate and all baseline risk factors in the total population. A Kaplan-Meier curve compared the total cardiovascular event rate for all four subgroups.

Results

In table 1, the baseline patient characteristics are shown. When significant differences between groups were found, the relation with the total cardiovascular event rate during follow-up is displayed. Differences between groups were found for the number of revascularised coronary arteries and plasma high-density lipoprotein levels ($P < 0.05$). However, there was no correlation with the total cardiovascular event rate ($P > 0.05$). In addition, the number of supervised exercise training sessions was significantly different between groups ($P < 0.05$) and related to the total cardiovascular event rate ($P < 0.05$). No other differences between groups were found.

Within the PCI patient subpopulations, the use of antiplatelets, ACE-inhibitors, digitalis, nitrates and statins was not different between the rehabilitation and control group. The use of anticoagulants (13% vs. 3%), beta-blockers (72% vs. 80%), and calcium antagonists (20% vs. 33%) was significantly different between the subgroups ($P < 0.05$). No significant correlation was found between the use of different medications and the total cardiovascular event rate. In the CABG patient subpopulations, between the rehabilitation and control group, the use of anticoagulants, beta-blockers, calcium antagonists, digitalis, and statins was not different. The use of antiplatelets (80% vs. 92%), ACE-inhibitors (20% vs. 7%), and nitrates (30% vs. 18%) was significantly different between groups ($P < 0.05$). No significant correlations were found between the use of medication and the total cardiovascular event rate.

In the total population, the incidence of total cardiovascular events (20.4% vs. 12.8%), acute myocardial infarction (3.9% vs. 0.9%) and death (4.2% vs. 0.6%) was significantly higher in the control group as compared to the rehabilitation group ($P < 0.05$) (see table 2). In the CABG patients, the incidence of total cardiovascular events (14.0% vs. 4.7%), acute

myocardial infarction (3.2% vs. 0.0%) and death (5.4% vs. 0.7%) was significantly higher in the control group, as compared to the rehabilitation group ($P < 0.05$). In the PCI patients, death and the need for repeat revascularisation of target coronary arteries were significantly higher in the control group (12.6% and 3.7%, respectively), as compared to the rehabilitation group (6.2% and 0.5%, respectively) ($P < 0.05$). No other differences were found between groups.

In figure 1, a Kaplan-Meier curve presents the total cardiovascular event rate in all subgroups during the follow-up period.

Discussion

In this study, the effects of a three-month cardiac rehabilitation programme in CABG and PCI patients were analysed during two years of follow-up. There are two major findings.

First, analysis of the total cardiovascular event rate showed that CABG patients benefit more from cardiac rehabilitation as compared to PCI patients, i.e. total cardiovascular event rate was not significantly lower as a result of the intervention in PCI patients.

Second, CABG and PCI patients who followed the rehabilitation programme show a significantly lower mortality as compared to the control patients. In accordance, other investigators have reported a mortality reduction as a result of exercise training in CABG and PCI patients¹⁶⁻¹⁹. Therefore, in agreement with previous studies, our findings suggest that cardiac rehabilitation has a profound impact on mortality in CABG and PCI patients. The reduction in mortality risk in these patients can be related, at least in part, to the increase in peak whole-body oxygen uptake capacity (VO_{2peak}). Previous studies have reported a correlation between changes of VO_{2peak} and mortality. Therefore, by selecting effective training modalities, the increase in VO_{2peak} might even be greater²², and contribute to larger survival benefits.

However, compared to CABG patients, total cardiovascular event rate is less reduced in PCI patients because of the high need for repeat coronary revascularisation. Because training modalities were equal in PCI and CABG patients, the difference in long-term outcome might be related to differences in post-rehabilitation lifestyle changes. After completion of the rehabilitation programme, previous studies have reported a lower physical activity level in PCI patients, as compared to CABG patients²³. This might be related to a worse illness perception in PCI patients. An increasing amount of evidence indicates that most PCI patients are not concerned about their medical condition²⁴. In recent studies, one out of three PCI patients considers their cardiac problem as nonexistent. At least 42% of PCI patients are

convinced they are completely cured^{25,26}. On the other hand, in CABG patients illness perception resulted in a higher treatment adherence²⁷. Therefore, a greater physical activity and lifestyle change promotion should be implemented in the care of PCI patients.

Besides changes in VO_{2peak} , rehabilitation may have favourable effects on blood circulation, endothelial function, collateral circulation, low-degree inflammation and coronary atherosclerosis⁻⁹. In addition, a multidisciplinary rehabilitation might also increase the awareness of health profile and lifestyle, resulting in a lower fat intake and/or increased physical activity²⁸.

From this study, an important implication emerges. Cardiac rehabilitation significantly reduces cardiovascular mortality in both CABG and PCI patients, and should therefore be a preferred intervention in both groups. However, for PCI patients, post-rehabilitation lifestyle changes should be addressed with emphasis on physical activity.

This study has some limitations. (i) We have compared groups of two different hospitals, with only one hospital having rehabilitation facilities. Our study tried to reduce bias by comparing populations undergoing CABG or PCI in one centre, by the same operators. In addition, baseline demographic and cardiovascular disease risk factors were closely matched between groups. (ii) Differences in medication have been found between groups. This should not bias the outcome results, as these differences were not in favour of one or another group. There was no correlation between the different medical regimens and cardiovascular event rate. (iii) We did not collect data on changes in daily physical activity and diet during the two-year follow-up. Changes between CABG and PCI patients might have contributed to and/or explain the different cardiovascular event reduction between subgroups. (iv) The number of training sessions –being different between CABG and PCI patients- was significantly related to the total cardiovascular event rate. However, it seems unlikely that a difference of five

training sessions would entirely contribute to the different cardiovascular event rate between CABG and PCI patients.

Conclusion

After two years of follow-up, cardiac rehabilitation significantly reduces mortality in CABG and PCI patients. However, with respect to the total cardiovascular event rate, the benefit seems to be greater in CABG patients.

Acknowledgements

We are grateful to Prof. dr. Luc Vanhees for reviewing this manuscript and offering valuable suggestions. In addition, we would like to thank dr. Didier Stroobants, dr. Benny Schoebrechts, and dr. Stephen Elshot for their kind assistance in the data collection for this study.

Sources of funding

This work was supported by an unrestricted grant from the Clinical Research Foundation Hartcentrum Hasselt.

Conflict of interest: none declared.

References

1. Clark AM, Hartling L, Vandermeer B, McAlister FA. Meta-analysis: secondary prevention programs for patients with coronary artery disease. *Ann Intern Med* 2005; **143**: 659-72.
2. Kavanagh T, Mertens DJ, Hamm L, Beyene J, Kennedy J, Corey P, Shephard RJ. Prediction of long-term prognosis in 12169 men referred for cardiac rehabilitation. *Circulation* 2002; **106**: 666-71.
3. Dorn J, Naughton J, Imamura D, Trevisan M. Results of a multicenter randomized clinical trial of exercise and long-term survival in myocardial infarction patients. The National Exercise and Heart Disease Project (NEHDP). *Circulation* 1999; **100**: 1764-9.
4. Vanhees L, Fagard R, Thijs L, Amery A. Prognostic value of training-induced change in peak exercise capacity in patients with myocardial infarcts and patients with coronary bypass surgery. *Am J Cardiol* 1995; **76**: 1014-9.
5. Erbs S, Linke A, Hambrecht R. Effects of exercise training on mortality in patients with coronary heart disease. *Coron Artery Dis* 2006; **17**: 219-25.
6. Church ST, Lavie CJ, Milani RV, Kirby GS. Improvements in blood rheology after cardiac rehabilitation and exercise training in patients with coronary heart disease. *Am Heart J* 2002; **143**: 349-55.
7. Gielen S, Schuler G, Hambrecht R. Exercise training in coronary artery disease and coronary vasomotion. *Circulation* 2001; **103**: E1-6.
8. Hambrecht R, Wolf A, Gielen S, Linke A, Hofer J, Erbs S, Schoene N, Schuler G. Effect of exercise on coronary endothelial function in patients with coronary artery disease. *N Engl J Med* 2000; **342**: 454-60.
9. Nicklas JN, You T, Pahor M. Behavioural treatments for chronic systemic inflammation: effects of dietary weight loss and exercise training. *CMAJ* 2005; **172**: 1199-1209.

10. Hoffman SN, TenBrook JA, Wolf MP, Pauker SG, Salem DN, Wong JB. A meta-analysis of randomized controlled trials comparing coronary artery bypass graft with percutaneous transluminal coronary angioplasty: one- to eight-year outcomes. *J Am Coll Cardiol* 2003; **41**: 1293-1304.
11. Boodwhani M, Rubens FD, Sellke FW, Mesana TG, Ruel M. Mortality and myocardial infarction following surgical versus percutaneous revascularisation of isolated left anterior descending artery disease: a meta-analysis. *Eur J Cardiothorac Surg* 2006; **29**: 65-70.
12. Mercado N, Wijns W, Serruys PW, Sigwart U, Flather MD, Stables RH, O'Neill WW, Rodriguez A, Lemos PA, Hueb WA, Gersh BJ, Booth J, Boersma E. One-year outcomes of coronary artery bypass graft surgery versus percutaneous coronary intervention with multiple stenting for multisystem disease: a meta-analysis of individual patient data from randomised clinical trials. *J Thorac Cardiovasc Surg* 2005; **130**: 512-9.
13. Bakhai A, Hill RA, Dundar Y, Dickson R, Walley T. Percutaneous transluminal coronary angioplasty with stents versus coronary artery bypass grafting for people with stable angina or acute coronary syndromes. *Cochrane Database Syst Rev* 2005; **25**: CD004588.
14. Lan C, Chen SY, Hsu CJ, Chiu SF, Lai JS. Improvement of cardiorespiratory function after percutaneous transluminal coronary angioplasty or coronary artery bypass grafting. *Am J Phys Med Rehabil* 2002; **81**: 336-41.
15. Shenkman Z, Shir Y, Weiss YG, Bleiberg B, Gross D. The effects of cardiac surgery on early and late pulmonary functions. *Acta Anaesthesiol Scand* 1997; **41**: 1193-9.
16. Lisspers J, Sundin O, Ohman A, Hofman-Bang C, Ryden L, Nygren A. Long-term effects of lifestyle behavior change in coronary artery disease: effects of recurrent coronary events after percutaneous coronary events. *Health Psychol* 2005; **24**: 41-8.

17. Belardinelli R, Paolini I, Cianci G, Piva R, Georgiou D, Purcaro A. Exercise training intervention after coronary angioplasty: the ETICA trial. *J Am Coll Cardiol* 2001; **37**: 1891-1900.
18. Dendale P, Berger J, Hansen D, Vaes J, Benit E, Weymans M. Cardiac rehabilitation reduces the rate of major adverse cardiac events after percutaneous coronary intervention. *Eur J Cardiovasc Nurs* 2005; **4**: 113-6.
19. Hedback B, Perk J, Hornblad M, Ohlsson U. Cardiac rehabilitation after coronary artery bypass surgery: 10-year results on mortality, morbidity and readmissions to hospital. *J Cardiovasc Risk* 2001; **8**: 153-8.
20. Kubo H, Yano K, Hirai H, Yabuki S, Machii K. Preventive effect of exercise training on recurrent stenosis after percutaneous transluminal coronary angioplasty (PCI). *Jpn Circ J* 1992; **56**: 413-21.
21. Hambrecht R, Walther C, Möbius-Winkler S, Gielen S, Linke A, Conradi K, Erbs S, Kluge R, Kendziorra K, Sabri O, Sick P, Schuler G. Percutaneous coronary angioplasty compared with exercise training in patients with stable coronary artery disease. A randomised trial. *Circulation* 2004; **109**: 1371-8.
22. Hansen D, Dendale P, Berger J, Meeusen R. Rehabilitation in cardiac patients: what do we know about training modalities? *Sports Med* 2005; **35**: 1063-84.
23. Reid RD, Morrin LI, Pipe AL, Dafoe WA, Higginson LA, Wielgosz AT, McDonald PW, Plotnikoff RC, Courneya KS, Oldridge NB, Beaton LJ, Papadakis S, Slovenic D'Angelo ME, Tulloch HE, Blanchard CM. Determinants of physical activity after hospitalization for coronary artery disease: the Tracking Exercise After Cardiac Hospitalization (TEACH) Study. *Eur J Cardiovasc Prev Rehabil* 2006; **13**: 529-37.
24. Gulanick M, Naito A. Patients' reactions to angioplasty: realistic or not? *Am J Crit Care* 1994; **3**: 368-73.

25. Campbell M, Torrance C. Coronary angioplasty: impact on risk factors and patients' understanding of the severity of their condition. *Aust J Adv Nurs* 2005; **22**: 26-31.
26. Fernandez RS, Griffiths R, Juergens C, Davidson P, Salamonson Y. Persistence of coronary risk factor status in participants 12 to 18 months after percutaneous coronary intervention. *J Cardiovasc Nurs* 2006; **21**: 379-87.
27. Worcester MU, Murphy BM, Mee VK, Roberts SB, Goble AJ. Cardiac rehabilitation programmes: predictors of non-attendance and drop-out. *Eur J Cardiovasc Prev Rehabil* 2004; **11**: 328-35.
28. Hofman-Bang C, Lisspers J, Nordlander R, Nygren A, Sundin O, Ohman A, Ryden L. Two-year results of a controlled study of residential rehabilitation for patients treated with percutaneous transluminal coronary angioplasty. *Eur Heart J* 1999; **20**: 1465-7.

Tables

Table 1. Presentation of the baseline patient characteristics in the four subgroups

	Rehabilitation		Control		Relation with MACE
	PCI	CABG	PCI	CABG	
Number	194	149	245	89	
Age (years)	63.4 ± 7.6	65.0 ± 9.0	64.2 ± 10.4	66.2 ± 8.3	
Men (%)	76.3	69.8	76.4	67.7	
<u>Cardiovascular disease risk factors</u>					
Body mass index (kg/m ²)	26.8 ± 3.3	26.9 ± 3.7	27.3 ± 4.2	26.8 ± 3.5	
Total cholesterol (mg/dl)	223 ± 41	225 ± 42	214 ± 40	217 ± 51	
HDL cholesterol (mg/dl)	51 ± 17*	50 ± 22	45 ± 12	46 ± 14	NS
LDL cholesterol (mg/dl)	144 ± 36	144 ± 41	136 ± 37	139 ± 44	
Systolic blood pressure (mm/Hg)	136 ± 19	141 ± 21	136 ± 22	138 ± 20	
Diastolic blood pressure (mm/Hg)	78 ± 11	78 ± 11	79 ± 11	80 ± 10	
Hyperglycaemia (>110 mg/dl) (%)	14.8	17.7	14.6	22.1	
Smoking (%)	28.3	29.6	34.6	24.7	
Family history of CVD (%)	54.6	64.2	56.0	58.1	
<u>Revascularised artery</u>					
Left descending artery (%)	50.0*	78.1	39.0	94.6	NS
Right ascending artery (%)	46.4*	59.2	37.8	36.6	NS
Circumflex artery (%)	30.4*	55.6	35.8	43.0	NS
Stenting (%)	79.9	x	78.8	x	
Supervised training sessions (n)	43 ± 19*	48 ± 25	0	0	<0.05

Abbreviations: CABG, coronary artery bypass graft surgery; NS, not significant; PCI, percutaneous coronary intervention.

*significantly different between four subgroups ($P < 0.05$)

Table 2. Effects of the rehabilitation programme on the **cardiovascular event rate**.

	Total population		PCI		CABG	
	Rehabilitation	Control (OR)	Rehabilitation	Control (OR)	Rehabilitation	Control (OR)
Number	343	334	194	245	149	89
Total cardiovascular event rate (%)	12.8*	20.4 (1.6)	19.1	22.4 (1.2)	4.7 [¶]	14.0 (3.0)
Acute myocardial infarction (%)	0.9*	3.9 (4.3)	1.5	4.2 (2.8)	0.0 [¶]	3.2
Repeat revascularisation with PCI (%)	10.2	12.0 (1.2)	14.9	13.8 (0.9)	4.0	6.5 (1.6)
*on target coronary artery (%)	x	x	6.2 [§]	12.6 (2.0)	x	x
Repeat revascularisation with CABG (%)	2.6	2.7 (1.0)	4.1	3.7 (0.9)	0.7	0.0
Death (%)	0.6*	4.2 (7.0)	0.5 [§]	3.7 (7.4)	0.7 [¶]	5.4 (7.7)

Abbreviations: CABG, coronary artery bypass graft surgery; OR, odds ratio; PCI, percutaneous coronary intervention

Odds ratios between rehabilitation and control groups.

*significant difference between total rehabilitation and control groups ($P < 0.05$)

[§]significant difference between PCI rehabilitation and control groups ($P < 0.05$)

[¶]significant difference between CABG rehabilitation and control groups ($P < 0.05$)

Figure legends

Figure 1. Kaplan-Meier curve presenting the total cardiovascular **event rate** in the four subgroups during a two-year follow-up.

*significantly different between control and rehabilitation group ($P < 0.05$)
MACE, major adverse cardiac event.