

## Air Pollution Stress and the Aging Phenotype: The Telomere Connection

Peer-reviewed author version

MARTENS, Dries & NAWROT, Tim (2016) Air Pollution Stress and the Aging Phenotype: The Telomere Connection. In: Current Environmental Health Reports, 3(3), p. 258-269.

DOI: 10.1007/s40572-016-0098-8

Handle: <http://hdl.handle.net/1942/23001>

**SpringerLink Header:** Early Life Environmental Health (J Sunyer, Section Editor)

**Air Pollution Stress and the Aging Phenotype: The Telomere Connection**

Dries S. Martens<sup>1</sup>, Tim S. Nawrot<sup>1,2</sup>

<sup>1</sup> Centre for Environmental Sciences, Hasselt University, 3500 Hasselt, Belgium

<sup>2</sup> Department of Public Health & Primary Care, Leuven University, 3000 Leuven, Belgium

**Corresponding author:** Tim S. Nawrot, PhD., Centre for Environmental Sciences, Hasselt University, Agoralaan gebouw D, 3590 Diepenbeek, Belgium. Phone: 0032-11 26 83 82, fax: 0032-11 26 82 99 e-mail: [tim.nawrot@uhasselt.be](mailto:tim.nawrot@uhasselt.be)

**Keywords:** Air pollution, telomere, aging, exposome

## **Abstract**

Aging is a complex physiological phenomenon. The question why some subjects grow old while remaining free from disease whereas others prematurely die remains largely unanswered. We focus here on the role of air pollution in biological aging. Hallmarks of aging can be grouped into three main categories: genomic instability, telomere attrition, and epigenetic alterations leading to altered mitochondrial function and cellular senescence. At birth, the initial telomere length of a person is largely determined by environmental factors. Telomere length shortens with each cell division and exposure to air pollution as well as low residential greens space exposure are associated with shorter telomere length. Recent studies show that the estimated effects of particulate air pollution exposure on the telomere mitochondrial axis of aging may play an important role in chronic health effects of air pollution. The exposome encompasses all exposures over the entire life. As telomeres can be considered as the cellular memories of exposure to oxidative stress and inflammation, telomere maintenance may be a proxy for assessing the “exposome”. If telomeres are causally related to the aging phenotype and environmental air pollution is an important determinant of telomere length, this might provide new avenues for future preventive strategies.

## **Introduction**

Extensive epidemiological studies support the link between ambient air pollution and adverse health outcomes, including cardiovascular and respiratory disease, both with short-term [1-3] and chronic exposure [4-8]. The WHO estimated the effects of combustion related particulate matter on life expectancy [9]. The results of this analysis indicate that current exposure to particulate matter from anthropogenic sources leads to an average loss of 8.6 months of life expectancy in Europe. The impacts vary from around 3 months in Finland to more than 13 months in Belgium. Along similar lines, cleaner air has contributed to increased life expectancy in the US [10]. A decrease of 10  $\mu\text{g}/\text{m}^3$  in the concentration of fine particulate matter was associated with an estimated increase in life expectancy of 7.3 months. Fifteen percent of the overall increase in life expectancy over a 2 decade period was attributed to reductions in air pollution. Public health impacts of nine major environmental exposures were evaluated in 6 European countries, and findings indicated that particulate air pollution clearly caused the greatest burden of disease of the nine factors investigated [11]. Based on the recent update of the Global Burden of Disease, Injuries, and Risk Factor study, air pollution is ranked 12<sup>th</sup> of a list of the most influential factors influencing health worldwide [12].

## **The Aging Phenotype**

Although aging is universal and unavoidable, the aging process does not occur in a uniform way. It is a complex phenotype responsive to a plethora of drivers. In the general population, aging represents a continuously distributed phenotype, in which genetic, behavioral and environmental factors interact with each other [13, 14]. Given the biological complexity of the aging process, there is no single, simple and reliable measure of an individual healthy aging process. Arbeeve and coworkers [15] analysed the relation between the risk of onset of “unhealthy life” (defined as the onset of cancer, cardiovascular disease, or diabetes) and longitudinal changes in body mass

index, diastolic pressure, hematocrit and pulse pressure, in the Framingham Heart Study. This process can be quantified already in young adults. Indeed, blood pressure, fasting glucose and glycated hemoglobin (HbA1C), bone mineral density, and blood lipids, appear to be predictive of biological age and of the rate of aging in younger healthy subjects [16]. Some of these factors including blood pressure [17-19], glucose [20] and diabetes risk [21] have been associated with exposure to air pollution. Further, the trajectory from healthy to unhealthy aging often comprises the microcirculation, which plays a role in diverse age related conditions such as hypertension, left ventricular dysfunction and chronic kidney disease [22]. Few studies of air pollution addressed the role of microcirculation. Among 4,607 participants of the Multi-Ethnic Study of Atherosclerosis (MESA), the retinal arteriolar diameter narrowed by 0.8  $\mu\text{m}$  in response to an interquartile increase in the 2-year exposure to  $\text{PM}_{2.5}$  (3  $\mu\text{g}/\text{m}^3$ ). This effect size was equal to a 7-year increase in age [23]. In a repeated measure study, including 84 healthy volunteers, each 10- $\mu\text{g}/\text{m}^3$  increase in  $\text{PM}_{10}$  was associated with 0.93 and 0.86  $\mu\text{m}$  decrease in the arteriolar and venular diameters, respectively [24].

Primary hallmarks of the aging process cause damage to cellular functions including genomic instability, telomere attrition, and epigenetic alterations. These are followed by critical responses including altered mitochondrial function and cellular senescence. Finally, these integrative hallmarks of the cellular aging process are possible culprits which ultimately contribute to the clinical effects of aging as seen in physiological loss of reserve, organ decline and reduced function. Air pollution might influence the aging phenotype through its action on the primary molecular hallmarks of aging.

### **Biological Underpinnings of Aging**

Aging begins at the very beginning of life, to accelerate at middle-age. The biological underpinnings of aging may begin before birth. Indeed, complications in adults often find their origin in risk factors operative in early life [13]. Some recently defined patterns of diseases that

begin in childhood include metabolic syndrome and cognitive aspects [14]. It is suggested that an adaptive response in the fetus to *in utero* exposures could result in persistent changes that influence health later in life [25].

### *Telomeres*

Telomeres are ribonucleoprotein complexes that cap the end of chromosomes and thereby protect it from degradation and end-to-end fusion to ensure genome stability and to prevent the loss of genetic information [26]. Human telomeres consists of several kb of tandem repeated TTAGGG sequences with a 3' G-rich single stranded overhang up to 200 nucleotides [27]. Telomeres shorten after each cellular division due to the end-replication problem and they are considered as a marker of biological aging [28]. During DNA replication on the lagging strand, DNA polymerase is not able to fully replicate the DNA strand, as the last RNA primer cannot be removed and fully replicated. Once telomeres reach a critical length (also known as the Hayflick limit), end-to end fusions are formed and genomic instability increases, leading to replicative senescence (mortality phase 1, M1) [29-31]. If human cells circumvent the M1 phase, more and more telomeres continue to shorten and reach a critical length, resulting in crisis (mortality phase 2, M2), characterized by end-to end fusions, chromosome breakage-fusion-bridge cycles, mitotic catastrophe and extensive genomic instability, eventually leading to cell death [32, 33]. To compensate for telomeric DNA loss, telomere length is maintained by the enzyme telomerase which is capable of adding the telomeric repeat sequences to the ends of the chromosomes [34]. Telomerase is a ribonucleoprotein containing a RNA template (TERC) and a reverse transcriptase (TERT) and is mainly active in germ, stem and immortal cells and is mainly repressed in somatic cells [35, 36]. DNA binding proteins are able to bind with telomeres to form the shelterin complex. Proteins of the shelterin complex that directly binds to the TTAGGG repeats includes the telomeric repeat binding factor 1 (TRF1) and 2 (TRF2) and protein protection of telomeres 1 (POT1). Other proteins of the shelterin complex that are connected are

the repressor activator protein 1 (RAP1), tripeptidyl peptidase 1 protein (POT1) and the TRF1 interacting nuclear factor 2 (TIN2) [37, 38]. This shelterin complex is able to affect the structure of telomeric DNA by the formation of a large duplex loop referred to as T-loop that protects the telomeres and regulates telomerase [39]. The single-stranded overhang invades into the DNA duplex of telomeric repeats and forms a displacement loop (D-loop) [40]. The amount of shelterins on telomeres can influence telomerase inhibition [37].

Telomere research in humans has explored whether telomere length provides information, over and above chronological age. Most of the large population based studies focused on leukocyte telomere lengths [41]. Leukocytes consists of the most replicative cell types, resulting in shorter telomeres compared with somatic tissues and results obtained from leukocyte telomere lengths research might not be generalized to other tissues [42]. Although leukocyte telomere length have showed to be highly correlated with other somatic tissues from the same individual such as muscle, fat, skin, synovial tissue , indicating that a clear intra-individual synchronization in telomere length exists in adults. Interestingly, Daniali *et al.* [43] showed that during adulthood the rate of telomere shortening in different tissues is equal regardless of their replicative activity. These studies suggest that leukocyte telomere length might be a proxy for telomere length in most somatic tissues.[44].

Studies among populations show that persons with shorter mean telomere length in leukocytes have increased risk for cardiovascular disease, [45-50] indices of obesity and insulin resistance [51, 47, 48, 52]. Excessive telomere shortening is associated with higher risk of age related diseases, such as heart failure [53] and cancer [46, 38]. However, whether mortality in elderly is associated with shorter telomere length is still a question with conflicting results among studies [54, 46, 55]. Telomere length is a complex trait, it is heritable (estimated between 36% and 82%) [56-60], longer in women than in men [61], and longer in offspring of older fathers [62]. At birth telomere length is highly variable among newborns [63], but as observed in adults, even at birth telomere lengths are synchronized across different cell types and tissues within the

newborn [64, 63, 65]. Telomere length during the *in utero* life and at birth sets the aging phenotype and might predict overall life expectancy. Data provided by telomere length assessment across different time-points during the entire life of zebra finches, indeed indicate that telomere length during early life predicts lifespan and these findings underline the importance of unravelling early life factors that sets telomere length [66]. During the first years of life, telomeres shorten at a much higher rate compared to adulthood [67, 68]. So in general, the main factors that influence telomere length are factors operative during the *in utero* life and/or childhood, and these early life influences on telomere length remain persistent later in life, with the least contribution during adulthood [43]. The environmental factors during *in utero* life that affect telomere length variability are largely unidentified. The natural shortening of telomeres associated with aging may be accelerated through oxidative stress and inflammation induced by environmental factors [69, 70].

### *Mitochondria*

Decreased mitochondrial function as exemplified by impaired ATP generation and increased reactive oxygen species (ROS) production are associated with aging, whereas conservation of mitochondrial function is an important mechanism of extending lifespan [71]. Mitochondria are involved in a variety of critical cell functions, including oxidative energy production, programmed cell death, growth, and redox signaling. By-products of mitochondrial electron transfer reactions in aerobic cells result in the production of reactive oxygen species (ROS), e.g. superoxide and hydrogen peroxide. Compared with nuclear DNA, mitochondrial DNA is more vulnerable to damage due to the lack of protective histones and insufficient repair capacity [72]. Thus, it has been proposed that mitochondrial damage, from any of a number of causes, might result in increased ROS that are causative agents in the development of genomic instability as observed in aging and cancer [73]. In this regard, genes that impact upon longevity have recently been characterized in *S. cerevisiae* and *C. elegans*, and the human homologs include the Sirtuin



family of protein deacetylases. Interestingly, three of the seven sirtuin proteins are localized into the mitochondria suggesting a connection between the mitochondrial sirtuins, the free radical theory of aging, and carcinogenesis [74, 75]. Based on these results it has been hypothesized that Sirt3 functions as a mitochondrial localized tumor suppressor [75]. Mitochondria have been linked with environmental interaction to an array of metabolic and age-related diseases, including cancer [76, 77], diabetes [78, 79] and cardiovascular illness [80, 81]. Endogenous mitochondrial oxidative stress is an important cardiovascular risk factor that can modulate atherogenesis by environmental inputs [82]. Mitochondrial superoxide dismutase 2 (SOD2)-specific activity declines in mice exposed to environmental tobacco smoke [83]. Until now, the focus has been on experimental research. However, biomarkers of mitochondria might be important in disease prediction and so far its association with environmental factors has barely been studied [84]. Moreover, observational studies in newborns indicate that low birth weight infants were prenatally subjected to conditions of oxidative stress and inflammation and this might be involved in vascular dysfunction [85] and atherosclerosis development and progression later in life [86].

#### *Assessment of telomere length and mitochondrial DNA content*

Different methods to assess telomere length have been extensively reviewed before [87-89]. The first available method to measure telomere lengths was by the use of terminal restriction fragment (TRF) length estimation using southern blotting. Throughout the years it became and remained the 'gold standard' to measure telomere lengths, but other methods to measure telomeres have been developed throughout recent years. TRF estimation is a robust method with low coefficients of variation, that can express average telomere lengths in absolute values (kb). Major disadvantages of this method is the high quantity of DNA that is required (up to several ug of DNA) which might be a major limitation in some epidemiological studies, besides it captures subtelomeric DNA which might bias absolute telomere length estimation [90]. In 2002 a

real-time qPCR method was described to measure telomere length, which was later adapted to a multiplex version with adapted telomere specific primers to generate a PCR product of a fixed length compared with the older version which generated PCR products of various lengths [91, 92]. Major advantages of the qPCR method over the TRF method is that it can be used for high-throughput measurements, it is less labor intensive, limited amount of DNA is required (less than 100ng), and relative low cost [90]. Disadvantages of the qPCR assay are that telomeres are expressed as average relative telomere length, by determining the ratio of telomere repeat copy number to a single-copy gene copy number (T/S ratio), and it has in general a higher coefficient of variation. Studies have shown modest to high correlations between the TRF and qPCR method ( $R^2$  up to 0.91) and both techniques are commonly used in epidemiological settings [93, 91, 92, 94]. Other methods available includes quantitative fluorescence *in situ* hybridization (Q-FISH), flow-FISH, and a PCR based method for single telomere length analysis (STELA), which are in general more labor intensive and are applied in context of specific research questions [95-97]. Mitochondrial DNA content is assessed by the use of a real-time qPCR method [98]. A ratio of a mitochondrial gene copy number to a single-copy nuclear gene copy number is determined to estimate the relative mtDNA content of human samples [99, 100].

## **Aging and Inflammation**

Long-term inflammation is a cause of aging-associated diseases [101]. A mechanistic role for immune activation and inflammation in the development of a plethora of age related disease emerged over the last decade. Vascular inflammation contributes to the pathogenesis of hypertension, heart failure and renal disease. An early step in the development of these diseases is the local production of cytokines that attract immune cells to the vascular wall. These immune cells subsequently cross the endothelial lining and migrate into the underlying tissue, where they produce an inflammatory response. Circulation levels of pro-inflammatory cytokines (e.g. TNF- $\alpha$ , IL1, IL6) are increased during the early onset of cardiovascular disease. Numerous

causes may contribute to aging-associated inflammation, such as pro-inflammatory tissue damages, a dysfunctional immune system [102], pro-inflammatory cytokines secreted by senescent cells. Mean leukocyte telomere length reflects the senescent status of circulating cells of the immune system and systemic influences on telomere maintenance in other tissues. Because of the pro-inflammatory processes instigated by immune cell senescence, telomere attrition in immune cells is relevant in inflammatory induced diseases.

### **Relation between inflammation, oxidative stress and air pollution**

Particulate air pollution can trigger multiple cellular responses in the lung such as cytotoxicity, inflammation, and mutagenesis. The adverse effects of particulate matter are not confined to the lung, but there are also extrapulmonary manifestations of exposure to particulate matter. The systemic consequences of particulate matter induced health effects may be due to a 'spill-over effect' of pulmonary inflammation or to translocation of the nano-sized particles into the circulation. Particulate matter has been described to induce oxidative stress, by the generation of reactive oxygen species (ROS). Indeed particles are able to generate ROS in different ways as reviewed by Risom and colleagues [103]. Firstly, ROS can directly be formed at the surface of particles. At the particle surface, soluble transition metals such as iron can be present that generates reactive hydroxyl radicals via a Fenton reaction ( $\text{Fe}^{2+} + \text{H}_2\text{O}_2 \rightarrow \text{Fe}^{3+} + \cdot\text{OH} + \cdot\text{OH}$ ) [104]. Higher levels of ROS can induce mitochondrial DNA damage and dysfunction leading towards more additional production of ROS by mitochondria itself. Exposure to  $\text{PM}_{2.5}$  has been associated with increased mitochondrial ROS production via the site III of the mitochondrial electron transport chain and subsequent produced ROS can activate apoptotic pathways through apoptosis signal-regulating kinase 1 (ASK1), c-Jun N-Terminal kinase (JNK) and tumor suppressor (p53) [105]. Other indirect sources of PM generated ROS are due to altered NADPH function and activation of inflammatory cells that are able to generate ROS [103].

## **The Air Pollution Telomere-Mitochondrial Connection**

### *Telomere Shortening and Oxidative Stress*

Besides cellular replication that imply constant telomere shortening, external factors and stress can interact with telomeres and may influence telomere attrition rates throughout the life course [68]. Interestingly, von Zglinicki was the first to show in 1995 experimentally, that cultivating human fibroblasts under hyperoxia conditions (represented as a state of oxidative stress) indeed shortened telomeres [106]. The underlying mechanism by which ROS induces DNA damage by which telomeres shorten is by the accumulation of single-strand nicks in telomeres which are less well repaired compared to other regions in the genome [107]. More specific, the G-rich parts of the telomere sequence (TTAGGG) and not the G-rich single stranded overhang are highly sensitive for DNA damage in human fibroblasts induced by oxidative stress conditions (after exposure to  $\text{H}_2\text{O}_2$ , an NO-generating agent and a  $\text{O}_2^-$  generating system) [108, 109]. Accelerated shortening of telomeres, and as such, senescence of cells may be an important pathway by which oxidative stress may accelerate biological aging and the resultant development of aging-related morbidity, including cardiovascular disease [47, 49].

### *The air pollution induced telomere-mitochondrial aging hypothesis*

Direct and indirect formation of ROS by air pollution as described previously, can explain a cascade by which both telomeres and mitochondria are targeted, leading to telomere shortening as well as mitochondrial dysfunction and this might explain the mechanism by which air pollution alters the aging phenotype (Fig 1).

Recently, Sahin *et al.* [71] unveiled a fascinating connection between the nuclear and mitochondrial aging processes. Telomere-deficient mice showed p53 activation which results in suppression of peroxisome proliferator-activated receptor gamma coactivator 1 alpha and beta (PGC-1) genes thereby revealing a direct link between telomere and mitochondrial biology [110]. Further evidence of the telomere-mitochondrial axis of aging was shown in sirtuin1 (SIRT1)

knock-out mice. SIRT1 functions as a metabolic sensor and its deacetylase activity is controlled by the cellular NAD<sup>+</sup> /NADH ratio [111]. SIRT1 expression was found to play a pivotal role in mitochondrial biogenesis [112, 113]. SIRT1 affects longevity in humans by influencing telomerase activity [114] and its function by inactivation of the 'guardian of the genome', TP53 [115]. In addition, SIRT1 stimulates PPARGC1A, a regulator of mitochondrial biogenesis [116]. Overexpression of SIRT1 in mice reduces the incidence of several aging related diseases, such as cardiovascular disease, metabolic disease and cancer [117].

The current accepted experimental model proposes that DNA damage to telomeres activates several signaling pathways and alters SIRT1 gene expression, which leads to mitochondrial dysfunction [110, 118]. This experimental model is in line with human observations. Among 166 elderly, telomere length and SIRT1 gene expression were found to be intermediate mechanisms between particulate matter air pollution exposure and mtDNA alterations. Formal mediation analysis indicated that the effect of long-term particulate matter air pollution on mtDNA content was mediated by SIRT1 expression [119].

#### *Current evidence of air pollution induced telomere-mitochondrial aging*

To date, epidemiological studies examining the effects of PM exposure and traffic related exposure on telomere length and mitochondrial DNA content have reported different responses after long-term or short-term exposure to PM. As the main focus here is on telomere biology, we summarized occupational and population based studies investigating the association between air pollution exposure and telomere length in Table 1 and Table 2 respectively. In several occupational studies a positive association is observed between PM exposure and telomere length. In 57 steel workers increased telomere length has been found in association with short-term (3 days of work) exposure to high levels of PM<sub>1</sub> and PM<sub>10</sub> [120]. This observation is in line with observations made by Hou *et al.* [121] in truck drivers compared to office workers in Beijing, China. In this repeated measure study, the participants were examined on two different work

days with 1 to 2 weeks interval. Exposure to PM<sub>2.5</sub> and elemental carbon during the last 24 hours were positively associated with blood telomere length [121]. In accordance with these studies at high exposure, variation in relatively low short-term exposure (1 month prior blood drawn) was also associated with a 17.5% increase in telomere length for a 5 µg/m<sup>3</sup> increase in PM<sub>2.5</sub> [119]. Although PM is hypothesized to induce oxidative stress which causes telomere shortening, the observed increase in telomere length due to short-term exposure might be explained by acute inflammatory processes that has been linked with increased telomerase activity in B-cells [122]. However, this observation could not yet be confirmed based on gene expression data of the human telomerase reverse transcriptase (*hTERT*) component of telomerase [120], and this lengthening of telomeres should be further evaluated by investigating both the involvement of telomerase and components of the shelterin complex.

Observations on long term-exposure to PM and short-term traffic related exposure in both occupational and population based studies, indicates shortening of telomeres and supports the role of PM induced oxidative stress in increased DNA damage at telomeres. Traffic officers compared with office workers had much lower leukocyte telomere length which corresponds to a telomeric year equivalence of 13 years [123]. Among 165 never-smoking men from the Normative Aging Study [124], an IQR increase in annual black carbon (0.25µg/m<sup>3</sup>) was associated with a 7.6% decrease in telomere length. Besides black carbon, it was shown in 166 elderly, that a 5µg/m<sup>3</sup> increment in annual PM<sub>2.5</sub> was associated with 16.8% decrease in telomere length, and a 25.7% decrease in mitochondrial content [119]. The estimated decrease in telomere length for a 5µg/m<sup>3</sup> higher annual PM<sub>2.5</sub> corresponded to a telomere-age equivalence loss of 4 years. Recently, in 211 newborn twins from the East Flanders Prospective Twin Survey, placental telomere lengths were inversely associated with traffic intensity within a buffer of 100m surrounding maternal residence, while placental telomeres were longer in association with maternal residential surrounding greenness [125].

Similar with the observed associations between telomere length and air pollution exposure, both positive and negative associations between short-term and chronic exposure to PM and mtDNA content have been reported. Hou *et al.* [84] found an increase in mtDNA content in relation to personal PM<sub>10</sub> and PM<sub>1</sub> in steelworkers while they found a decrease in mtDNA content in office workers and truck drivers in association with short-term PM<sub>10</sub> and elemental carbon exposure. These opposite results indicate that mtDNA may react in a different way under different environmental circumstances [99]. A study in newborns [100] revealed that a 10 µg/m<sup>3</sup> increment in PM<sub>10</sub> exposure during the last trimester of pregnancy was associated with a 17.4% decrease in placental mtDNA content. A significantly decrease in mtDNA content was observed after long-term (1 year) exposure to PM<sub>2.5</sub> whereas a significantly increase in mtDNA was observed for short-term (1 month and 1 week) exposure windows of PM<sub>2.5</sub> in an elderly population [119]. Recently, in 675 elderly men from the Normative Aging Study an increase of mtDNA was observed in association with BC exposure (5days to 28days) [126].

### **Telomeres and the Exposome**

The exposome encompasses all exposures over the entire life [127, 128]. Aside from air pollution, telomere length is associated with a broad spectrum of environmental factors including life stress [129], mediterranean diet [130], education [131, 132], smoking status [48, 52], obesity [133], sleep quality [134], childhood maltreatment [135]. Therefore, telomere maintenance may be a proxy for assessing the “exposome”. Telomere research has largely been focused on studies in adults, although, even at birth, newborn cord and placental telomere lengths have been associated with environmental factors such as prenatal stress [136, 137], maternal residential traffic exposure [125], intrauterine tobacco exposure [138], and *in utero* cadmium

exposure [139], indicating that from early-life (even intrauterine life) onwards these environmental factors change telomere biology. Since factors of the exposome are already present during the intrauterine life and may contribute to telomere length setting of newborns, this might have a large impact on later life expectations and may impact overall life quality and the development of age-related diseases as well.

## **Conclusions**

In addressing the epidemic of non-communicable disease, identification of factors that influence healthy or unhealthy aging is pivotal in prevention. Aging is a complex phenotype. Cellular aging, as exemplified by shorter telomere length, is associated with higher exposure to particulate air pollution during adulthood and in newborns with lower maternal residential greenness.

## **Acknowledgments**

This research was funded by the EU Program “Ideas” (ERC-2012-StG 310898) and by the Flemish Scientific Fund (FWO, G073315N/G.0880.13).

## **Compliance with Ethics Guidelines**

## **Conflict of Interest**

Dries S. Martens and Tim S. Nawrot declare that they have no conflict of interest.

## **Human and Animal Rights and Informed Consent**

This article does not contain any studies with human or animal subjects performed by any of the authors.



## References

Papers of particular interest, published recently, have been highlighted as:

•Of importance

••Of outstanding importance

1. Levy JJ, Hammitt JK, Spengler JD. Estimating the mortality impacts of particulate matter: what can be learned from between-study variability? *Environ Health Perspect.* 2000;108(2):109-17.
2. Nawrot TS, Perez L, Kunzli N et al. Public health importance of triggers of myocardial infarction: a comparative risk assessment. *Lancet.* 2011;377(9767):732-40.
3. Provost EB, Louwies T, Cox B et al. Short-term fluctuations in personal black carbon exposure are associated with rapid changes in carotid arterial stiffening. *Environ Int.* 2016;88:228-34.
4. Beelen R, Raaschou-Nielsen O, Stafoggia M et al. Effects of long-term exposure to air pollution on natural-cause mortality: an analysis of 22 European cohorts within the multicentre ESCAPE project. *Lancet.* 2014;383(9919):785-95.
5. Beelen R, Stafoggia M, Raaschou-Nielsen O et al. Long-term exposure to air pollution and cardiovascular mortality: an analysis of 22 European cohorts. *Epidemiology.* 2014;25(3):368-78.
6. Hamra GB, Guha N, Cohen A et al. Outdoor particulate matter exposure and lung cancer: a systematic review and meta-analysis. *Environ Health Perspect.* 2014;122(9):906-11.
7. Laden F, Schwartz J, Speizer FE et al. Reduction in fine particulate air pollution and mortality: Extended follow-up of the Harvard Six Cities study. *Am J Respir Crit Care Med.* 2006;173(6):667-72.
8. Provost EB, Madhloum N, Int Panis L et al. Carotid intima-media thickness, a marker of subclinical atherosclerosis, and particulate air pollution exposure: the meta-analytical evidence. *PLoS One.* 2015;10(5):e0127014.
9. WHO. Health Risks of Particulate Matter from Long-range Transboundary Air Pollution (World Health Organization). 2006.  
[http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0006/78657/E88189.pdf](http://www.euro.who.int/__data/assets/pdf_file/0006/78657/E88189.pdf).
10. Pope CA, 3rd, Ezzati M, Dockery DW. Fine-particulate air pollution and life expectancy in the United States. *N Engl J Med.* 2009;360(4):376-86.
11. Hanninen O, Knol AB, Jantunen M et al. Environmental burden of disease in Europe: assessing nine risk factors in six countries. *Environ Health Perspect.* 2014;122(5):439-46.
12. Collaborators GBDRF, Forouzanfar MH, Alexander L et al. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2015;386(10010):2287-323.
13. Barker DJ. Fetal origins of coronary heart disease. *BMJ.* 1995;311(6998):171-4.
14. Dietert RR, DeWitt JC, Germolec DR et al. Breaking patterns of environmentally influenced disease for health risk reduction: immune perspectives. *Environ Health Perspect.* 2010;118(8):1091-9.
15. Arbeevev KG, Ukraintseva SV, Akushevich I et al. Age trajectories of physiological indices in relation to healthy life course. *Mech Ageing Dev.* 2011;132(3):93-102.
16. Belsky DW, Caspi A, Houts R et al. Quantification of biological aging in young adults. *Proc Natl Acad Sci U S A.* 2015;112(30):E4104-10.
17. Jacobs L, Buczynska A, Walgraeve C et al. Acute changes in pulse pressure in relation to constituents of particulate air pollution in elderly persons. *Environ Res.* 2012;117:60-7.
18. Liang R, Zhang B, Zhao X et al. Effect of exposure to PM2.5 on blood pressure: a systematic review and meta-analysis. *J Hypertens.* 2014;32(11):2130-40; discussion 41.

19. Pieters N, Koppen G, Van Poppel M et al. Blood Pressure and Same-Day Exposure to Air Pollution at School: Associations with Nano-Sized to Coarse PM in Children. *Environ Health Perspect.* 2015;123(7):737-42.
20. Calderon-Garciduenas L, Franco-Lira M, D'Angiulli A et al. Mexico City normal weight children exposed to high concentrations of ambient PM<sub>2.5</sub> show high blood leptin and endothelin-1, vitamin D deficiency, and food reward hormone dysregulation versus low pollution controls. Relevance for obesity and Alzheimer disease. *Environ Res.* 2015;140:579-92.
21. Eze IC, Schaffner E, Fischer E et al. Long-term air pollution exposure and diabetes in a population-based Swiss cohort. *Environ Int.* 2014;70:95-105.
22. Struijker-Boudier HA, Heijnen BF, Liu YP et al. Phenotyping the microcirculation. *Hypertension.* 2012;60(2):523-7.
23. Adar SD, Klein R, Klein BE et al. Air Pollution and the microvasculature: a cross-sectional assessment of in vivo retinal images in the population-based multi-ethnic study of atherosclerosis (MESA). *PLoS Med.* 2010;7(11):e1000372.
24. Louwies T, Panis LI, Kicinski M et al. Retinal microvascular responses to short-term changes in particulate air pollution in healthy adults. *Environ Health Perspect.* 2013;121(9):1011-6.
25. Wild CP, Kleinjans J. Children and increased susceptibility to environmental carcinogens: evidence or empathy? *Cancer Epidemiol Biomarkers Prev.* 2003;12(12):1389-94.
26. Blackburn EH. Structure and function of telomeres. *Nature.* 1991;350(6319):569-73.
27. Wright WE, Tesmer VM, Huffman KE et al. Normal human chromosomes have long G-rich telomeric overhangs at one end. *Genes Dev.* 1997;11(21):2801-9.
28. Levy MZ, Allsopp RC, Futcher AB et al. Telomere end-replication problem and cell aging. *J Mol Biol.* 1992;225(4):951-60.
29. Hayflick L. The Limited in Vitro Lifetime of Human Diploid Cell Strains. *Exp Cell Res.* 1965;37:614-36.
30. Shay JW, Wright WE. Hayflick, his limit, and cellular ageing. *Nat Rev Mol Cell Biol.* 2000;1(1):72-6.
31. Campisi J, d'Adda di Fagagna F. Cellular senescence: when bad things happen to good cells. *Nat Rev Mol Cell Biol.* 2007;8(9):729-40.
32. Shay JW, Wright WE. Senescence and immortalization: role of telomeres and telomerase. *Carcinogenesis.* 2005;26(5):867-74.
33. Shay JW, Wright WE, Werbin H. Defining the molecular mechanisms of human cell immortalization. *Biochim Biophys Acta.* 1991;1072(1):1-7.
34. Blackburn EH, Epel ES, Lin J. Human telomere biology: A contributory and interactive factor in aging, disease risks, and protection. *Science.* 2015;350(6265):1193-8.
35. Kim NW, Piatyszek MA, Prowse KR et al. Specific association of human telomerase activity with immortal cells and cancer. *Science.* 1994;266(5193):2011-5.
36. Hiyama K, Hirai Y, Kyoizumi S et al. Activation of telomerase in human lymphocytes and hematopoietic progenitor cells. *J Immunol.* 1995;155(8):3711-5.
37. de Lange T. Shelterin: the protein complex that shapes and safeguards human telomeres. *Genes Dev.* 2005;19(18):2100-10.
38. O'Sullivan RJ, Karlseder J. Telomeres: protecting chromosomes against genome instability. *Nat Rev Mol Cell Biol.* 2010;11(3):171-81.
39. Griffith JD, Comeau L, Rosenfield S et al. Mammalian telomeres end in a large duplex loop. *Cell.* 1999;97(4):503-14.
40. Greider CW. Telomeres do D-loop-T-loop. *Cell.* 1999;97(4):419-22.
41. Aviv A, Valdes AM, Spector TD. Human telomere biology: pitfalls of moving from the laboratory to epidemiology. *Int J Epidemiol.* 2006;35(6):1424-9.

42. Sanders JL, Newman AB. Telomere length in epidemiology: a biomarker of aging, age-related disease, both, or neither? *Epidemiol Rev.* 2013;35:112-31.
43. Daniali L, Benetos A, Susser E et al. Telomeres shorten at equivalent rates in somatic tissues of adults. *Nat Commun.* 2013;4:1597.
44. Mather KA, Jorm AF, Parslow RA et al. Is telomere length a biomarker of aging? A review. *J Gerontol A Biol Sci Med Sci.* 2011;66(2):202-13.
45. Benetos A, Okuda K, Lajemi M et al. Telomere length as an indicator of biological aging: the gender effect and relation with pulse pressure and pulse wave velocity. *Hypertension.* 2001;37(2 Pt 2):381-5.
46. Cawthon RM, Smith KR, O'Brien E et al. Association between telomere length in blood and mortality in people aged 60 years or older. *Lancet.* 2003;361(9355):393-5.
47. Fitzpatrick AL, Kronmal RA, Gardner JP et al. Leukocyte telomere length and cardiovascular disease in the cardiovascular health study. *Am J Epidemiol.* 2007;165(1):14-21.
48. Nawrot TS, Staessen JA, Holvoet P et al. Telomere length and its associations with oxidized-LDL, carotid artery distensibility and smoking. *Front Biosci (Elite Ed).* 2010;2:1164-8.
49. Samani NJ, Boulton R, Butler R et al. Telomere shortening in atherosclerosis. *Lancet.* 2001;358(9280):472-3.
50. Haycock PC, Heydon EE, Kaptoge S et al. Leucocyte telomere length and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ.* 2014;349:g4227.
51. Demissie S, Levy D, Benjamin EJ et al. Insulin resistance, oxidative stress, hypertension, and leukocyte telomere length in men from the Framingham Heart Study. *Aging Cell.* 2006;5(4):325-30.
52. Valdes AM, Andrew T, Gardner JP et al. Obesity, cigarette smoking, and telomere length in women. *Lancet.* 2005;366(9486):662-4.
53. Kuznetsova T, Codd V, Brouillette S et al. Association between left ventricular mass and telomere length in a population study. *Am J Epidemiol.* 2010;172(4):440-50.
54. Bischoff C, Petersen HC, Graakjaer J et al. No association between telomere length and survival among the elderly and oldest old. *Epidemiology.* 2006;17(2):190-4.
55. Martin-Ruiz CM, Gussekloo J, van Heemst D et al. Telomere length in white blood cells is not associated with morbidity or mortality in the oldest old: a population-based study. *Aging Cell.* 2005;4(6):287-90.
56. Hjelmborg JB, Dalgard C, Moller S et al. The heritability of leucocyte telomere length dynamics. *J Med Genet.* 2015;52(5):297-302.
57. Slagboom PE, Droog S, Boomsma DI. Genetic determination of telomere size in humans: a twin study of three age groups. *Am J Hum Genet.* 1994;55(5):876-82.
58. Vasa-Nicotera M, Brouillette S, Mangino M et al. Mapping of a major locus that determines telomere length in humans. *Am J Hum Genet.* 2005;76(1):147-51.
59. Nawrot TS, Staessen JA, Gardner JP et al. Telomere length and possible link to X chromosome. *Lancet.* 2004;363(9408):507-10.
60. Andrew T, Aviv A, Falchi M et al. Mapping genetic loci that determine leukocyte telomere length in a large sample of unselected female sibling pairs. *Am J Hum Genet.* 2006;78(3):480-6.
61. Gardner M, Bann D, Wiley L et al. Gender and telomere length: systematic review and meta-analysis. *Exp Gerontol.* 2014;51:15-27.
62. Kimura M, Cherkas LF, Kato BS et al. Offspring's leukocyte telomere length, paternal age, and telomere elongation in sperm. *PLoS Genet.* 2008;4(2):e37.
63. Okuda K, Bardeguet A, Gardner JP et al. Telomere length in the newborn. *Pediatr Res.* 2002;52(3):377-81.

64. Kimura M, Gazitt Y, Cao X et al. Synchrony of telomere length among hematopoietic cells. *Exp Hematol*. 2010;38(10):854-9.
65. Youngren K, Jeanclos E, Aviv H et al. Synchrony in telomere length of the human fetus. *Hum Genet*. 1998;102(6):640-3.
66. Heideringer BJ, Blount JD, Boner W et al. Telomere length in early life predicts lifespan. *Proc Natl Acad Sci U S A*. 2012;109(5):1743-8.
67. Aubert G, Baerlocher GM, Vulto I et al. Collapse of telomere homeostasis in hematopoietic cells caused by heterozygous mutations in telomerase genes. *PLoS Genet*. 2012;8(5):e1002696.
68. Frenck RW, Jr., Blackburn EH, Shannon KM. The rate of telomere sequence loss in human leukocytes varies with age. *Proc Natl Acad Sci U S A*. 1998;95(10):5607-10.
69. Mitchell C, Hobcraft J, McLanahan SS et al. Social disadvantage, genetic sensitivity, and children's telomere length. *Proc Natl Acad Sci U S A*. 2014;111(16):5944-9.
70. von Zglinicki T. Oxidative stress shortens telomeres. *Trends Biochem Sci*. 2002;27(7):339-44.
71. Sahin E, DePinho RA. Axis of ageing: telomeres, p53 and mitochondria. *Nat Rev Mol Cell Biol*. 2012;13(6):397-404.
72. Liu P, Demple B. DNA repair in mammalian mitochondria: Much more than we thought? *Environ Mol Mutagen*. 2010;51(5):417-26.
73. Singh KK. Mitochondria damage checkpoint, aging, and cancer. *Ann N Y Acad Sci*. 2006;1067:182-90.
74. Kim HS, Patel K, Muldoon-Jacobs K et al. SIRT3 is a mitochondria-localized tumor suppressor required for maintenance of mitochondrial integrity and metabolism during stress. *Cancer Cell*. 2010;17(1):41-52.
75. Park SH, Ozden O, Jiang H et al. Sirt3, mitochondrial ROS, ageing, and carcinogenesis. *Int J Mol Sci*. 2011;12(9):6226-39.
76. Hochhauser D. Relevance of mitochondrial DNA in cancer. *Lancet*. 2000;356(9225):181-2.
77. Kuo JH, Chu YL, Yang JS et al. Cantharidin induces apoptosis in human bladder cancer TSGH 8301 cells through mitochondria-dependent signal pathways. *Int J Oncol*. 2010;37(5):1243-50.
78. Cree LM, Patel SK, Pyle A et al. Age-related decline in mitochondrial DNA copy number in isolated human pancreatic islets. *Diabetologia*. 2008;51(8):1440-3.
79. Gianotti TF, Sookoian S, Dieuzeide G et al. A decreased mitochondrial DNA content is related to insulin resistance in adolescents. *Obesity (Silver Spring)*. 2008;16(7):1591-5.
80. Ballinger SW, Patterson C, Knight-Lozano CA et al. Mitochondrial integrity and function in atherogenesis. *Circulation*. 2002;106(5):544-9.
81. Roy Chowdhury SK, Sangle GV, Xie X et al. Effects of extensively oxidized low-density lipoprotein on mitochondrial function and reactive oxygen species in porcine aortic endothelial cells. *Am J Physiol Endocrinol Metab*. 2010;298(1):E89-98.
82. Harrison CM, Pompilius M, Pinkerton KE et al. Mitochondrial oxidative stress significantly influences atherogenic risk and cytokine-induced oxidant production. *Environ Health Perspect*. 2011;119(5):676-81.
83. Knight-Lozano CA, Young CG, Burow DL et al. Cigarette smoke exposure and hypercholesterolemia increase mitochondrial damage in cardiovascular tissues. *Circulation*. 2002;105(7):849-54.
84. Hou L, Zhu ZZ, Zhang X et al. Airborne particulate matter and mitochondrial damage: a cross-sectional study. *Environ Health*. 2010;9:48.
85. Halvorsen CP, Andolf E, Hu J et al. Discordant twin growth in utero and differences in blood pressure and endothelial function at 8 years of age. *J Intern Med*. 2006;259(2):155-63.
86. Leeson CP, Kattenhorn M, Morley R et al. Impact of low birth weight and cardiovascular risk factors on endothelial function in early adult life. *Circulation*. 2001;103(9):1264-8.

87. Aubert G, Hills M, Lansdorp PM. Telomere length measurement-caveats and a critical assessment of the available technologies and tools. *Mutat Res.* 2012;730(1-2):59-67.
88. Montpetit AJ, Alhareeri AA, Montpetit M et al. Telomere length: a review of methods for measurement. *Nurs Res.* 2014;63(4):289-99.
89. Vera E, Blasco MA. Beyond average: potential for measurement of short telomeres. *Aging (Albany NY).* 2012;4(6):379-92.
90. Kimura M, Stone RC, Hunt SC et al. Measurement of telomere length by the Southern blot analysis of terminal restriction fragment lengths. *Nat Protoc.* 2010;5(9):1596-607.
91. Cawthon RM. Telomere measurement by quantitative PCR. *Nucleic Acids Res.* 2002;30(10):e47.
92. Cawthon RM. Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Res.* 2009;37(3):e21.
93. Aviv A, Hunt SC, Lin J et al. Impartial comparative analysis of measurement of leukocyte telomere length/DNA content by Southern blots and qPCR. *Nucleic Acids Res.* 2011;39(20):e134.
94. Elbers CC, Garcia ME, Kimura M et al. Comparison between southern blots and qPCR analysis of leukocyte telomere length in the health ABC study. *J Gerontol A Biol Sci Med Sci.* 2014;69(5):527-31.
95. Baird DM, Rowson J, Wynford-Thomas D et al. Extensive allelic variation and ultrashort telomeres in senescent human cells. *Nat Genet.* 2003;33(2):203-7.
96. Canela A, Vera E, Klatt P et al. High-throughput telomere length quantification by FISH and its application to human population studies. *Proc Natl Acad Sci U S A.* 2007;104(13):5300-5.
97. Rufer N, Dragowska W, Thornbury G et al. Telomere length dynamics in human lymphocyte subpopulations measured by flow cytometry. *Nat Biotechnol.* 1998;16(8):743-7.
98. Miller FJ, Rosenfeldt FL, Zhang C et al. Precise determination of mitochondrial DNA copy number in human skeletal and cardiac muscle by a PCR-based assay: lack of change of copy number with age. *Nucleic Acids Res.* 2003;31(11):e61.
99. Hou L, Zhang X, Dioni L et al. Inhalable particulate matter and mitochondrial DNA copy number in highly exposed individuals in Beijing, China: a repeated-measure study. *Part Fibre Toxicol.* 2013;10:17.
100. Janssen BG, Munters E, Pieters N et al. Placental mitochondrial DNA content and particulate air pollution during in utero life. *Environ Health Perspect.* 2012;120(9):1346-52.
101. Perry VH, Cunningham C, Holmes C. Systemic infections and inflammation affect chronic neurodegeneration. *Nat Rev Immunol.* 2007;7(2):161-7.
102. Deeks SG. HIV infection, inflammation, immunosenescence, and aging. *Annu Rev Med.* 2011;62:141-55.
103. Risom L, Moller P, Loft S. Oxidative stress-induced DNA damage by particulate air pollution. *Mutat Res.* 2005;592(1-2):119-37.
104. Ghio AJ, Carraway MS, Madden MC. Composition of air pollution particles and oxidative stress in cells, tissues, and living systems. *J Toxicol Environ Health B Crit Rev.* 2012;15(1):1-21.
105. Soberanes S, Urich D, Baker CM et al. Mitochondrial complex III-generated oxidants activate ASK1 and JNK to induce alveolar epithelial cell death following exposure to particulate matter air pollution. *J Biol Chem.* 2009;284(4):2176-86.
106. von Zglinicki T, Saretzki G, Docke W et al. Mild hyperoxia shortens telomeres and inhibits proliferation of fibroblasts: a model for senescence? *Exp Cell Res.* 1995;220(1):186-93.
107. Petersen S, Saretzki G, von Zglinicki T. Preferential accumulation of single-stranded regions in telomeres of human fibroblasts. *Exp Cell Res.* 1998;239(1):152-60.
108. Kawanishi S, Oikawa S. Mechanism of telomere shortening by oxidative stress. *Ann N Y Acad Sci.* 2004;1019:278-84.

109. von Zglinicki T, Pilger R, Sitte N. Accumulation of single-strand breaks is the major cause of telomere shortening in human fibroblasts. *Free Radic Biol Med*. 2000;28(1):64-74.
- 110.●● **Sahin E, Colla S, Liesa M et al. Telomere dysfunction induces metabolic and mitochondrial compromise. *Nature*. 2011;470(7334):359-65. These authors made a substantial contribution in unravelling the link between telomeres, mitochondria and aging.**
111. Yamamoto H, Schoonjans K, Auwerx J. Sirtuin functions in health and disease. *Mol Endocrinol*. 2007;21(8):1745-55.
112. Lagouge M, Argmann C, Gerhart-Hines Z et al. Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1alpha. *Cell*. 2006;127(6):1109-22.
113. Nemoto S, Fergusson MM, Finkel T. SIRT1 functionally interacts with the metabolic regulator and transcriptional coactivator PGC-1{alpha}. *J Biol Chem*. 2005;280(16):16456-60.
114. Narala SR, Allsopp RC, Wells TB et al. SIRT1 acts as a nutrient-sensitive growth suppressor and its loss is associated with increased AMPK and telomerase activity. *Mol Biol Cell*. 2008;19(3):1210-9.
115. Vaziri H, Dessain SK, Ng Eaton E et al. hSIR2(SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell*. 2001;107(2):149-59.
116. Aquilano K, Vigilanza P, Baldelli S et al. Peroxisome proliferator-activated receptor gamma co-activator 1alpha (PGC-1alpha) and sirtuin 1 (SIRT1) reside in mitochondria: possible direct function in mitochondrial biogenesis. *J Biol Chem*. 2010;285(28):21590-9.
117. Donmez G, Guarente L. Aging and disease: connections to sirtuins. *Aging Cell*. 2010;9(2):285-90.
118. Sahin E, Depinho RA. Linking functional decline of telomeres, mitochondria and stem cells during ageing. *Nature*. 2010;464(7288):520-8.
- 119.● **Pieters N, Janssen BG, Dewitte H et al. Biomolecular Markers Within the Core Axis of Aging and Particulate Air Pollution Exposure in the Elderly: A Cross-Sectional Study. *Environ Health Perspect*. 2015. This study integrates different aging related molecular markers in elderly in association with long-term air pollution exposure.**
120. Dioni L, Hoxha M, Nordio F et al. Effects of short-term exposure to inhalable particulate matter on telomere length, telomerase expression, and telomerase methylation in steel workers. *Environ Health Perspect*. 2011;119(5):622-7.
121. Hou L, Wang S, Dou C et al. Air pollution exposure and telomere length in highly exposed subjects in Beijing, China: a repeated-measure study. *Environ Int*. 2012;48:71-7.
122. Weng NP, Granger L, Hodes RJ. Telomere lengthening and telomerase activation during human B cell differentiation. *Proc Natl Acad Sci U S A*. 1997;94(20):10827-32.
123. Hoxha M, Dioni L, Bonzini M et al. Association between leukocyte telomere shortening and exposure to traffic pollution: a cross-sectional study on traffic officers and indoor office workers. *Environ Health*. 2009;8:41.
- 124.● **McCracken J, Baccarelli A, Hoxha M et al. Annual ambient black carbon associated with shorter telomeres in elderly men: Veterans Affairs Normative Aging Study. *Environ Health Perspect*. 2010;118(11):1564-70. First study on telomere attrition and long-term air pollution exposure in elderly.**
- 125.● **Bijnens E, Zeegers MP, Gielen M et al. Lower placental telomere length may be attributed to maternal residential traffic exposure; a twin study. *Environ Int*. 2015;79:1-7. This study provides evidence that maternal traffic exposure is linked with telomere length in early life.**
126. Zhong J, Cayir A, Trevisi L et al. Traffic-Related Air Pollution, Blood Pressure, and Adaptive Response of Mitochondrial Abundance. *Circulation*. 2016;133(4):378-87.
127. Rappaport SM, Smith MT. Epidemiology. Environment and disease risks. *Science*. 2010;330(6003):460-1.
128. Wild CP. The exposome: from concept to utility. *Int J Epidemiol*. 2012;41(1):24-32.

129. Epel ES, Blackburn EH, Lin J et al. Accelerated telomere shortening in response to life stress. *Proc Natl Acad Sci U S A*. 2004;101(49):17312-5.
130. Crous-Bou M, Fung TT, Prescott J et al. Mediterranean diet and telomere length in Nurses' Health Study: population based cohort study. *BMJ*. 2014;349:g6674.
131. Steptoe A, Hamer M, Butcher L et al. Educational attainment but not measures of current socioeconomic circumstances are associated with leukocyte telomere length in healthy older men and women. *Brain Behav Immun*. 2011;25(7):1292-8.
132. Adler N, Pantell MS, O'Donovan A et al. Educational attainment and late life telomere length in the Health, Aging and Body Composition Study. *Brain Behav Immun*. 2013;27(1):15-21.
133. Muezzinler A, Zaineddin AK, Brenner H. Body mass index and leukocyte telomere length in adults: a systematic review and meta-analysis. *Obes Rev*. 2014;15(3):192-201.
134. Liang G, Schernhammer E, Qi L et al. Associations between rotating night shifts, sleep duration, and telomere length in women. *PLoS One*. 2011;6(8):e23462.
135. Surtees PG, Wainwright NW, Pooley KA et al. Life stress, emotional health, and mean telomere length in the European Prospective Investigation into Cancer (EPIC)-Norfolk population study. *J Gerontol A Biol Sci Med Sci*. 2011;66(11):1152-62.
136. Entringer S, Epel ES, Lin J et al. Maternal psychosocial stress during pregnancy is associated with newborn leukocyte telomere length. *Am J Obstet Gynecol*. 2013;208(2):134 e1-7.
137. Marchetto NM, Glynn RA, Ferry ML et al. Prenatal stress and newborn telomere length. *Am J Obstet Gynecol*. 2016.
138. Salihu HM, Pradhan A, King L et al. Impact of intrauterine tobacco exposure on fetal telomere length. *Am J Obstet Gynecol*. 2015;212(2):205 e1-8.
139. Lin S, Huo X, Zhang Q et al. Short placental telomere was associated with cadmium pollution in an electronic waste recycling town in China. *PLoS One*. 2013;8(4):e60815.
140. Wong JY, De Vivo I, Lin X et al. Cumulative PM(2.5) exposure and telomere length in workers exposed to welding fumes. *J Toxicol Environ Health A*. 2014;77(8):441-55.
141. Xia Y, Chen R, Wang C et al. Ambient air pollution, blood mitochondrial DNA copy number and telomere length in a panel of diabetes patients. *Inhal Toxicol*. 2015;27(10):481-7.
142. Shan M, Yang X, Ezzati M et al. A feasibility study of the association of exposure to biomass smoke with vascular function, inflammation, and cellular aging. *Environ Res*. 2014;135:165-72.

**Table 1. Occupational studies describing the association between air pollution exposure and telomere length**

Authors	Study Population	n	Age, y	% Men	Telomere method and tissue	Exposure time	Exposure, concentration in $\mu\text{g}/\text{m}^3$	% change or $\beta$ , (95%CI), p value	Adjustments
Wong et al., 2014 [140]	Boilermakers Study: Population of males, Massachusetts, USA	48	39.3 $\pm$ 12.8 <sup>a</sup> at baseline	100%	qPCR, WBC	Career exposure, last year and last month exposure	Career cumulative PM <sub>2.5</sub> : 713.7 $\pm$ 1457.9 <sup>a</sup> mg/m <sup>3</sup> hr Year cumulative PM <sub>2.5</sub> : 74.0 $\pm$ 77.3 <sup>a</sup> mg/m <sup>3</sup> hr Month cumulative PM <sub>2.5</sub> : 8.5 $\pm$ 10.2 <sup>a</sup> mg/m <sup>3</sup> hr	Career: $\beta$ : -0.021 (-0.048 to 0.006), p>0.05 Year: $\beta$ : -0.022 (-0.053 to 0.009), p>0.05 Month: $\beta$ : -0.04 (-0.08 to 0.001), p≤0.05	% neutrophils, % lymphocytes, % monocytes, % eosinophils, current smoking status, age at baseline blood drawn, BMI, years as boilermaker
Hou et al., 2012 [121]	The Beijing Truck Driver Air Pollution Study: Population of truck drivers and indoor office workers, Beijing, China	120	Truck drivers: 33.5 $\pm$ 5.7 <sup>a</sup> Office workers: 30.27 $\pm$ 7.96 <sup>a</sup>	67%	qPCR, WBC	Personal PM <sub>2.5</sub> and EC exposure for 8-h  Ambient PM <sub>10</sub> exposure: Examination day, 1day, 1-2 days, 1-5 days, 1-7 days, 1-10 days, 1-14 days	Personal PM <sub>2.5</sub> for truck drivers vs office workers: 126.8 $\pm$ 68.8 <sup>a</sup> vs 94.6 $\pm$ 64.9 <sup>a</sup>  Personal EC for truck drivers vs office workers: 17.3 $\pm$ 6.7 <sup>a</sup> vs 13.1 $\pm$ 4.0 <sup>a</sup>  Ambient PM <sub>10</sub> for truck drivers vs office workers: - Examination day: 116.7 $\pm$ 50.2 <sup>a</sup> vs 123.5 $\pm$ 50.1 <sup>a</sup> - 1-day: 121.5 $\pm$ 47.8 <sup>a</sup> vs 119.5 $\pm$ 51.2 <sup>a</sup> - 1-2 days: 121.6 $\pm$ 38 <sup>a</sup> vs 119.3 $\pm$ 40.3 <sup>a</sup> - 1-5 days: 119.5 $\pm$ 26.9 <sup>a</sup> vs 118.2 $\pm$ 25.6 <sup>a</sup> - 1-14 days: 119.9 $\pm$ 18.7 <sup>a</sup> vs 121.7 $\pm$ 17.8 <sup>a</sup>	PM <sub>2.5</sub> : 5.2% (1.5% to 9.1%), p=0.007 for each IQR increase  Personal EC: 4.9% (1.2% to 8.8%), p=0.01 for each IQR increase  PM <sub>10</sub> : - Examination day: 7.7% (3.7% to 11.9%), p<0.001 - 1-day: 8.4% (4.0% to 13.0%), p<0.001 - 1-2 days: 8.1% (3.1% to 13.3%), p=0.002 - 1-5 days: -1.3% (-6.3% to 4.1%), p=0.64 - 1-14 days: -9.9% (-17.6%, -1.5%), p=0.02	Age, gender, BMI, number cigarettes smoked during examination time, day of the week, usage central heating, time used for commuting to work, temperature, dew point
Dioni et al., 2011 [120]	Population of steel workers, Brescia, Italy	57	44 (27-55) <sup>b</sup>	100%	qPCR, WBC	PM <sub>10</sub> and PM <sub>1</sub> exposure for 3 days	PM <sub>10</sub> <sup>a</sup> : 262 $\pm$ 272 PM <sub>1</sub> <sup>a</sup> : 8.0 $\pm$ 7.7	PM <sub>10</sub> : $\beta$ : 0.30 (0.11 to 0.49), p=0.002 for each increase of 285 $\mu\text{g}/\text{m}^3$ (10 <sup>th</sup> -90 <sup>th</sup> perc)  PM <sub>1</sub> : $\beta$ : 0.29 (0.01 to 0.57), p=0.042 for each increase of 11.05 $\mu\text{g}/\text{m}^3$ (10 <sup>th</sup> -90 <sup>th</sup> perc)	Age, BMI, pack-years of smoking, % lymphocytes



Hoxha et al., 2009 [123]	Population of street traffic officers and office workers (referents), Milan, Italy	134	< 30 y: 39% 30-40 y: 37% > 40 y: 24%	63%	qPCR, WBC	One work shift (7 hours)	Airborne benzene: Referents: 13.0 (2.0-115.1) <sup>b</sup> Traffic officers: 31.8 (9.0- 315.7) <sup>b</sup>  Airborne toluene: Referents: 43.4 (6.0-368.0) <sup>b</sup> Traffic officers: 128.7 (24.4- 1710.7)	Benzene: -6.4% (-10.4% to - 2.1%), p=0.004 for each IQR (11.2 µg/m <sup>3</sup> ) increase  Toluene: -6.2% (-10.4% to - 1.7%), p= 0.008 for each IQR (25.7 µg/m <sup>3</sup> ) increase	Age, gender, smoking, pack years of smoking
-----------------------------	---	-----	--	-----	-----------	--------------------------	--	---	--

---

<sup>a</sup> Mean±SD


<sup>b</sup> Mean (range)

**Table 2. Population-based studies describing the association between air pollution exposure and telomere length**

Authors	Study Population	n	Age, y	% Men	Telomere method and tissue	Exposure time	Exposure, concentration in $\mu\text{g}/\text{m}^3$	% change or $\beta$ (95%CI), p value	Adjustments
Pieters <i>et al.</i> 2015 [119]	Population of elderly, all non-smokers, Flanders, Belgium	166	70.6 $\pm$ 4.7 <sup>a</sup>	46%	qPCR, WBC	One year exposure, last month and last week	Annual $\text{PM}_{2.5}$ : 21.1 $\pm$ 1.76 <sup>a</sup>	<p>Last year: -16.8% (-26.0% to -7.4%), <math>p=0.0005</math> for each <math>5\mu\text{g}/\text{m}^3</math> increment</p> <p>Last month: 17.5% (9.6% to 26.0%), <math>p=0.0001</math> for each <math>5\mu\text{g}/\text{m}^3</math> increment</p> <p>Last week: 1% (-2.4% to 4.9%), <math>p=0.5</math> for each <math>5\mu\text{g}/\text{m}^3</math> increment</p>	Gender, age, BMI, socio-economic status, statin use, past smoking status, WBC count, % neutrophils
Xia <i>et al.</i> , 2015 [141]	Type 2 diabetes patients, Shanghai, China	35	65 $\pm$ 8 <sup>a</sup>	49%	qPCR, WBC	Lag 0, 1, 2, 3 and 4-7 days prior blood drawn	<p>24-h PMC:</p> <p>PMC &lt; <math>1\mu\text{m}</math>: 39.6<math>\pm</math>21.7<sup>a</sup></p> <p>PMC 1-2.5<math>\mu\text{m}</math>: 6.6<math>\pm</math>4.5<sup>a</sup></p> <p>PMC 2.5-10<math>\mu\text{m}</math>: 21.1<math>\pm</math>18.5<sup>a</sup></p>	<p>All lags not significant</p> <p><math>\text{PM}_{2.5}</math>: 0.11% (-0.97% to 1.19%), <math>p=\text{NS}</math> for each IQR increment in 24-h average air pollutant</p>	Temperature and relative humidity on concurrent day and previous 3 days, day of the week, month, age, gender, BMI, income, duration T2DM, statin use, random intercept for correlation among multiple measurements
Shan <i>et al.</i> , 2014 [142]	Population of females, Sichuan, China	21	59 (38-85) <sup>b</sup>	0%	qPCR, Buccal cells	24-h exposure	<p>24-h personal <math>\text{PM}_{2.5}</math> in low vs high exposed: 39<math>\pm</math>11<sup>a</sup> vs 101<math>\pm</math>37<sup>a</sup></p> <p>24-h personal BC in low vs high exposed: 2.6<math>\pm</math>1.8<sup>a</sup> vs 14.9<math>\pm</math>11.2<sup>a</sup></p>	<p>Low vs high exposed group</p> <p>-43% (-113% to 28%) <math>p=\text{NS}</math></p>	Age, BMI
McCracken <i>et al.</i> , 2010 [124]	Normative Aging Study (NAS): Population of males, all never smokers, Massachusetts, USA	165	73.6 $\pm$ 7.1 <sup>a</sup>	100%	qPCR, WBC	365 days before each blood drawn	Annual BC <sup>a</sup> : 0.32 <sup>a</sup> $\pm$ 0.2 <sup>a</sup>	-7.6 % (-12.8% to -2.1%) for each IQR (0.25 $\mu\text{g}/\text{m}^3$ ) increase. $p=0.008$	Age at baseline, age change between measurements, year, BMI, WBC count, % neutrophils, % lymphocytes, statin use, diabetes, fasting blood glucose, education, urbanity, census-tract socioeconomic status, inverse probability of follow-up response weighting

<sup>a</sup> Mean±SD

<sup>b</sup> Mean (range)



**Fig 1. The air-pollution induced telomere-mitochondrial aging hypothesis.** Particulate matter exposure induces the formation of ROS at particle surfaces by Fenton like reaction. Besides direct ROS formation from the particle surface, PM induces elevated ROS levels due to altered function of NADPH-oxidase, mitochondria and activation of inflammatory cells. The cellular presence of ROS induces DNA damage and induces single-strand breaks at the G rich telomeres leading to telomere shortening and induces cellular aging. Subsequently DNA damage and telomere shortening is associated with the increase of p53 production. Elevated levels of p53 leads to increased mitochondrial dysfunction, leading to accelerated cellular aging. Under normal conditions high levels of SIRT1 reduces the production of p53. Telomere damage and shortening have been associated with the suppression of SIRT1 which is associated with high p53 levels. Besides the direct effects of ROS production by particulate matter on telomeres and subsequently the p53 pathways, particulate matter has also direct effects on mitochondria by inducing mitochondrial dysfunction, which leads to mitochondrial ROS production. This mitochondrial induced ROS production may alternatively also influences telomere shortening and again inducing the p53 pathway cascade. These mechanisms indicates a close relationship between mitochondrial and telomere function in the aging phenotype affected by exposure to particulate matter.