

Occurrence and effect of trematode metacercariae in two endangered killifishes from Greece

Peer-reviewed author version

Kalogianni, Eleni; Kmentová, Nikol; Harris, Eileen; Zimmerman, Brian; Giakoumi, Sofia; Chatzinikolaou, Yorgos & VANHOVE, Maarten (2017) Occurrence and effect of trematode metacercariae in two endangered killifishes from Greece. In: PARASITOLOGY RESEARCH, 116(11), p. 3007-3018.

DOI: 10.1007/s00436-017-5610-z

Handle: <http://hdl.handle.net/1942/25486>

1 Occurrence and effect of trematode metacercariae in two endangered killifishes from Greece

12 Eleni Kalogianni¹, Nikol Kmentová², Eileen Harris³, Brian Zimmerman⁴, Sofia Giakoumi¹, Yorgos

3 Chatzinikolaou¹, Maarten P.M. Vanhove^{1,2,5,6,7*}

5 ¹*Institute of Marine Biological Resources and Inland Waters, Hellenic Centre for Marine*

6 *Research, 46.7 km Athinon – Souniou Av., 190 13, P.O. Box 712, Anavyssos, Greece*

7 ²*Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2, 611 37*

8 *Brno, Czech Republic*

9 ³*Division of Parasites and Vectors, Department of Life Sciences, The Natural History Museum,*

10 *Cromwell Road, London, SW7 5BD, United Kingdom*

11 ⁴*Zoological Society of London, Regent's Park, London NW1 4RY, United Kingdom*

12 ⁵*Laboratory of Biodiversity and Evolutionary Genomics, Department of Biology, University of*

13 *Leuven, Ch. Deberiotstraat 32, B-3000 Leuven, Belgium*

14 ⁶*Capacities for Biodiversity and Sustainable Development, Operational Directorate Natural*

15 *Environment, Royal Belgian Institute of Natural Sciences, Vautierstraat 29, B-1000 Brussels,*

16 *Belgium*

17 ⁷*Hasselt University, Centre for Environmental Sciences, Research Group Zoology: Biodiversity*

18 *and Toxicology, Agoralaan Gebouw D, B-3590 Diepenbeek, Belgium*

19 * *Corresponding author email: maarten.vanhove@kuleuven.be; telephone: +32 (0)2 627 44 52*

20 **Running title:**

21 **Digenean infection of endangered Greek killifishes**

22 **Abstract**

23 We report digeneans (Diplostomidae, Crassiphialinae) in the endangered freshwater fishes *Valencia*

24 *letourneuxi* and *Valencia robertae*, endemics of Western Greece. Digenean metacercariae occurred

in two forms in the abdominal cavity, excysted and encysted, the latter attached to gonads, liver and alimentary tract. Parasites were, using morphological and molecular techniques, identified as two representatives of Crassiphalinae, specifically part of the *Posthodiplostomum-Ornithodiplostomum* clade. The spatial, seasonal and age class variation in parasite prevalence was examined. Autumn parasite prevalence varied between the six populations sampled (18.2% to 100%). Seasonal prevalence at the two sites sampled quadannually peaked in autumn and reached its lowest value in spring; prevalence increased with size to 100% in young adult fish. We did not find a correlation between prevalence and host sex. Overall parasites' weight averaged 0.64% of the host's, while parasite weight increased with host weight. A comparison of relative condition, and hepatosomatic and gonadosomatic indices of infected and metacercariae-free specimens showed that infection did not have a significant effect on host body condition and reproduction. Regarding the parasite's life cycle, planorbid gastropods are proposed as potential first intermediate hosts in view of the host's diet and occurrence data of molluscs in the ecosystem. This is the first record of a diplostomid digenean in valenciid fishes and of representatives of the *Posthodiplostomum-Ornithodiplostomum* clade in a native Greek freshwater fish. Our findings are discussed in conjunction to fish conservation interventions, since parasites may contribute to the decline of endangered species.

Keywords

Digenea; parasite prevalence; host body condition; Planorbidae; Platyhelminthes; Valenciidae

Introduction

The critically endangered freshwater fish *Valencia letourneuxi* (Sauvage, 1880) (Valenciidae) is an endemic of Western Greece and southern Albania (Crivelli 2006). Its sister species, the recently described *Valencia robertae* Freyhof, Kärst and Geiger, 2014, endemic of Central Greece, encompasses the most southern populations previously included in *V. letourneuxi*. Both species are

characterized by a fragmented geographical distribution, narrow ecological requirements and low population densities (Barbieri et al. 2000; Kalogianni et al. 2010a). In the last 30 years, both species' geographical range has been reduced, with some populations now being extinct and others in a precarious state (2005 survey data, Kalogianni et al. 2010b; see also Economidis 1991; Bianco et al. 1996; Barbieri et al. 2002a, b, 2015). Their decline has been attributed to the degradation of their habitats due to anthropogenic activities, as well as to predation and competition with the introduced Eastern mosquitofish *Gambusia holbrooki* Girard, 1859 (Bianco and Miller 1989; Barbieri et al. 2000, 2002b; Kottelat and Freyhof 2007). The parasite fauna of these Greek killifishes has never been studied. Parasitization, however, of individuals of a native fish species, either by non-native parasites transmitted from introduced fish (Prenter et al. 2004) or by native parasites, can also potentially contribute to its decline, by influencing host behaviour, survival, growth and fecundity, as well as host population dynamics (Marcogliese 2004). Parasitic organisms are often neglected in the management and conservation of biological resources and ecosystems (Marcogliese 2004). Research on the parasites of endangered species can, however, provide information about their host organisms and the ecological interactions between these organisms (e.g. Whiteman et al. 2007). Parasite community composition can provide valuable information for the management and conservation of aquatic species and habitats, e.g. by contributing to understanding introduction routes (e.g. Huyse et al. 2015). Furthermore, the assemblage of parasites within a host organism potentially reflects that host's trophic position in the food web, as well as the presence in the ecosystem of various other organisms that participate in the life cycles of these parasites. Parasite populations and communities could also be useful indicators of environmental stress, such as eutrophication or acidification, as well as of food web structure and biodiversity (Marcogliese 2005; Vidal-Martínez et al. 2010).

A preliminary examination of the abdominal cavity of *V. letourneuxi* and *V. robertae* revealed the presence of digenean trematode metacercariae, possibly belonging to Diplostomidae Poirier, 1886. Among parasites, trematodes are the dominant group that causes retarded growth, morbidity

and mortality, especially in juvenile fishes (Shareef and Abidi 2015). Digenean trematodes are widespread around the globe and are characterized by a complex life cycle, often involving a mollusc as first intermediate host, a fish, an amphibian or occasionally a mammal as second intermediate host and piscivorous birds or mammals as definitive hosts (Niewiadomska 2002; Cribb et al. 2003). Digeneans have been shown to induce behavioural changes in their fish secondary host, such as decreasing swimming performance (Coleman 1993) or decreasing predator avoidance (Poulin 1993) resulting in increased predation of the host (Ondráčková et al. 2006). They have also been shown to cause damage to fish host tissues resulting in blindness, inflammatory reactions and perforations in some cases (Shariff et al. 1980; Niewiadomska 2002; Vianna et al. 2005), though there are also studies reporting no effect of digenean parasitism on fish host condition (e.g. Silva-Souza and Ludwig 2005).

In this study, we further examined the abovementioned metacercariae retrieved from *V. letourneuxi* and *V. robertae* in order to identify these parasites using morphological and molecular techniques, to examine the variation in prevalence between seasons, locations and host sexes and size-classes, and to assess the effect of the metacercariae on host condition and reproduction.

Materials and methods

Sampling methodology and phenotypic characterisation of hosts and parasites

Samples were collected from six sites in Western Greece, in the autumn of 2005, 2006 and 2009; one site hosts *V. robertae* and the other five sites *V. letourneuxi* (Fig. 1 and Table 1). To explore seasonal variation in parasitization, seasonal samplings (July, September, January, May) were conducted at two sites, Mornos and Acheron (sites 4 and 5, hosting *V. robertae* and *V. letourneuxi* respectively, Table 1). All sampling sites were located at lowland semi-lotic streams or canals (elevation range 0-6 m), associated with springs, while distance from sea ranged from 0.4 to 13.5

105 km. Site depth ranged between 0.9 and 1.7 m and site width between 4 and 13 m. All sites were
106 thickly vegetated (surface aquatic vegetation ranged between 40 and 95%). Salinity varied between
107 0.1 and 6.5 ppt and temperature between 14.5 and 25 °C.

108 This study was conducted within the frame of a wide scope research programme during 2005-2009,
109 targeting *V. letourneuxi* (*V. robertae* had not yet been described as a separate species at the time).

110 This research included dietary studies, studies on the effect of *G. holbrooki* Girard, 1859 on *V.*
111 *letourneuxi* and genetic studies. Fish were collected with a D-shaped net (2 mm mesh) from the
112 stream banks at the six sites described above (Fig.1 and Table 1). In five of the six sites sampled
113 (sites 1, 2, 3, 5, 6, Table 1), representatives of *Valencia* are found in association with the introduced
114 *G. holbrooki* (Kalogianni et al., 2010a). After identification, a total of 296 specimens of *Valencia*
115 were anesthetized with quinaldine and preserved in 4% formaldehyde for further laboratory
116 analyses. In the laboratory, for each fish total and standard length (TL and SL, nearest 0.1 mm)
117 were measured, total and net host weight, before and after evisceration respectively (HW and NW,
118 nearest 0.01 mg) as well as liver and gonad weight (LW and GW, nearest 0.01 mg after blotting
119 dry) were also recorded. External surfaces, viscera and musculature of the fish were examined
120 under a Olympus SZX7 stereo microscope for parasitic infection. Parasites were removed, blotted
121 dry and their weight was recorded (PW, nearest 0.01 mg, weighed for all metacercariae of one host
122 specimen together).

123 Parasite prevalence (number of infected fish per total fish examined) was calculated for all
124 metacercariae together (i.e. no species-level morphological identification of the digeneans could be
125 made and these prevalence values are hence not parasite species-specific). Differences in parasite
126 prevalence between sexes and 2 mm size classes (larval size range < 13 mm, juvenile range ≥ 13
127 and ≤ 17 mm and adults > 17 mm, see Barbieri et al. 2000 and Kalogianni et al. 2010b) were
128 examined pooling data from all the autumn samples. The effect of host sex on the prevalence of
129 parasites was tested by applying chi-square test. Differences in mean standard length between

infected and metacercariae-free specimens were examined using a t-test. The relationship between total PW and HW was examined with Pearson's correlation. Finally, ANCOVA was used to investigate potential effects of parasitation on the condition and reproduction of the female and male fish hosts, after calculation of the relative condition (NW/SL), the hepatosomatic (LW/NW) and the gonadosomatic (GW/NW) indices of infected and metacercariae-free specimens. Fish standard length was used as a covariate to account for possible size effects. Prior to statistical analysis, values of the above indices were log10 transformed. ANCOVA was conducted with PASW 17 software.

Parasite identification

Parasite identification was done using combined morphological and molecular methods. The digenean metacercariae, excysted when necessary, were stained using paracarmine, dehydrated through a series of alcohols and cleared in beechwood creosote. They were mounted in Canada balsam on glass microscope slides and examined under a Olympus BH2 high-power microscope with interference phase.

Additional samples, used for the genetic characterization of the parasites, were collected from the Acheron site (site 5) in June 2008. We extracted DNA from individual (artificially or naturally) excysted metacercariae found in the abdominal cavity with the DNeasy Blood and Tissue Kit (Qiagen) according to the manufacturer's instructions. Polymerase Chain Reaction was performed using a GeneAmp PCR system 9700 thermocycler (Applied Biosystems) and Illustra PuReTaq Ready-To-Go PCR Beads (GE Healthcare), adding 1 µL of each primer (20 µM) (Sigma Aldrich), 2 µL of template DNA and 21 µL of double distilled, autoclaved and filter sterilized water, for a total reaction volume of 25 µL. We amplified fragments of the mitochondrial cytochrome *c* oxidase subunit I (COI) gene, and of the nuclear rDNA region (ITS-1, 5.8, ITS-2). Primer combinations were MplatCOX1dF (5'-TGTAACGACGGCCAGTTTWCITTRGATCATAAG-3') and MplatCOX1dR (5'-CAGGAAACAGCTATGACTGAAAYAAAYAIIGGATCICCACC-3')

(Moszczyńska et al. 2009) for COI and the combinations of D1 (5'-AGGAATTCCTGGTAAGTGCAAG-3') with D2 (5'-CGTTACTGAGGGAATCCTGG-3') (Hillis and Dixon 1991) and 81_f (5'-GTAACAAGGTTTCCGTAGGTGAA-3') (Gustinelli et al. 2010) with ITS2.S_r (5'-CCTGGTTAGTTTCTTTTCCTCCGC-3') (Cribb et al. 1998) for rDNA. These regions (ITS-1, 5.8 and ITS-2) are commonly used for species identification in flatworms (Vanhove et al. 2013; Stoyanov et al. 2017). After an initial denaturation of 2 min at 94 °C, samples were subjected to 35 cycles (40 for 81f – ITS2sr) of 30 s at 94 °C, 30 s (40 s for 81f – ITS2sr) at 50 °C and 60 s (90s for 81f – ITS2sr) at 72 °C. After a final elongation of 10 min (5 min for 81f – ITS2sr) at 72 °C, samples were cooled to 4 °C. PCR products were purified using the QIAquick PCR Purification Kit (Qiagen) following the manufacturer's protocol. Sequencing of both strands was carried out using the same primers as above with an Applied Biosystems 3730 DNA analyser and BigDye version 1.1. Sequences were deposited in NCBI GenBank under accession numbers KY320571-3. Voucher specimens for the genetically characterized parasite population were deposited in the Natural History Museum (London, United Kingdom) (NHMUK 2015.12.2.1) (parasite) and the Natural History Museum Rijeka (Croatia) (PMR VP 3140-2) (host). Sequences were visually corrected and aligned in MEGA v6 (Tamura et al. 2013) with the MUSCLE algorithm and UPGMB clustering method (Edgar 2004) under default conditions. The best fitting substitution model describing molecular evolution of the sequences was selected by TOPALi v2.5 (Milne et al. 2009) based on the Bayesian information criterion. The GTR model (Rodriguez et al. 1990) was used for the rDNA region. Pairwise deletion was used to construct a distance matrix. The phylogenetic position of the collected parasite haplotypes within Diplostomidae (GenBank accession numbers in Table 2, representatives of available diplostomid genera were selected) was inferred based on the rDNA combining the results of a maximum likelihood tree search performed in RAxML 8.7.4. (Stamatakis 2014) with bootstrap values calculated using 1,000 replicates and Bayesian interference performed in MrBayes 3.2 (Ronquist et al. 2011). Posterior probabilities were approximated for 10,000,000 generations, sampled at each 1,000th generation and with a burn-in of

10% in two separate runs. Chain stationarity and parameter convergence were checked in Tracer 1.6 (Rambaut et al. 2014). Because of its position in a different but related digenean family (Clinostomidae), *Clinostomum complanatum* (Rudoplhi, 1814) collected from *Barbus barbus* (L.) was used as an outgroup. File conversion was carried out using ALTER (Glez-Peña et al. 2010). Phylogenetic trees were rendered by FigTree 1.4.2 (Rambaut and Drummond 2009) and edited in Adobe Photoshop CS6.

Results

Parasite identification

Digenean metacercariae were found in the abdominal cavity of *V. letourneuxi* and *V. robertae*, while isolated cysts were also found in some fishes beneath the lens of the eye. Parasites were identified as members of Diplostomidae, subfamily Crassiphialinae. This was based on the morphological characters of the ‘neascus’ (a characteristic type of diplostomid metacercariae, see Niewiadomska 2002), such as the presence of a bipartite body, with a reserve bladder consisting of a ramified median and two lateral canals forming a net-like structure in the forebody, and a developed hindbody with unconnected excretory canals. Since only immature gonads were present, further identification to genus or species level was not possible, because the classification is based on the size of testes and the absence/presence of an ejaculatory pouch (Niewiadomska 2002).

For the genetic identification of the parasites, sequences from nine metacercariae (only specimens from the abdominal cavity were available for molecular work) were obtained and two different rDNA haplotypes were recorded, with a length of 1159 base pairs and a pairwise difference of 4.8%. These haplotypes did not correspond with the distinction between encysted and excysted metacercariae. Only one corresponding COI haplotype was recorded with a length of 531 base pairs, due to low amplification success. The pairwise distances in the entire dataset ranged from 0.5–37.8 % in the rDNA regions and from 0.5–33.2 % in the COI region (pairwise deletion). Phylogenetic analyses of rDNA did not cluster the two haplotypes of metacercariae infecting Greek

killifishes together. Genetic distances between both haplotypes surpassed those between other sequences considered to belong to different species. This indicates the presence of two parasite species. They clustered with representatives of *Posthodiplostomum* Dubois, 1936 and *Ornithodiplostomum* Dubois, 1936 placed among other basal lineages of this clade. Both methods produced the same tree topology for rDNA (Fig. 2). The analyses confirmed the previously observed polyphyly of *Posthodiplostomum*.

Host-parasite ecology

Metacercariae occurred in two forms (encysted and excysted, Fig. 3), the encysted form usually in groups attached to the anterior end of the gonad, adjacent to the mesentery and at the area between the liver and the digestive tract, with parasites varying in size. Of a total of 296 fishes, 219 were found to be parasitized, corresponding to a high overall metacercaria prevalence of 73.99% (not separated between parasite forms).

Parasite prevalence, in the six *Valencia* populations studied, varied in autumn from 18.18% to 100% (Table 1; maximum prevalence was 100% for both *V. robertae* and *V. letourneuxi*). Seasonal prevalence remained high throughout the year at the two sites sampled quadannually (Table 1) ranging between 74.3% and 100% (Fig. 4). In these two sites, prevalence peaked in autumn with all fish being parasitized (100%) and then decreased in the winter to 82.3%, due to the appearance of a group of metacercariae-free fish, ranging in size from 13 to 36 mm SL (juveniles and adults, see Kalogianni et al. 2010b). In the spring, prevalence further decreased to 74.3%, reaching its lowest value. Parasite prevalence in function of the host's sex (ratio 5:4 in favor of females), was 52.88% and 63.15% in host females (n=52) and males (n=42) respectively, but this difference was not statistically significant ($\chi^2 = 0.792$; $df = 1$; $P = 0.374$).

Parasite prevalence increased gradually with host size (SL, Fig. 5), from a prevalence of 0% for the larval length classes 6-8 and 8-10 mm to a maximum prevalence value of 100% in the 22-24 mm adult length class (larval size range < 13 mm, juvenile range ≥ 13 and ≤ 17 mm and adults > 17

mm, see Barbieri et al. 2000 and Kalogianni et al. 2010b). For fish lengths > 24 mm these prevalence values remained stable, with one exception for the adult length class 24-26 mm (prevalence 75%, Fig. 5). Mean SL values for infected and metacercariae-free fish differed, i.e. mean SL of infected fish was 19.70 ± 0.92 S.E. ($n = 58$); mean SL of metacercariae-free fish was 13.64 ± 0.55 S.E. ($n = 36$); this difference was statistically significant ($t = -4.846$, $P < 0.0001$). Mean parasite:host weight ratio was 0.0064 ± 0.0010 S.E., with parasite weight averaging 0.64% of the host weight. Parasite weight and host weight were positively correlated ($n = 9$, rho correlation = 0.882, $P = 0.002$, see Fig. 6) for fish < 600 mg, with a mean parasite:host weight ratio of 0.0076 ± 0.0009 S.E. and percentage mean 0.76%; however, there was no correlation for the larger specimens (mean \pm S.E. parasite:host weight ratio of 0.0009 ± 0.0007 and percentage mean 0.09%, $n = 2$).

The study of the condition, gonadosomatic and hepatosomatic indices showed no statistically significant differences between infected and metacercariae-free specimens, in either males or females ($p > 0.05$).

Discussion

Parasite identification

This study is the first record of valenciids as second intermediate hosts for diplostomid parasites, being heavily infected by their metacercariae. No other endoparasites were found in this study. (Dactylogyridean monogenean gill parasites were also found – at very low prevalence and infection intensity – but fall outside of the scope of this study.) Dominance of (immature) endoparasites in general, and of trematode metacercariae in particular, in killifishes has been previously observed. It was suggested to demonstrate the importance of these fishes as intermediate or paratenic hosts (Nezhybová et al. 2017). Pairwise uncorrected genetic distances between the retrieved rDNA haplotypes (4.8%) suggest the existence of two different diplostomid species (Georgieva et al.

2013; Stoyanov et al. 2017). Unfortunately, the metacercarial lifestage does not allow morphology-based species-level distinction. Therefore, parasite haplotypes could not be linked with specific phenotypic characters; neither was there a link with being encysted or not. The collected parasite species are considered as representatives of *Posthodiplostomum* or *Ornithodiplostomum* based on the results of phylogenetic analysis (Fig. 2). Interestingly, haplotype 1 takes a basal position within the *Posthodiplostomum*-*Ornithodiplostomum* clade and most likely represents a hitherto unsequenced phylogenetic lineage. Low maximum likelihood bootstrap values of deeper nodes compared to the Bayesian inference posterior probabilities highlight the unresolved position of several diplostomid genera and an insufficient number of already published sequences. Moreover, the tree also indicates that *Posthodiplostomum* is not monophyletic and in need of revision (Locke et al. 2010b; Athokpam and Tandon 2014; García-Varela et al. 2016). However, according to Stoyanov et al. (2017) the uncertain phylogenetic positions of *Posthodiplostomum* species is often the result of incorrect morphological identification due to low quality of specimens, creating confusion even in available molecular data. Although previous studies based on barcoding approaches reported cryptic diversity in the family (Locke et al. 2010a, b; Georgieva et al. 2013), complicated morphological identification and the lack of information about adult stages makes formal species description challenging as it depends on high quality stained material. Four species of *Posthodiplostomum* (*P. brevicaudatum* (von Nordmann, 1832); *P. centrarchi* Hoffman, 1958; *P. cuticola* (von Nordmann, 1832); *P. minimum* (MacCallum, 1921)) and one of *Ornithodiplostomum* (*O. scardinii* (Schulman in Dubinin, 1952)) respectively, have been reported from more than 70 freshwater fish hosts, mostly cyprinids and cobitids, in Europe (Sonin 1986; Stoyanov et al. 2017). To the best of our knowledge this is the first report of a freshwater species native to Greece found infected by either of both abovementioned diplostomid genera (see e.g. Stoyanov et al., 2017). Our results enrich the list of the known digenean fauna in Greece, following the record of *Diplostomum spathaceum* (Rudolphi, 1819) from ten different fish host species in Lakes Volvi and Vistonis (Kalfa-Papaioannou and Sinis 1985).

286

287 *Host-parasite ecology*

2

288

3

4

5

289

6

290

7

8

291

9

10

292

11

12

293

13

294

14

295

15

296

16

297

17

298

18

299

19

300

20

301

21

302

22

303

23

304

24

305

25

306

26

307

27

308

28

309

29

310

30

311

31

312

32

313

33

314

34

315

35

316

36

317

37

318

319

The linear correlation between parasite weight and host weight indicates that infection occurs early in the life of the fish and then the parasites grow with the host and/or fish accumulate metacercariae as they grow. The positive relation between the length of the host and the prevalence of the parasite observed in the target species is most likely a result of temporal accumulation of parasitisation, as larger fish could be exposed repeatedly to infection for a longer time than younger fish (Saad-Fares and Combes 1992; Paes et al. 2010). This is also supported by the presence of different stages of digenean metacercariae in the target species, as suggested also in various cyprinids or cobitids (Ondráčková et al. 2004a). Finally, there were no significant differences in the susceptibility of infection between males and females, as reported also for other freshwater fish species infected with diplostomid digeneans (Flores and Semenas 2002; Machado et al. 2005). Digenean metacercariae have been also found, in various freshwater species, subcutaneously in the trunk region and head, in fins, gills, the eyes and muscle tissue, as well as in viscera (Sonin 1986; Niewiadomska 2002). In the two host species of this study, metacercariae were found mostly in the visceral cavity, in association with the gonads and the digestive tract and liver, but also beneath the lens of the eye. Hence, it seems a systemic infection. This distribution of the parasite therefore corresponds with previous studies and it is correlated with the high infection level reported (Kvach et al. 2017).

Previous studies have shown that the natural definitive hosts of *Posthodiplostomum* and *Ornithodiplostomum* are piscivorous birds, with planorbid or lymnaeid gastropods the most common first intermediate hosts (Miller 1954; Niewiadomska 2002; Faltýnková et al. 2008; Nguyen et al. 2012). Planorbids and lymnaeids were the only gastropods that were found at both Louros, Mornos and Acheron *Valencia* habitats during a benthic faunal study conducted in summer 2009 (unpublished data). However, only planorbid availability reflected the variation of parasite prevalence between these three populations, with the Louros habitat having both lower planorbid availability and lower parasite prevalence. Furthermore, benthic macroinvertebrate data collected seasonally at the Mornos habitat to assess food availability for a dietary study on *V. robertae* (then *V. letourneuxi*) showed that planorbids were the only gastropod taxon available throughout the year

(Kalogianni et al. 2010b). In addition, that study showed that the target species consumed only three gastropod prey categories (Planorbidae, Valvatidae and Physidae), with Lymnaeidae being absent from its diet and Planorbidae being the most frequent and abundant gastropod in the species diet. Since the planorbids reflect variation in metacercariae prevalence and figure as preferable prey of valenciid fishes, we assume high contact rates. We therefore suggest planorbids as the first intermediate host of the collected digenean species.

Parasite effects on the host species and conservation implications

The results of the current study show that there was no correlation between the diplostomid infection of the target species and the condition and reproduction of the hosts, as reported also elsewhere (Paes et al. 2010; Gholami et al. 2011), though there are studies that have shown a negative correlation between the abundance of *Posthodiplostomum* sp. and the relative condition factor of its hosts (Lucký 1970).

The absence of any detectable differences on fish condition and reproduction between infected and metacercariae-free *Valencia* specimens leads us to tentatively assume that this endoparasite is not pathogenic to its host. Given the short lifespan of the target species (two or three years in the wild, Barbieri et al. 2002b) and the fact that trematode species may live for more than a year in the fish host and even for the whole lifespan of the host (Kalantan et al. 1987; Dias et al. 2006), it appears that this host-parasite relationship bears the characteristics of a strategy in which the parasite does not affect fish survival, fitness and reproduction. However, the observed eye infection by diplostomid metacercariae could affect fish vision and thus increase its predation by birds, as reported for other diplostomids (Seppälä et al. 2005).

Parasites are a potential risk factor in conservation initiatives targeting native species, such as population enhancement, assisted migration or reintroduction actions. Digenean colonization of non-native areas depends on the strategy of larval stages, highly productive asexual reproduction, host specificity, level of virulence in intermediate hosts and measure of similarity of environmental

conditions between source or recipient localities (see Bauer 1991; Kennedy 1993). Based on the enemy release hypothesis, introduced endangered host species could profit of parasite loss (Genner et al. 2008) or be affected by spill-back of parasites from alien hosts in an introduction locality (McCallum and Dobson 1995; Daszak et al. 2000; Holt et al. 2003). On the other hand, a scenario suggesting a greater pathogenetic effect of co-introduced parasites on native hosts was also documented (naïve host hypothesis) (Anderson and May 1992; McCallum and Dobson 1995; Hudson et al. 1998). In this respect, translocations of endangered species to reestablish or to help recover populations could introduce parasites, harmless to the reintroduced population, but pathogenic to the already present naïve conspecifics or other sympatric species (see Daszak et al. 2000; Britt et al. 2004). Therefore, even the seemingly harmless diplostomid digeneans reported here should be considered carefully in the context of the conservation-related release or translocation of *Valencia* populations. Finally, no specimens of the introduced mosquitofish *G. holbrooki*, examined in two water systems where the species is sympatric with *Valencia* species (Acheron and Louros, unpublished data), were found infected by the same diplostomid metacercariae. This indicates that the parasites were not introduced locally through the mosquitofish nor that it could pose a threat to the native species, acting as a reservoir for these parasites. A similar absence of metacercariae of digenean parasites has been also reported in the only available study on mosquitofish parasites in Europe, namely in *G. holbrooki* from eight Mediterranean river mouths in Spain and France (Benejam et al. 2009).

Conclusion

This study showed no negative effect of metacercaria infection on *Valencia* species. Furthermore, seasonal differences in digenean prevalence and parasite accumulation over fish age were documented, with no differences in infection parameters between host sexes. It also confirmed the need for a revision of the complicated taxonomy of diplostomids and their unresolved phylogenetic classification. While we suggest planorbids to be potential first intermediate hosts, further

investigations reconstructing the life cycle of the here reported parasites are required, in order to understand the ecological parameters of infection of their secondary host, as well as to identify the other host taxa. Such information is important to the understanding of parasite-host interactions, as well as to the planning or implementation of appropriate conservation measures for the endangered fish species, targeted in this study, as well as other vulnerable fish hosts of these parasites.

Acknowledgements

This research was conducted in the frame of a wider study on the ecology, biology, genetics and diet of *V. letourneuxi* undertaken by scientists of HCMR and was partially funded by the European Union of Aquarium Curators (EUAC) in the frame of the project “Rapid Assessment of the Status of *Valencia letourneuxi*, the Greek Killifish” (2005-2009). This research received support from the SYNTHESYS Project (<http://www.synthesys.info/>) (GB-TAF-2984) which is financed by European Community Research Infrastructure Action under the FP7 Integrating Activities Programme. N.K. and M.P.M.V. are supported by the Czech Science Foundation [P505/12/G112 (ECIP)]. The authors wish to thank Andrea Waeschenbach and D. Tim J. Littlewood for advice regarding molecular work, Rod A. Bray for advice regarding parasite identification, the staff of the Wolfson Wellcome Biomedical Laboratories at the Natural History Museum for their technical support, Nicholas Koutsikos for producing the map with the sampling sites' location and Marcelo Kovačić for curatorial services. For this study, HCMR had secured all necessary permits for fish collection from the Greek Ministry of Environment, Energy and Climate change (permit numbers 97429/4350 and 85404/130).

Compliance with ethical standards

Conflict of interest: The authors declare that they have no conflict of interest.

Ethical approval: All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.

417 **References**

- 418 Alvarez-Pellitero P (2008) Fish immunity and parasite infections: from innate immunity to
 419 immunoprophylactic prospects. *Vet Immunol Immunop* 126(3-4):171-98
- 420 Anderson RM, May RM (1992) *Infectious diseases of humans: Dynamics and control*. Oxford
 421 University Press, Oxford
- 422 Athokpam VD, Tandon V (2014) Morphological and molecular characterization of
 423 *Posthodiplostomum* sp. (Digenea: Diplostomidae) metacercaria in the muscles of
 424 snakeheads (*Channa punctata*) from Manipur, India. *Helminthologia* 51(2):141-152
- 425 Barbieri R, Daoulas Ch, Psarras Th, Stoumboudi MTh, Economou AN (2000) The biology and
 426 ecology of *Valencia letourneuxi* Sauvage, 1880 (Valenciidae) - Prospects for conservation.
 427 *Mediterr Mar Sci* 1:75-90
- 428 Barbieri R, Economou AN, Stoumboudi MTh, Economidis PS (2002a) Freshwater fishes of
 429 Peloponnese (Greece): distribution, ecology and threats. In: Collares-Pereira MJ, Cowx IG,
 430 Coelho MM (eds) *Conservation of freshwater fishes: options for the future*, Fishing News
 431 Books, Blackwell Science, Oxford, pp 55-64
- 432 Barbieri R, Stoumboudi MTh, Economou AN (2002b) Threatened fishes of the world: *Valencia*
 433 *letourneuxi* Sauvage, 1880 (Valenciidae). *Environ Biol Fish* 65:46
- 434 Barbieri R, Zogaris S, Kalogianni E, Stoumboudi M, Chatzinikolaou Y, Giakoumi S, Kapakos Y,
 435 Kommatas D, Koutsikos N, Tachos V, Vardakas L, Economou AN (2015) Freshwater fishes
 436 and lampreys of Greece: An annotated checklist. *HCMR Monographs on Marine Sciences* 8,
 437 Anavyssos
- 438 Bauer ON (1991) Spread of parasites and diseases of aquatic organisms by acclimatization: a short
 439 review. *J Fish Biol* 39:679-686

- Benejam L, Alcaraz C, Sasal P, Simon-Levert G, García-Berthou E (2009) Life history and parasites of an invasive fish (*Gambusia holbrooki*) along a latitudinal gradient. Biol Invasions 11:2265-2277
- Bianco PG, Miller RR (1989) First record of *Valencia letourneuxi* (Sauvage, 1880) in Peloponnese (Greece) and remarks on the Mediterranean family Valenciidae (Cyprinodontiformes). Cybium 13:385-387
- Bianco PG, Ahnelt H, Economidis P (1996) The freshwater fishes from eastern, and large Mediterranean islands with comments on their safety status. Acta Univ Carol Biol 40:45-60
- Britt A, Welch C, Katz A, Iambana B, Porton I, Junge R, Crawford G, Williams C, Haring D (2004) The re-stocking of captive-bred ruffed lemurs (*Varecia variegata variegata*) into the Betampona Reserve, Madagascar: methodology and recommendations. Biodivers Conserv 13(3):635-657
- Chubb JC (1979) Seasonal occurrence of helminths in freshwater fishes. Part II. Trematoda. Adv Parasitol 17:141-313
- Coleman FC (1993) Morphological and physiological consequences of parasites encysted in the bulbus arteriosus of an estuarine fish, the sheepshead minnow, *Cyprinodon variegatus*. J Parasitol 79:247-254
- Cribb TH, Anderson GR, Adlard RD, Bray RA (1998) A DNA-based demonstration of a three-host life-cycle for the Bivesiculidae (Platyhelminthes: Digenea). Int J Parasitol 28:1791-1795
- Cribb TH, Bray RA, Olson PD, Littlewood DTJ (2003) Life cycle evolution in the digenea: A new perspective from phylogeny. Adv Parasitol 54:197-254
- Crivelli AJ (2006) *Valencia letourneuxi*. The IUCN Red List of Threatened Species 2006: e.T22830A9393054. <http://dx.doi.org/10.2305/IUCN.UK.2006.RLTS.T22830A9393054.en>. Accessed 27 June 2017
- Daszak P, Cunningham AA, Hyatt AD (2000) Emerging infectious diseases of wildlife – Threats to biodiversity and human health. Science 287(5452):443-449

- 466 Dias MLGG, Minte-Vera CV, Eiras JC, Machado MH, Souza GTR, Pavanelli GC (2006) Ecology
467 of *Clinostomum complanatum* Rudolphi, 1814 (Trematoda: Clinostomidae) infecting fish
2 from the floodplain of the high Paraná River, Brazil. Parasitol Res 99:675-681
468
4 from the floodplain of the high Paraná River, Brazil. Parasitol Res 99:675-681
5
469 Economidis PS (1991) Check list of freshwater fishes of Greece. Recent status of threats and
7 protection. Hellenic Society for the Protection of Nature, Athens
470
9
10
471 Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput.
12 Nucleic Acids Res 32:1792-1797
472
14
15
473 Faltýnková A, Našincová V, Kablášková L (2008) Larval trematodes (Digenea) of planorbid snails
17 (Gastropoda: Pulmonata) in Central Europe: a survey of species and key to their
474 identification. Syst Parasitol 69:155-178
19
20
475 Flores V, Semenas L (2002) Infection patterns of *Tylodelphys bariloensis* and *T. crubensis*
22 (Trematoda: Diplostomatidae) metacercariae in *Galaxias maculatus* (Osmeriformes:
476 Galaxiidae) from two Patagonian lakes and observations on their geographical distribution
24 in the Southern Andean Region, Argentina. J Parasitol 88(6):1135-1139
25
477
26
27
478
28
29
479
31
32
480 García-Varela M, Sereno-Urbe AL, Pinacho-Pinacho CD, Domínguez-Domínguez O, Pérez-Ponce
34 de León G (2016) Molecular and morphological characterization of *Austrodiplostomum*
481 *ostrowskiae* Dronen, 2009 (Digenea: Diplostomatidae), a parasite of cormorants in the
36 Americas. J Helminthol 90(2):174-185
37
482
38
39
483
41
42
484 Genner MJ, Michel E, Todd JA (2008) Resistance of an invasive gastropod to an indigenous
43 trematode parasite in Lake Malawi. Biol Invasions 10:41-49
44
485
46
47
486 Georgieva S, Soldánová M, Pérez-del-Olmo A, Dangel DR, Sitko J, Sures B, Kostadinova A (2013)
48 Molecular prospecting for European *Diplostomum* (Digenea: Diplostomidae) reveals cryptic
49 diversity. Int J Parasitol 43:57-72
50
51
488
53
54
489 Gholami Z, Mobedi I, Esmaeili HR, Kia EB (2011) Occurrence of *Clinostomum complanatum* in
55 *Aphanius dispar* (Actinopterygii: Cyprinodontidae) collected from Mehran River, Hormuzgan
56 Province, South of Iran. Asian Pac J Trop Biomed 1(3):189-192
57
58
490
59
60
61
62
63
64
65

- 492 Glez-Peña D, Gómez-Blanco D, Reboiro-Jato M, Fdez-Riverola F, Posada D (2010) ALTER:
493 program-oriented format conversion of DNA and protein alignments. Nucleic Acids Res
2 38(suppl. 2):W14-W18
- 494 38(suppl. 2):W14-W18
- 495 Gustinelli A, Caffara M, Florio D, Otachi EO, Wathuta EM, Fioravanti ML (2010) First description
7 of the adult stage of *Clinostomum cutaneum* Paperna, 1964 (Digenea: Clinostomidae) from
496 grey herons *Ardea cinerea* L. and a redescription of the metacercaria from the Nile tilapia
10
497 *Oreochromis niloticus niloticus* (L.) in Kenya. Syst Parasitol 76(1):39-51
11
12
- 498 Hillis DM, Dixon MT (1991) Ribosomal DNA: molecular evolution and phylogenetic inference. Q
15
499 Rev Biol 66:411-426
16
17
- 500 Holt RD, Dobson AP, Begon M, Bowers RG, Schauber EM (2003) Parasite establishment in host
19
501 communities. Ecol Lett 6:837-842
20
21
22
- 502 Hudson PJ, Dobson AP, Newborn D (1998) Prevention of population cycles by parasite removal.
25
503 Science 282(5397): 2256-2258
26
27
28
- 504 Huyse T, Vanhove MPM, Mombaerts M, Volckaert FAM, Verreycken H (2015) Parasite
31
505 introduction with an invasive goby in Belgium: double trouble? Parasitol Res 114(7):2789-
32
506 2793
33
34
35
- 507 Kalantan AMN, Arfin M, Nizami WA (1987) Seasonal incidence and pathogenicity of the
37
508 metacercariae of *Clinostomum complanatum* in *Aphanius dispar*. Jap J Parasitol 36:17-23
38
39
- 509 Kalfa-Papaioannou AM, Sinis AI (1985) Metacercariae of *Diplostomurn spathaceum* in freshwater
42
510 fish of the Lake System Koronia and Volvi (Macedonia, Greece). Zbl Vet Med B 32:65-70
43
44
45
- 511 Kalogianni E, Giakoumi S, Zogaris S, Chatzinikolaou Y, Zimmerman B, Economou AN (2010a)
47
512 Current distribution and ecology of the critically endangered *Valencia letourneuxi* in
48
513 Greece. Biologia 65:128-139.
49
50
51
52
- 514 Kalogianni E, Giakoumi S, Andriopoulou A, Chatzinikolaou Y (2010b) Feeding ecology of the
54
515 critically endangered *Valencia letourneuxi* (Valenciidae). Aquat Ecol 44(1):289-299
55
56
57
58
59
60
61
62
63
64
65

- Kennedy CR (1993) Introductions, spread and colonization of new localities by fish helminth and crustacean parasites in the British Isles: a perspective and appraisal. *J Fish Biol* 43:287-301
- Kondo Y, Ohtsuka S, Hirabayashi T, Okada S, Ogawa NO, Ohkouchi N, Shimazu T, Nishikawa J (2016) Seasonal changes in infection with trematode species utilizing jellyfish as hosts: evidence of transmission to definitive host fish via medusivory. *Parasite* 23:16
- Kottelat M, Freyhof J (2007) Handbook of European freshwater fishes. Kottelat, Cornol and Freyhof, Berlin
- Kvach Y, Jurajda P, Bryjová A, Trichkova T, Ribeiro F, Přikrylová I, Ondračková M (2017) Distribution of North American digenean, *Posthodiplostomum* cf. *minimum centrarchi* (Strigeiformes: Diplostomidae) metacercariae in Europe. *Parasitol Int* doi: 10.1016/j.parint.2017.06.003
- Locke SA, McLaughlin JD, Dayanandan S, Marcogliese DJ (2010a) Diversity and specificity in *Diplostomum* spp. metacercariae in freshwater fishes revealed by cytochrome *c* oxidase I and internal transcribed spacer sequences. *Int J Parasitol* 40:333-343
- Locke SA, McLaughlin JD, Marcogliese DJ (2010b) DNA barcodes show cryptic diversity and a potential physiological basis for host specificity among Diplostomoidea (Platyhelminthes: Digenea) parasitizing freshwater fishes in the St. Lawrence River, Canada. *Mol Ecol* 19:2813-2827
- Lucký Z (1970) Pathological changes with posthodiplostomosis of fish fry. *Acta Vet Brno [Suppl]* 1:51-66
- Machado PM, Takemoto RM, Pavanelli GC (2005) *Diplostomum* (*Austrodiplostomum*) *compactum* (Lutz, 1928) (Platyhelminthes, Digenea) metacercariae in fish from the floodplain of the Upper Paraná River, Brazil. *Parasitol Res* 97(6):436-444
- Marcogliese DJ (2004) Parasites: Small players with crucial roles in the ecological theatre. *Ecohealth* 1:151-164

- Marcogliese DJ (2005) Parasites of the superorganism: Are they indicators of ecosystem health? *Int J Parasitol* 35:705–716
- Mbokane EM, Theron J, Luus-Powell WJ (2015) Seasonal occurrence of some larval stages of endoparasites in three cyprinids from the Nwanedi-Luphephe dams, the Limpopo River System, South Africa. *Helminthologia* 52(3):229-235
- McCallum H, Dobson A (1995) Detecting disease and parasite threats to endangered species and ecosystems. *Trends Ecol Evol* 10:190-194
- Miller JH (1954) Studies on the life history of *Posthodiplostomum minimum* (MacCallum 1921). *J Parasitol* 3(40):255-270
- Milne I, Lindner D, Bayer M, Husmeier D, McGuire G, Marshall DF, Wright F (2009) TOPALi v2: a rich graphical interface for evolutionary analyses of multiple alignments on HPC clusters and multi-core desktops. *Bioinformatics* 25:126-127
- Moszczyńska A, Locke SA, McLaughlin JD, Marcogliese DJ, Crease TJ (2009) Development of primers for the mitochondrial cytochrome *c* oxidase I gene in digenetic trematodes (Platyhelminthes) illustrates the challenge of barcoding parasitic helminths. *Mol Ecol Resour* 9:75-82
- Nezhybová V, Reichard M, Blažek R, Ondračková M (2017) Metazoan parasites of African annual killifish (Nothobranchiidae): abundance, diversity, and their environmental correlates. *Biotropica* 49(2):229-238
- Nguyen TC, Li Y-C, Makouloutou P, Jimenez LA, Sato H (2012) *Posthodiplostomum* sp. metacercariae in the trunk muscle of northern snakeheads (*Channa argus*) from the Fushinogawa River, Yamaguchi, Japan. *J Vet Med Sci* 74(10):1367-1372
- Niewiadomska K (2002) Superfamily Diplostomoidea Poirier, 1886. In: Gibson DI, Jones A, Bray RA (eds) *Keys to the Trematoda, Volume 1*, The Natural History Museum, London, pp 159-166

- Ondráčková M, Dávidová M, Gelnar M, Jurajda P (2006) Susceptibility of Prussian carp infected by metacercariae of *Posthodiplostomum cuticola* (v. Nordmann, 1832) to fish predation. Ecol Res 21:526-529
- Ondráčková M, Reichard M, Jurajda P, Gelnar M (2004a) Seasonal dynamics of *Posthodiplostomum cuticola* (Digenea, Diplostomatidae) metacercariae and parasite-enhanced growth of juvenile host fish. Parasitol Res 93:131-136
- Ondráčková M, Šimková A, Gelnar M, Jurajda P (2004b) *Posthodiplostomum cuticola* (Digenea: Diplostomatidae) in intermediate fish hosts: factors contributing to the parasite infection and prey selection by the definitive bird host. Parasitology 129:761-770
- Paes JVK, Carvalho ED, da Silva RJ (2010) Infection levels of *Austrodiplostomum compactum* (Digenea, Diplostomidae) metacercariae in *Plagioscion squamosissimus* (Teleostei, Sciaenidae) from the Nova Avanhandava reservoir, Sao Paulo State, Brazil. J Helminthol 84:284–291.
- Poulin R (1993) Age-dependent effects of parasites on anti-predator responses in two New Zealand freshwater fish. Oecologia 96:431-438
- Prenter J, MacNeil C, Dick JTA, Dunn AM (2004) Roles of parasites in animal invasions. Trends Ecol Evol 19:385-390
- Rambaut A, Drummond A (2009) FigTree: Molecular evolution, phylogenetics and epidemiology. Available at <http://tree.bio.ed.ac.uk/>
- Rambaut A, Suchard MA, Xie D, Drummond AJ (2014) Tracer v1.6. Available at <http://beast.bio.ed.ac.uk/Tracer>
- Rodríguez F, Olivier JL, Marín A, Medina JR (1990) The general stochastic model of nucleotide substitution. J Theor Biol 142:485-501
- Ronquist F, Teslenko M, van der Mark P, Ayres D, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2011) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Syst Biol 61(3):539-542

- 593 Saad-Fares A, Combes C (1992) Abundance/host size relationship in a fish trematode community. J
594 Helminthol 66:187–192.
- 595 Seppälä O, Karvonen A, Valtonen ET (2005) Manipulation of fish host by eye flukes in relation to
596 cataract formation and parasite infectivity. Anim Behav 70:89-894
- 597 Sharriff M, Richards RH, Sommerville C (1980) The histopathology of acute and chronic infections
598 of rainbow trout *Salmo gairdneri* Richardson with eye flukes, *Diplostomum* spp. J Fish Dis
599 3:455-465
- 600 Shareef PAA, Abidi SMA (2015) Studies on the epidemiology and histopathology of
601 *Euclinostomum heterostomum* (Trematoda; Digenea) infection in *Channa punctata* from
602 North India. Arch Pol Fish 23(3):133-140
- 603 Silva-Souza AT, Ludwig G (2005) Parasitism of *Cichlasoma paranaense* Kullander, 1983 and
604 *Gymnotus carapo* Linnaeus, 1814 by *Clinostomum complanatum* (Rudolphi, 1814)
605 metacercariae in the Taquari river. Braz J Biol 65(3):513-519
- 606 Sonin MD (1986) Key to the trematode parasites of fish-eating birds in Palearctic region. Nauka,
607 Moscow
- 608 Stamatakis A (2014) RAxML Version 8: A tool for phylogenetic analysis and post-analysis of large
609 phylogenies. Bioinformatics 30(9):1312-1313
- 610 Stoyanov B, Georgieva S, Pankov P, Kudlai O, Kostadinova A, Georgiev BB (2017) Morphology
611 and molecules reveal the alien *Posthodiplostomum centrarchi* Hoffman, 1958 as the third
612 species of *Posthodiplostomum* Dubois, 1936 (Digenea: Diplostomidae) in Europe. Syst
613 Parasitol 94:1–20
- 614 Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular Evolutionary
615 Genetics Analysis version 6.0. Mol Biol Evol 30: 2725-2729
- 616 Vanhove MPM, Tessens B, Schoelinck C, Jondelius U, Littlewood DTJ, Artois T, Huyse T (2013)
617 Problematic barcoding in flatworms: A case-study on monogeneans and rhabdocoels
618 (Platyhelminthes). ZooKeys 365:355-379

- 619 Vianna RT, Pereira Júnior J, Brandão DA (2005) *Clinostomum complanatum* (Digenea,
620 Clinostomidae) density in *Rhamdia quelen* (Siluriformes, Pimelodidae) from South Brazil.
2 Braz Arch Biol Techn 48(4):635-642
3
4
5
622 Vidal-Martinez VM, Pech D, Sures B, Purucker ST, Poulin R (2010). Can parasites really reveal
7
8
9
10
624 Whiteman NK, Kimball RT, Parker PG (2007) Co-phylogeography and comparative population
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 Features of the six sites sampled in autumn, number of fish examined, Standard length range and parasite prevalence (%) in the total sample, in females and males. The two sites that were also sampled seasonally are marked with an asterisk. Sites 1, 2, 3, 5 and 6 host *V. letourneuxi*; Site 4 hosts *V. robertae*. Veg denotes Vegetation cover.

Site	Date	Latitude	Longitude	Veg (%)	Temp (°C)	Salinity (ppt)	n fish	SL range (mm)	% Total	Parasite prevalence (%) % ♀ % ♂
1	Oct 2005	38° 54' 54" N	20° 52' 32" E	50	21.4	3.5	20	8.8 - 24.8	20.00	27.27 12.50
2	Oct 2005	39° 10' 31" N	20° 45' 53" E	95	15.8	0.2	16	9.7 - 24.8	50.00	33.33 71.43
3	Oct 2005	39° 05' 30" N	21° 02' 13" E	70	17.8	0.4	11	10.0 - 24.5	18.18	16.67 20.00
4	Sep 2006	38° 24' 26" N	21° 55' 03" E	80	25.0	0.3	23	16.5 - 56.0	100.00	100.00 100.00
5	Sep 2006	39° 14' 55" N	20° 28' 50" E	40	17.2	3.2	16	12.7 - 20.0	100.00	100.00 100.00
6	Sep 2009	39° 17' 15" N	20° 32' 35" E	60	16.9	0.1	9	7.0 - 29.2	55.56	40.00 75.00

Table 2 List of digenean species (Diplostomatidae) obtained from GenBank with their accession numbers for the ITS1-5.8S-ITS2 rDNA region and for the COI sequences retrieved, their host species and the country where the species were collected.

Parasite species	Host species	Country	ITS1-5.8S-ITS2 rDNA	COI mtDNA
<i>Alaria mustelae</i> Bosma, 1931	<i>Mustela frenata</i> Lichtenstein, 1831	USA	JF820609.1	KT254032.1
<i>Austrodiplostomum ostrowskiae</i> Dronen, 2009	<i>Dorosoma cepedianum</i> (Lesueur, 1818)	USA	KT728782.1	KR271028.1
<i>Bolbophorus confusus</i> (Krause, 1914)	<i>Pelecanus onocrotalus</i> Linné, 1758	Israel	AY242851.1	-
<i>Clinostomum complanatum</i> Rudolphi, 1814	<i>Triturus carnifex</i> (Laurenti, 1768)	Italy	KM518257.1	JF718595.1
<i>Diplostomum mergi</i> Dubois, 1932	<i>Radix auricularia</i> Linnaeus, 1758	Czech Republic	KR149499.1	KR149528.1
<i>Diplostomum paracaudum</i> (Iles, 1959)	<i>Gadus morhua</i> Linnaeus, 1758	Denmark/Germany	KJ889013.1	JQ639176.1
<i>Diplostomum spathaceum</i> Rudolphi, 1819	<i>Larus ridibundus</i> Linné, 1766	Czech Republic	KR269765.1	JX986895.1
<i>Ornithodiplostomum</i> sp. 1 Dubois, 1936	<i>Percina caprodes</i> (Rafinesque, 1818)	Canada	HM064937.1	HM064748.1
<i>Ornithodiplostomum</i> sp. 2 Dubois, 1936	<i>Notemigonus crysoleucas</i> (Mitchill, 1814)	Canada	HM064939.1	HM064764.1
<i>Ornithodiplostomum</i> sp. 3 Dubois, 1936	<i>Pimephales promelas</i> (Rafinesque, 1820)	Canada	HM064942.1	HM064784.1
<i>Ornithodiplostomum</i> sp. 4 Dubois, 1936	<i>Pimephales promelas</i>	Canada	HM064945.1	HM064785.1
<i>Ornithodiplostomum</i> sp. 5 Dubois, 1936	<i>Notemigonus crysoleucas</i> (Mitchill, 1814)	Canada	FJ469595.1	KT831368.1
<i>Ornithodiplostomum</i> sp. 6 Dubois, 1936	<i>Pimephales promelas</i> (Rafinesque, 1820)	Canada	HM064946.1	HM064790.1
<i>Ornithodiplostomum scardinii</i> (Schulman in Dubinin, 1952)	<i>Scardinus erythrophthalmus</i> (Linnaeus, 1758)	Czech Republic	KX931443.1	KX931425.1
<i>Posthodiplostomum brevicaudatum</i> 1(von Nordmann, 1832)	<i>Perca fluviatilis</i> (Linnaeus, 1758)	Czech Republic	KX931428.1	KX931418.1
<i>Posthodiplostomum brevicaudatum</i> 2	<i>Gasterosteus aculeatus</i> (Linnaeus, 1758)	Bulgaria	KX931429.1	KX931419.1
<i>Posthodiplostomum centrarchi</i> 1 (Hoffman, 1958)	<i>Lepomis gibbosus</i> (Linnaeus, 1758)	Bulgaria	KX931441.1	KX931421.1
<i>Posthodiplostomum centrarchi</i> 2	<i>Lepomis gibbosus</i>	Slovakia	KX931442.1	KX931423.1
<i>Posthodiplostomum</i> sp. 1 Dubois, 1936	<i>Channa punctata</i> (Bloch, 1793)	India/Canada	KF738447.1	HM064795.1
<i>Posthodiplostomum</i> sp. 2 Dubois, 1936	<i>Channa argus</i> (Cantor, 1842)	Japan/Canada	AB693170.1	HM064798.1
<i>Posthodiplostomum</i> sp. 3 Dubois, 1936	<i>Lepomis gibbosus</i>	Canada	HM064957.1	HM064821.1
<i>Posthodiplostomum</i> sp. 4 Dubois, 1936	<i>Morone americana</i> (Gmelin, 1789)	Canada	HM064960.1	HM064844.1
<i>Posthodiplostomum</i> sp. 5 Dubois, 1936	<i>Lepomis gibbosus</i>	Canada	HM064958.1	HM064857.1
<i>Posthodiplostomum</i> sp. 6 Dubois, 1936	<i>Micropterus salmoides</i> Lacepède, 1802	Canada	HM064962.1	HM064864.1
<i>Posthodiplostomum</i> sp. 7 Dubois, 1936	<i>Perca flavescens</i> Mitchill, 1814	Canada	HM064961.1	HM064865.1
<i>Tetracotyle xenentodon</i> Chakrabarti, 1970	<i>Tetracotyle xenentodon</i> (Hamilton, 1822)	India/Canada	KU316948.1	HM064876.1
<i>Tylodelphys azteca</i> García-Varela, Sereno-Uribe, Pinacho-Pinacho, Hernández-Cruz, Pérez-Ponce de León, 2015	<i>Podilymbus podiceps</i> (Linnaeus, 1758)	Mexico	KT175388.1	KT175369.1

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tylodelphys mashonensis (Beverley-Burton, 1963)

Clarias gariepinus Burchell, 1822

Tanzania

KC685363.1

KR863382.1

Figure Captions

Figure 1 Location of sampling sites in Western Greece. Mornos hosts *Valencia robertae*, while Vlychos, Arachthos, Louros, Kypseli and Acheron host *V. letourneuxi*.

Figure 2 Phylogenetic tree based on rDNA fragments from 29 haplotypes of Diplostomatidae. Posterior probabilities for Bayesian inference, (before slash) and bootstrap percentages for maximum likelihood (behind slash) are shown. Clades that neither yield a support value higher than 80 nor of 50 under BI or ML, respectively, are collapsed. The haplotypes obtained in this study are called Haplotype 1 and 2.

Figure 3 Encysted and excysted metacercariae of diplostomatid digenean from the abdominal cavity of *V. letourneuxi* and *V. robertae* (Scale bar 250 μ m), OS – oral sucker; PS – pseudosuckers; GP – genital pore; VS – ventral sucker.


Figure 4 Seasonal variation of the size frequency distribution of infected and metacercariae-free specimens in Mornos and Acheron habitats that host *V. robertae* and *V. letourneuxi* respectively. Number of specimens and parasite prevalence values (%) are also shown. The two groups of metacercariae-free fish of the winter sample are marked with asterisks.

Figure 5 Size frequency distribution of metacercariae-free and infected *V. robertae* and *V. letourneuxi* specimens of the autumn samplings (n = 94). Maximum parasite prevalence (100%) was first observed at the 22-24 mm length class (marked with asterisk).

Figure 6 Parasite weight (PW) and host weight (HW, i.e. total fish host weight) positive relationship for fish < 600 mg.



Figure1






Figure4




Figure5




Figure6

