Black carbon reduces beneficial effect of physical activity on lung function

Peer-reviewed author version

LAEREMANS, Michelle; DONS, Evi; Avila-Palencia, Ione; Carrasco-Turigas, Glòria; Orjuela-Mendoza, Juan Pablo; Anaya-Boig, Esther; Cole-Hunter, Tom; De Nazelle, Audrey; Nieuwenhuijsen, Mark; Standaert, Arnout; Van Poppel, Martine; DE BOEVER, Patrick & INT PANIS, Luc (2018) Black carbon reduces beneficial effect of physical activity on lung function. In: MEDICINE AND SCIENCE IN SPORTS AND EXERCISE, 50(9), p. 1875-1881.

DOI: 10.1249/MSS.0000000000001632

Handle: <http://hdl.handle.net/1942/27574>

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/324431512>

Black Carbon Reduces the Beneficial Effect of Physical Activity on Lung Function

Article in *Medicine & Science in Sports & Exercise* · April 2018

DOI: 10.1249/MSS.0000000000001632

CITATIONS

3

READS

247

13 authors, including:

Michelle Laeremans

Flemish Institute for Technological Research

38 PUBLICATIONS 74 CITATIONS

[SEE PROFILE](#)

Evi Dons

Hasselt University

100 PUBLICATIONS 1,661 CITATIONS

[SEE PROFILE](#)

Ione Avila-Palencia

Instituto de Salud Global de Barcelona

40 PUBLICATIONS 75 CITATIONS

[SEE PROFILE](#)

Glòria Carrasco-Turigas

Instituto de Salud Global de Barcelona

55 PUBLICATIONS 200 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:

BlueHealth [View project](#)

AVENUE [View project](#)

Medicine & Science IN Sports & Exercise

The Official Journal of the American College of Sports Medicine
www.acsm-msse.org

. . . Published ahead of Print

Black Carbon Reduces the Beneficial Effect of Physical Activity on Lung Function

Michelle Laeremans^{1,2}, Evi Dons^{1,3}, Ione Avila-Palencia^{4,5,6}, Glòria Carrasco-Turigas^{4,5,6},
Juan Pablo Orjuela-Mendoza⁷, Esther Anaya-Boig⁷, Tom Cole-Hunter^{4,5,6,8}, Audrey
de Nazelle⁷, Mark Nieuwenhuijsen^{4,5,6}, Arnout Standaert¹, Martine Van Poppel¹,
Patrick De Boever^{1,3}, and Luc Int Panis^{1,2}

¹Flemish Institute for Technological Research (VITO), Mol, Belgium; ²Transportation Research Institute (IMOB), Hasselt University, Diepenbeek, Belgium; ³Centre for Environmental Sciences, Hasselt University, Diepenbeek, Belgium; ⁴ISGlobal, Centre for Research in Environmental Epidemiology, Barcelona, Spain; ⁵Universitat Pompeu Fabra, Barcelona, Spain; ⁶CIBER Epidemiology and Public Health, Madrid, Spain; ⁷Centre for Environmental Policy, Imperial College London, London, United Kingdom; ⁸Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, CO

Accepted for Publication: 26 March 2018

Medicine & Science in Sports & Exercise® Published ahead of Print contains articles in unedited manuscript form that have been peer reviewed and accepted for publication. This manuscript will undergo copyediting, page composition, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered that could affect the content.

Black Carbon Reduces the Beneficial Effect of Physical Activity on Lung

Function

Michelle Laeremans^{1,2}, Evi Dons^{1,3}, Ione Avila-Palencia^{4,5,6}, Glòria Carrasco-Turigas^{4,5,6},
Juan Pablo Orjuela-Mendoza⁷, Esther Anaya-Boig⁷, Tom Cole-Hunter^{4,5,6,8}, Audrey de Nazelle⁷,
Mark Nieuwenhuijsen^{4,5,6}, Arnout Standaert¹, Martine Van Poppel¹,
Patrick De Boever^{1,3}, and Luc Int Panis^{1,2}

¹Flemish Institute for Technological Research (VITO), Mol, Belgium; ²Transportation Research Institute (IMOB), Hasselt University, Diepenbeek, Belgium; ³Centre for Environmental Sciences, Hasselt University, Diepenbeek, Belgium; ⁴ISGlobal, Centre for Research in Environmental Epidemiology, Barcelona, Spain; ⁵Universitat Pompeu Fabra, Barcelona, Spain; ⁶CIBER Epidemiology and Public Health, Madrid, Spain; ⁷Centre for Environmental Policy, Imperial College London, London, United Kingdom; ⁸Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, CO

Correspondence to: Prof dr Luc Int Panis, Flemish Institute for Technological Research (VITO), Boeretang 200, 2400 Mol, Belgium,

e-mail: luc.intpanis@vito.be

This research received funding from the European Union's Seventh Framework Program to the PASTA project under grant agreement No. 602624 (FP7-HEALTH-2013-INNOVATION-1) (Physical Activity through Sustainable Transport Approaches). Duration 1 November 2013 to 31

October 2017. Michelle Laeremans is supported by a VITO PhD scholarship. Evi Dons is supported by a postdoctoral scholarship from FWO-Research Foundation Flanders. Juan Pablo Orjuela-Mendoza received a beneficiary grant from Colciencias (Administrative Department of Science, Technology and Innovation of the Colombian Government) (grant number 646). All authors declare that they have no conflicts of interest and that the results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The results of the present study do not constitute endorsement by ACSM.

Abstract

Introduction: When physical activity is promoted in urban outdoor settings (e.g. walking and cycling), individuals are also exposed to air pollution. It has been reported that short-term lung function increases as a response to physical activity, but this beneficial effect is hampered when elevated air pollution concentrations are observed. Our study assessed the long-term impact of air pollution on the pulmonary health benefit of physical activity. **Methods:** Wearable sensors were used to monitor physical activity levels (SenseWear) and exposure to black carbon (microAeth) of 115 healthy adults during one week in three European cities (Antwerp, Barcelona, London). The experiment was repeated in three different seasons to approximate long-term behavior. Spirometry tests were performed at the beginning and end of each measurement week. All results were averaged on a participant level as a proxy for long-term lung function. Mixed effect regression models were used to analyze the long-term impact of physical activity, black carbon and their interaction on lung function parameters FEV₁, FVC, FEV₁/FVC, FEF₂₅₋₇₅ and PEF. Interaction plots were used to interpret the significant interaction effects. **Results:** Negative interaction effects of physical activity and black carbon exposure on FEV₁ (p=0.07), FEV₁/FVC (p=0.03) and FEF₂₅₋₇₅ (p=0.03) were observed. For black carbon concentrations up to approximately 1 µg/m³, an additional METhour per week resulted in a trend towards lung function increases (FEV₁, FEV₁/FVC and FEF₂₅₋₇₅ increased 5.6 mL, 0.1% and 14.5 mL/s, respectively). **Conclusion:** We found that lung function improved with physical activity at low black carbon levels. This beneficial effect decreased in higher air pollution concentrations. Our results suggest a greater need to reduce air pollution exposures during physical activity.

Key words: air pollution, active mobility, Tiffeneau, FVC, FEV₁

Introduction / Background

Today's global physical inactivity pandemic is responsible for approximately five million premature deaths annually (1). However, when physical activity in urban outdoor settings (e.g. walking and cycling) is promoted, air quality may need to be taken into account.

Long-term air pollution exposure contributes to increased prevalence of cardiovascular and respiratory diseases (2). Globally, many people live in urbanized regions, which are hotspots of air pollution due to poor urban and transport planning (3). Consequently, walking, cycling and other forms of both incidental and intentional physical activity in urban environments often involve exposure to high concentrations of air pollution.

Higher ventilation rates during physical activity result in an increased pollutant dose and deeper penetration of particles into the lungs (4,5). Nevertheless, various health impact assessments have been carried out and reported that the overall benefits of physical activity outweigh the risks of air pollution (6–8). Also, the Danish Diet, Cancer and Health Cohort did not observe a modification effect of air pollution on the relationship between physical activity and all-cause mortality (9). However, short-term respiratory responses to physical activity have been shown to alter with increasing air pollution concentrations (10,11). A recent study in older adults, with and without pre-existing chronic obstructive pulmonary disease (COPD) or ischaemic heart disease, found that the beneficial effects of walking on lung function were lost after a two-hour walk in a polluted environment (11). In addition, lung function responses to physical activity in healthy participants also differed when air pollution was taken into account (10,12). Evidence on the chronic impact of air pollution on the health benefit of physical activity for specific physiological systems remains absent (5). Therefore, the main aims of the current study were: (1) to assess the chronic effects of physical activity and air pollution on lung function and (2) to assess whether

the pulmonary responses to physical activity and air pollution interact in a sample of healthy adults across three European cities.

Methods

Study design and participants

The framework of this study was the PASTA project (Physical Activity through Sustainable Transport Approaches). In this project, data on physical activity and travel behavior among over 12,500 volunteers living or working in seven European cities (Antwerp, Barcelona, London, Örebro, Rome, Vienna, Zurich) was collected through an online survey (13,14). From this sample, 122 participants were selected to participate in a real-life monitoring study in three cities (Antwerp: 41 participants, Barcelona: 41 participants, London: 40 participants). High-resolution data on physical activity and black carbon (BC) exposure was collected during seven consecutive days using wearable sensors. Lung function was assessed with a spirometry test at the beginning and end of the measurement week. Each participant repeated the measurement week three times: in the mid-season (spring or autumn), in the summer and in the winter, between February 2015 and March 2016. Eligible participants were non-smoking, 18-65 year olds with a BMI below 30 and no self-reported cardiovascular, respiratory or neurological condition. The study was approved by the Ethics Committee of the University hospital in Antwerp (UZA), the Clinical Research Ethical Committee of the Parc de Salut Mar in Barcelona and the Imperial College Research Ethics Committee in London. All participants gave written informed consent prior to participation.

Physical activity assessment

Physical activity was objectively measured with the SenseWear armband (model MF-SW, BodyMedia, USA). This multi-sensor body monitor measures heat flux, galvanic skin response, skin temperature and 3-axis accelerometry. The sensor was worn on the triceps muscle of the left arm and was only removed when there was contact with water (bathing, showering, etc.); wearing time was on average 96 ± 4 % of the total time. Age, sex, body weight and height of the participants were entered in the SenseWear professional software (version 8.0). Energy expenditure and METs were calculated on a one-minute basis using proprietary algorithms based on pattern recognition (15). For each measurement week, the amount of METhours was calculated in R version 3.3.1. Only bouts of at least 10 consecutive minutes with an intensity ≥ 3 METs were considered. This is in accordance with the WHO recommendations on physical activity for health (16). To clarify, an activity that requires 3 METs is walking the dog and doing this for one hour results in 3 METhours (according to the compendium of physical activities (17)).

BC exposure assessment

Personal exposure to BC was assessed with the microAeth (model AE51, Aethlabs, USA). Ambient air was drawn over a Teflon-coated borosilicate glass fibre filter at a flow rate of 100 mL/min, resulting in BC accumulation on the filter. Light attenuation at a wavelength of 880 nm was measured and converted into a BC concentration (ng/m^3). The device was set to estimate the average BC concentration every five minutes. Participants always carried the device with them and replaced the filter every two days to prevent saturation. For indoor activities, participants left the microAeth in the room where they spent most of their time. A short tube was attached to the inlet of the microAeth which enabled participants to carry the device in a bag.

Raw BC data were smoothened with the ONA algorithm (Optimized Noise-Reduction Algorithm; developed by the Environmental Protection Agency), and data tagged with an error code that could potentially impact concentration levels were excluded.

Lung function measurement

Spirometry (EasyOne, ndd Medical Technologies) was performed according to the European Respiratory Society and the American Thoracic Society guidelines (18,19). Lung function parameters selected for this study are forced expiratory volume in the first second (FEV_1), forced vital capacity (FVC), the Tiffeneau index (FEV_1/FVC), peak expiratory flow (PEF) and forced expiratory flow at 25% to 75% of the FVC (FEF_{25-75}). Percentage predicted values of FEV_1 and FVC were calculated based on the Global Lung Function 2012 equations.

Analysis

For each measurement week, the participants' physical activity level was calculated as total METhours, and their BC exposure was averaged. We calculated the mean of the three measurements per participant as a proxy for long-term physical activity behavior and long-term personal BC exposure. This is justified as we objectively measured physical activity and BC concentrations during a whole week in three different seasons. Consequently, differences due to seasonal changes were accounted for. Three participants did not complete all three measurement weeks and were excluded from the analysis. Also, only non-smokers ($n=100$) and former smokers ($n=19$) were recruited. Pack years of former smokers were low with an average of 2.7 ± 2.9 . This corresponds to 20 cigarettes per day during 2.7 years. Four of those former smokers were excluded from the analysis since they quit smoking less than 5 years ago. This resulted in a total of 115 eligible participants out of 122. Long-term lung function parameters

were approximated by calculating the average FEV₁, FVC, FEV₁/FVC, FEF₂₅₋₇₅ and PEF per participant.

The study population was described by sex, age, ethnicity, height, BMI, and education level. For categorical variables the amount (n) and percentage was reported. Arithmetic mean and standard deviation (SD) were used for continuous variables and pulmonary outcomes. In addition, the overall and city-specific median and interquartile range (IQR) of physical activity and BC concentrations were calculated.

The effects of long-term physical activity, air pollution and their interaction were tested using mixed effect regression models where city was introduced as a random variable. First, we tested the effects of physical activity and BC on lung function separately. Then the effects of physical activity and BC were estimated together in one single model. Finally, models including the interaction term were tested. All models included sex, age and height based on the European Respiratory Society and the American Thoracic Society guidelines (18). To account for socio-economic differences, education level was also added as a covariate.

The significance level was set at $p < 0.05$. When trends towards significant interaction effects were observed ($p < 0.1$), regression estimates of physical activity were plotted with their 95% confidence interval as a function of BC concentrations in an interaction plot.

R software version 3.3.1 was used for data processing and analysis. Random intercept models were tested with the lme4 and lmerTest R-packages for mixed effect regression analysis. All marginal residuals were normally distributed. Interaction plots were made using the R-package Interplot.

Results

A total of 115 healthy adults were included in the analysis. Almost half of the study population (44%) were males; 94% were Caucasian and 90% had a higher education degree (Table 1). They were on average 36.6 ± 10 years old, 1.7 ± 0.1 m tall with a BMI of 23.7 ± 3 kg/m². Long-term FEV₁ ranged from 2.28 to 5.13 L with an average of 3.53 ± 0.70 L (mean % predicted FEV₁: $93.9 \pm 10.6\%$); FVC ranged from 2.63 to 6.78 L with an average of 4.44 ± 0.92 L (mean % predicted FVC: $96.8 \pm 10.7\%$). The median, long-term FEV₁/FVC ratio was 80.7%. Hence, we recruited a sample of participants with normal lung function (18).

Table 2 shows the participants' physical activity level and BC exposure aggregated over all three measurement weeks per individual. In general, we recruited a physically active sample: the median amount of METhours per week was 42, which exceeds the WHO recommendation of 10 METhours. The least active population was recruited in Barcelona, while a similar physical activity level was observed for participants in Antwerp and London. On the other hand, BC concentrations were highest in Barcelona. Overall, median BC exposures were 1.4 µg/m³.

Mixed effect regression analysis was performed using models with and without the interaction term of physical activity and BC exposure (Table 3). There were no significant effects of physical activity and BC on the lung function parameters in models excluding the interaction term. This changed when the interaction term was included. These models showed significant, negative interaction effects of physical activity and BC exposure on FEV₁ ($p=0.07$), the FEV₁/FVC ratio ($p=0.03$) and FEF₂₅₋₇₅ ($p=0.03$).

The effect of physical activity increased to borderline significant levels when the interaction term was included. According to these models, FEV₁, FEV₁/FVC and FEF₂₅₋₇₅ would gain 5.6 mL ($p=0.13$), 0.1 % ($p=0.08$) and 14.5 mL/s ($p=0.06$) per additional weekly METhour. However, the

interaction term also shows trends towards significance which means that the value and significance of the main physical activity effect should be interpreted for specific BC concentrations.

None of the models could detect a significant effect of physical activity and BC on FVC and PEF.

The effects of physical activity on FEV_1 , FEV_1/FVC and FEF_{25-75} were plotted as a function of BC concentration. Figure 1 visualizes how increasing BC concentrations modify the effect of physical activity on FEV_1 , FEV_1/FVC and FEF_{25-75} . The dashed vertical line represents the BC concentrations where the 95% confidence interval of the estimated effect of physical activity on FEV_1 , FEV_1/FVC and FEF_{25-75} only included negative values. Consequently, based on our results, FEV_1 , FEV_1/FVC and FEF_{25-75} improves with physical activity at lower BC concentrations. At high BC levels, additional physical activity may reduce these lung function parameters.

Discussion

This is the first study to examine the combined long-term effects of air pollution and physical activity on lung function using personal measurements obtained with wearable devices. We found that physical activity is associated with an improved pulmonary function at low BC concentrations. This respiratory benefit decreased when BC concentrations increased.

BC is a fraction of particulate matter with most particles having an aerodynamic diameter smaller than 1 μm . Such small particles are believed to be more harmful than particles of larger sizes (20). This makes BC a valuable marker to study physiological effects of air pollution (21). Overall, we measured a median BC concentration of 1.4 $\mu g/m^3$, with the highest concentrations

in Barcelona ($1.7 \mu\text{g}/\text{m}^3$). These concentrations were similar to those measured in the ESCAPE project, which covered a broad range of areas in Europe ($\text{PM}_{2.5}$ absorbance was converted to BC based on the formulas in Dons (24) and Gan et al. (25)). The ESCAPE cohort found a negative association between long-term NO_2 and PM_{10} exposure and lung function in adults (24). However, the relationship was not significant for $\text{PM}_{2.5}$ absorbance (a marker for BC) which is consistent with our results.

At low BC concentrations up to approximately $1 \mu\text{g}/\text{m}^3$, the effect of an additional METhour per week on FEV_1 , FEV_1/FVC and FEF_{25-75} was an increase of 5.6 mL, 0.1 % and 14.5 mL/s respectively. This benefit decreased with increasing BC concentrations. Our findings are in line with observations from a study by Cheng et al. (2003), executed in Texas. In this study, self-reported exercise patterns of healthy adults between 25 and 55 years old were associated with spirometry results (25). These participants were categorized according to their physical activity level where the highly active participants were approximately 16 METhour per week more active than those with a low physical activity level (based on the formulas in Ainsworth et al. (28)). Highly active men had a 130 mL higher FEV_1 ; for women, the difference was 80 mL. This corresponded to a 8 mL and 5 mL FEV_1 increase per METhour for men and women respectively with an annual BC concentration $0.7 \mu\text{g}/\text{m}^3$ in Texas, similar to our results (26). Other studies have also observed correlations between physical activity related variables and FEV_1 or FVC (27,28).

Little research has been done on the chronic impact of air pollution on the respiratory benefits of physical activity. A review by Giles et al. (2014) suggested that chronic exposure during exercise has important consequences for respiratory function (29). Hence, a higher prevalence of exercise-induced bronchoconstriction, asthma and lower lung function has been observed in

athletes who train in environments with high particulate matter emissions (5). Moreover, reductions in respiratory mortality associated with cycling were greater for participants with low to moderate compared to high air pollution exposure in the Danish Diet, Cancer and Health Cohort (9).

It is well documented that inhalation of air pollutants causes oxidative damage and inflammation in the respiratory system resulting in airway obstruction and a reduced FEV_1 (5,30,31). Exercising in polluted air also results in inhalation of a higher pollutant dose which may enhance its adverse respiratory effects (4,5,31). Contrary, regular physical activity improves endurance and strength of the respiratory muscles resulting in an increased FEV_1 (30). However, compared to the cardiovascular system, respiratory adaptations to physical activity are rather small because the lungs are already equipped to deal with the demands of high intensity exercise. Consequently, exercise in elevated air pollution concentrations may lead to a level of obstruction potentially outweighing the beneficial effect of physical activity on FEV_1 and FEF_{25-75} . Therefore, chronic exposure to high air pollution doses during physical activity may result in a decreased FEV_1 , FEV_1/FVC and FEF_{25-75} . Since FEF_{25-75} specifically informs us about small airway function, this may indicate that physical activity and BC interact at this level (32,33).

Contrary, various health impact assessment reported that the overall physical activity benefits outweigh the negative physiological effects of air pollution (8). However, the modification effects on physiological systems are potentially masked in previous studies as they did not investigate such subclinical reactions. Our results suggest that there may be a greater need to reduce air pollution exposures during physical activity than was previously thought.

Our sample consisted of healthy, highly educated adults with an average FEV_1/FVC ratio of $80.1 \pm 6.4\%$. The level of physical activity measured in our sample meets the WHO

recommendation of 600 METminutes to maintain health. Overall, the IQR of measured METhours falls within light to moderate physical activity levels (2). Our findings aren't entirely representative of the general population as we recruited mostly active and highly educated volunteers of Caucasian ethnicity. We covered, however, multiple European cities which is unique in this field of research. We also recruited a larger sample compared to previous studies where air pollution exposure was assessed on a personal level (12,34). Moreover, measuring air pollution exposure and physical activity with wearable sensors reduces the risk of exposure misclassification. This is of particular importance when studying traffic-related pollutants such as BC of which the concentrations rapidly change in space and time (5). Accurately assessing both physical activity and air pollution exposure remains a major challenge in epidemiological research. In addition, we were able to use our personal measurements as a proxy for long-term behavior as all participants collected exposure data in three different seasons.

It has been reported that the SenseWear underestimates energy expenditure during high intensity activities which is a limitation of our study design (35). However, especially MET values above 10 METs are underestimated which our participants only achieved during 0.2 % of their time awake (35). In addition, the most recent version of the SenseWear provides the best estimation compared to previous SenseWear versions and the Actigraph, the most widely used wearable physical activity tracker in research. Another limitation is that only a limited number of participants was exposed to the highest BC concentrations where we found a negative relationship between physical activity and respiratory health. Consequently, we were unable to robustly characterize the association at these BC concentrations. Also, the addition of an interaction term to our models resulted in wider confidence intervals. This may be due to the limited amount of data available for each combination of physical activity and BC levels. While

our results are of potential interest to public health, the study needs to be replicated in a larger study sample to confirm them. Finally, high average physical activity levels of our participants may have affected the results. Potentially, the respiratory benefit of physical activity in low BC concentrations is larger in a less active sample. In addition, the interaction effect may be smaller since the baseline amount of inhaled particles during physical activity would be lower, also lowering the chances to provide a respiratory hazard.

Due to the potentially high impact on public health, we recommend future research to further characterize the modification effect of air pollution on the relationship between physical activity and long-term respiratory health. We suggest to (1) complement spirometry with other measures providing information on different parts of the respiratory system (e.g. frequency of self-reported respiratory symptoms of both the upper and lower tract), and (2) to investigate the combined effects of air pollution and physical activity in vulnerable sub-populations such as children, athletes and pulmonary patients.

Conclusion

This is the first study executed in multiple European cities simultaneously that integrated effects of air pollution and physical activity to assess their combined impact on pulmonary function. We found a long-term beneficial effect of long-term physical activity on respiratory at low BC concentrations. The beneficial effect decreased with increasing yearly, average BC concentrations in a healthy adult population. This illustrates the need for an integrated approach to better understand the health impact of strategies and policy measures that try to tackle physical inactivity and pollution.

Acknowledgements

We are grateful to all study participants, and to Eva Govarts for her excellent support in the statistical analysis.

Conflicts of Interest and Source of Funding:

This research received funding from the European Union's Seventh Framework Program to the PASTA project under grant agreement No. 602624 (FP7-HEALTH-2013-INNOVATION-1) (Physical Activity through Sustainable Transport Approaches). Duration 1 November 2013 to 31 October 2017. Michelle Laeremans is supported by a VITO PhD scholarship. Evi Dons is supported by a postdoctoral scholarship from FWO-Research Foundation Flanders. Juan Pablo Orjuela-Mendoza received a beneficiary grant from Colciencias (Administrative Department of Science, Technology and Innovation of the Colombian Government) (grant number 646). All authors declare that they have no conflicts of interest and that the results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The results of the present study do not constitute endorsement by ACSM.

Ethics committee approval

The study was approved by the Ethics Committee of the University hospital in Antwerp (UZA), the Clinical Research Ethical Committee of the Parc de Salut Mar in Barcelona and the Imperial College Research Ethics Committee in London.

References

1. Lee I, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012;380(9838):219–29.
2. Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions , 1990 – 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2224–2060.
3. Nieuwenhuijsen MJ. Urban and transport planning, environmental exposures and health- new concepts, methods and tools to improve health in cities. *Environ Heal*. 2016;15(Suppl 1):38;161-71.
4. Dons E, Laeremans M, Orjuela JP, et al. Wearable sensors for personal monitoring and estimation of inhaled traffic-related air pollution: evaluation of methods. *Environ Sci Technol*. 2017;51(3):1859–67.
5. Rundell KW. Effect of air pollution on athlete health and performance. *Br J Sports Med*. 2012;46(6):407–12.
6. Johan de Hartog J, Boogaard H, Nijland H, Hoek G. Do the health benefits of cycling outweigh the risks? *Environ Health Perspect*. 2010 Aug;118(8):1109–16.
7. Buekers J, Dons E, Elen B, Int Panis L. Health impact model for modal shift from car use to cycling or walking in Flanders: application to two bicycle highways. *J Transp Heal*. 2015;2(4):549–62.
8. Mueller N, Rojas-rueda D, Cole-hunter T, et al. Health impact assessment of active

- transportation: A systematic review. *Prev Med (Baltim)*. 2015;76:103–14.
9. Andersen ZJ, de Nazelle A, Mendez MA, et al. A study of the combined effects of physical activity and air pollution on mortality in elderly urban residents: The Danish Diet, Cancer, and Health Cohort. *Environ Heal Perspect Heal Perspect*. 2015;123(6):557–63.
 10. Matt F, Cole-hunter T, Donaire-Gonzalez D, et al. Acute respiratory response to traffic-related air pollution during physical activity performance. *Environ Int*. 2016;97:45–55.
 11. Sinharay R, Gong J, Barratt B, et al. Respiratory and cardiovascular responses to walking down a traffic-polluted road compared with walking in a traffic-free area in participants aged 60 years and older with chronic lung or heart disease and age-matched healthy controls: a randomised, crossover. *Lancet*. 2017;6736(17):1–11.
 12. Kubesch N, de Nazelle A, Westerdahl D, et al. Respiratory and inflammatory responses to short-term exposure to traffic-related air pollution with and without moderate physical activity. *Occup Environ Med*. 2015;72(4):284–93.
 13. Dons E, Götschi T, Nieuwenhuijsen M, et al. Physical Activity through Sustainable Transport Approaches (PASTA): protocol for a multi-centre, longitudinal study. *BMC Public Health*. 2015;15(1126):1–11.
 14. Gerike R, de Nazelle A, Nieuwenhuijsen M, et al. Physical Activity through Sustainable Transport Approaches (PASTA): a study protocol for a multicentre project. *BMJ Open*. 2016;6(1):1–11.
 15. Bassett DRJ, Rowlands A V, Trost SG. Calibration and validation of wearables monitors. *Med Sci Sports Exerc*. 2012;44(1):1–13.
 16. WHO. Global recommendations on physical activity for health. 2010. 58 p.

17. Ainsworth BE, Haskell WIL, Whitt MC, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc.* 2000;32(9 Suppl):S498–504.
18. Quanjer PH, Stanojevic S, Cole TJ, et al. Multi-ethnic reference values for spirometry for the 3-95 year age range: the global lung function 2012 equations. *Eur Respir J.* 2012;40(6):1324–43.
19. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J.* 2005;26(2):1–22.
20. Seaton A, Godden D, MacNee W, Donaldson K. Particulate air pollution and acute health effects. *Lancet.* 1995;345(8943):176–8.
21. Janssen NAH, Hoek G, Simic-Lawson M, et al. Black carbon as an additional indicator of the adverse health effects of airborne particles compared with PM₁₀ and PM_{2.5}. *Environ Health Perspect.* 2011;119(12):1691–9.
22. Dons E. Air pollution exposure assessment through personal monitoring and activity-based modeling [dissertation]. Hasselt University; 2013. 302 p.
23. Gan WQ, Koehoorn M, Davies HW, Demers PA, Tamburic L, Brauer M. Long-term exposure to traffic-related air pollution and the risk of coronary heart disease hospitalization and mortality. *Environ Health Perspect.* 2011;119(4):501–7.
24. Adam M, Schikowski T, Carsin AE, et al. Adult lung function and long-term air pollution exposure. ESCAPE: a multicentre cohort study and meta-analysis. *Eur Respir J.* 2015;45(1):38–50.
25. Cheng YJ, Macera CA, Addy CL, Sy FS, Wieland D, Blair SN. Effects of physical activity on exercise tests and respiratory function. *Br J Sports Med.* 2003;37(6):521–8.

26. Barrett TE, Sheesley RJ. Urban impacts on regional carbonaceous aerosols: case study in central Texas. *J Air Waste Manage Assoc.* 2014;64(8):917–26.
27. Camarri B, Eastwood PR, Cecins NM, Thompson PJ, Jenkins S. Six minute walk distance in healthy subjects aged 55 – 75 years. *Respir Med.* 2006;100(4):658–65.
28. Jakes RW, Day NE, Patel B, et al. Physical inactivity is associated with lower forced expiratory volume in 1 second - European Prospective Investigation into Cancer-Norfolk prospective population study. *Am J Epidemiol.* 2002;156(2):139–47.
29. Giles L V, Koehle MS. The health effects of exercising in air pollution. *Sport Med.* 2014;44(2):223–49.
30. McKenzie DC. Respiratory physiology: adaptations to high-level exercise. *Br J Sports Med.* 2012;46(6):381–4.
31. Int Panis L, de Geus B, Vandenbulcke G, et al. Exposure to particulate matter in traffic: a comparison of cyclists and car passengers. *Atmos Environ.* 2010;44(19):2263–70.
32. Rao DR, Gaffin JM, Baxi SN, Sheenan WJ, Hoffman EB, Phipatanakul W. The utility of forced expiratory flow between 25% and 75% of vital capacity in predicting childhood asthma morbidity and severity. *J asthma.* 2012;49(6):586–92.
33. Marseglia GL, Cirillo I, Vizzaccaro A, et al. Role of forced expiratory flow at 25–75% as an early marker of small airways impairment in subjects with allergic rhinitis. *J Allergy Clin Immunol.* 2007;28(1):74–9.
34. Weichenthal S, Kulka R, Dubeau A, Martin C, Wang D, Dales R. Traffic-related air pollution and acute changes in heart rate variability and respiratory function in urban cyclists. *Environ Health Perspect.* 2011;119(10):1373–8.
35. Lopez GA, Brønd JC, Dencker M, Arvidsson D. Validation of SenseWear Armband in children, adolescents, and adults. *Scand J Med Sci Sports.* 2017;28(2):487–495.

Figure captions

Figure 1 Regression estimates of physical activity effects (95% confidence interval) on FEV₁ (A), FEV₁/FVC (B) and FEF₂₅₋₇₅ (C) as a function of BC. Barplots on the x-axis represent the number of participants exposed to the corresponding BC concentration. The dashed vertical line represents the BC concentration where the 95% confidence interval of the effect of physical activity only includes negative values. This means that the lung function parameters decrease with additional physical activity at BC concentrations above the highlighted levels.

Figure 1

Table 1 Characteristics of the study population and pulmonary outcomes aggregated over all sessions per individual (n=115). Categorical characteristics are reported as n (%); continuous variables are reported as mean±SD. The percentage predicted values of FEV1 and FVC were calculated based on the 2012 Global Lung Function equations.

Personal characteristics	
Men	51 (44%)
Age (years)	36.6±10
Ethnicity (Caucasian)	108 (94%)
Height (m)	1.7±0.1
BMI (kg/m ²)	23.7±3
Higher education	104 (90%)
Pulmonary outcomes (mean±SD)	
FEV ₁	3.53±0.69 L 93.9±10.6 % pred.
FVC	4.44±0.92 L 96.8±10.7 % pred.
FEV ₁ /FVC	80±6.3 %
PEF	8.58±1.87 L/s
FEF ₂₅₋₇₅	3.41 ± 0.95 L/s

Table 2 Median and IQR of the average physical activity level (PA) and BC concentrations per participant (n=115).

	Overall	Antwerp	Barcelona	London
PA level (METhours/week)	42 (28-74)	56 (36-89)	32 (17-59)	51 (29-82)
BC ($\mu\text{g}/\text{m}^3$)	1.4 (1.1-1.7)	1.3 (1.1-1.6)	1.7 (1.5-1.9)	1.3 (1.0-1.5)

Table 3 Estimated effects of physical activity (METhour/week), BC ($\mu\text{g}/\text{m}^3$) and their interaction on lung function parameters : FEV1, FVC, FEV1/FVC, PEF and FEF₂₅₋₇₅. Estimates were obtained using mixed effect regression analysis (n=115). The first column specifies the main effects that were included in the associated models. All models included sex, age, height and education levels as covariates. City was introduced as a random effect. PA = physical activity; BC = Black carbon

Outcome	PA (METhour/week)		BC ($\mu\text{g}/\text{m}^3$)		PA x BC	
	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p
FEV1 (mL)						
PA	-1.01 (-3.14;1.12)	0.35				
BC			-74.30 (-248.21;96.62)	0.40		
PA+BC	-0.95 (-3.09;1.19)	0.39	-68.87 (-243.40;105.66)	0.44		
PA+BC+PAxBC	5.61 (-1.65;12.87)	0.13	189.90 (-133.77;513.57)	0.25	-4.73 (-9.74;0.28)	0.07
FVC (mL)						
PA	0.10 (-2.40;2.60)	0.94				
BC			19.59 (-184.81;223.99)	0.85		
PA+BC	0.08 (-2.44;2.61)	0.95	19.11 (-186.74;224.95)	0.86		
PA+BC+PAxBC	2.01 (-6.67;10.70)	0.65	95.22 (-292.22;482.65)	0.63	-1.39 (-7.39;4.60)	0.65
FEV1/FVC (%)						
PA	-0.03 (-0.06;0.004)	0.09				
BC			-2.57 (-5.10;-0.04)	0.05		
PA+BC	-0.02 (-0.05;0.01)	0.23	-2.18 (-4.73;0.36)	0.10		
PA+BC+PAxBC	0.09 (-0.009;0.19)	0.08	2.21 (-2.30;6.72)	0.34	-0.08 (-0.15;-0.01)	0.03
PEF (mL/s)						
PA	-3.90 (-9.66;1.86)	0.19				
BC			-208.10 (-698.29;282.08)	0.41		
PA+BC	-3.54 (-9.41;2.34)	0.24	-144.61 (-641.56;352.34)	0.57		
PA+BC+PAxBC	5.60 (-13.61;24.82)	0.57	225.03 (-663.26;1131.32)	0.62	-6.64 (-19.90;6.63)	0.33
FEF₂₅₋₇₅ (mL/s)						
PA	-2.37 (-6.78;2.04)	0.30				
BC			-336.44 (-718.72;45.83)	0.10		
PA+BC	-1.60 (-6.04;2.85)	0.49	-306.46 (-695.13;82.20)	0.14		
PA+BC+PAxBC	14.54 (-0.20;29.27)	0.06	344.86 (-337.35;1027.07)	0.33	-11.70(-21.89;-1.51)	0.03