Biophysical skin measurements to evaluate the effectiveness of photobiomodulation therapy in the prevention of acute radiation dermatitis in breast cancer patients

Peer-reviewed author version

ROBIJNS, Jolien; Censabella, S.; Claes, S.; Pannekoeke, L.; Bussé, L.; Colson, D.; Kaminski, I.; LODEWIJCKX, Joy; BULENS, Paul; MAES, Annelies; Noé, L.; Brosens, M.; Timmermans, A.; LAMBRICHTS, Ivo; SOMERS, Veerle & MEBIS, Jeroen (2018) Biophysical skin measurements to evaluate the effectiveness of photobiomodulation therapy in the prevention of acute radiation dermatitis in breast cancer patients. In: Supportive care in cancer, 27(4), p. 1245-1254.

DOI: 10.1007/s00520-018-4487-4

Handle: <http://hdl.handle.net/1942/28110>

Dear Author

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style.
- Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s00520-018-4487-4>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

1	Article Title	Biophysical skin measurements to evaluate the effectiveness of photobiomodulation therapy in the prevention of acute radiation dermatitis in breast cancer patients	
2	Article Sub- Title		
3	Article Copyright - Year	Springer-Verlag GmbH Germany, part of Springer Nature 2018 (This will be the copyright line in the final PDF)	
4	Journal Name	Supportive Care in Cancer	
5	Corresponding Author	Family Name	Robijns
6		Particle	
7		Given Name	Jolien
8		Suffix	
9		Organization	Hasselt University
10		Division	Faculty of Medicine & Life Sciences
11		Address	Martelarenlaan 42, Hasselt 3500, Belgium
12		e-mail	jolien.robijns@uhasselt.be
13	Author	Family Name	Censabella
14		Particle	
15		Given Name	Sandrine
16		Suffix	
17		Organization	Jessa Hospital
18		Division	Department of Medical Oncology
19		Address	Stadsomvaart 11, Hasselt, Belgium
20		e-mail	
21	Author	Family Name	Claes
22		Particle	
23		Given Name	Stefan
24		Suffix	
25		Organization	Jessa Hospital
26		Division	Limburg Oncology Centre
27		Address	Stadsomvaart 11, Hasselt, Belgium
28		e-mail	
29	Author	Family Name	Pannekoeke
30		Particle	

31		Given Name	Luc
32		Suffix	
33		Organization	Jessa Hospital
34		Division	Limburg Oncology Centre
35		Address	Stadsomvaart 11, Hasselt, Belgium
36		e-mail	
37		Family Name	Bussé
38		Particle	
39		Given Name	Lore
40		Suffix	
41	Author	Organization	Hasselt University
42		Division	Faculty of Medicine & Life Sciences
43		Address	Martelarenlaan 42, Hasselt 3500, Belgium
44		e-mail	
45		Family Name	Colson
46		Particle	
47		Given Name	Dora
48		Suffix	
49	Author	Organization	Hasselt University
50		Division	Faculty of Medicine & Life Sciences
51		Address	Martelarenlaan 42, Hasselt 3500, Belgium
52		e-mail	
53		Family Name	Kaminski
54		Particle	
55		Given Name	Iris
56		Suffix	
57	Author	Organization	Hasselt University
58		Division	Faculty of Medicine & Life Sciences
59		Address	Martelarenlaan 42, Hasselt 3500, Belgium
60		e-mail	
61		Family Name	Lodewijckx
62		Particle	
63		Given Name	Joy
64		Suffix	
65	Author	Organization	Hasselt University
66		Division	Faculty of Medicine & Life Sciences
67		Address	Martelarenlaan 42, Hasselt 3500, Belgium
68		e-mail	

69		Family Name	Bulens
70		Particle	
71		Given Name	Paul
72		Suffix	
73		Organization	Jessa Hospital
74	Author	Division	Department of Medical Oncology
75		Address	Stadsomvaart 11, Hasselt, Belgium
76		Organization	Jessa Hospital
77		Division	Limburg Oncology Centre
78		Address	Stadsomvaart 11, Hasselt, Belgium
79		e-mail	
80		Family Name	Maes
81		Particle	
82		Given Name	Annelies
83		Suffix	
84		Organization	Jessa Hospital
85	Author	Division	Department of Medical Oncology
86		Address	Stadsomvaart 11, Hasselt, Belgium
87		Organization	Jessa Hospital
88		Division	Limburg Oncology Centre
89		Address	Stadsomvaart 11, Hasselt, Belgium
90		e-mail	
91		Family Name	Noé
92		Particle	
93		Given Name	Leen
94		Suffix	
95		Organization	Jessa Hospital
96	Author	Division	Department of Medical Oncology
97		Address	Stadsomvaart 11, Hasselt, Belgium
98		Organization	Jessa Hospital
99		Division	Limburg Oncology Centre
100		Address	Stadsomvaart 11, Hasselt, Belgium
101		e-mail	
102		Family Name	Brosens
103		Particle	
104	Author	Given Name	Marc
105		Suffix	
106		Organization	Jessa Hospital

107		Division	Department of Medical Oncology
108		Address	Stadsomvaart 11, Hasselt, Belgium
109		Organization	Jessa Hospital
110		Division	Limburg Oncology Centre
111		Address	Stadsomvaart 11, Hasselt, Belgium
112		e-mail	
113		Family Name	Timmermans
114		Particle	
115		Given Name	An
116		Suffix	
117	Author	Organization	Jessa Hospital
118		Division	Department of Dermatology
119		Address	Stadsomvaart 11, Hasselt, Belgium
120		e-mail	
121		Family Name	Lambrichts
122		Particle	
123		Given Name	Ivo
124		Suffix	
125	Author	Organization	Hasselt University
126		Division	Faculty of Medicine & Life Sciences
127		Address	Martelarenlaan 42, Hasselt 3500, Belgium
128		e-mail	
129		Family Name	Somers
130		Particle	
131		Given Name	Veerle
132		Suffix	
133	Author	Organization	Hasselt University
134		Division	Faculty of Medicine & Life Sciences
135		Address	Martelarenlaan 42, Hasselt 3500, Belgium
136		e-mail	
137		Family Name	Mebis
138		Particle	
139		Given Name	Jeroen
140		Suffix	
141	Author	Organization	Hasselt University
142		Division	Faculty of Medicine & Life Sciences
143		Address	Martelarenlaan 42, Hasselt 3500, Belgium
144		Organization	Jessa Hospital

145	Division	Department of Medical Oncology
146	Address	Stadsomvaart 11, Hasselt, Belgium
147	Organization	Jessa Hospital
148	Division	Limburg Oncology Centre
149	Address	Stadsomvaart 11, Hasselt, Belgium
150	e-mail	
151	Received	30 April 2018
152	Schedule	Revised
153	Accepted	24 September 2018
154	Abstract	<p>Purpose: The purpose of this study was to evaluate objectively the effectiveness of photobiomodulation therapy (PBMT) for the prevention of acute radiation dermatitis (ARD) by using biophysical skin measurements.</p> <p>Methods: A randomized, placebo-controlled trial with 120 breast cancer patients who underwent an identical radiotherapy (RT) regimen post-lumpectomy was performed (TRANSDERMIS trial). Patients were randomized to receive PBM (808 nm CW/905 nm pulsed, 168 mW/cm², spot size 19.6 cm², fluence 4 J/cm²) or placebo treatments from the first day of RT (2×/week). Biophysical skin measurements were collected to assess the skin pigmentation and barrier function. Measurements were collected at the first day of RT, a RT dose of 40 Gray (Gy), and the end of RT (66 Gy).</p> <p>Results: The incidence of moist desquamation was significantly higher in the control than in the PBMT group at the end of RT (30 vs. 7%, respectively, odds ratio = 6, <i>p</i> = 0.004). The biophysical skin measures showed that the mean percentage change from the baseline transepidermal water loss (TEWL), erythema, and melanin values was significantly higher in the control than in the PBMT group at the end of RT (<i>ps</i> < 0.05). Logistic regression analysis revealed that the risk on moist desquamation was significantly increased for patients with a large (> 800 cc) breast volume (odds ratio = 4, <i>p</i> = 0.017).</p> <p>Conclusions: This is the first randomized controlled trial demonstrating by objective measurements that PBMT is effective in reducing the incidence of moist desquamation in breast cancer patients undergoing RT. Additionally, a large breast volume is an important risk factor for the development of moist desquamation.</p>
155	Keywords separated by ' - '	Breast cancer - Photobiomodulation therapy - Radiotherapy - Skin toxicity - Radiation dermatitis - Objective skin evaluation
156	Foot note information	<p>This research was orally presented at the International Symposium of the Multinational Association of Supportive Care in Cancer and the International Society for Oral Oncology (MASCC/ISOO; June 28–30, 2018, Vienna, Austria).</p> <p>The online version of this article (https://doi.org/10.1007/s00520-018-4487-4) contains supplementary material, which is available to authorized users.</p>

Electronic supplementary material

ESM 1
(DOCX 130 kb)

Biophysical skin measurements to evaluate the effectiveness of photobiomodulation therapy in the prevention of acute radiation dermatitis in breast cancer patients

Jolien Robijns¹ · Sandrine Censabella² · Stefan Claes³ · Luc Pannekoeke³ · Lore Bussé¹ · Dora Colson¹ · Iris Kaminski¹ · Joy Lodewijckx¹ · Paul Bulens^{2,3} · Annelies Maes^{2,3} · Leen Noé^{2,3} · Marc Brosens^{2,3} · An Timmermans⁴ · Ivo Lambrechts¹ · Veerle Somers¹ · Jeroen Mebis^{1,2,3}

Received: 30 April 2018 / Accepted: 24 September 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Purpose The purpose of this study was to evaluate objectively the effectiveness of photobiomodulation therapy (PBMT) for the prevention of acute radiation dermatitis (ARD) by using biophysical skin measurements.

Methods A randomized, placebo-controlled trial with 120 breast cancer patients who underwent an identical radiotherapy (RT) regimen post-lumpectomy was performed (TRANSDERMIS trial). Patients were randomized to receive PBM (808 nm CW/905 nm pulsed, 168 mW/cm², spot size 19.6 cm², fluence 4 J/cm²) or placebo treatments from the first day of RT (2×/week). Biophysical skin measurements were collected to assess the skin pigmentation and barrier function. Measurements were collected at the first day of RT, a RT dose of 40 Gray (Gy), and the end of RT (66 Gy).

Results The incidence of moist desquamation was significantly higher in the control than in the PBMT group at the end of RT (30 vs. 7%, respectively, odds ratio = 6, *p* = 0.004). The biophysical skin measures showed that the mean percentage change from the baseline transepidermal water loss (TEWL), erythema, and melanin values was significantly higher in the control than in the PBMT group at the end of RT (*ps* < 0.05). Logistic regression analysis revealed that the risk on moist desquamation was significantly increased for patients with a large (> 800 cc) breast volume (odds ratio = 4, *p* = 0.017).

Conclusions This is the first randomized controlled trial demonstrating by objective measurements that PBMT is effective in reducing the incidence of moist desquamation in breast cancer patients undergoing RT. Additionally, a large breast volume is an important risk factor for the development of moist desquamation.

Keywords Breast cancer · Photobiomodulation therapy · Radiotherapy · Skin toxicity · Radiation dermatitis · Objective skin evaluation

This research was orally presented at the International Symposium of the Multinational Association of Supportive Care in Cancer and the International Society for Oral Oncology (MASCC/ISOO; June 28–30, 2018, Vienna, Austria).

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00520-018-4487-4>) contains supplementary material, which is available to authorized users.

✉ Jolien Robijns
jolien.robijns@uhasselt.be

¹ Faculty of Medicine & Life Sciences, Hasselt University, Martelarenlaan 42, 3500 Hasselt, Belgium

² Department of Medical Oncology, Jessa Hospital, Stadsomvaart 11, Hasselt, Belgium

³ Limburg Oncology Centre, Jessa Hospital, Stadsomvaart 11, Hasselt, Belgium

⁴ Department of Dermatology, Jessa Hospital, Stadsomvaart 11, Hasselt, Belgium

Q3 33

Introduction

Acute radiation dermatitis (ARD) is a severe side effect occurring in about 90–95% of the cancer patients undergoing radiotherapy (RT) [1]. This is a cutaneous reaction that is caused by direct damage of ionizing radiation, which manifests 2–4 weeks after the first RT session [2].

In normal healthy skin, the superficial cells of the epidermis (i.e., upper skin layer) are shed through normal desquamation and replaced by stem cells from the underlying basal layer. From the first RT dose, stem cells within the basal layer of the epidermis are destroyed, leading to a disruption in the self-renewing property of the skin. During RT, this process continues which will negatively affect the skin barrier function and the wound healing process. This ultimately results in changes of the skin structure and vasculature, clinically characterized by erythema, dryness, flaking skin, pruritus, folliculitis (i.e., skin rash), and hyperpigmentation. Due to the compromised skin barrier function and cutaneous immune system, the skin will become more susceptible to water loss, chemical substances, allergens, ultraviolet radiation (UV), and infections [3, 4].

Clinically, ARD is evaluated by the criteria of the Radiation Therapy Oncology Group (RTOG) into three grades: mild erythema and dry desquamation (grade 1), bright erythema and moist desquamation in skin folds (grade 2), and confluent moist desquamation (grade 3). However, this grading system lacks objectivity [5].

A variety of biophysical skin techniques are available to measure the skin pigmentation, hydration, pH, blood flow, and sebum level in order to investigate the underlying physiological mechanism of ARD [6].

Up to now, the evidence for a general consensus on the prevention and management of ARD is limited. Nevertheless, the Multinational Association of Supportive Care in Cancer (MASCC) developed skin care guidelines concerning the prevention and treatment of RD in 2013. Still, many RT centers develop their own skin care protocol [7].

Photobiomodulation therapy (PBMT) is the application of visible and/or (near-) infrared light at a low power on tissue to stimulate the wound healing process and reduce inflammation and pain [8]. There is evidence that PBMT could be used as a new preventive and therapeutic tool in the management of ARD [9–12]. Recently, our research group performed two clinical trials in which we demonstrated that PBMT is able to prevent the development of ARD grade 2 or higher in breast cancer patients by clinically evaluating the skin reactions by the RTOG grading [13, 14].

In this project, we evaluated the effectiveness of PBMT in the prevention of ARD in breast patients by objectively

assessing the skin hydration, transepidermal water loss (TEWL), and pigmentation.

Material and methods

Study design and setting

This was a secondary analysis of the TRANSDERMIS trial, a monocentric, prospective, placebo-controlled, randomized controlled trial (RCT) [14], to evaluate objectively the effectiveness of PBMT in breast cancer patients undergoing RT. Female patients with unilateral breast cancer who were treated at the RT Department of the Limburg Oncology Centre (Jessa Hospital, Hasselt, Belgium) were screened on eligibility between April 2015 and June 2017. The study was approved by the ethics committees of the Jessa Hospital and the University of Hasselt (B243201524443) and was conducted according to the Declaration of Helsinki. The study was registered at [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02443493) (NCT02443493).

Study population

To be eligible for the study, patients needed to fulfill the following criteria: female, diagnosed with primary unilateral breast cancer, underwent lumpectomy, scheduled to undergo a RT regimen consisting of 25 fractions of 2 Gray (Gy) to the whole breast and 8 fractions of 2 Gy to the tumor region (total RT dose 66 Gy). Patients were excluded when they met the following criteria: irradiation to the same breast in the past, hypofractionated RT, mastectomy, metastatic disease, concomitant chemotherapy, and infection of the to-be-irradiated zone. Eligible patients were recruited during the CT simulation session, approximately 2 weeks before the start of the RT. Written informed consent of all patients was collected before study participation.

Randomization and blinding

The planning target volume (PTV) of the eligible patients was used to stratify them into three groups: small (< 450 cc), medium (450–800 cc), and large breasts (> 800 cc) [15]. Patients were randomly assigned to the control or PBMT group in a 1:1 ratio based on a computer-generated random number list, which was held by a researcher who was not involved in the clinical part of the study. Allocation was concealed to the PBM operator until the first treatment session. Both the participating patient and the outcome assessor were blinded until the last treatment session.

128	Interventions	
129	Radiotherapy	
130	The Eclipse™ treatment planning system was used to plan	synchronously with coincident propagation axes (average
131	the RT sessions (version 11.0, Varian Medical System,	radiant power 3.3 W). The energy density (fluence) was
132	Palo Alto, CA). The standard RT regimen consisted of	set at 4 J/cm ² based on earlier recommendations and on
133	25 daily fractions (2 Gy/fraction, 5 fractions/week) to	our clinical experience [13, 16]. During the PBMT ses-
134	the whole breast followed by boost of 8 fractions (2 Gy/	sions, the whole irradiated area was treated (whole breast,
135	fraction, 5 fractions/week) to the tumor bed during a pe-	inframammary fold, and axilla). The complete list of
136	riod of 6 to 7 weeks (total RT dose of 66 Gy). The whole	PBMT parameters can be found in Table 1. The PBMT
137	breast was irradiated with two tangential photon (half)	parameters were selected based on the successful results
138	beams set up isocentrically using a 6-MV or a 6 + 15-	of our previous trial (DERMIS trial) [14] and on the
139	MV linear accelerator (Clinac® DHX, Varian Medical	guidelines of Zecha et al. [17].
140	Systems, Palo Alto, CA) and the tumor region with a	During the sham treatments of the control group, the PBM
141	two-field conformal photon (4–15 MV) or a one-field	device did not emit light but made the same sound as an active
142	vertical electron (6–15 MeV) beam. A selected ¹ group	device. All patients, independently of their treatment group,
143	of patients were irradiated using the deep inspiration	wore safety glasses and eye shields to avoid any perceived risk
144	breath-hold (DIBH) in order to reduce the mean heart	of eye damage and to blind them during the PBM or sham
145	dose (MHD).	sessions.
		Outcome measures
146	Topical skin care treatment	
147	Each patient was individually advised to follow the gen-	Patient data
148	eral skin care guidelines (e.g., wear loose fit clothing,	
149	gentle washing with or without mild soap, patting dry	Clinical information regarding the patient's personal and
150	with a soft towel instead of rubbing). Further, the patients	disease- and treatment-related characteristics was collected
151	were instructed to apply a topical, hydroactive colloid gel	via patient questionnaires and the patient's medical charts.
152	(Flamigel®, Flen Pharma, Kontich, Belgium) on the irra-	
153	diated zone (3×/day), starting at the first day of RT. Foam,	RTOG grading
154	absorbent, self-adhesive silicone dressings (Mepilex®,	
155	Mölnlycke Health Care, Gothenburg, Sweden) were used	Clinically, the severity of ARD was evaluated by the criteria of
156	in the case of painful skin reactions and/or moist	the Radiation Therapy Oncology Group/European
157	desquamation.	Organization for Research and Treatment of Cancer
		(RTOG/EORTC [5]). Two experienced RT nurses performed
158	PBMT	this in a blinded manner.
159	PBMT was applied from the first until the last day of RT	Objective skin measurements
160	(2×/week, 14 sessions) by a trained operator using the	
161	class IV MLS® M6 laser (ASA Srl, Vicenza, Italy), as	In order to assess the impact of RT on the skin barrier
162	described previously [14]. This device is commercially	function, the transepidermal water loss (TEWL) and the
163	available, built in compliance with EC/EU rules, received	skin hydration level were determined. TEWL was mea-
164	FDA approval, and is CE certified. It consists of two laser	sured by the Tewameter® TM300 (Courage-Khazaka,
165	diodes with different wavelengths (808–905 nm), peak	Cologne, Germany), according to the guidelines published
166	powers (1.1–25 W), and emission modes (continuous	both by the standardization group of the European
167	and pulsed). Both diodes work simultaneously and	Contact Dermatitis Society [18] and by the European
		group on Efficacy Measurements of Cosmetics and
		Other topical products [19]. The skin hydration was mea-
		sured with the Corneometer® (Courage-Khazaka,
		Cologne, Germany) according to Heinrich et al. [20]. A
		reflectance spectrophotometer, Mexameter® MX18
		(Courage-Khazaka, Cologne, Germany), was used to mea-
		sure the pigmentation of the skin (e.g., melanin and ery-
		thema) as previously described by Clarys et al. [21].
		All four measurements (e.g., TEWL, hydration, erythe-
		ma, and melanin) were taken at the four quadrants of each
		breast (irradiated and non-irradiated), with three

¹ DIBH was used when the patients matched the following criteria: bilateral breast cancer; left-sided breast cancer and lymph node metastases under the age of 70 years; left-sided breast cancer and lymph node metastases above the age of 70 years and undergoing chemotherapy; left-sided breast cancer without lymph node metastasis but with a MHD ≥ 35 Gy. DIBH was applied using the Varian Real-Time Position Management (RPM) gating system (Varian Medical System, Palo Alto, CA).

214 measurements per quadrant (see Online Resource 1). The
 215 average values of these measurements were taken as a
 216 value for the whole breast. The measurements were car-
 217 ried out after a 30-min acclimatization period at room
 218 temperature (20–22 °C) and 40–60% humidity. The final

219 objective measurements were described as percentages in
 220 order to calculate deviations from pre-treatment baseline
 221 values, also termed as indexes. Therefore, the following
 222 formula was used:

$$\left[\left(\frac{\text{Obj.measure irradiated breast at indicated time/Obj.measure control breast at indicated time}}{\text{Obj.measure irradiated breast at baseline/Obj.measure control breast at baseline}} \right) - 1 \right] \times 100\%$$

226
 228
 229

Q4 t1.1 **Table 1** Photobiomodulation parameters

t1.2	PBMT parameters			
t1.3	Device information	Manufacturer	ASA Srl	
t1.4		Model identifier	MLS® laser M6	
t1.5		Year produced	2012	
t1.6		Number of emitters	3	
t1.7		Emitter type	IR laser diodes	
t1.8		Spatial distribution of emitters	Three emitters spaced 2 cm apart in a triangle pattern	
t1.9		Beam delivery system	Scanning head (five pre-settled directions)	
t1.10	Irradiation parameters		Laser diode 1	Laser diode 2
t1.11		Center wavelength	808 nm	905 nm
t1.12		Number of emitters	1	2
t1.13		Spectral bandwidth	± 5 nm	± 5 nm
t1.14		Operating mode	Continuous pulsed wave mode	
t1.15		Peak radiant power	1.1 W	25 W
t1.16		Average radiant power	3.3 W	
t1.17		Maximum frequency (frequency range)		90 kHz (1–2000 Hz)
t1.18		- Pulse on duration		- 100-ns single pulse width
t1.18		- Duty cycle		- 50%
t1.19		Aperture diameter	5 cm	
t1.20		Irradiance at aperture	0.168 W/cm ²	
t1.21		Beam divergence at 60%	42.8 mrad	59.2 mrad
t1.22		Beam profile	Two laser beams work simultaneously and synchronously with coincident propagation axes	
t1.23	Treatment parameters	Beam spot size at target area	19.625 cm ²	
t1.24		Irradiance at target	0.168 W/cm ²	
t1.25		Radiant exposure (fluence)	4 J/cm ²	
t1.26		Number of points irradiated	-Breast: Whole breast, inframammary fold and/or axilla, depending on the location of radiodermatitis	
t1.27		Exposure duration	- Whole breast: ± 420–720 s - Inframammary fold: ± 103 s - Axilla: ± 68 s	
t1.28		Application technique	5 cm above the skin	
t1.29		Timing	After the RT session	
t1.30		Number and frequency of treatment sessions	14 sessions in total, delivered biweekly from the first until the last day of RT over a period of 7 weeks	

IR, infrared; MLS, Multiwave Locked System; PBMT, photobiomodulation therapy; RD, radiodermatitis; RT, radiotherapy

231 Measurement collection schedule

232 All the previously described measurements were collected on
233 three time points: at the first day of RT, at a RT dose of 40 Gy,
234 and at the last day of RT (66 Gy).

235 Statistical analysis

236 Differences in patient- and therapy-related characteristics be-
237 tween both groups were analyzed by means of chi-square tests
238 (χ^2), Fisher's exact tests, Student's *t* tests, or Mann-Whitney
239 *U* tests, as appropriate. RTOG scores were analyzed by means
240 of χ^2 or Fisher's exact tests, as appropriate. The objective skin
241 measurements at each time point were analyzed by Mann-
242 Whitney *U* tests. Longitudinal analysis of the biophysical skin
243 measurements was performed by mixed analyses of variance
244 (ANOVAs) with time (between the RT dose of 40 Gy and
245 66 Gy) as within-subject factor and group (control vs.
246 PBMT group) as between-subject factor. To determine the risk

on moist desquamation, univariate logistic regressions with,
as predictor variables, treatment group and breast size (based
on the PTV) were performed. The level of statistical signifi-
cance for all analyses was set assuming a significance level of
5% ($p < 0.05$, two-tailed). SPSS 24.0 (IBM, Chicago, IL) was
used for all analyses.

Results

Patient characteristics

A total of 139 patients were randomized into the placebo or
PBMT group between April 2015 and June 2017. During the
course of RT, 2 patients of the control group withdrew their
consent. Further 17 patients were excluded due to a RT regi-
men change or a RT interruption (5 and 8 in the control and
PBMT, resp.). For the final analysis data of 120 patients, 60
patients in each group were used (Fig. 1). Both groups were

Fig. 1 CONSORT flow chart [14]

matched for all the patient- and treatment-related characteristics (Table 2).

Clinical evaluation of ARD

Patients' RT-induced skin reactions were evaluated by the criteria of the RTOG, as shown in Table 3. Our results demonstrated that the incidence of moist desquamation (ARD grade 2 or higher) was significantly lower in the PBMT group in comparison with the control group at the end of RT ($p = 0.004$). This was confirmed by the univariate logistic regression analysis demonstrating that patients only receiving the standard skin care were six times more likely to develop moist desquamation in comparison with patients that also were treated with PBMT ($p = 0.003$, 95% CI [OR] 1.881–19.82). Further, the risk on moist desquamation rose with an increasing breast volume. As such, patients with large breasts (> 800 cc) had a four times higher risk to develop moist desquamation than patients with small breast volumes ($p = 0.017$, 95% CI [OR] 1.290–12.936).

Objective evaluation of ARD

Erythema

The mixed 2×2 ANOVAs revealed a significant main time effect and group by time interaction ($ps < 0.05$) for the erythema index. However, the main group effect was

Table 3 RTOG grading at a RT dose of 40 and 66 Gy (end RT)

RTOG grading	Control group ($n = 60$) N (%)	PBMT group ($n = 60$) N (%)	p^a
40 Gray			0.562
Grade 1	1 (1.7)	3 (5)	
Grade 2	55 (91.7)	54 (90)	
Grade 3	4 (6.7)	3 (5)	
66 Gray (end RT)			0.004
Grade 1	42 (70)	56 (93.3)	
Grade 2	16 (26.7)	4 (6.7)	
Grade 3	2 (3.3)	0 (0)	

PBMT, photobiomodulation therapy; RTOG, Radiation Therapy Oncology Group (grade 0: no change; grade 1: follicular, dull, or faint erythema, dry desquamation; grade 2: tender or bright erythema, patchy moist desquamation; grade 3: confluent moist desquamation other than skin folds)

^a Chi-square tests (two-tailed)

not significant. As depicted in Fig. 2a, the degree of erythema in both groups increased during the course of RT. At the RT dose of 40 Gy, the percentage change in erythema from baseline did not significantly differ between the control group and the PBMT group. However, at the end of RT, the percentage change from baseline in erythema was significantly higher in the control group in comparison with the PBMT group ($p = 0.016$).

Table 2 Patient and treatment characteristics

	Control group ($n = 60$)	PBMT group ($n = 60$)	p^a
Mean age (SD), years	56.92 (10.34)	56.52 (10.54)	0.88
Mean body mass index (SD)	25.03 (4.47)	25.27 (3.87)	0.63
Mean breast size (SD) ^b , cc	796.27 (439.67)	742.55 (353.92)	0.67
Breast size ^b , n (%)			0.97
Small (< 450 cc)	11 (18.3)	12 (20)	
Medium (450–800 cc)	26 (43.3)	26 (43.3)	
Large (> 800 cc)	23 (38.3)	22 (36.7)	
Prior chemo, n (%)	46 (76.6)	44 (73.3)	0.83
RT energy level, n (%)			0.19
6 MV	43 (71.7)	50 (83.3)	
6 MV + 15 MV	17 (28.3)	10 (16.7)	
Boost type, n (%)			0.86
Photons	31 (51.7)	29 (48.3)	
Electrons	29 (48.3)	31 (51.7)	
DIBH	17 (28.3)	11 (18.3)	0.28

DIBH, deep inspiration breath-hold; PBMT, photobiomodulation therapy; RT, radiotherapy; SD, standard deviation

^a Student's t test, Wilcoxon-Mann-Whitney U test, chi-square tests, or Fisher's exact tests, as appropriate (two-tailed)

^b Radiotherapy target volume that consists of the macroscopic primary tumor, the surrounding microscopic tumor spread and a margin to account for patient and/or organ movement, shape changes of the tumor, and daily setup variations. PTV was measured via treatment planning system by contouring manually each slice of breast tissue on planning CT

Fig. 2 Evaluation of the skin pigmentation (erythema (a) and melanin (b)) and barrier function (hydration (c) and TEWL (d)) by biophysical measurements. Data are shown as mean percentage change from baseline ($\pm$ SEM). *Significant difference between the two groups at the indicated

time point ($p < 0.05$; Mann-Whitney U test, two-tailed). TEWL, transepidermal water loss; PBMT, photobiomodulation therapy; RT, radiotherapy

Pigmentation

Concerning the melanin index, there were both a significant main time effect and group by time interaction ($p < 0.05$), but no significant main group effect. Figure 2b demonstrates that the degree of pigmentation increased during the progression of RT in both groups. The increase in pigmentation started off slowly, with no significant difference in percentage change over baseline in melanin between the two groups at the RT dose of 40 Gy. Towards the end of RT, the melanin index was significantly higher in the control than in the PBMT group ($p = 0.019$).

Hydration

The mixed 2×2 ANOVAS revealed a significant main time and group effect ($p < 0.05$), but no significant group by time interaction for the skin moisture level. As shown in Fig. 2c, during the course of RT, the skin hydration level decreased in both groups in comparison with the baseline values. The skin hydration level was significantly lower at the RT dose of 40 Gy in the PBMT group in comparison with the control group ($p = 0.036$). However, at the end of RT, both groups showed a comparable skin moisture index.

Transepidermal water loss

Regarding the TEWL, there was a significant main time and group effect ($p < 0.05$), but no significant group by time interaction. The TEWL decreased in comparison with the baseline value in both the control and PBMT groups at the RT dose of 40 Gy, to a comparable level (Fig. 2d). Towards the end of RT, the TEWL level increased in both groups, although the final TEWL index was significantly lower in the PBMT group in comparison with the control group ($p = 0.05$).

Discussion

Results of this trial show that PBMT is an effective tool to prevent the development of moist desquamation. This was confirmed by objectively evaluating the skin's biophysical condition. Our results demonstrated that PBMT was able to reduce the increase in the skin's pigmentation level and improve the skin barrier function. Additionally, the main risk factor for the development of severe ARD is the breast volume, which implies that patients with large breasts (> 800 cc) have an increased risk on moist desquamation.

The erythema index progressively increased during RT in both treatment arms. These findings are in line with previous studies [22–25]. This increase in erythema is caused by the

RT-induced inflammatory reaction leading to vasodilation and leaking of the blood vessels [6, 26, 27]. However, the increase was significantly lower in the PBMT than in the control group at the end of RT. This proves that PBMT is able to reduce the degree of erythema. These results are consistent with earlier in vivo studies and clinical trials on various erythematous skin disorders (e.g., acne vulgaris, UV damage, laser resurfacing wounds, burn wounds) [28–31]. The anti-inflammatory effect of PBMT, correlated with a decrease in inflammatory cytokine production, might explain this observation [28, 32].

Further, our results also showed a significant increase in skin pigmentation in both groups during the course of RT. This is explained by post-inflammatory hyperpigmentation (PIH) after the RT-induced skin reaction [6, 27]. PIH is caused by the stimulation of melanocytes due to an inflammatory skin reaction leading to an increased melanin production and transport to the surrounding keratinocytes. Remarkably, our results demonstrated that at the end of RT, the increase in melanin content of the skin was significantly lower in the PBMT than in the control group. As such, PBMT was able to stabilize the hyperpigmentation reaction of the patients' skin during RT. Several in vitro studies showed that PBMT can inhibit the melanin synthesis in human melanocyte cultures [33]. Also, clinical trials demonstrated that PBMT is able to reduce hyperpigmentation in numerous skin conditions (e.g., acne vulgaris, photoaging, melasma) [34, 35].

In healthy skin, a low TEWL and a high hydration value correlate with a good barrier function [36]. Ionizing radiation deregulates the cellular function and causes apoptosis of the epidermal cells, resulting into an affected skin barrier function, correlated with a high TEWL and a low skin moisture level [6, 27, 37, 38]. The findings in our control group are in line with these studies. However, in the PBMT group, both the TEWL and hydration index were significantly decreased at the end of RT. The epidermal thickening effect might explain these conflicting results. This effect is characterized by epidermal hyperproliferation leading to a thickened stratum corneum (outermost layer of the epidermis) caused by repetitive exposure to external stimuli. The thickening of the stratum corneum improves the skin barrier function and thereby it is correlated with a decrease in TEWL [38, 39]. Several studies, both in vitro and in vivo, have demonstrated that PBMT can stimulate the proliferation of several types of cells, including keratinocytes. PBMT seems to be able to stimulate the epidermal thickening effect in the skin caused by RT and thereby it can improve the skin barrier function [40–43].

The results of the logistic regression analysis demonstrated that patients who were treated with standard skin care had a six time higher risk to develop moist desquamation in comparison with the patients treated with PBMT. This implies that the preventive application of

PBMT can seriously lower the severity of the RT-induced skin reactions, as previously published by our study group [14]. Further, our results showed that patients with large breasts developed more severe skin reactions. These findings are consistent with those of earlier published studies [44, 45].

The main limitation of the study was the enrolled patient population, which was confined to breast cancer patients post-lumpectomy, who underwent a standard fractionated RT regimen. In the future, more clinical trials in a broader patient population with different cancer types and RT regimens need to be conducted, which will increase the generalizability of the study results.

Conclusion

This is the first RCT demonstrating by an objective approach that the preventive application of PBMT is effective in reducing the incidence of moist desquamation in breast cancer patients. The biophysical skin measurements showed that PBMT is able to stabilize the degree of pigmentation (both erythema and melanin) and improve the skin barrier function during the course of RT. Interestingly; patients with a large breast volume have an increased risk on moist desquamation. In conclusion, we can state that PBMT is an effective tool to prevent the development of severe ARD in breast cancer patients. Further, screening patients on breast volume before the start of RT can allow the radiotherapist to optimize the skin management during the course of RT.

Acknowledgments The corresponding author, Mrs. Jolien Robijns, received the Young Investigator Award of the MASCC for her scientific abstract.

Funding information This research is part of the Limburg Clinical Research Program (LCRP) UHasselt-ZOL-Jessa, financially supported by the foundation Limburg Sterk Merk, province of Limburg, Flemish government, Hasselt University, Ziekenhuis Oost-Limburg, and Jessa Hospital. Additionally, this research is supported by Kom op Tegen Kanker, Limburgs Kankerfonds, and ASA Srl.

Compliance with ethical standards

Conflict of interest The authors declare that they have no competing interests.

Ethical approval The ethics committees of the Jessa Hospital and the University of Hasselt approved the study (B243201524443). All procedures performed in the study were in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

1. Wells M, MacBride S (2003) Radiation skin reactions. Elsevier, New York
2. Seite S, Bensadoun RJ, Mazer JM (2017) Prevention and treatment of acute and chronic radiodermatitis. *Breast Cancer* (Dove Med Press) 9:551–557. <https://doi.org/10.2147/BCTT.S149752>
3. Robijns J, Laubach H-J (2018) Acute and chronic radiodermatitis: clinical signs, pathophysiology, risk factors and management options. *Journal of the Egyptian Women's Dermatologic Society* 15(1):2–9. <https://doi.org/10.1097/01.EWX.0000529960.52517.4c>
4. Proksch E, Brandner JM, Jensen JM (2008) The skin: an indispensable barrier. *Exp Dermatol* 17(12):1063–1072
5. Cox JD, Stetz J, Pajak TF (1995) Toxicity criteria of the Radiation Therapy Oncology Group (RTOG) and the European Organization for Research and Treatment of Cancer (EORTC). *Int J Radiat Oncol Biol Phys* 31(5):1341–1346. [https://doi.org/10.1016/0360-3016\(95\)00060-C](https://doi.org/10.1016/0360-3016(95)00060-C)
6. Hu SC, Hou MF, Luo KH, Chuang HY, Wei SY, Chen GS, Chiang W, Huang CJ (2014) Changes in biophysical properties of the skin following radiotherapy for breast cancer. *J Dermatol* 41(12):1087–1094. <https://doi.org/10.1111/1346-8138.12669>
7. Wong RK, Bensadoun RJ, Boers-Doets CB, Bryce J, Chan A, Epstein JB, Eaby-Sandy B, Lacouture ME (2013) Clinical practice guidelines for the prevention and treatment of acute and late radiation reactions from the MASCC Skin Toxicity Study Group. *Support Care Cancer* 21(10):2933–2948. <https://doi.org/10.1007/s00520-013-1896-2>
8. Hamblin M, Ferraresi C, Huang YY, de Freitas LF, Carroll J (2018) Low-Level Light Therapy: Photobiomodulation. *Tutorial Texts in Optical Engineering*, vol TT115. SPIE Press, Bellingham 390
9. Schindl M, Kerschman K, Schindl A, Schon H, Heinzl H, Schindl L (1999) Induction of complete wound healing in recalcitrant ulcers by low-intensity laser irradiation depends on ulcer cause and size. *Photodermatol Photoimmunol Photomed* 15(1):18–21
10. DeLand MM, Weiss RA, McDaniel DH, Geronemus RG (2007) Treatment of radiation-induced dermatitis with light-emitting diode (LED) photomodulation. *Lasers Surg Med* 39(2):164–168. <https://doi.org/10.1002/lsm.20455>
11. Fife D, Rayhan DJ, Behnam S, Ortiz A, Elkeeb L, Aquino L, Eduardo Roa D, Ramsinghani N, Kuo J, Newcomb R, Zachary CB, Kelly KM (2010) A randomized, controlled, double-blind study of light emitting diode photomodulation for the prevention of radiation dermatitis in patients with breast cancer. *Dermatol Surg* 36(12):1921–1927. <https://doi.org/10.1111/j.1524-4725.2010.01801.x>
12. Strouthos I, Chatzikonstantinou G, Tselis N, Bon D, Karagiannis E, Zoga E, Ferentinos K, Maximenko J, Nikolettou-Fischer V, Zamboglou N (2017) Photobiomodulation therapy for the management of radiation-induced dermatitis : a single-institution experience of adjuvant radiotherapy in breast cancer patients after breast conserving surgery. *Strahlenther Onkol* 193(6):491–498. <https://doi.org/10.1007/s00066-017-1117-x>
13. Censabella S, Claes S, Robijns J, Bulens P, Mebis J (2016) Photobiomodulation for the management of radiation dermatitis: the DERMIS trial, a pilot study of MLS® laser therapy in breast cancer patients. *Support Care Cancer* 24:1–9. <https://doi.org/10.1007/s00520-016-3232-0>
14. Robijns J, Censabella S, Claes S, Pannekoek L, Bussé L, Colson D, Kaminski I, Bulens P, Maes A, Noé L, Brosens M, Timmermans A, Lambrechts I, Somers V, Mebis J (2018) Prevention of acute radiodermatitis by photobiomodulation: a randomized, placebo-controlled trial in breast cancer patients (TRANSDERMIS trial). *Lasers Surg Med* 50(7):763–771. <https://doi.org/10.1002/lsm.22804>
15. Burnet NG, Thomas SJ, Burton KE, Jefferies SJ (2004) Defining the tumour and target volumes for radiotherapy. *Cancer Imaging* 4(2):153–161. <https://doi.org/10.1102/1470-7330.2004.0054>
16. Nair RG, Bensadoun RJ (2016) Mitigation of cancer therapy side-effects with light. Morgan & Claypool Publishers, San Rafael
17. Zecha JA, Raber-Durlacher JE, Nair RG, Epstein JB, Elad S, Hamblin MR, Barasch A, Migliorati CA, Milstein DM, Genot MT, Lansaet L, van der Brink R, Arnabat-Dominguez J, van der Molen L, Jacobi I, van Diessen J, de Lange J, Smeele LE, Schubert MM, Bensadoun RJ (2016) Low-level laser therapy/ photobiomodulation in the management of side effects of chemoradiation therapy in head and neck cancer: part 2: proposed applications and treatment protocols. *Support Care Cancer* 24(6):2793–2805. <https://doi.org/10.1007/s00520-016-3153-y>
18. Pinnagoda J, Tupker RA, Agner T, Serup J (1990) Guidelines for transepidermal water loss (TEWL) measurement. A report from the Standardization Group of the European Society of Contact Dermatitis. *Contact Dermatitis* 22(3):164–178
19. Rogiers V, E. Group (2001) EEMCO guidance for the assessment of transepidermal water loss in cosmetic sciences. *Skin Pharmacol Appl Ski Physiol* 14(2):117–128 56341
20. Heinrich U, Koop U, Leneveu-Duchemin MC, Osterrieder K, Bielfeldt S, Chkarnat C, Degwert J, Hantschel D, Jaspers S, Nissen HP, Rohr M, Schneider G, Tronnier H (2003) Multicentre comparison of skin hydration in terms of physical-, physiological- and product-dependent parameters by the capacitive method (Comeometer CM 825). *Int J Cosmet Sci* 25(1–2):45–53. <https://doi.org/10.1046/j.1467-2494.2003.00172.x>
21. Clarys P, Alewaeters K, Lambrecht R, Barel AO (2000) Skin color measurements: comparison between three instruments: the Chromameter(R), the DermaSpectrometer(R) and the Mexameter(R). *Skin Res Technol* 6(4):230–238
22. Yamazaki H, Yoshida K, Kobayashi K, Tsubokura T, Kodani N, Aibe N, Ikeno H, Nishimura T (2012) Assessment of radiation dermatitis using objective analysis for patients with breast cancer treated with breast-conserving therapy: influence of body weight. *Jpn J Radiol* 30(6):486–491. <https://doi.org/10.1007/s11604-012-0073-8>
23. Primavera G, Carrera M, Berardesca E, Pinnaro P, Messina M, Arcangeli G (2006) A double-blind, vehicle-controlled clinical study to evaluate the efficacy of MAS065D (XClair), a hyaluronic acid-based formulation, in the management of radiation-induced dermatitis. *Cutan Ocul Toxicol* 25(3):165–171. <https://doi.org/10.1080/15569520600860009>
24. Yoshida EJ, Chen H, Torres MA, Curran WJ, Liu T (2011) Spectrophotometer and ultrasound evaluation of late toxicity following breast-cancer radiotherapy. *Med Phys* 38(10):5747–5755. <https://doi.org/10.1118/1.3633942>
25. Iacovelli NA, Naimo S, Bonfantini F, Cavallo A, Bossi P, Fallai C, Pignoli E, Alfieri S, Bergamini C, Favale F, Orlandi E (2017) Preemptive treatment with Xonrid(R), a medical device to reduce radiation induced dermatitis in head and neck cancer patients receiving curative treatment: a pilot study. *Support Care Cancer* 25(6):1787–1795. <https://doi.org/10.1007/s00520-017-3569-z>
26. Nystrom J, Geladi P, Lindholm-Sethson B, Rattfelt J, Svensk AC, Franzen L (2004) Objective measurements of radiotherapy-induced erythema. *Skin Res Technol* 10(4):242–250
27. Yamazaki H, Yoshida K, Kotsuma T, Kuriyama K, Masuda N, Nishimura T, Kobayashi K, Tsubokura T, Nishimura T (2009) Longitudinal practical measurement of skin color and moisture during and after breast-conserving therapy: influence of neoadjuvant systemic therapy. *Jpn J Radiol* 27(8):309–315. <https://doi.org/10.1007/s11604-009-0345-0>
28. Hamblin MR (2017) Mechanisms and applications of the anti-inflammatory effects of photobiomodulation. *AIMS Biophys* 4(3):337–361. <https://doi.org/10.3934/biophys.2017.3.337>

- 569 29. Gupta A, Keshri GK, Yadav A, Gola S, Chauhan S, Salhan AK, 607
570 Bala Singh S (2015) Superpulsed (Ga-As, 904 nm) low-level laser 608
571 therapy (LLLT) attenuates inflammatory response and enhances 609
572 healing of burn wounds. *J Biophotonics* 8(6):489–501. <https://doi.org/10.1002/jbio.201400058> 610
- 573 30. Khoury JG, Goldman MP (2008) Use of light-emitting diode 611
574 photomodulation to reduce erythema and discomfort after intense 612
575 pulsed light treatment of photodamage. *J Cosmet Dermatol* 7(1): 613
576 30–34. <https://doi.org/10.1111/j.1473-2165.2008.00358.x> 614
- 577 31. Barolet D, Boucher A (2008) LED photoprevention: reduced MED 615
578 response following multiple LED exposures. *Lasers Surg Med* 616
579 40(2):106–112. <https://doi.org/10.1002/lsm.20615> 617
- 580 32. Chen CH, Wang CZ, Wang YH, Liao WT, Chen YJ, Kuo CH, Kuo 618
581 HF, Hung CH (2014) Effects of low-level laser therapy on M1- 619
582 related cytokine expression in monocytes via histone modification. 620
583 *Mediat Inflamm* 2014:625048. [https://doi.org/10.1155/2014/](https://doi.org/10.1155/2014/625048) 621
584 [625048](https://doi.org/10.1155/2014/625048) 622
- 585 33. Kim HK, Min KO, Choi JH, Kim SH (2016) Effects of low-level 623
586 laser therapy, electroacupuncture, and radiofrequency on the pig- 624
587 mentation and skin tone of adult women. *J Phys Ther Sci* 28(5): 625
588 1407–1411. <https://doi.org/10.1589/jpts.28.1407> 626
- 589 34. Lee SY, You CE, Park MY (2007) Blue and red light combination 627
590 LED phototherapy for acne vulgaris in patients with skin phototype 628
591 IV. *Lasers Surg Med* 39(2):180–188. [https://doi.org/10.1002/lsm.](https://doi.org/10.1002/lsm.20412) 629
592 [20412](https://doi.org/10.1002/lsm.20412) 630
- 593 35. Barolet D (2018) Photobiomodulation in dermatology: harnessing 631
594 light from visible to near infrared. *Medical Research Archives* 6(1). 632
595 <https://doi.org/10.18103/mra.v6i1.1610> 633
- 596 36. Baroni A, Buommino E, De Gregorio V, Ruocco E, Ruocco V, Wolf 634
597 R (2012) Structure and function of the epidermis related to barrier 635
598 properties. *Clin Dermatol* 30(3):257–262. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.clindermatol.2011.08.007) 636
599 [clindermatol.2011.08.007](https://doi.org/10.1016/j.clindermatol.2011.08.007) 637
- 600 37. Schmutz M, Wimmer MA, Hofer S, Sztankay A, Weinlich G, 638
601 Linder DM, Elias PM, Fritsch PO, Fritsch E (2002) Topical corti- 639
602 costeroid therapy for acute radiation dermatitis: a prospective, ran- 640
603 domized, double-blind study. *Br J Dermatol* 146(6):983–991 641
604 38. Jensen JM, Gau T, Schultze J, Lemnitz G, Folster-Holst R, May T, 642
605 Abels C, Proksch E (2011) Treatment of acute radiodermatitis with 643
606 an oil-in-water emulsion following radiation therapy for breast can- 607
608 cer: a controlled, randomized trial. *Strahlenther Onkol* 187(6):378– 609
610 384. <https://doi.org/10.1007/s00066-011-2224-8> 611
612 39. Park SY, Kim JH, Cho SI, Kim KI, Cho HJ, Park CW, Lee CH, Kim 613
614 HO (2014) Induction of a hardening phenomenon and quantitative 615
615 changes of ceramides in stratum corneum. *Ann Dermatol* 26(1):35– 616
616 42. <https://doi.org/10.5021/ad.2014.26.1.35> 617
- 617 40. Hawkins DH, Abrahamse H (2006) The role of laser fluence in cell 618
618 viability, proliferation, and membrane integrity of wounded human 619
619 skin fibroblasts following helium-neon laser irradiation. *Lasers* 620
620 *Surg Med* 38(1):74–83. <https://doi.org/10.1002/lsm.20271> 621
621 41. Lanzafame RJ, Stadler I, Kurtz AF, Connelly R, Peter TA Sr, 622
622 Brondon P, Olson D (2007) Reciprocity of exposure time and irra- 623
623 diance on energy density during photoradiation on wound healing 624
624 in a murine pressure ulcer model. *Lasers Surg Med* 39(6):534–542. 625
625 <https://doi.org/10.1002/lsm.20519> 626
- 626 42. Rizzi M, Migliario M, Tonello S, Rocchetti V, Reno F (2018) 627
627 Photobiomodulation induces in vitro re-epithelialization via nitric 628
628 oxide production. *Lasers Med Sci* 33:1003–1008. [https://doi.org/](https://doi.org/10.1007/s10103-018-2443-7) 629
629 [10.1007/s10103-018-2443-7](https://doi.org/10.1007/s10103-018-2443-7) 630
- 630 43. Sperandio FF, Simoes A, Correa L, Aranha AC, Giudice FS, 631
631 Hamblin MR, Sousa SC (2015) Low-level laser irradiation pro- 632
632 motes the proliferation and maturation of keratinocytes during epi- 633
633 thelial wound repair. *J Biophotonics* 8(10):795–803. [https://doi.](https://doi.org/10.1002/jbio.201400064) 634
634 [org/10.1002/jbio.201400064](https://doi.org/10.1002/jbio.201400064) 635
- 635 44. Censabella S, Claes S, Orlandini M, Braekers R, Bulens P (2017) 636
636 Efficacy of a hydroactive colloid gel versus historical controls for 637
637 the prevention of radiotherapy-induced moist desquamation in 638
638 breast cancer patients. *Eur J Oncol Nurs* 29:1–7. [https://doi.org/](https://doi.org/10.1016/j.ejon.2017.04.003) 639
639 [10.1016/j.ejon.2017.04.003](https://doi.org/10.1016/j.ejon.2017.04.003) 640
- 640 45. De Langhe S, Mulliez T, Veldeman L, Remouchamps V, van 641
641 Greveling A, Gilsoul M, De Schepper E, De Ruyck K, De Neve 642
642 W, Thierens H (2014) Factors modifying the risk for developing 643
643 acute skin toxicity after whole-breast intensity modulated radiother- 644
644 apy. *BMC Cancer* 14:711. [https://doi.org/10.1186/1471-2407-14-](https://doi.org/10.1186/1471-2407-14-711) 645
645 [711](https://doi.org/10.1186/1471-2407-14-711) 646

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES.

- Q1. Please check if the affiliations are presented correctly.
- Q2. Please check if the article note is presented correctly.
- Q3. Please check if the section headings are assigned to appropriate levels.
- Q4. Tables 1, 2, 3, and 4 were slightly modified. Please check if data are presented correctly.
- Q5. Please supply/verify the standard abbreviation of the journal name in References Robijns and Laubach (2018) and Barolet (2018).

UNCORRECTED PROOF