

Salmonella in Belgian laying hens: An identification of risk factors

Peer-reviewed author version

NAMATA, Harriet; Meroc, Estelle; AERTS, Marc; FAES, Christel; CORTINAS ABRAHANTES, Jose; Imberechts, H & Mintiens, K. (2008) Salmonella in Belgian laying hens: An identification of risk factors. In: PREVENTIVE VETERINARY MEDICINE, 83(3-4). p. 323-336.

DOI: 10.1016/j.prevetmed.2007.09.002

Handle: <http://hdl.handle.net/1942/8181>

***Salmonella* in Belgian laying hens: an Identification of Risk Factors**

¹Harriet Namata, ²Estelle Méroc ¹Marc Aerts, ¹Christel Faes, ¹José Cortiñas

Abrahantes, ²Hein Imberechts and ²Koen Mintiens.

¹ Hasselt University, Center for Statistics, Campus Diepenbeek, Agoralaan, Gebouw D,

B 3590 Diepenbeek, Belgium Belgium. ²Veterinary and Agrochemical Research Centre, Groeselenberg 99,

B1180 Brussels, Belgium.

Correspondence:

Harriet Namata

Hasselt University, Center for Statistics,

Campus Diepenbeek, Agoralaan, Gebouw D,

B 3590 Diepenbeek, Belgium

Tel: +32 11 26 82 57 Fax: +32 11 26 82 99 Email: harriet.namata@uhasselt.be

Abstract

Since the 1980s, the prevalence of *Salmonella* in Belgian poultry layers and broilers has greatly fluctuated with a rise observed in 2003 and a significant decrease in 2005. In order to alleviate the risk at egg consumer level, it is crucial to understand the factors which influence the contamination and the spread of *Salmonella* in laying hens. To study such determinants we explored the Belgian data from the 2005 baseline study on the prevalence of *Salmonella* in laying flocks of *Gallus gallus* in the European Union. The response variables corresponded to presence or absence of *Salmonella* from dust and faecal samples taken from the environment of a Belgian layer flock. The explanatory variables included: region of Belgium, sampling time (month the flock was sampled), production type (cage or barn/ free range), *Salmonella* vaccination status, flock age and flock size. Analyses of these data were performed using a bivariate logistic regression model assuming independence between the

two responses and bivariate Generalized Estimating Equations model, which incorporates the correlation between the two responses on the same flock. The main risk factor that was identified was rearing flocks in cages compared to barns and free-range systems. The results also showed a significant higher risk for *Salmonella* for a one week increase in flocks' age as well as with a unit increase in the size of the flock.

Key words: *Salmonella*; laying hens; production type; within-flock correlation.

Introduction

Salmonellosis constitutes a major public health burden and represents a significant cost in many countries. In Belgium, the disease ranks high among the reported food borne illnesses (Collard *et al.*, 2004). Even if the incidence of human salmonellosis has diminished since 1999, in 2004, 9545 cases were reported in the country (EFSA, 2006a). As in most of the countries around the world, Belgian *Salmonella* outbreaks in humans are very often linked to the consumption of contaminated eggs (Davies and Breslin, 2001a; Van Immerseel *et al.*, 2005). The most frequently isolated serotype in layer flocks in the EU as well as in Belgium is *Salmonella* Enteritidis which is a non-typhoid non-host adapted serotype with a very wide host range (Baird-Parker, 1990; Gast *et al.*, 2005; Quinet, 2005; VAR, 2005; EFSA, 2004). The bacterium infects the eggs by two processes: first by vertical transmission during the development of the egg within the ovary or its passage through the oviduct and secondly by horizontal transmission through trans-shell contamination (Kinde *et al.*, 2000; WHO FAO, 2002; Davies and Breslin, 2003a; Van Immerseel *et al.*, 2005). Vertical transmission is considered to be the major route of egg contamination and should be controlled by applying sanitary measures at the breeders level (that is, hygiene practices and eventually vaccination) while horizontal transmission should be reduced by preventing contacts between the layer

1 hens and by cleaning and disinfecting the flock's environment. *Salmonella* is known for its
2 ability to asymptotically infect the hen's oviduct (De Buck *et al.*, 2004a; 2004b).
3 Therefore detection of infected flocks depends entirely on laboratory analysis. An infected
4 hen may contaminate one egg out of 200 (Quinet, 2005). Reducing *Salmonella* flock
5 prevalence results in a directly proportional reduction in human health risk (Altekruse *et al.*,
6 1993). This suggests that sanitary measures at the flock level contribute to a significant
7 reduction of the risk for salmonellosis due to egg consumption. In Belgium, the layer
8 breeders are not significantly infected, probably due to the many years' efforts of control at
9 this level and therefore, it is reasonable to assume that most day-old chicks are free from
10 *Salmonella* when placed on farms (Davies and Breslin, 2001; AFSCA, 2004). The majority
11 of the infections in layer hens seem to be attributed to the persistent contamination of the
12 farm. Indeed, the presence of *Salmonella* in the laying house environment has been strongly
13 correlated with the probability that hens will lay contaminated eggs. Chicken are infected
14 after oral ingestion of the bacteria from the environmental sources (for example,
15 contaminated fluff, dust, feed etc) invasion of the mucosal epithelial cells, which leads to
16 systemic dissemination and colonization of the ovary and oviduct (Henzler *et al.*, 1998;
17 Davies and Breslin, 2003b). The primary control should focus at farm level. Control
18 measures include preventing contacts with contaminated feed and visitors, wearing house-
19 specific clothing, thorough cleaning and disinfection of the layer houses, vaccination, rodent
20 control programs. In Belgium, every holding housing more than 5000 hens is required to be
21 sampled for *Salmonella* diagnosis 3 weeks before slaughter time. This measure probably has
22 contributed to a reduction of the risk for food-borne salmonellosis. However, in 2004, still
23 27% of the layer flocks analysed remained positive for *Salmonella* (AFSCA, 2004). Several
24 risk factors have been described, but in order to advise the Belgian competent authority
25 (Federal Agency for the safety of the Food Chain) with detailed, practical guidance, an

understanding of possible causal factors is essential. The objective of the study reported here is to investigate the risk factors which are associated with the occurrence of *Salmonella* in laying hens in Belgium using data collected for the Baseline Study on the Prevalence of *Salmonella* in laying flocks of *Gallus gallus* f. domestica in the European Union (SANCO/34/2004 and Commission decision 2004/665/EG). Although it would be worthwhile to utilize data from earlier years, the 2005 data set contained flock information, particularly on some demographic factors and *Salmonella* vaccination status, which were unavailable for earlier databases.

Materials and methods

Data collection

The Belgian part of the Baseline Study on the Prevalence of *Salmonella* in egg laying flocks of *Gallus gallus* in the European Union consisted of a cross-sectional study that covered the year 2005 from February to September in Belgium. The primary sample size providing the number of holdings which had to be tested was calculated on the basis of a target prevalence of 20%, a confidence level of 95% and an accuracy of 3% (Commission decision 2004/665/EG). The population of laying hens was stratified according to holding size (below 1000, 1000-2999, 3000-4999, 5000-9999, 10000-29999, 30000 and more). The number of holdings to be sampled was subsequently distributed proportionally to the number of holdings in each class. In all cases, only one flock per holding was sampled. Seven different samples, two dust samples and five faecal samples were collected from each selected flock. The dust samples were any of these types: 1) dust from different places in case of barn or free range flocks, 2) dust from egg belts, 3) dusty material from beneath cages. Faecal samples were any of these types: 1) boot swabs which are indeed socks placed over the boots. They are sufficiently absorptive to collect faecal or moist litter samples from the floor

1 surfaces (SANCO/34/2004 and Commission decision 2004/665/EG). 2) Pooled faecal
2 samples from dip pits, 3) pooled faecal samples from dropping belts, 4) pooled faecal
3 samples from scrapers. The collection of these samples was as follows: There had to be five
4 pooled faecal samples taken per selected flock. For the pooled faecal samples in cages, there
5 are normally several stacks of cages within a henhouse. The material from each stack picked
6 up using a new pair of plastic gloves for each individual sample was included in each of the
7 five pooled faecal samples of 200-300 grams. For the boot swabs in barns and free range
8 flocks, each henhouse was divided in sectors of at least 100m that were walked on with new
9 boot swabs, five pairs of boot swabs per henhouse. Each of the five pooled samples
10 comprised of faecal material fixed to a pair of boot swabs. The dust material from beneath
11 cages was obtained from 20 separate locations within a henhouse using a new pair of plastic
12 gloves for each sample. Finally for the dust from different places from barns and free range,
13 each dust sample was collected in a 250ml plastic jar or bag ensuring that all parts of the
14 henhouse like from exhaust fan, ledges, beams etcetera were covered. In order to maximise
15 sensitivity both faecal material (5 out of 7) and dust material from the environment (2 out of
16 7) were sampled, depending on whether the birds were reared in cages or barns or free-
17 range, in such a way that the complete farm was represented. The hens were sampled at the
18 end of their laying period, within a maximum of 9 weeks before depopulation. Samples were
19 sent within 24 hours to the laboratory. The detection method was as recommended by the
20 Community Reference Laboratory for Salmonella in Bilthoven, The Netherlands, that is, a
21 modification of ISO 6579:2002. *Salmonella* isolates were serotyped following the
22 Kaufmann-White scheme (Popoff, 2001; VAR, 2005).

Data description

Although the proportion of flocks infected with *Salmonella* may significantly differ depending on the type of sample that was used (Kinde *et al.*, 2005), for the analyses in this study we grouped the three dust-type samples to form the ‘dust material’ and the four faecal-type samples formed the ‘faecal material’ thus reducing the seven sample types to two for all flocks. The ‘dust material’ and ‘faecal material’ were considered as positive (outcome=1) when at least one of the dust or faecal samples respectively was positive. They were considered as negative (outcome=0) when all of the dust or faecal samples respectively were negative. The frequencies of infected flocks were obtained based on the ‘dust material’ and ‘faecal material’ separately. Since the two outcomes, one from the ‘dust material’ and one from the ‘faecal material’, occurred on each flock, it was important to examine if an association existed between them. This was done using the Pearson Chi-square test of independence (FREQUENCY procedure in SAS). Also the measure of this association was explored using the Pearson correlation coefficient with the SAS CORRELATION procedure. The existence of an association signalled the necessity for the two outcomes to be modelled jointly.

The explanatory variables used include: region (1= Walloon or 0= Flanders), sampling time (month the flock was sampled: February to September), production type (cage or barn/ free range), age (in weeks), flock size (number of hens in the flock considered) and vaccination status against *Salmonella* (yes, unknown, or no). The flocks were vaccinated against *Salmonella enterica*, serovar Enteritidis during the rearing period (one day to 18-20 weeks) with either a live or inactivated vaccine type although for some flocks the vaccine type was not known. The last dose was administered a few weeks before the onset of laying eggs. The pullets were kept in separated installations on the laying farm considering special conditions like temperature and light among others. The associations between presence and absence of

Salmonella and each of the categorical variables was investigated using the Pearson chi-Square test of independence (FREQUENCY procedure in SAS). To explore the relation of the outcomes with the continuous explanatory variables we used the mean.

Data analysis

In this study we modelled the probability of infection of a flock. Therefore, we carried out analyses for the dichotomized bivariate response where a flock was infected if at least one of the samples of the ‘dust material’ or ‘faecal material’ tested positive otherwise the flock was considered not infected. Since the ‘dust material’ and ‘faecal material’ responses were binary outcomes, a natural assumption for their distribution was the binomial distribution. Various approaches and models were used to model these data. The first approach to analyse these data was to perform separate analyses for the two outcomes, for example, by fitting a logistic model, $\text{logit}(P(Y_D = 1)) = \mathbf{X}^T \boldsymbol{\beta}_D$ for the dust outcome variable and another logistic model, $\text{logit}(P(Y_F = 1)) = \mathbf{X}^T \boldsymbol{\beta}_F$ for the faecal outcome variable. The probabilities of the presence of *Salmonella*, $P(Y_D = 1)$ in dust and $P(Y_F = 1)$ in faeces, were predicted as functions of explanatory variables contained in the \mathbf{X} design matrix using the logit link function. The estimates of the model parameters, $\boldsymbol{\beta}_D$ and $\boldsymbol{\beta}_F$, were obtained using maximum likelihood estimation. This however would ignore the correlation between the two outcomes. Moreover the Pearson correlation coefficient showed a tendency for the two outcomes to relate positively, meaning that when the dust outcome was positive the faecal outcome tended to be positive as well or vice versa.

In the second approach we modelled both outcomes jointly as (Y_D, Y_F) , for example; by fitting the Generalized Estimating Equations (GEE) model, introduced by Liang and Zeger (1986). In order to use maximum likelihood estimation the joint

probabilities: $P(Y_D = 1, Y_F = 1)$, $P(Y_D = 1, Y_F = 0)$, $P(Y_D = 0, Y_F = 1)$, and $P(Y_D = 0, Y_F = 0)$ must be assumed at each combination of explanatory variables. However, when there are many explanatory variables this is not practical especially if some are continuous (Agresti, 2002). An alternative to maximum likelihood fitting uses the quasi-likelihood. Instead of assuming a bivariate binomial distribution for (Y_D, Y_F) , the quasi-likelihood method specifies a model for the means of the marginal distributions of Y_D and Y_F ; a variance function describing how the variance of Y_D and Y_F depend on their means; and a pairwise correlation, $\text{corr}(Y_D, Y_F) = \rho$ between the outcomes. Therefore the model was applied to two sets of marginal binomial parameters $\{P(Y_D = 1)\}$ and $\{P(Y_F = 1)\}$. The marginal logit model is then of the form

$$\begin{bmatrix} \log \text{it}\{P(Y_D = 1)\} \\ \log \text{it}\{P(Y_F = 1)\} \end{bmatrix} = \begin{bmatrix} \mathbf{X}^T \boldsymbol{\beta}_D \\ \mathbf{X}^T \boldsymbol{\beta}_F \end{bmatrix}$$

The estimates to the model parameters were obtained as solutions of quasi-likelihood equations called generalized estimating equations. In this study we assumed an exchangeable working correlation structure. Essentially the correlation between the outcomes was estimated and then used to re-estimate the regression parameters and adjust the standard errors. An advantage of the GEE model is that the estimates are valid even if one misspecifies the variance-covariance structure (Agresti, 2002; Molenberghs and Verbeke, 2005). In addition the GEE model estimates the magnitude of the correlation between the outcomes taking into account the explanatory variables.

The models were fitted using the SAS GENMOD procedure. A parsimonious model was built based on the ordinary logistic model where the probability of *Salmonella* positivity was modelled by including one explanatory variable (two continuous and four categorical) at a time and the variables that had a p-value less than 0.25 were introduced in the multiple

logistic regression models. A stepwise automatic selection procedure was also used to supplement the model selection. The two criteria led to the same model. Along with the selected main factors, their two-way interactions were added to the model. Higher interactions were not considered in order to keep a reasonable number of parameters in regard to estimation. However, two-way interactions between categorical variables, for instance, production type by vaccination status resulted into observations with only one type of the outcomes causing difficulties in estimation. The interactions between categorical and continuous variables posed no estimation problems but were found to be non-significant. Therefore the final model considered eliminated the 'region' variable and the interactions. By fitting the final model the results for the risk factors were expressed as odds ratios along with their corresponding 95% confidence intervals and probability values. A probability value of less than 0.05 indicated a statistically significant result.

Results

Data description

In total, data was recorded for 148 flocks. In Figures 1 and 2, we show the number of flocks with positive or negative results for *Salmonella* for dust and faecal samples. The numbers at the top of the bars indicate the number of flocks in each category on the horizontal axis. Specific to the dust sample type, Figure 1 also shows that in 102 flocks no dust sample was *Salmonella* positive whereas 22 flocks had one positive dust sample and 24 flocks had both dust samples positive. A similar interpretation follows for the faecal sample type. Grouping the results from Figure 1 into *Salmonella* positive flocks (if at least one sample was *Salmonella* positive) and *Salmonella* negative flocks (if all samples were *Salmonella* negative) produces Figure 2. Considering the dust sample type, for instance, 102 out of 148 flocks were *Salmonella* negative while the 46 were positive for *Salmonella*. The frequencies

for the faecal sample type are interpreted in a similar manner. The Pearson chi-square statistic for the association between the two outcome variables was estimated at 66.60 ($p < 0.001$) which rejects the null hypothesis of no association between the dust and faecal outcomes. The Pearson correlation coefficient between the two outcomes was obtained as 0.6708 giving an indication of moderate to strong positive association.

Figure 1 here

Figure 2 here

Table 1 shows the distribution of the number of *Salmonella* positive and negative flocks per each categorical explanatory variable. The percentages of all flocks that were positive or negative and the associations between presence and absence of *Salmonella* and each of the categorical variables using Pearson chi-Square test of independence are also shown. For both sample types there seems to be significant (p -values < 0.05) associations of production type and *Salmonella* vaccination status on the presence and absence of *Salmonella*.

Table 1 here

For the *Salmonella* positive group, the flocks' mean age (in weeks) was 74.87 and 76.15 while the mean flock size was 21929.22 and 22156.6 for dust and faecal materials respectively. Similarly, for the *Salmonella* negative group, the mean age was 70.75 and 70.11 while the mean flock size was 13912.28 and 13727.1 for dust and faecal materials respectively. The mean age and mean flock size were higher for the *Salmonella* infected flocks than for the uninfected ones, suggesting an increased risk for *Salmonella* as the hens get older and as the flock size increases.

Data analysis

Knowing that the two outcomes of *Salmonella* were from the same flock meant that analyses which take into account the dependence between the responses from the dust material and faecal material outcomes were more appropriate. However, in order to explore the changes in effects with the complexity of a model, we analysed the two outcomes separately, first with univariate simple logistic models (USLM) shown in column 2 of Table 2 and secondly using univariate multiple logistic models (UMLM) presented in column 3 of Table 2. In these separate analyses the dust and faecal datasets were assumed to be independent. These findings were then compared with the findings from the appropriate model, the bivariate multiple logistic model (BMLM) in column 4 of Table 2. Column 2 shows the odds ratios estimated from the univariate simple logistic model analyses where one covariate was entered in the model while column 3 gives the estimated odds ratios from the univariate multiple logistic regression model where other covariates were controlled for. The USLMs identified that rearing flocks in cages compared to barns and free-range, not vaccinating flocks, a unit increase of flock size and a one week increase in flock age as significant risk factors for *Salmonella* in Belgian layer flocks. Controlling for other factors in the UMLMs showed that rearing layer flocks in cages is a significant risk factor in both dust and faecal data sets whereas a one week increase in flock age and a unit increase of flock size were significant risk factors for the faecal dataset alone.

However in column 4, which presents the appropriate analysis for the data, the joint analysis of the two datasets with the correlation between dust and faecal outcomes modelled as an exchangeable working correlation using GEE we observed that, controlling for other variables, rearing layer flocks in cages was still a significant risk factor but flock age, flock size and the month of July became borderline significant while *Salmonella* vaccination status

turned non-significant. A working correlation of 0.7384 was estimated which indicates a strong positive association between the responses that was ignored by USLMs and UMLMs, which modelled the two responses separately. Therefore the bivariate GEE confirmed the strong association between the outcomes and estimated this association even higher compared to the exploratory measure, the Pearson correlation coefficient of 0.6708 that did not account for other factors. About the risk factors for *Salmonella* found in this study, the exploratory data analysis, which gave an indication of risk factors using Pearson chi-square test of independence for categorical variables (Table 1) and using means for the continuous variables, and the confirmatory analysis via modelling (Table 2) led to similar conclusions.

Table 2 here

Discussion

The prevalence of *Salmonella* in commercial holdings of laying hens in Belgium is relatively high, especially when compared to the northern European countries (EFSA, 2006b). However, it should be mentioned that Belgium has many laying hens compared to neighbouring countries (Quinet, 2005). The European survey was based on environmental sampling which is considered to be an accurate and representative indicator for the presence of *Salmonella* in layer flocks and for the probability that hens would lay contaminated eggs (Henzler *et al.*, 1994; Kinde *et al.*, 2005). The persistence of the pathogen in the intestinal tract is more important when infection occurs in young chicks, since bacterial clearance occurs more efficiently in adults. Genetically distinct lines of hens and various breeds can also be responsible for differences in the presence of *Salmonella* in the faeces of a contaminated animal. It is important to take these factors into account as the duration of this shedding can influence the detection of *Salmonella* in the threatening flocks (Kinde *et al.*, 2000; Gast *et al.*, 2005). Environmental sampling is not entirely reliable as it can miss flocks

1 which passed the peak of infection but which are still producing contaminated eggs (Kinde *et*
2 *al.*, 1996; Davies and Breslin, 2004; Van Immerseel *et al.*, 2005). The fact that one specific
3 type of sample would be more contaminated than others helped identify risk factors, for
4 example, a high level of the bacteria in dust (two dust samples positive instead of one) could
5 point out a problem due to the ventilation system in the hen house or may be associated with
6 cleaning and disinfection of the house, or with insufficient rodent control. A study from Gast
7 *et al.*, 1998 suggested that infection could, among other things, occur by oral ingestion of
8 external surfaces contaminated by airborne movement of *Salmonella* during the feeding or
9 pecking. From our findings, we saw differences in the statistical relations between the
10 response variable and the predictors. For instance, the age factor was statistically associated
11 to *Salmonella* status in the faecal dataset ($p=0.05$), while not significantly associated
12 ($p=0.439$) in the dust dataset. The risk for *Salmonella* in cages versus barn and free range
13 was twice as high in the dust dataset as in the faecal dataset (OR= 20.11 versus 10.27).
14 The major risk factor identified from the analysis was rearing flocks in cages compared to
15 rearing in barns and free-range systems. The risk of contamination with *Salmonella* is
16 thought to be higher when eggs are produced in non-cage systems, because of the greater
17 exposure of layers to environmental contamination (Kinde *et al.*, 1996; EFSA, 2004).
18 However, in practice, control is not easier in cage layer houses; due to the difficulty to
19 efficiently disinfect the cages and the higher densities of birds which produce a larger
20 volume of contaminated faeces and dust (Davies and Breslin, 2004). The result of the current
21 study clearly corroborates this finding. In addition, a clear difference was noticed in the
22 proportions of vaccinated hens in the two types of production systems: 88% of the barn and
23 free-range birds were vaccinated, while only 53% for the cage system poultry. The
24 vaccination variable can act here as a confounding factor on the apparent association
25 between production type and *Salmonella* status. However, in the description of the sampled

population of this present study, we noticed that the proportion of the “barn and free-range” category is relatively small (23%). Moreover, the very wide confidence intervals suggest that there might be a problem due to sample size.

Most of the studies have proven vaccination to be an important aid to reduce or possibly eliminate *Salmonella* Enteritidis from laying flocks (Davies and Breslin, 2001b; 2003b; 2004). In the United Kingdom for instance, most of the laying flocks which have been implicated in the recent outbreaks of *Salmonella* Enteritidis in human beings were unvaccinated (Davies and Breslin, 2001). In the present analysis vaccination seemed not to have a significant protective effect. In the cases when *Salmonella* serovars other than *Salmonella enteritidis* are present concurrently in flocks vaccinated for *Salmonella enteritidis*, then considerably more contamination with these other *Salmonella* serovars may occur (Davies and Breslin, 2004). Another explanation why vaccination was less effective than expected, is that hens might have been infected before the vaccination was completed. Therefore it would have been interesting to exploit the period when the flock had been vaccinated as an explanatory variable. Such a variable was indeed available in the initial database but we chose to leave it aside for two main reasons. First, since the variables “vaccination status” and “vaccination period” were related to each other, we used only one of them to avoid multicollinearity problems. Second, from the description of the “vaccination period” variable, we had 88 holdings where vaccination was performed at rearing out of the 90 holdings where hens were vaccinated, leaving us with nothing to properly compare these findings with. Furthermore, effective protection owed to vaccination might occur only when the challenge dose is low. It is crucial to keep in mind that for vaccination to work effectively, an efficient cleaning and disinfection of laying houses between successive flocks is compulsory (Davies and Breslin, 2003b; Van den Bosch 2003). In this study, other factors like hygiene practices or pest control and their potentially

1 confounding effects on the association between vaccination and the probability of being
2 infected by *Salmonella*, were not taken into account.

3 The influence of temperature on the growth of *Salmonella* in food has been well
4 documented. It is known that in all countries the incidence of human salmonellosis is highest
5 during the summer (Baird-Parker, 1990; CNRSS, 2004; Kovats *et al.*, 2004). Even though a
6 statistically significant effect of the “month” variable is reported from our study, it is
7 difficult to show the direction of the influence as only the month of July had borderline
8 significance. Mollenhorst *et al.*, 2005 came to the same conclusion. During the summer
9 season of the year 2003, a large increase of *Salmonella* infections was observed in Belgium
10 and in The Netherlands. This increase could probably be attributed to the extremely hot
11 weather during the summer of 2003. The Dutch study (Van Pelt *et al.*, 2004) showed that a
12 concomitant outbreak of *Salmonella* and avian influenza led to a shortage of eggs on the
13 Dutch market, which was to be compensated for with imports, providing a reasonable
14 explanation for this apparent seasonal trend.

15 This present study showed no evidence of significant differences in the distribution of
16 *Salmonella* among laying flocks according to regional repartition, and the odds ratios were
17 very close to 1 in both faecal and dust samples. Again we should note that the sample
18 repartition is not really equitable, the Walloon holdings representing only 18%. On the other
19 hand, the number of human salmonellosis cases across the country is clearly much higher in
20 Flanders. Although the eggs produced in Belgium do not necessarily tend to be consumed
21 locally, the food practices vary between both regions (CNRSS, 2004; AFSCA, 2006).

22 The impact of the age factor on the occurrence of *Salmonella* among egg laying flocks
23 cannot really be established here, as the odds ratios and the confidence intervals were all
24 close to 1.

At last, other risk factors which were not considered in the present study are important to mention. For example, it could be useful to build a model taking into account flock characteristics (type of breed, number of flocks on the farm, multi-age farm or not), farm management (control of pest access, visitors allowed or not, feed composition and feeding practices, drinking water), cleaning and disinfecting practices related with the contamination status of the previous flock in the same hen house ((Henzler & Opitz, 1992; Kinde *et al.*, 1996; Shirota *et al.*, 2000; Garber *et al.*, 2003; Liebana *et al.*, 2003; Kinde *et al.*, 2005). Knowing that non-typhoid *Salmonellae* have very wide host ranges, it is important to take into consideration all various potential vectors surrounding the flock.

References

AFSCA, Agence Fédérale de la Sécurité de la Chaîne Alimentaire, 2004. Salmonella, rapport annuel 2004.

AFSCA, Agence Fédérale pour la Sécurité de la Chaîne Alimentaire, 2006. Bulletin de l'Agence Fédérale pour la Sécurité de la Chaîne Alimentaire, Mai 2006, 4-5.

Agresti, A., 2002. Categorical Data Analysis. (Second edition). Wiley, New York.

Altekruse, S., Koehler, J., Hickman-Brenner, F., Tauxe, R.V., Ferris, K. A., 1993. Comparison of Salmonella enteritidis phage types from egg-associated outbreaks and implicated laying flocks. Epidemiology and Infection 110 (1), 17-22.

Baird-Parker, A.C., 1990. Foodborne salmonellosis. Lancet 336, 1231-1235.

Collard, J.M., Bertrand, S., Willems, L., Baeyens, D., De Cooman, F., Steenhaut, H., Lattuca, M., Mairiaux, E., Dupont, Y., Godard, C., Wildenauwe, C., Vrints, M., 2004. Human Salmonellosis in Belgium: recent trends and outbreaks in 2003. Proceedings of Belgian symposium on Salmonella research and control in poultry.

CNRSS, Centre National de Référence des *Salmonella* et *Shigella*, 2004. Rapport Annuel 2004. Institut scientifique de Santé Publique.

Davies, R., Breslin, M., 2001. Environmental contamination and detection of Salmonella enterica serovar enteritidis in laying flocks. Veterinary Record 149 (23), 699-704.

Davies, R., Breslin, M., 2003a. Observations on *Salmonella* contamination of commercial laying farms before and after cleaning and disinfection. Veterinary Record 152 (10), 283-287.

Davies, R., Breslin, M., 2003b. Effects of vaccination and other preventive methods for Salmonella enteritidis on commercial laying chicken farms. Veterinary Record 153 (22), 673-677.

Davies, R. and Breslin, M., 2004. Observations on *Salmonella* contamination of eggs from infected commercial laying flocks where vaccination for Salmonella enterica serovar Enteritidis had been used. Avian Pathology 33 (2), 133-144.

De Buck, J., Pasmans, F., Van Immerseel, F., Haesebrouck, F. and Ducatelle, R., 2004a. Tubular glands of the isthmus are the predominant colonization site of Salmonella Enteritidis in the upper oviduct of laying hens. Poultry Science 83, 352-358.

De Buck, J., Van Immerseel, F., Haesebrouck, F., Ducatelle, R., 2004b. Recent insights on egg contamination and control. Proceedings of Belgian symposium on Salmonella research and control in poultry.

EFSA, European Food Safety Authority, 2004. Opinion of the AHW Panel related to the Welfare aspects of various systems of keeping laying hens. EFSA Journal 197, 1-23.

EFSA, European Food Safety Authority, 2006a. Trends and sources of zoonoses, zoonotic agents and antimicrobial resistance in the European Union in 2004. The EFSA Journal 2005, 23-95.

EFSA, European Food Safety Authority, 2006b. Preliminary Report. Analysis of the baseline study on the prevalence of *Salmonella* in laying hen flocks of *Gallus gallus*. EFSA Journal 81, 1-71.

Garber, L., Smeltzer, M., Fedorka-Cray, P., Ladely, S., Ferris, K., 2003. *Salmonella* enterica serotype enteritidis in Table Egg Layer House Environments and in Mice in U.S. Layer Houses and Associated Risk Factors. Avian Diseases 47, 134-143.

Gast, R.K., Mitchell, B.W., Holt, P.S., 1998. Airborne transmission of *Salmonella* enteritidis infection between groups of chicks in controlled-environment isolation cabinets. Avian Diseases 42 (2), 315-320.

Gast, R.K., Guard-Bouldin, J., and Holt, P.S., 2005. The relationship between the duration of faecal shedding and the production of contaminated eggs by laying hens infected with strains of *Salmonella* Enteritidis and *Salmonella* Heidelberg. Avian Diseases 49, 382-386.

Henzler, D. J., Opitz , H. M., 1992. The role of mice in the epizootiology of *Salmonella* enteritidis infection on chicken layer farms. Avian Diseases 36(3), 625-631.

Henzler, D.J., Ebel, E., Sanders, J., Kradel, D., Mason, J., 1994. *Salmonella* enteritidis in eggs from commercial chicken layer flocks implicated in human outbreaks. Avian Diseases 38 (1), 37-43.

Henzler, D.J., Kradel, D.C. and Sischo, W.M., 1998. Management and environmental risk factors for *Salmonella* enteritidis contamination of eggs. American journal of veterinary research 59, 824-829.

Kinde, H., Read, D.H., Chin, R.P., Bickford, A.A., Walker, R.L., Ardans, A., Breitmeyer, R.E., Willoughby, D., Little, H.E., Kerr, D., Gardner, I.A., 1996. *Salmonella* enteritidis, phase type 4 infection in a commercial layer flock in southern California: bacteriologic and epidemiologic findings. Avian Diseases 40 (3), 665-671.

Kinde, H., Shivaprasad, H.L., Daft, B.M., Read, D.H., Ardans, A., Breitmeyer, R., Rajashekara, G., Nagaraja, K.V., Gardner, I.A., 2000. Pathologic and bacteriologic findings in 27-week-old commercial laying hens experimentally infected with *Salmonella* enteritidis, phage type 4. Avian Disease 44 (2), 239-248.

Kinde, H., Castellan, D.M., Kerr, D., Campbell, J., Breitmeyer, R., Ardans, A., 2005. Longitudinal Monitoring of Two Commercial Layer Flocks and Their Environments for *Salmonella* Enterica Serovar Enteritidis and Other *Salmonellae*. Avian Diseases 49, 189-194.

Kovats, R.S., Edwards, S.J., Hajat, S., Armstrong, B.G., Ebi, K.L., Menne, B., 2004. The effect of temperature on food poisoning: a time-series analysis of salmonellosis in ten European countries. Epidemiology and Infection. 132 (3), 443 – 453.

Liang, K.Y., Zeger, S.L., 1986. Longitudinal data analysis using generalized linear models. *Biometrika* 73, 13-22.

Liebana, E., Garcia-Migura, L., Clouting, C., Clifton-Hadley, F.A., Breslin, M., Davies, R. H., 2003. Molecular fingerprinting evidence of the contribution of wildlife vectors in the maintenance of *Salmonella* Enteritidis infection in layer farms. *Journal of Applied Microbiology* 94 (6), 1024-1029.

Molenberghs, G. and Verbeke, G., 2005. *Models for Discrete Longitudinal Data*. Springer, New York.

Mollenhorst, H., van Woudenberg, C.J., Bokkers, E.G., de Boer, I.J.M., 2005. Risk factors for *Salmonella* enteritidis infections in laying hens. *Poultry Science* 84 (8), 1308-1313.

Popoff, M.Y., 2001. *Antigenic Formulas of the Salmonella Serovars*. 8th edition. WHO Collaborating Centre for Reference and Research on *Salmonella*. Institut Pasteur. Paris, France.

Quinet, C., 2005. La Salmonellose aviaire, état des lieux. *Arsia infos* 20, 1-2.

Shirota, K., Katoh, H., Ito, T., and Otsuki, K., 2000. *Salmonella* Contamination in Commercial Layer Feed in Japan. *Journal of Veterinary Medicine Science* 62 (7), 789-791.

Van den Bosch., G., 2003. Vaccination versus treatment: How Europe is tackling the eradication of *Salmonella*. *Asian Poultry Magazine*, 2-4.

Van Immerseel, F., De Buck, J., Boyen, F., Pasmans, F., Bertrand, S., Collard, J.M., Saegerman, C., Hooyberghs, J., Haesebrouck, F., Ducatelle, R., 2005. *Salmonella* dans la viande de volaille et dans les oeufs, un danger pour le consommateur qui demande la mise en place d'un programme de lutte efficace. *Annales de Médecine Vétérinaire* 149, 34-48.

Van Pelt W., Mevius D., Stoelhorst H.G., Kovats S., Van de Giessen A.W., Wannet W., Duynhoven Y.T.H.P., 2004. A large increase of *Salmonella* infections in 2003 in the Netherlands: hot summer or side effect of the avian influenza outbreak? *Eurosurveillance* 9 (7), 17-19.

VAR, Veterinary and Agrochemical Research Centre, Laboratory of General Bacteriology, 2005. *Salmonella* Serotypes analysed at the CODA-CERVA in 2005. Evolution among Poultry, Cattle and Pig isolates from 1992 to 2005 with results of Antimicrobial Resistance Testing. VAR Report data 2005.

WHO FAO, World Health Organization, Food and Agriculture Organization of the United Nations, 2002. Risk Assessments of *Salmonella* in eggs and broiler chickens, an interpretative summary. *Microbiological Risk Assessment Series* 1.

List of Tables

Table 1: Frequency of *Salmonella* positive/negative flocks (percentage of all 148 flocks) by categorical independent variables and sample type. Association P-values between each categorical variable and the presence/absence of *Salmonella* using Pearson Chi-Square test are shown.

Variable	Dust sample type			Faecal sample type		
	Positive (%)	Negative(%)	Assoc χ^2 P-value	Positive(%)	Negative(%)	Assoc χ^2 P-value
<u>region</u>			0.9698			0.5598
<i>Flanders</i>	38 (25.68)	84 (56.76)		40 (27.03)	82 (55.41)	
<i>Walloon</i>	8 (5.41)	18 (12.16)		7 (4.73)	19 (12.84)	
<u>SamplingTime</u>			0.6570			0.4347
<i>Feb</i>	2 (1.35)	4 (2.70)		3 (2.03)	3 (2.03)	
<i>Mar</i>	4 (2.70)	14 (9.46)		7 (4.73)	11 (7.43)	
<i>Apr</i>	5 (3.38)	16 (10.81)		4 (2.70)	17 (11.49)	
<i>May</i>	7 (4.73)	15 (10.14)		9 (6.08)	13 (8.78)	
<i>Jun</i>	12 (8.11)	16 (10.81)		7 (4.73)	21 (14.19)	
<i>Jul</i>	7 (4.73)	10 (6.76)		8 (5.41)	9 (6.08)	
<i>Aug</i>	4 (2.70)	8 (5.41)		3 (2.03)	9 (6.08)	
<i>Sep</i>	5 (3.38)	19 (12.84)		6 (4.05)	18 (12.16)	
<u>Production Type</u>			<0.0001			0.0002
<i>Cage</i>	45 (30.41)	69 (46.62)		45 (30.41)	69 (46.62)	
<i>barn & free range</i>	1 (0.67)	33 (22.30)		2 (1.35)	32 (21.62)	
<u>Vaccination Status</u>			0.0260			0.0573
<i>Yes</i>	22 (14.86)	68 (45.95)		22 (14.86)	68 (45.95)	
<i>No</i>	22 (14.86)	26 (17.57)		21 (14.19)	27 (18.24)	
<i>Unknown</i>	2 (1.35)	8 (5.41)		4 (2.70)	6 (4.05)	

Table 2: Estimated *Salmonella* infection odds ratios (95% confidence interval limits) and p-values from univariate simple logistic models (USLM) and univariate multiple logistic models (UMLM) under Independency and the bivariate multiple logistic model (BMLM) using GEE approach assuming an exchangeable working correlation between the outcomes. Significant risk factors ($p < 0.05$) are denoted by (*) while the borderline ($0.05 \leq p < 0.1$) risk factors with (+).

1	2		3		4	
	USLM	USLM	UMLM	UMLM	BMLM using GEE	
COVARIATE	DUST	FAECES	DUST	FAECES	DUST	FAECES
Sampling time						
February vs september	1.90 (0.27-13.52)	3.00 (0.47-19.04)	0.95 (0.11-8.39)	1.60 (0.20-12.78)	0.61 (0.03-10.89)	1.11 (0.13-9.22)
March vs september	0.522 (0.25-4.79)	0.244 (0.51-7.17)	0.965 (0.14-3.61)	0.660 (0.27-5.63)	0.738 (0.12-4.14)	0.925 (0.27-6.13)
April vs september	1.09 (0.25-4.79)	1.91 (0.51-7.17)	0.70 (0.14-3.61)	1.24 (0.27-5.63)	0.70 (0.12-4.14)	1.28 (0.27-6.13)
May vs september	0.914 (0.29-4.85)	0.338 (0.17-2.94)	0.669 (0.15-3.35)	0.780 (0.07-1.76)	0.697 (0.15-3.70)	0.758 (0.06-1.97)
June vs September	1.19 (0.29-4.85)	0.71 (0.17-2.94)	0.71 (0.15-3.35)	0.35 (0.07-1.76)	0.73 (0.15-3.70)	0.35 (0.06-1.97)
July vs september	0.811 (0.47-6.72)	0.633 (0.59-7.29)	0.662 (0.28-7.20)	0.203 (0.45-10.92)	0.709 (0.36-7.60)	0.232 (0.44-15.03)
August vs September	1.77 (0.47-6.72)	2.08 (0.59-7.29)	1.41 (0.28-7.20)	2.22 (0.45-10.92)	1.65 (0.36-7.60)	2.56 (0.44-15.03)
September vs September	0.399 (0.83-9.82)	0.254 (0.28-3.52)	0.677 (0.67-12.01)	0.326 (0.17-3.21)	0.520 (0.69-13.35)	0.298 (0.14-3.47)
October vs September	2.85 (0.83-9.82)	1.00 (0.28-3.52)	2.83 (0.67-12.01)	0.74 (0.17-3.21)	3.04 (0.69-13.35)	0.70 (0.14-3.47)
November vs September	0.097 (0.67-10.57)	1.00 (0.71-10.05)	0.157 (0.66-18.75)	0.691 (0.65-17.05)	0.142 (0.83-28.96)	0.664 (0.58-23.45)
December vs September	2.66 (0.67-10.57)	2.67 (0.71-10.05)	3.53 (0.66-18.75)	3.32 (0.65-17.05)	4.89+ (0.83-28.96)	3.68 (0.58-23.45)
January vs September	0.165 (0.40-8.98)	0.147 (0.20-4.95)	0.139 (0.32-11.39)	0.150 (0.10-4.00)	0.080 (0.27-14.21)	0.169 (0.09-4.28)
February vs September	1.90 (0.40-8.98)	1.00 (0.20-4.95)	1.92 (0.32-11.39)	0.64 (0.10-4.00)	1.97 (0.27-14.21)	0.62 (0.09-4.28)
March vs September	0.418 (0.40-8.98)	1.000 (0.20-4.95)	0.475 (0.32-11.39)	0.634 (0.10-4.00)	0.503 (0.27-14.21)	0.627 (0.09-4.28)
Production type						
Cage vs barn and free-range	21.52* (2.84-162.98)	10.43* (2.38-45.70)	16.38* (1.92-139.99)	7.88* (1.47-42.12)	20.11* (2.52-160.49)	10.27* (2.13-49.57)
Free-range vs cage	0.003 (0.000-0.004)	0.002 (0.000-0.004)	0.011 (0.000-0.004)	0.016 (0.000-0.004)	0.005 (0.000-0.004)	0.004 (0.000-0.004)
Vaccination status						
Vaccination vs no vaccination	0.38* (0.18-0.80)	0.42* (0.20-0.88)	0.50 (0.20-1.28)	0.70 (0.27-1.83)	0.49 (0.18-1.31)	0.69 (0.23-2.02)
Vaccination unknown vs no vaccination	0.011 (0.06-1.54)	0.021 (0.21-3.43)	0.148 (0.05-2.23)	0.463 (0.34-9.65)	0.154 (0.02-2.06)	0.494 (0.37-7.89)
Unknown vs no vaccination	0.30 (0.06-1.54)	0.86 (0.21-3.43)	0.35 (0.05-2.23)	1.81 (0.34-9.65)	0.23 (0.02-2.06)	1.70 (0.37-7.89)
Unknown vs vaccination	0.148 (0.06-1.54)	0.828 (0.21-3.43)	0.266 (0.05-2.23)	0.486 (0.34-9.65)	0.188 (0.02-2.06)	0.498 (0.37-7.89)
Age						
Age	1.03+ (1.00-1.06)	1.04* (1.01-1.08)	1.02 (0.98-1.05)	1.04* (1.00-1.08)	1.02 (0.98-1.06)	1.03+ (1.00-1.07)
Age	0.067 (0.000-0.004)	0.008 (0.000-0.004)	0.291 (0.000-0.004)	0.027 (0.000-0.004)	0.439 (0.000-0.004)	0.050 (0.000-0.004)
Flocksize						
Flocksize	1.00* (1.00-1.00)	1.00* (1.00-1.00)	1.00+ (1.00-1.00)	1.00* (1.00-1.00)	1.00+ (1.00-1.00)	1.00+ (1.00-1.00)
Flocksize	0.002 (0.000-0.004)	0.001 (0.000-0.004)	0.081 (0.000-0.004)	0.030 (0.000-0.004)	0.049 (0.000-0.004)	0.071 (0.000-0.004)

List of Figures




Figure 1: number of flocks and the frequency of samples, out of the two for dust and out of five for faeces, which tested positive for *Salmonella*.




Figure 2: number of flocks and their state of salmonella, 0 for absence and 1 for presence of *Salmonella* after collapse of figure 1.