

Functions of some finger joints while handling the PC mouse, and their possible relevance for computer aided learning

Non Peer-reviewed author version

VAN ZWIETEN, Koos Jaap; Hotterbeekx, An; Thywissen, Carlo; Helder, Paul; LIPPENS, Peter; SCHMIDT, Klaus; ZOUBOVA, Irina; Piskùn, Oleg; Varzin, Sergey & Zinkovsky, Anatoly (2010) Functions of some finger joints while handling the PC mouse, and their possible relevance for computer aided learning. In: Auer, Michael E. & Schreurs, Jeanne (Ed.) Academic and Corporate E-Learning In a Global Context. p. 1098-1101..

Handle: <http://hdl.handle.net/1942/11210>

# Functions of some finger joints while handling the PC mouse, and their possible relevance for computer aided learning

K.J. van Zwieten<sup>1</sup>, A. Hotterbeekx<sup>1</sup>, C. Thywissen<sup>2</sup>, P. Helder<sup>3</sup>, P.L. Lippens<sup>1</sup>, K.P. Schmidt<sup>1</sup>, I.A. Zoubova<sup>4</sup>, O.E. Piskun<sup>4</sup>, S.A. Varzin<sup>4</sup>, A.V. Zinkovsky<sup>4</sup>

<sup>1</sup>Functional Morphology, Anatomy, Universiteit Hasselt, BioMed, Diepenbeek, Belgium,

<sup>2</sup>Department of Radiology, Jessa Hospital, Hasselt, Belgium, <sup>3</sup>HandShoe Mouse, Hippus NV, Rotterdam, The Netherlands, <sup>4</sup>Department of Health Sciences and Biomechanics, St. Petersburg State Polytechnic University, IMOP, St. Petersburg, Russia

**Key words:** *e-learning, computer mouse use, finger motion, functional analyses*

## Abstract:

*Highly intense use of the finger, while handling the computer mouse, as in e-learning, may lead to hand- or finger complaints, e.g. in the elderly. To understand the finger positions concerned, we analysed some of its joints by functional anatomical research. A statically and dynamically stabilised finger arch is needed to prevent complaints.*

## 1 Introduction, material and methods

After the development of several distance learning courses, including study routes, on Human Anatomy for Students of Biomechanics and Health Sciences in Russian-speaking countries (1,2), the present authors became aware of ever-growing health-problems worldwide by the continuous use of personal computers in persons of all ages, including our current e-learning students (3,4). Many of such problems are upper extremity movement disorders, as in hands and fingers (3). As finger movement studies related tot PC use mainly concentrate on the second joint of the finger, known as the proximal interphalangeal (PIP-) joint, this study deals with mobility and stability of this PIP-joint in e.g. handling the PC mouse (5,6). Structures governing the PIP-joint's (and other finger-joints') stabile and coordinated movements were therefore observed by *in vitro* anatomical specimens, supple enough to simulate motion *in vivo*. Collateral ligaments of the PIP-joint, revealed by micro-dissection, were then compared to current literature (7). Motion-behaviour of their various parts was interpreted kinematically. By understanding which elements underlie finger coordinated motion, movement problems by intense use of the PC mouse, possibly occurring in e-learning too, may probably be prevented.

## 2 Results

### 2.1 Observations

By their finely tuned positions relative to the axes of each finger's proximal and distal interphalangeal joints, the various tendinous bands within the so-called extensor assembly of the finger, purely mechanically help to guarantee the ratios of coordinated flexion as well as extension in both joints simultaneously. Along the PIP-joint each lateral band is supported by the joint's proper collateral ligament (PCL) (14). This PCL shows its more superficial part, which crosses its deeper part in extension, to shift over the latter in PIP-flexion however. Both parts of the PCL then become taut (7). These two parts, plus the finger's proximal and middle phalanx to which they attach, are thus acting as elements of a crossed four-bar linkage system.

## **2.2 Proximal interphalangeal joint in man and other primates**

At the level of the proximal interphalangeal joint, the disposition in the extended human finger is as follows. The medial tendon or band of the extensor assembly lies on the dorsum of the first phalanx's trochlea, while its lateral bands lie practically dorsally over the strong collateral ligaments of the proximal interphalangeal joint (8). This disposition is determined in man by the trapezoid shape of the trochlea (9). The same shape on cross-section is found in higher primates, whereas in lower primates the trochlea is squarer, becoming even rectangular in primate predecessors like the opossum (10). As a consequence, the lateral bands will mainly stay on the shoulders of the trochlea in these species, impairing a more independent use of their separate fingers as compared to higher primates' fingers, man included (11,12,13).

## **2.3 Flexion of the proximal interphalangeal joint**

In flexion of the human finger, the lateral bands of the extensor assembly, fusing distally into one extensor tendon for the distal interphalangeal joint, glide alongside the flexed proximal interphalangeal joint, thereby acquiring more palmar positions. Thus, by a purely mechanical coupling, their displacements favour the simultaneous flexion of the distal interphalangeal joint, whose extension they normally control (14, 15). In PIP-flexion, their displacements are in turn enhanced by the PIP-joint's collateral ligaments, *viz.* the proper collateral ligaments (PCL) becoming taut whereas the PIP accessory collateral ligaments (ACL) become slack (7).

## **2.4 Imaging results**

In the extended proximal interphalangeal joint, HR-imaging techniques revealed characteristic features of the positions of PIP collateral ligaments, in cross-sections as well as in coronal slices (16,17,18). Undulating trajectories of the PCL fibres in particular, at finger extension, contribute to certain cushioning effects with respect to the extensor assembly's lateral bundles which they support. Thus the lateral bundles maintain dorsal positions in PIP extension (8,14).

## **2.5 Proximal interphalangeal joint ligaments behaviour**

Manipulating anatomical specimens of the finger during micro-dissection enabled us to point out the positions as well as the mechanical behaviour of the proper collateral ligaments (PCL) controlling PIP-flexion. On both sides of the joint, either PCL consists of a superficial part and a deep part that cross each other (7). Together with the bone of the proximal phalanx and the bone of the middle phalanx on which they both insert, these parts (being taut in PIP-flexion) may be interpreted as behaving mechanically like a crossed four-bar mechanism (19).

## **2.6 Kinematical aspects**

A crossed four-bar mechanism in the finger has been suggested, but mainly theoretically (20). With respect to hand prosthetics however, this concept was recently successfully applied (21). The movement pattern of the PIP-joint *in se*, in fact more complicated than that of a simple hinge-joint, was recently described in terms of differing geometries of its joint surfaces (22). Regarding the latter analysis, the paths of motion of the proximal interphalangeal joint, being "generated" by crossed four-bar mechanisms like the one proposed here, do unmistakably fit.

### 3 Practical applications

Because of the key-role of the proximal interphalangeal joint of the finger (together with its enigmatic complexity) in relation to finger flexor tendons and the extensor tendon apparatus or extensor assembly, in creating a stabilised arch of the moving finger, functional-anatomical analyses should use *in vitro* micro-dissections as well as *in vivo* HR-imaging techniques (23). After applying these methods, we now could present a concise description of the initial results

Besides, the functional demand of a stabilised arch of the finger will be met, by designing and using those PC mice that are pre-shaped to prevent disorders by intense use of the mouse (24). In current e-learning practice, each student's fingers, hand, and even whole upper extremity, may profit from ergonomically safe working conditions, thus using the computer successfully.

### 4 Conclusion and discussion

The stabilised arch of the freely moving finger, in contact with the PC mouse, largely depends on positions of its flexor- and its extensor-tendons (the so-called extensor assembly) relative to the proximal interphalangeal joint. We could point out these positions, by micro-dissection *in vitro*, simulating finger movements *in vivo*. The latter, visualised in two healthy individuals (AH; KJvZ) by HR-imaging techniques, showed the extensor assembly's precise positions. PIP-joint collateral ligaments moreover act as a crossed four-bar linkage system, guaranteeing this joint's movements in static and dynamic situations contributing to a stabilised finger arch. To prevent motion complaints, computer mouse designers should take into account these data.

### References:

- [1] Van Zwieten, K. J.; Wouters, I.; Curvers B. (2001): Osnovii Anatomii dlja Izuchayauschich Biomechaniku i Valeologiyu, Konspekt lektsij - Base Course Anatomy for Students of Biomechanics and Health Sciences. Editors: Van Zwieten, K.J.; Zinkovsky, A.V.; Schmidt, K.P. Russian translation and redaction: I.A. Zoubova, A.A. Ivanov, S.A. Varzin, 155 pp., also CD-ROM. Diepenbeek - St. Petersburg, 2001.
- [2] Van Zwieten, K.J.; Wouters, I.; Curvers B. (2001): Osnovii Anatomii dlja Izuchayauschich Biomechaniku i Valeologiyu, Rukovodstvo dlja Samostayatelnaya Isuchenya - Base Course Anatomy for Students of Biomechanics and Health Sciences, Study Route. Russian translation and redaction: I. A. Zoubova, A.A. Ivanov, S.A. Varzin, 47 pp., also CD-ROM. Diepenbeek - St. Petersburg, 2001.
- [3] Van Boxtel, M.P.J.; Slegers, K.; Jolles, J.; Ruijgrok, J.M. (2007): Risk of upper limb complaints due to computer use in older persons: a randomized study. *BioMed Central Geriatrics*, 7, 21.
- [4] Blackstone, J. M.; Karr, C.; Camp, J.; Johnson, P. W. (2008): Physical exposure differences between children and adults when using standard and small computer input devices. *Ergonomics*, 51, 6, 872-889.
- [5] Lee, D.L.; McLoone, H.; Dennerlein, J.T. (2008): Observed finger behaviour during computer mouse use. *Applied Ergonomics*, 39, 107-113.
- [6] Chivapornthip, P.; Bohez, E.L.J.; Nanthavanij, S.; Siththiseripratip, K.; Lorprayoon, E. (2010): Biomechanics of Index Finger during Mouse Click. Proceedings of the 3rd International Conference on the Development of Biomedical Engineering in Vietnam, 11-14th Jan 2010, 51-54.
- [7] Schmidt, H.-M.; Lanz, U. (2004): *Surgical Anatomy of the Hand*. Thieme, Stuttgart New York.
- [8] Van Zwieten, K. J.; Lippens, P.L. (2003): Some Factors Influencing 3-D Modelling of the Human Finger. Brussels International Symposium: Advances in Anatomy and Clinical Examination of the Hand and Wrist, March 28-29, 2003. Department of Orthopedics, University of Brussels <http://www.ulb.ac.be/assoc/chorus/orthogenva/2003/abstracts/session4.pdf>
- [9] Landsmeer, J.M.F.; Van Zwieten, K.J. (1974): Observations on the extensor assembly in some primate species. *Journal of Anatomy*, 117, 204-205.

- [10] Van Zwieten, K.J.; Lippens, P.L.; Hauglustaine, S.; Lemelin, P. (1994): Interphalangeal flexion in the opossum. 1st Benelux-congress of Zoology, University of Leuven, 4-5 November 1994, Book of Abstracts, 39.
- [11] Landsmeer, J.M.F. (1979): The extensor assembly in two species of opossum, *Philander opossum* and *Didelphis marsupialis*. *Journal of Morphology*, 161, 373–346.
- [12] Van Zwieten, K.J. (1980): The extensor assembly of the finger in man and non-human primates. PhD Thesis, University of Leiden, J.J. Groen en Zoon, Leiden.
- [13] Christel, M. (1993): Grasping techniques and hand preferences in Hominoidea. In : *Hands of Primates*. H. Preuschoft, D.J. Chivers (Editors). Springer-Verlag, Wien New York.
- [14] Van Zwieten, K.J.; Lippens, P.L.; Gelan, J.; Adriaensens, P.; Schmidt, K.P.; Thywissen, C.; Duyvendak, W. (2008): Coordination of interphalangeal flexion in the human finger. *Journal of Hand Surgery - British and European Volume*, 33(1), 170-171.
- [15] Landsmeer, J.M.F. (1976): *Atlas of Anatomy of the Hand*. Churchill Livingstone, Edinburgh London and New York.
- [16] Kanavel, A.B. (1921): *Infections of the Hand*. Lea & Febiger, Philadelphia and New York.
- [17] Lewis, A.R.; Nolan, M.J.; Hodgson, R.J.; Ralphs, J.R.; Archer, C.W.; Tyler, J.A.; Hall, L.D. (1996): High Resolution Magnetic Resonance Imaging of the Proximal Interphalangeal Joints. *Journal of Hand Surgery (British and European Volume)*, 21B, 4, 488-495.
- [18] Lewis, A.R.; Ralphs, J.R.; Kneafsey, B.; Benjamin, M. (1998): Distribution of Collagens and Glycosaminoglycans in the Joint Capsule of the Proximal Interphalangeal Joint of the Human Finger. *The Anatomical Record*, 250, 281-291.
- [19] Van Zwieten, K.J.; Hauglustaine, S.; Curvers, L.; Lippens, P.L.; Sholukha, V.A.; Ivanov, A.A. (2001): Jaw opening and closing movements, comparative and biomechanical aspects. *European Journal of Morphology*, 39, 4, 246.
- [20] Ngalé Haulin, E.; Lakis, A.; Vinet, R. (2001): Optimal synthesis of a planar four-link mechanism used in a hand prosthesis. *Mechanism and Machine Theory*, 36, 1203-1214.
- [21] Ramirez, I.A.; Lusk, C.P.; Dubey, R.; Highsmith, M.J.; Maitland, M.E. (2009): Crossed four-bar mechanism for improved prosthetic grasp. *Journal of Rehabilitation Research & Development*, 49, 8, 1011-1020.
- [22] Dumont, C.; Albus, G.; Kubein-Meesenburg, D.; Fanghänel, J.; Stürmer, K.M.; Nägerl, H. (2008): Morphology of the Interphalangeal Joint Surface and its Functional Relevance. *Journal of Hand Surgery*, 33 A, 9-18.
- [23] Lippens, P.L.; Sholukha, V.A.; Hauglustaine, S.; Adriaensens, P.; Van Zwieten, K.J.; Gelan, J. (2001): Computer reconstruction of human tendon architecture based on NMR images. *Journal of Morphology*, 148, 255.
- [24] Chen, H.-M.; Leung, C.-L. (2007): The effect on forearm and shoulder muscle activity in using different slanted computer mice. *Clinical Biomechanics*, 22, 518-523.

## Corresponding Author:

Van Zwieten, Koos Jaap, MD PhD

Department of Anatomy, Functional Morphology Research Group, Faculty of Medicine, BioMed Institute, University of Hasselt, University Campus, Agoralaan, Bldg. D, Room D 50 B 3590 Diepenbeek, Belgium

Email [koosjaap.vanzwieten@uhasselt.be](mailto:koosjaap.vanzwieten@uhasselt.be)