

Muscles involved in foot eversion movements during the swing phase of gait - some comparative-anatomical and functional-morphological aspects

Non Peer-reviewed author version

NARAIN, Faridi; Brauns, Annelien; Lemmens, Karolien; VAN ZWIETEN, Koos Jaap; DUYSSENS, Jacques; HAUGLUSTAINE, Stephan; VANDERSTEEN, Marjan; LIPPENS, Peter; ADRIAENSENS, Peter; THEUNISSEN, Evi; GELAN, Jan & Lamur, Kenneth (2011) Muscles involved in foot eversion movements during the swing phase of gait - some comparative-anatomical and functional-morphological aspects. In: Wallner, Christian & Secretary N.A.V. - Nederlandse Anatomen Vereniging (Ed.) Proceedings of 173rd Scientific Meeting Nederlandse Anatomen Vereniging (N.A.V.): vol. 1. p. 28-28..

Handle: <http://hdl.handle.net/1942/11478>

# **Muscles involved in foot eversion movements during the swing phase of gait - some comparative-anatomical and functional-morphological aspects**

Faridi Narain (1), Annelien Brauns (2), Karolien Lemmens (2), Koos Jaap van Zwieten (2), Jacques Duysens (3), Stephan Hauglustaine (2), Marjan Vandersteen (2), Peter Lippens (2), Peter Adriaensens (2), Evi Theunissen (2), Jan Gelan (2), Kenneth Lamur (1)

(1) Department of Anatomy, Faculty of Medical Sciences, Medisch Wetenschappelijk Instituut (MWI), Anton de Kom University of Suriname, Paramaribo, Suriname (2) Department of Anatomy, Functional Morphology, BioMed Research Institute, Faculty of Medicine, Universiteit Hasselt, Diepenbeek, Belgium (3) Department of Biomedical Kinesiology, Faculty of Kinesiology and Rehabilitation Sciences (FaBeR), Katholieke Universiteit Leuven, Belgium

Surinam opossum *Didelphis marsupialis*, considered as a predecessor of primates including man, shows extreme inversion of hindfoot at late stance during quadrupedal walking (Narain et al., 2003). In inversion, the sole of the foot turns inward, in eversion it turns outward. Repositioning of the opossum foot occurs in swing, eversion starting at toe-off (Narain et al., 2009). In midswing, *Didelphis* keeps its foot horizontally, as video-grabbed images reveal. Tendons of extensor digitorum brevis, and of peroneus muscles, passing behind the fibular malleolus “pulley”, may contribute to foot eversion in *Didelphis* (Lewis, 1966). Pronator profundus muscle, revealed by microdissection, may account for rotating the crus medially during swing (Narain et al., 2005). Various muscles might therefore be involved in guaranteeing opossum foot eversion at touchdown.

With regard to eversion in bipedal primates like man, the strength of peroneus muscles may be trained, to guarantee stable foot positioning at touchdown. Therefore, after approval by our medical-ethical committee, a group of nine youthful gymnasts without a medical history was trained by warm-up exercises four times weekly intended to reinforce peroneus muscles. After four weeks this group performed the single-leg standing balance on board longer than a control group exercising with non-specific warm-ups. Evertor muscles reinforcement might therefore be used to prevent inversion traumatism.

## **References**

- Narain F. H. M. et al. (2003) Aspects of arthrology in the lower leg of the opossum. *European Journal of Morphology* 41, 1, 68.
- Narain F. H. M. et al. (2009) Human foot inversion prior to toe-off. *Journal of Vibroengineering*, 11, 3, 530-535.
- Lewis O. J. (1966) The phylogeny of the cruropedal extensor musculature, with special reference to the primates. *Journal of Anatomy*, 100, 4, 865-880.
- Narain F. H. M. et al. (2005) Deep muscles in the lower leg of the opossum. *European Journal of Morphology*, 42, 3, 147-148.