Made available by Hasselt University Library in https://documentserver.uhasselt.be

Some New Examples of Nondegenerate Quiver Potentials Non Peer-reviewed author version

de Volcsey, Louis de Thanhoffer & VAN DEN BERGH, Michel (2013) Some New Examples of Nondegenerate Quiver Potentials. In: INTERNATIONAL MATHEMATICS RESEARCH NOTICES (20), p. 4672-4686.

DOI: 10.1093/imrn/rns182 Handle: http://hdl.handle.net/1942/15969

SOME NEW EXAMPLES OF NON-DEGENERATE QUIVER POTENTIALS

L. DE THANHOFFER DE VÖLCSEY AND M. VAN DEN BERGH

ABSTRACT. We prove a technical result which allows us to establish the nondegeneracy of potentials on quivers in some previously unknown cases. Our result applies to McKay quivers and also to potentials derived from geometric helices on Del Pezzo surfaces.

On the other hand we also give an example of a skew group ring with a degenerate potential. This shows that for 3-CY orders Iyama-Reiten mutations cannot always be iterated indefinitely.

1. INTRODUCTION

In [8] Derksen, Weyman and Zelevinsky describe how to "mutate" at a vertex a pair (Q, W) consisting of a quiver Q and a potential $W \in (kQ)^{/}[(kQ)^{,}(kQ)^{]}$. This construction produces a new such pair (Q', W'). The Jacobian (or Ginzburg) algebras of the pairs (Q, W), (Q', W') share many homological properties [7, 9, 10].

One peculiarity of the mutation process is that it is only defined if the vertex is not incident to a loop or two-cycle. Even if all vertices in Q have this property then this is not necessarily the case for Q'. If the property of having no loops or two-cycles persists under iterated mutations then we say that (Q, W) (or W) is non-degenerate.

In this paper we give a technical result (Theorem 3.1) which allows us to establish the non-degeneracy of potentials in some previously unknown cases. Our theorem applies for example to the McKay quiver associated to a group $G \subset SL(V)$ where dim V = 3 and G acts freely on $V^* - \{0\}$ (this implies that G is cyclic). See Corollary 4.1.2 below.

¹⁹⁹¹ Mathematics Subject Classification. 16T30.

Key words and phrases. Quiver potentials, mutations.

The second author is a senior researcher at the FWO.

Let us give an explicit example considered in [9]. Assume that Q is the McKay quiver of the pair $(\mathbb{Z}/5\mathbb{Z}, kx_1 + kx_2 + kx_3)$ where the generator of $\mathbb{Z}/5\mathbb{Z}$ acts diagonally by (ξ, ξ^2, ξ^2) . The quiver Q looks as follows



The potential W is the signed sum of all three-cycles containing x_1, x_2 and x_3 . The Jacobian algebra of (Q, W) is the skew group ring $k[[x_1, x_2, x_3]] \# \mathbb{Z}/5\mathbb{Z}$. Our main theorem implies that W is non-degenerate.

On the other hand if G does not act freely on $V^* - \{0\}$ then the associated potential will generally be degenerate. Consider the McKay quiver associated to the cyclic group $\mathbb{Z}/6\mathbb{Z}$ acting with (ξ^2, ξ^5, ξ^5) .



A mutation at $\overline{0}$ produces a 2-cycle between $\overline{2}$ and $\overline{4}$. See Example 4.1.3 below. Hence this potential is degenerate. Since for 3-CY orders the mutations from [8] coincide with the Iyama-Reiten mutations from [9] (see [7, 10]) this implies that Iyama-Reiten mutations cannot always be iterated indefinitely.

Our theorem also implies that potentials derived from geometric helices on Del Pezzo surfaces (see below) are non-degenerate. In this way we recover part of the main result of [6].

The main technical tool in this paper is the lifting of mutations to the graded setting (see [1]).

Throughout all quivers are finite. k is a base field.

2. Graded and ungraded mutations

Let Q be a quiver. A potential on Q is an element $W \in (kQ)^{\hat{}}([kQ)^{\hat{}}, (kQ)^{\hat{}}]$ containing no paths of length one or zero. We write W as a sum of oriented cycles in Q. The corresponding Jacobian algebra is defined in the usual way

$$\mathcal{P}(Q,W) = (kQ)^{\hat{}} / (\partial_a W)_a$$

The space of paths of length zero in kQ will be denoted by R. This is a semisimple k-algebra.

A potential is said to be reduced if it contains no two-cycles. Such two-cycles lead to relations which express some arrows in terms of others and hence such arrows can be eliminated in the Jacobian algebra. This observation is refined in the splitting theorem [8, Thm 4.6] which asserts that (Q, W) is "right equivalent" (see [8, Def. 4.2]) to a direct sum decomposition

(2.1)
$$(Q, W) \sim (Q^{\text{red}}, W^{\text{red}}) \oplus (Q^{\text{triv}}, W^{\text{triv}})$$

where W^{red} is reduced and W^{triv} contains only two-cycles and its associated Jacobian algebra is equal to R.

The decomposition theorem implies in particular

(2.2)
$$\mathcal{P}(Q, W) \cong \mathcal{P}(Q^{\text{red}}, W^{\text{red}})$$

If G is an abelian group then we say that Q is G-graded if we have assigned a G-degree |a| to all arrows a in Q. If W is homogeneous of degree r the partially completed Jacobian algebra

$$\mathcal{P}^{\mathrm{gr}}(Q,W) = (kQ)^{\mathrm{gr}}/(\partial_a W)_a$$

(where we complete only with respect to sequences of paths having ascending length but constant degree), is naturally a G-graded algebra. It is observed in [1] that the graded analogue of the splitting theorem holds and hence the decomposition (2.1) can be performed on the graded level.

The following lemma is very useful.

Lemma 2.1. Let Q' be obtained from Q by repetively deleting pairs of arrows $i \xrightarrow{a} j \xrightarrow{b} i$ with $i \neq j$ and |a| + |b| = r. If Q' contains a two-cycle then so does Q^{red} .

Proof. By the degree constraint on W we can onl eliminate two-cycles of the form ab with |a| + |b| = r. This implies the result.

Assume that Q does not have loops or two-cycles and let i be a vertex of Q. The mutation $\mu_i(Q, W) = (Q', W')$ at i is defined in [8] as follows.

• For any sequence of arrows $u \xrightarrow{a} i \xrightarrow{b} v$ we add an arrow $u \xrightarrow{[ba]} v$.

• All arrows a starting or ending in i are replaced with opposite arrows a^* . We have

$$W' = [W] + \sum b^* a^*[ab]$$

where [W] is obtained from W by replacing all compositions ba through the vertex i by the new arrows [ba]. Following [8] we put $\tilde{\mu}_i(Q, W) = (Q'^{\text{red}}, W'^{\text{red}})$. The operations $\tilde{\mu}_i$ is an operation on right equivalence classes of quivers with potential (see [8, Thm 5.2]).

Now assume that Q is in addition G-graded and that W is homogeneous of degree r. Then we make Q' into a G-graded quiver by fixing the degrees of the arrows as follows [1]

• Arrows which are both in Q and Q' do not change degrees.

- |[ab]| = |a| + |b|.
- $|a^*| = -|a| + r$ if a ends in i.
- $|a^*| = -|a|$ if a starts in *i*.

With this grading W' is homogeneous of degree r. If we perform mutation on the graded level we write $\mu_i^{\text{gr}}(Q, W) = (Q', W')$ and $\tilde{\mu}_i^{\text{gr}}(Q, W) = (Q'^{\text{red}}, W'^{\text{red}})$. Such graded mutations are compatible with forgetting the grading.

3. Main theorem

Let $S = \bigoplus_{n \ge 0} S_n$ be a commutative noetherian \mathbb{Z} -graded ring and let Λ be a graded S-order (i.e. Λ is a graded S-algebra, finite as a module over S). We will say that Λ/S is almost Azumaya if for all $P \in \operatorname{Spec} S$ such that $P \not\supseteq S_{\ge 1}$ the algebra Λ_P is Azumaya over S_P . This implies in particular that for any non-zero idempotent $e \in \Lambda_0$ the quotient $\Lambda/\Lambda e \Lambda$ is a finite S_0 -module. The latter property is in fact the only one we will use below.

Theorem 3.1. Assume the following assumptions hold

- (1) (Q, W) is a \mathbb{Z} -graded connected quiver with a reduced homogeneous potential W of degree r.
- (2) Q has at least three vertices.
- (3) Put $\Lambda = \mathcal{P}^{\mathrm{gr}}(Q, W)$. Then dim $\Lambda_i < \infty$ for all i and dim $\Lambda_i = 0$ for $i \ll 0$.
- (4) Λ is a 3-dimensional CY-order over a noetherian center which is almost Azumaya.
- (5) $\operatorname{HH}_0(\Lambda) = \Lambda/[\Lambda, \Lambda]$ contains no homogeneous elements with degree lying in the interval [1, r/2].

Then Q has no loops or two-cycles.

Proof. Assume first that Q contains a loop $i \xrightarrow{a} i$ and let e be the sum of the idempotents corresponding to the vertices different from i. Let Q_0 be the subquiver of Q consisting of all loops at i. Put $\overline{\Gamma} = \Gamma/\Gamma e\Gamma$. Then $\overline{\Gamma}$ is the graded Jacobian algebra $\mathcal{P}^{\mathrm{gr}}(Q_0, W_0)$ where W_0 is obtained from \overline{W} by dropping all cycles passing through vertices other than i.

If $W_0 = 0$ then $\overline{\Gamma}$ is infinite dimensional which is impossible by the discussion preceding the theorem. Hence W_0 is a sum of terms of degree r which are products of loops

with $n \geq 3$ (since W_0 is reduced). Assume that (3.1) is the shortest such term. Then there must be some a_i which has degree $\leq r/3 \leq r/2$. As the paths occurring in the relations of $\Gamma/\Gamma e\Gamma$ are products of at least two arrows we have $a_i \neq 0$. Furthermore it is also clear that a_i is not in the image of $[kQ_0, kQ_0]$. Thus a_i represents a non-zero element of $HH_0(\bar{\Gamma})$. Since there is a surjection $HH_0(\Gamma) \to HH_0(\bar{\Gamma})$, we obtain a non-trivial element of $HH_0(\Gamma)$ as well. Since $|a_i| \leq r/2$ we obtain from the hypotheses that $|a_i| \leq 0$. But then a_i must be nilpotent and hence by [13] a_i is a sum of commutators, which is a contradiction.

We now assume that Q has no loops. The proof in the case of a two-cycle $i \to j \to i$ is similar. Let Q_0 be the subquiver of Q consisting of all arrows between i and j and vice versa and let e be the sum of the idempotents corresponding to the vertices different from i or j. Put $\overline{\Gamma} = \Gamma/\Gamma e\Gamma$. Then $\overline{\Gamma} = \mathcal{P}^{\mathrm{gr}}(Q_0, W_0)$ where

 W_0 is obtained from W by dropping all cycles passing through vertices other than i or j.

The case $W_0 = 0$ is again impossible. Hence W_0 is a sum of terms of degree r which are products of two-cycles

$$a_1b_1a_2b_2\cdots a_nb_n$$

with $i \xrightarrow{a_l} j$ and $j \xrightarrow{b_m} i$ with $n \ge 2$. As above this leads to a contradiction. \Box

Proposition 3.2. Let (Q, W) be as in Theorem 3.1 and let *i* be a vertex of *Q*. Then up to right equivalence we may assume that $\tilde{\mu}_i(Q, W) = (Q', W')$ also satisfies the conditions of Theorem 3.1. In particular Q' has no loops or two-cycles.

Proof. Since ungraded and graded mutations are compatible and since the decomposition (2.1) can be performed at the graded level, we can up to right equivalence, assume that Q' is graded and W' is homogeneous.

It has been proved in [7, Thm 5.2] and in [10, §6] that for Jacobian algebras which are Calabi-Yau of dimension three, Derksen-Weyman-Zelevinsky mutations correspond to Iyama-Reiten mutations. With some work these proofs can be carried over to the graded context. Hence in particular we obtain that $\Gamma = \mathcal{P}^{\text{gr}}(Q', W')$ is 3-CY. Furthermore $Z(\Gamma) = Z(\Lambda)$, Γ is almost Azumaya over its center and Γ is derived equivalent to Λ [9]. Finally by the derived invariance of Hochschild homology we have an isomorphism of graded vector spaces

 $\Lambda/[\Lambda,\Lambda] = \mathrm{HH}_0(\Lambda) = \mathrm{HH}_0(\Gamma) = \Gamma/[\Gamma,\Gamma] \quad \Box$

Corollary 3.3. If (Q, W) is as in Theorem 3.1 then (Q, W) is non-degenerate.

Proof. This is clear from Proposition 3.2.

4. Examples and counterexamples

4.1. Skew group rings. In this section we assume that k is algebraically closed of characteristic zero. Let V be a three dimensional vector space and let G be a finite subgroup of SL(V) of order n. Then it is well-known that the skew group algebra $\Lambda = SV \# G$ is 3-Calabi-Yau. Furthermore Λ is graded Morita equivalent to a graded Jacobian algebra $\mathcal{P}^{gr}(Q, W)$ where Q is the McKay quiver of (G, V) and W has degree three. The McKay quiver is the quiver with vertices the irreducible representations $(V_i)_i$ of G and arrows $i \to j$ with multiplicity p if V_j occurs p times in $V \otimes V_i$. We refer to [3, Thm 3.2] for a detailed discussion on how to construct the potential.

In the case of a cyclic group $G \cong \mathbb{Z}/n\mathbb{Z}$, the situation is particularly simple. The irreducible representations of G are given by characters. After the choice of a primitive *n*-th root of unity ξ a character χ may be identified with an element \bar{a} of $\mathbb{Z}/n\mathbb{Z}$ through the rule $\chi_{\bar{a}}(\bar{m}) = \xi^{am}$.

Let \bar{a}_1 , \bar{a}_2 , \bar{a}_3 be the weights of V. The fact that $G \subset SL(V)$ is equivalent to $\sum_i \bar{a}_i = \bar{0}$. The vertices of the McKay quiver of (G, V) are indexed by the elements of $\mathbb{Z}/n\mathbb{Z}$. If $\bar{l} \in \mathbb{Z}/n\mathbb{Z}$ then there are three arrows x_1, x_2, x_3 leaving \bar{l} and ending respectively in $\bar{l} + \bar{a}_1$, $\bar{l} + \bar{a}_2$, $\bar{l} + \bar{a}_3$. The potential is given by

(4.1)
$$W = \sum \pm x_{i_1} x_{i_2} x_{i_3}$$

where the sum runs over all three-cycles such that $\{i_1, i_2, i_3\} = \{1, 2, 3\}$ and the sign is positive iff (i_1, i_2, i_3) is equal to (1, 2, 3), up to cyclic permutation.

Lemma 4.1.1. The hypotheses of Theorem 3.1 hold for (Q, W) if and only if $G \cong \mathbb{Z}/n\mathbb{Z}$ is cyclic and acts with weights $(\bar{a}_i)_i$ on V such that $gcd(n, a_i) = 1$.

Proof. Assume that the hypotheses for Theorem 3.1 hold. It is well-known that Λ is almost Azumaya over its center if and only if SV^G has an isolated singularity which is equivalent to G acting freely on $V^* - \{0\}$ and hence that G is cyclic (see [12]). The fact that G acts freely is equivalent to the stated condition on weights.

Conversely assume that G is cyclic and acts with weights relatively prime to n. We verify the hypotheses of Theorem 3.1: we have already mentioned (1). (2) follows from the fact that the conditions imply $G \neq \mathbb{Z}/2\mathbb{Z}$. (3) is clear. (4) has been mentioned previously. It remains to check (5). We must show $\text{HH}(\Lambda)_1 = 0$. If $H(\Lambda)_1 \neq 0$ then Q must have a loop. This implies that one of the weights is zero and thus not relatively prime to n. Contradiction!

We obtain:

Corollary 4.1.2. Assume that Q is the McKay quiver of a cyclic group $\mathbb{Z}/n\mathbb{Z}$ acting with weights $\bar{a}_1, \bar{a}_2, \bar{a}_3$ such that $gcd(a_i, n) = 1$ and $\sum_i \bar{a}_i = 0$. Let W be the potential (4.1). Then W is non-degenerate.

In [9] Iyama and Reiten consider the case where $\mathbb{Z}/5\mathbb{Z}$ acts with weights $(\overline{1}, \overline{2}, \overline{2})$ and mention that they do not know if the mutations of the associated McKay quiver do not create loops or two-cycles. Corollary 4.1.2 establishes that this does indeed not happen.

On the other hand the hypotheses for Corollary 4.1.2 can not be weakened as shown by the following example:

Example 4.1.3. Let $G = \mathbb{Z}/6\mathbb{Z}$ and assume that G acts with weights $(\overline{2}, \overline{5}, \overline{5})$. Since 2|6 the hypotheses for Corollary 4.1.2 do not hold.

The McKay quiver corresponding to this case has been given in the introduction. We note that the McKay quiver of a cyclic groups is naturally \mathbb{Z}^3 -graded. It is convenient to use the grading by the monomials in x_1, x_2, x_3 such that $|x_i| = x_i$. Then W is homogeneous of degree $x_1x_2x_3$. With these conventions we see that performing a mutation at the vertex $\overline{0}$ results in a two-cycle $[x_1x_1]x_1$ between vertices $\overline{2}$ and $\overline{4}$. This two-cycle has degree x_1^3 and cannot be eliminated for degree reasons (see Lemma 2.1).

4.2. **Del Pezzo surfaces.** Let Y be a Del-Pezzo surface and let $(E_i)_{i=1,...,n}$ be a full exceptional collection on Y. Put $\mathbb{E} = \bigoplus_{i=1}^{n} E_i$ and define $A(\mathbb{E}) = \text{End}(\mathbb{E})$. Assume that

$$\mathbb{H}(E) = (E_i)_{i \in \mathbb{Z}} = (\dots, \omega_Y \otimes E_n, E_1, \dots, E_n, \omega_Y^{-1} \otimes, E_1, \dots)$$

is a geometric helix (see [4, 6]). I.e. every slice $(E_{i+1}, \ldots, E_{i+n})$ is an exceptional collection and furthermore

 $\forall i < j, \forall k > 0 : \operatorname{Ext}^k(E_i, E_j) = 0$

The rolled up helix algebra is the Z-graded ring

$$B(\mathbb{H}) = \bigoplus_{k \in \mathbb{Z}} \operatorname{Hom}(\mathbb{E}, \omega_Y^{-k} \otimes \mathbb{E})$$

with obvious multiplication.

Theorem 4.2.1. $B(\mathbb{H})$ is an N-graded Jacobian algebra derived from a graded super potential of degree one which is Calabi-Yau of dimension three (as ungraded algebra) and which is almost Azumaya over its center.

This result is folklore although we don't know a good reference. For the convenience of the reader give a proof in Appendix A using some results from [11]. The fact that $B(\mathbb{H})$ is Calabi-Yau of dimension dimension three is [6, Thm 3.6]. The fact that $B(\mathbb{H})$ is Jacobian does not follow immediately from [2, Thm 3.1] as the results in loc. cit. are for algebras of the form kQ/(R) where Q is a quiver with arrows of degree one.

Theorem 4.2.2. Let (Q, W) be such that W is reduced and $\mathcal{P}^{\mathrm{gr}}(Q, W) = B(\mathbb{H})$. Then (Q, W) satisfies the hypotheses of Theorem 3.1.

In particular it follows from Corollary 3.3 that (Q, W) is non-degenerate. This is a consequence of [6, Thm 1.7]. In loc. cit. the authors deduce this fact from their result that a mutation of $B(\mathbb{H})$ is always of the form $B(\mathbb{H}')$ for another geometric helix \mathbb{H}' .

Proof of Theorem 4.2.2. (1) is true with r = 1. (2) is true since $\operatorname{rk} K_0(\operatorname{coh}(Y)) \geq 3$. (3) is obvious. (4) follows from Theorem 4.2.1. (5) is vacuous since the potential has degree one.

APPENDIX A. POTENTIALS ASSOCIATED TO DEL PEZZO SURFACES

Let the notations be as in §4.2. In this appendix we will prove Theorem 4.2.1. We keep the setting of the theorem although with trivial changes most of the proof applies to arbitrary Fano varieties. Our arguments are related to those appearing in recent papers by Minamoto and Mori [14, 15].

Lemma A.1. $B(\mathbb{H})$ is the derived tensor algebra of $B(\mathbb{H})_1$ over $B(\mathbb{H})_0 = A(\mathbb{E})$.

Proof. For k < 0 we have

$$\operatorname{Hom}(\mathbb{E}, \omega_V^{-k} \otimes \mathbb{E}) = \operatorname{Ext}^2(\mathbb{E}, \omega_V^{k+1} \otimes \mathbb{E})^* = 0$$

For $k \geq 0$ we have

$$\operatorname{Hom}(\mathbb{E}, \omega_Y^{-k} \otimes \mathbb{E}) = R \operatorname{Hom}(\mathbb{E}, \omega_Y^{-k} \otimes \mathbb{E})$$

For $k, l \ge 0$ we have to prove that the canonical map

$$\operatorname{RHom}(\mathbb{E}, \omega_Y^{-k} \overset{L}{\otimes} \mathbb{E}) \overset{L}{\otimes}_{A(\mathbb{E})} \operatorname{RHom}(\mathbb{E}, \omega_Y^{-l} \otimes \mathbb{E}) \to \operatorname{RHom}(\mathbb{E}, \omega_Y^{-k-l} \otimes \mathbb{E})$$

is an isomorphism

One verifies that this is equivalent to showing that the composition map

$$\operatorname{RHom}(\mathbb{E}, \omega_Y^{-k} \overset{L}{\otimes} \mathbb{E}) \overset{L}{\otimes}_{A(\mathbb{E})} \operatorname{RHom}(\omega_Y^l \otimes \mathbb{E}, \mathbb{E}) \to \operatorname{RHom}(\omega_Y^l \otimes \mathbb{E}, \omega_Y^{-k} \otimes \mathbb{E})$$

is an isomorphism.

Since \mathbb{E} is a classical generator for $D^b(\operatorname{coh}(\mathcal{O}_Y))$ it suffices to prove the result with $\omega_Y^{-k} \otimes \mathbb{E}$ and $\omega_Y^l \otimes \mathbb{E}$ replaced by \mathbb{E} . Then the statement is obvious. \Box

Lemma A.2. Consider $A(\mathbb{E}) \otimes A(\mathbb{E})$ as $A(\mathbb{E})$ -bimodule with the outer bimodule structure. We have an equality of $A(\mathbb{E})$ -bimodules

$$B(\mathbb{H})_1 = \operatorname{RHom}_{A(\mathbb{E})^e}(A(\mathbb{E}), A(\mathbb{E}) \otimes A(\mathbb{E}))[2]$$

where on the righthandside the $A(\mathbb{E})$ -bimodule structure comes from the surviving inner bimodule structure on $A(\mathbb{E}) \otimes A(\mathbb{E})$. In other words $B(\mathbb{H})_1$ is a shifted inverse dualizing complex of $A(\mathbb{E})$ ([11, §3.3]).

Proof. We have to compute

 $\operatorname{RHom}_{\operatorname{End}(\mathbb{E})^e}(\operatorname{End}(\mathbb{E}), \operatorname{End}(\mathbb{E}) \otimes \operatorname{End}(\mathbb{E}))$

Working on $Y \times Y$ we have

$$\operatorname{End}(\mathbb{E})^e \cong \operatorname{End}(\mathbb{E} \boxtimes \mathbb{E}^*)$$

It is easy to see that $\mathbb{E}\boxtimes\mathbb{E}^*$ is a classical generator for $D^b(\operatorname{coh}(Y \times Y))$ (e.g. because it is derived from an exceptional collection, or else invoke [5]). Under the ensuing derived equivalence $D^b(\operatorname{End}(\mathbb{E}\boxtimes\mathbb{E}^*)) \cong D^b(\operatorname{coh}(Y \times Y))$ it is easy to see that $\operatorname{End}(\mathbb{E})$ corresponds to \mathcal{O}_Y and $\operatorname{End}(\mathbb{E}) \otimes \operatorname{End}(\mathbb{E})$ to $\mathbb{E}\boxtimes\mathbb{E}^*$.

We now have to compute

$$\operatorname{RHom}_{Y \times Y}(\mathcal{O}_{\Delta}, \mathbb{E} \boxtimes \mathbb{E}^*)$$

where Δ is the diagonal. We use the well-known formula

$$\mathcal{RHom}_{\mathcal{O}_{Y\times Y}}(\mathcal{O}_{\Delta},\mathcal{O}_{Y\times Y}) = \omega_{\Delta}^{-1}[-2$$

Thus

$$\operatorname{RHom}_{Y \times Y}(\mathcal{O}_{\Delta}, \mathbb{E} \boxtimes \mathbb{E}^{*}) = R\Gamma(Y \times Y, \operatorname{R}\mathcal{H}om_{Y \times Y}(\mathcal{O}_{\Delta}, \mathcal{O}_{Y \times Y}) \overset{L}{\otimes}_{Y \times Y} \mathbb{E} \boxtimes \mathbb{E}^{*})$$
$$= R\Gamma(Y \times Y, \omega_{\Delta}^{-1} \overset{L}{\otimes}_{Y \times Y} \mathbb{E} \boxtimes \mathbb{E}^{*})[-2]$$
$$= \operatorname{RHom}_{Y}(\mathbb{E}, \omega_{Y}^{-1} \otimes \mathbb{E})[-2] \quad \Box$$

Lemma A.3. Assume that B is an N-graded ring such that dim $B_i < \infty$ for all i. Then

$$\operatorname{gl}\dim B_0 \leq \operatorname{gl}\dim B$$

If the category of graded finite length B-modules is Ext-finite and has a Serre functor given by ?(d)[n] for $d \neq 0$ then

$$\operatorname{gl}\dim B_0 < \operatorname{gl}\dim B = n$$

Proof. Let S, T be simple B_0 -modules which we view as B-modules concentrated in degree zero. The part of degree zero of a graded projective B-resolution of S is a projective B_0 -resolution of S. This implies the first inequality. We also get

$$\operatorname{Ext}^{i}_{B_{0}}(S,T) = \operatorname{Ext}^{i}_{B}(S,T)_{0}$$

Assume now that the category of graded finite length *B*-modules is Ext-finite and has a Serre functor given by ?(d)[n] for $d \neq 0$. Then it is standard that gl dim B = n. For the second inequality we note that $\operatorname{Ext}_B^n(S,T)_0 = \operatorname{Hom}_B(T,S(d))_0^* = 0$. \Box

Proof of Theorem 4.2.1. Since $A(\mathbb{E})$ is finite dimensional and has finite global dimension the same is true for $A(\mathbb{E})^e$. Hence $A(\mathbb{E})$ is perfect as a bimodule and as such homologically smooth.

By Lemmas A.1,A.2 we obtain that $B(\mathbb{H})$ is a (graded) 3-Calabi-Yau completion of $A(\mathbb{E})$ (see [11, §4]). By [11, Thm 4.8] we obtain that $B(\mathbb{H})$ is 3-Calabi-Yau and the proof shows that the Serre functor is given by ?(-1). By Lemma A.3 we then find that gl dim $A(\mathbb{E}) \leq 2$. We may then use [11, Thm 6.10] to obtain that $B(\mathbb{H})$ is quasi-isomorphic to a Ginzburg algebra derived from a super potential of degree one. Since $B(\mathbb{H})$ is concentrated in degree zero it is also a Jacobian algebra.

References

- C. Amiot and S. Oppermann, Cluster equivalence and graded derived equivalence, arXiv:1003.4916.
- [2] R. Bocklandt, Graded Calabi Yau algebras of dimension 3, J. Pure Appl. Algebra 212 (2008), no. 1, 14–32.
- [3] R. Bocklandt, T. Schedler, and M. Wemyss, Superpotentials and higher order derivations, arXiv:0802.0162.
- [4] A. Bondal and A. Polishchuk, Homological properties of associative algebras: the method of helices, Russian Acad. Sci. Izv. Math 42 (1994), 219–260.
- [5] A. Bondal and M. Van den Bergh, Generators and representability of functors in commutative and noncommutative geometry, Mosc. Math. J. 3 (2003), no. 1, 1–36, 258.
- [6] T. Bridgeland and D. Stern, Helices on del Pezzo surfaces and tilting Calabi-Yau algebras, arXiv:0909.173.
- [7] A. B. Buan, O. Iyama, I. Reiten, and D. Smith, Mutation of cluster-tilting objects and potentials, arXiv:0804.3813.
- [8] H. Derksen, J. Weyman, and A. Zelevinsky, Quivers with potentials and their representations. I. Mutations, Selecta Math. (N.S.) 14 (2008), no. 1, 59–119.
- [9] O. Iyama and I. Reiten, Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras, Amer. J. Math. 130 (2008), no. 4, 1087–1149.
- [10] B. Keller and D. Yang, Derived equivalences from mutations of quivers with potential, arXiv:0906.0761.
- [11] B. Keller, Deformed Calabi-Yau completions, arXiv:0908.349.
- [12] K. Kurano and S. Nishi, Gorenstein isolated quotient singularities of odd prime dimension are cyclic, arXiv:0903.3270v1.
- [13] H. Lenzing, Nilpotente Elemente in Ringen von endlicher globaler Dimension, Math. Z. 108 (1969), 313–324.
- [14] H. Minamoto, Ampleness of two-sided tilting complexes, to appear.
- [15] H. Minamoto and I. Mori, Structures of AS-regular algebra, to appear.

E-mail address: louicious@mac.com, michel.vandenbergh@uhasselt.be

DEPARTEMENT WNI, UNIVERSITEIT HASSELT, 3590 DIEPENBEEK, BELGIUM.