

REGULAR GRAPHS AND THE SPECTRA OF TWO-VARIABLE LOGIC WITH COUNTING*

ERYK KOPCZYŃSKI[†] AND TONY TAN[‡]

Abstract. The *spectrum* of a first-order logic sentence is the set of natural numbers that are cardinalities of its finite models. In this paper we show that when restricted to using only two variables, but allowing counting quantifiers, the class of spectra of first-order logic sentences is exactly the class of semilinear sets, and hence, closed under complement. At the heart of our proof are semilinear characterisations for the existence of regular and biregular graphs, the class of graphs in which there are a priori bounds on the degrees of the vertices. Our proof also provides a simple characterisation of models of two-variable logic with counting – that is, up to renaming and extending the relation names, they are simply a collection of regular and biregular graphs.

Key words. two-variable logic with counting, first-order spectra, regular graphs, semi-linear, Presburger arithmetic

AMS subject classifications. 15A15, 15A09, 15A23

1. Introduction. The spectrum of a first-order sentence ϕ , denoted by $\text{SPEC}(\phi)$, is the set of natural numbers that are cardinalities of finite models of ϕ . Or, more formally, $\text{SPEC}(\phi) = \{n \mid \text{there is a model of } \phi \text{ with universe of cardinality } n\}$. A set is a *spectrum*, if it is the spectrum of a first-order sentence.

In this paper we consider the logic \mathcal{C}^2 , the class of first-order sentences using only two variables and allowing counting quantifiers $\exists^k z \phi(z)$, where $k \geq 1$. Semantically $\exists^k z \phi(z)$ means there exist at least k number of z 's such that $\phi(z)$ holds. We prove that the spectra of \mathcal{C}^2 are precisely semilinear sets. In fact, our proof also shows that the family of models of a \mathcal{C}^2 formula can be viewed as a collection of regular graphs.

Related works. The notion of spectrum was introduced by Scholz [36] where he also asked whether there exists a necessary and sufficient condition for a set to be a spectrum. Since its publication, Scholz's question and many of its variants have been investigated by many researchers for the past 60 years. One of the arguably main open problems in this area is the one asked by Asser [1], formally known as *Asser's conjecture*, whether the complement of a spectrum is also a spectrum.

The notion of spectrum has a deep connection with complexity theory as shown by Jones and Selman [22], as well as Fagin [5] independently that a set of integers is a spectrum if and only if its binary representation is in NE. Hence, Asser's conjecture is equivalent to asking whether $\text{NE} = \text{co-NE}$. It also immediately implies that if Asser's conjecture is false, i.e., there is a spectrum whose complement is not a spectrum, then $\text{NP} \neq \text{co-NP}$, hence $\text{P} \neq \text{NP}$. The converse implication is still open. An interesting result by Woods [40] states that if spectra are precisely rudimentary sets, then $\text{NE} = \text{co-NE}$ and $\text{NP} \neq \text{co-NP}$.* There are a number of interesting connections between spectrum and various models of computation such as RAM as well as intrinsic computational behavior [9, 10, 11, 28, 35]. We refer the reader to a

*The same manuscript has been published in ArXiv and available online in <http://arxiv.org/abs/1304.0829>.

[†]University of Warsaw, erykk@mimuw.edu.pl

[‡]Hasselt University and Transnational University of Limburg, ptony.tan@gmail.com

*It should be noted that the class of rudimentary sets corresponds precisely to *linear time hierarchy* – the linear time analog of polynomial time hierarchy [41].

recent survey by Durand, Jones, Makowsky and More [3] for a more comprehensive treatment on the spectra problem and its history.

The logic C^2 is not the first logic known to have semilinear spectra. A well known Parikh theorem states the spectra of context-free languages are semilinear, and closed under complementation. Using the celebrated composition method, Gurevich and Shelah [17] showed that the spectra of monadic second order logic with *one unary function* are semilinear. Fischer and Makowsky [7] showed that the *many-sorted spectra* of the monadic second-order logic with *modulo counting* over structures with *bounded tree-width* are semilinear.[†]

These two results are orthogonal to ours here. Structures expressible in C^2 do not have bounded tree-width. An example is d -regular graphs for $d \geq 3$. Hence, it is very unlikely that one can apply some sort of “pumping” or automata theoretic argument as used by Fischer and Makowsky [7] to obtain the semilinearity of C^2 spectra. Moreover, it should also be noted that in C^2 one can express a few unary functions, hence our result does not follow from the result of Gurevich and Shelah [17], and neither theirs from ours since we are restricted to using only two variables. As far as we know, C^2 is the first logic known to have its spectra closed under complement without any restriction on the vocabulary nor in the interpretation.

The result closest to ours is the one by Grandjean [11] where he considers the spectra of first-order sentences using only one variable. A similar result due to Grohe [3] says that for every Turing machine M , there exists a first-order sentence ϕ_M using only *three variables* such that $\text{SPEC}(\phi_M) = \{t^2 \mid t \text{ is the length of an accepting run of } M\}$.

To end our study of related work, we should mention that the two-variable logic and many of its variants have been extensively studied, with the focus mainly being on the satisfiability problem [31, 39, 32, 15, 14, 12, 19, 33, 26, 37, 34]. Some other works by van den Broeck, Meert and Darwiche [2] consider the problem of counting the number of models of a given size on two-variable logic.

Sketch of our proof. Consider the following instances of structures expressible in C^2 .[‡]

(Ex.1) (c, d) -biregular graphs: the bipartite graphs on the vertices $U \cup V$, where the degree of each vertex in U and V is c and d , respectively.

(Ex.2) (c, d) -regular digraphs: the directed graphs in which the in-degree and the out-degree of each vertex are c and d , respectively.

An observation from basic graph theory tells us that for “big enough” M and N ,[§]

(C1) there is a (c, d) -biregular graph in which M vertices are of degree c and N vertices are of degree d if and only if $Mc = Nd$;

(C2) there is a (c, d) -regular digraph of N vertices if and only if $Nc = Nd$, and hence, $c = d$.

These characterisations immediately imply that the spectra of the C^2 sentences that describe (c, d) -biregular graphs and (c, d) -regular digraphs are linear sets. It is from these observations that we draw our inspiration to prove the semilinearity of the spectra of C^2 .

[†]Intuitively, the many-sorted spectrum of a formula is the spectrum that not only counts the sizes of the models, but also the cardinality of the unary predicates in the models of the formula.

[‡]Though the result in this paper holds for arbitrary structures, it helps to assume that the structures of C^2 are graphs in which the vertices and the edges are labelled with a finite number of colours.

[§]“Big enough” means M and N are greater than a constant K which depends only on c and d .

More precisely, we show that given a \mathcal{C}^2 sentence φ , one can construct a *Presburger* formula ψ that expresses precisely the spectrum of φ . Presburger formulas are first-order formulas with the relation symbols $+$ and \leq and constants 0 and 1 interpreted over the domain \mathbb{N} in the natural way. It is shown by Ginsburg and Spanier in [8] that Presburger formulas express precisely the class of semilinear sets. That is, if $\psi(\bar{X})$ is a Presburger formula with free variables $\bar{X} = (X_1, \dots, X_k)$, the set of the solutions of $\psi(\bar{X})$, i.e. \bar{N} such that $\psi(\bar{N})$ holds, is semilinear.

The crux of our construction of the Presburger formula is a generalisation of the characterisations (C1) and (C2) above to the following setting. Let \mathcal{C} be a set of ℓ colors, denoted by $\text{col}_1, \text{col}_2, \dots, \text{col}_\ell$, and let C and D be $(\ell \times m)$ - and $(\ell \times n)$ -matrices whose entries are all non-negative integers. We say that a bipartite graph $G = (U, V, E)$ is (C, D) -biregular, if we can color its edges with colors from \mathcal{C} such that there are a partition $U_1 \cup \dots \cup U_m$ of U and a partition $V_1 \cup \dots \cup V_n$ of V where

- for every integer $1 \leq i \leq m$, for every vertex $u \in U_i$, for every $1 \leq j \leq \ell$, the number of edges with color col_j adjacent to u is *precisely* $C_{j,i}$; and
- for every integer $1 \leq i \leq n$, for every vertex $v \in V_i$, for every $1 \leq j \leq \ell$, the number of edges with color col_j adjacent to v is *precisely* $D_{j,i}$.

Our setting also allows us to say that the number of edges with color col_j adjacent to v is *at least* $D_{j,i}$. In Theorem 5.1 we effectively construct a Presburger formula that characterises the set $\{N \mid \text{there is a } (C, D)\text{-biregular graph of } N \text{ vertices}\}$.

In a similar manner, we can define (C, D) -regular digraphs, where C and D control the number of incoming and outgoing edges of each vertex, respectively. Likewise, we obtain a similar Presburger formula that characterises the set $\{N \mid \text{there is a } (C, D)\text{-regular digraph of } N \text{ vertices}\}$.[¶] We then proceed to observe that the relations in every model of a \mathcal{C}^2 formula can be partitioned in such a way that every part forms a (C, D) -regular digraph, and every two parts a (C, D) -biregular graph. In a sense this shows that each model of a \mathcal{C}^2 sentence is simply a collection of regular graphs. Applying the Presburger formula that characterises the existence of these regular graphs, we obtain the semilinearity of the spectra of \mathcal{C}^2 sentences.

For the converse direction, it is not that difficult to show that every semilinear set is a spectrum of a \mathcal{C}^2 sentence. Since semilinear sets are closed under complement, this establishes the fact that the class of spectra of \mathcal{C}^2 sentences are closed under complement. It can also be deduced immediately from our proof that the many-sorted spectra of \mathcal{C}^2 are also semilinear. Moreover, our result extends trivially to the class $\exists\text{SOC}^2$, the class of sentences of the form: $\exists R_1 \dots \exists R_m \phi$, where R_1, \dots, R_m are second-order variables and ϕ is a \mathcal{C}^2 formula. We simply regard R_1, \dots, R_m as part of the signature.

Outline of the paper. This paper is organised as follows. In Section 2 we review the logic \mathcal{C}^2 and state our main result: Theorem 2.1 which states that every \mathcal{C}^2 spectrum can be expressed as the set of solutions of an effectively constructed Presburger formula. The proof of Theorem 2.1 is rather complex. So we present its outline in Section 3, before its details in Sections 4–8. Finally we conclude with a few observations and future directions in Section 9.

[¶]Closely related to our result is the work by S. L. Hakimi [18] which deals with the question: given a vector (d_1, \dots, d_m) , is there a graph with vertices v_1, \dots, v_m whose degrees are precisely d_1, \dots, d_m , respectively? Another related result concerns the notion of score sequence obtained by H. G. Landau [23] which deals with the question: given a vector (d_1, \dots, d_m) , is there a tournament with vertices v_1, \dots, v_m whose outdegrees are precisely d_1, \dots, d_m , respectively? These questions are evidently different from our characterisations provided in Section 5.

2. The logic \mathcal{C}^2 . We fix $\mathcal{P} = \{P_1, P_2, \dots\}$ to be the set of predicate symbols of arity 1; and $\mathcal{R} = \{R_1, R_2, \dots\}$ the set of predicate symbols of arity 2. Two-variable logic with counting, denoted by \mathcal{C}^2 , is defined by the following syntax.

$$\phi ::= z = z \mid R(z, z) \mid P(z) \mid \neg\phi \mid \phi \wedge \phi \mid \exists^k z \phi,$$

where the variable z ranges over x, y , and the symbols R and P over \mathcal{R} and \mathcal{P} , respectively.

The quantifier $\exists^k z \phi$ means *there are at least k elements z such that ϕ holds*. Note that $\exists^1 z \phi$ is the standard $\exists z \phi$, and $\forall z \phi$ is equivalent to $\neg\exists^1 z \neg\phi$. By default, we assume that $\exists^0 z \phi$ always holds.

As usual, we write $\mathfrak{A} \models \phi$ to denote that the structure \mathfrak{A} is a model of ϕ and $\text{SPEC}(\phi)$ to denote the spectrum of ϕ . Theorem 2.1 below is the main result in this paper. Its proof spans over Sections 4–8.

THEOREM 2.1. *For every $\phi \in \mathcal{C}^2$, there exists a Presburger formula $\text{PREB}(x)$ such that $\text{SPEC}(\phi) = \{n \mid \text{PREB}(n) \text{ holds}\}$. Moreover, the formula $\text{PREB}(x)$ can be constructed effectively.*

We should remark that Theorem 2.1 also holds for arbitrary vocabulary. Since \mathcal{C}^2 uses only two variables, relations of greater arity can be viewed as unary or binary relations [13].

An immediate consequence of Theorem 2.1 is that the spectrum of a \mathcal{C}^2 sentence is a semilinear set.

COROLLARY 2.2. *For every sentence $\phi \in \mathcal{C}^2$, the spectrum $\text{SPEC}(\phi)$ is semilinear.*

On the other hand, it is not that difficult to show that every semilinear set is a spectrum of a \mathcal{C}^2 sentence, as formally stated below.

PROPOSITION 2.3. *For every semilinear set $\Lambda \subseteq \mathbb{N}$, there exists a sentence $\phi \in \mathcal{C}^2$ such that $\text{SPEC}(\phi) = \Lambda$.*

Proof. For a linear set $\Gamma_{m,n} = \{m + in \mid i = 0, 1, 2, \dots\}$, consider the vocabulary $\tau_{m,n} = \{A, B_0, B_1, \dots, B_{n-1}, E\}$, where A, B_0, \dots, B_{n-1} are unary and E binary. Consider the \mathcal{C}^2 sentence $\phi_{m,n}$ which states that $A \cup B_0 \cup \dots \cup B_{n-1}$ partition the whole universe, the predicate A contains exactly m elements, and for every x , if $B_i(x)$ holds,

- there is exactly one y such that $x \neq y$ and $B_{i+1 \bmod n}(y)$ and $E(x, y)$ hold;
- there is exactly one y such that $x \neq y$ and $B_{i-1 \bmod n}(y)$ and $E(y, x)$ hold.

It is straightforward that $\text{SPEC}(\phi_{m,n}) = \Gamma_{m,n}$. For a semilinear set, we simply takes the finite disjunction of such $\phi_{m,n}$'s. This completes our proof of Proposition 2.3. \square

Now, take Corollary 2.2, apply the fact that the class of semilinear sets is closed under complement, and then Proposition 2.3, we obtain the following corollary.

COROLLARY 2.4. *The class of spectra of \mathcal{C}^2 sentences is closed under complement.*

3. The plan for the proof of Theorem 2.1. As mentioned earlier, the proof of Theorem 2.1 is rather complex and spans over Sections 4–8. We give its outline here.

- In Section 4 we define the logic QMLC (Quantified Modal Logic with Counting), which for our purpose, will be easier to work with. In particular, we will show that \mathcal{C}^2 and QMLC are equivalent in terms of spectra.
- In Section 5 we define the class of biregular graphs and regular digraphs. The main theorems in this section are Theorems 5.2 and 5.4. Theorem 5.2 gives




FIG. 3.1. The skeleton for the proof of Theorem 2.1. Theorem 5.2 gives us the Presburger characterisations for the existence of biregular graphs, while Theorem 5.4 for regular digraphs.

us the Presburger characterisations of the existence of biregular graphs, while Theorem 5.4 the same characterisations for regular digraphs.

- Equipped with Theorems 5.2 and 5.4, we construct the formula $\text{PREB}(x)$ as required in Theorem 2.1 in Section 6.
- However, the proofs of Theorems 5.2 and 5.4 themselves are rather long and involved. So we postpone their proofs until Sections 7 and 8.

Figure 3.1 illustrates the interdependence among Sections 4–8.

4. Quantified modal logic with counting. In this section we present *quantified modal logic with counting* (QMLC), which for our purpose, will be easier to work with. We are going to show that \mathcal{C}^2 and QMLC are equivalent in terms of spectra. In fact, our proof shows that \mathcal{C}^2 and QMLC are equivalent up to renaming/deleting/adding relational symbols, when QMLC are restricted to “complete” structures defined as follows. A structure \mathfrak{A} is a complete structure, if it satisfies the following properties.

- (N1) \mathfrak{A} is a clique over A . That is, for every $a, b \in A$, either $a = b$ or $R(a, b)$ for some $R \in \mathcal{R}$.
- (N2) Every binary relation in \mathcal{R} does not intersect identity relation. That is, for every $R \in \mathcal{R}$, if $R(a, b)$ holds, $a \neq b$.
- (N3) \mathcal{R} is closed under inverse. That is, for every $R \in \mathcal{R}$, there exists $\overleftarrow{R} \in \mathcal{R}$ such that $\overleftarrow{R} \neq R$ and for every $a, b \in A$, $R(a, b)$ if and only if $\overleftarrow{R}(b, a)$.
- (N4) The binary predicates in \mathcal{R} are pairwise disjoint.

Our proof is an adaptation of the proof by Lutz, Sattler and Wolter [29] which shows that similar equivalence holds between two-variable logic and modal logic.

The class MLC of *modal logic with counting* is defined with the following syntax.

$$\phi ::= \neg\phi \mid \alpha \mid \phi \wedge \phi \mid \diamond_{\beta}^k \phi$$

where α ranges over \mathcal{P} and β over \mathcal{R} .

The semantics of MLC is as follows. Let \mathfrak{A} be a structure of τ and $a \in A$ and ϕ be an MLC formula. That \mathfrak{A} satisfies ϕ from a , denoted by $\mathfrak{A}, a \models \phi$, is defined as follows.

- $\mathfrak{A}, a \models P$, where $P \in \mathcal{P}$, if $P(a)$ holds in \mathfrak{A} .
- $\mathfrak{A}, a \models \neg\phi$, if $\mathfrak{A}, a \not\models \phi$.
- $\mathfrak{A}, a \models \phi_1 \wedge \phi_2$, if $\mathfrak{A}, a \models \phi_1$ and $\mathfrak{A}, a \models \phi_2$.
- $\mathfrak{A}, a \models \diamond_R^k \phi$, if there exist at least k elements $b_1, \dots, b_k \in A$ such that $R(a, b_i)$ holds in \mathfrak{A} and $\mathfrak{A}, b_i \models \phi$ for $i = 1, \dots, k$.

We define the class of quantified modal logic with counting, denoted by QMLC with the following syntax.

$$\psi ::= \neg\psi \mid \psi_1 \wedge \psi_2 \mid \exists^k \phi$$

where the formula $\phi \in \text{MLC}$. A QMLC formula ψ is called a *basic* QMLC, if it is of the form $\exists^k \phi$, where $\phi \in \text{MLC}$.

The semantics of QMLC is as follows. Let \mathfrak{A} be a structure of τ and $\psi \in \text{QMLC}$. That \mathfrak{A} satisfies ψ , denoted by $\mathfrak{A} \models \psi$, is defined as follows.

- $\mathfrak{A} \models \neg\psi$, if it is not the case that $\mathfrak{A} \models \psi$.
- $\mathfrak{A} \models \psi_1 \wedge \psi_2$, if $\mathfrak{A} \models \psi_1$ and $\mathfrak{A} \models \psi_2$.
- $\mathfrak{A} \models \exists^k \phi$, if there exist at least k elements $a_1, \dots, a_k \in A$ such that $\mathfrak{A}, a_i \models \phi$ for $i = 1, \dots, k$.

We denote by $\text{SPEC}(\psi)$ the set of the cardinalities of the universes of complete structures of ψ . Note that for QMLC, the notion of spectrum is restricted to complete structures.

In the following we are going to show that from spectral point of view, \mathcal{C}^2 and QMLC are equivalent. The intuitive explanation for the requirement of complete structure is as follows. Notice that in QMLC we cannot express the negation of a binary relation $\neg R(x, y)$. Rather, to “express” $\neg R(x, y)$ in QMLC, we introduce a new relation symbol to capture $\neg R(x, y)$, hence, the requirement (N1) and (N4) in the complete structure. Similarly, in QMLC from an element x , we cannot express the “inverse” direction $R(y, x)$. So we introduce a new relation \overline{R} that captures the “inverse” of R , and $R(y, x)$ will be simulated by $\overline{R}(x, y)$, instead, hence the requirement (N3). We require (N2) simply for technical convenience.

Theorem 4.1 below states the spectral equivalence between \mathcal{C}^2 and QMLC, when QMLC is restricted to complete structures.

THEOREM 4.1. *For every $\varphi \in \mathcal{C}^2$, there is a QMLC formula ϕ such that*

- *for every structure $\mathfrak{A} \models \varphi$, there is a complete structure $\mathfrak{B} \models \phi$ where $|A| = |B|$;*
- *for every complete structure $\mathfrak{B} \models \phi$, there is a structure $\mathfrak{A} \models \varphi$ and $|A| = |B|$.*

Proof. Let $\varphi \in \mathcal{C}^2$. By extending/renaming/deleting the relations, and by modifying the sentence φ , if necessary, we can obtain another \mathcal{C}^2 sentence φ' such that

- *for every structure $\mathfrak{A} \models \varphi$, there is a complete structure $\mathfrak{A}' \models \varphi'$ where $|A| = |A'|$;*
- *for every complete structure $\mathfrak{A}' \models \varphi'$, there is a structure $\mathfrak{A} \models \varphi$ and $|A| = |A'|$.*

The details of the construction of φ' is straightforward, hence, omitted. For example, to achieve (N1) and (N4) we can introduce a new binary relation for each Boolean combination of relations in φ . We can do similar trick to achieve (N2) and (N3).

From this formula φ' , we are going to construct the desired QMLC formula ϕ . It consists of the following two steps.

1. Convert the sentence φ' into its “normal form” ψ such that for every complete structure \mathfrak{A} , we have $\mathfrak{A} \models \varphi'$ if and only if $\mathfrak{A} \models \psi$.^{||}
2. Convert the sentence ψ into a “quantified modal logic” (QMLC) sentence ϕ such that for every complete structure \mathfrak{A} , we have $\mathfrak{A} \models \psi$ if and only if $\mathfrak{A} \models \phi$.

In the following paragraphs we are going to describe formally these two steps.

A \mathcal{C}^2 sentence is in *normal form*, if all the quantifiers are either of form

$$\exists^k y \left(R(x, y) \wedge \theta(y) \right), \quad \text{or} \quad \exists^k x \theta(x)$$

and all other applications of variables are of form $P(x)$, where P is a unary predicate.

The \mathcal{C}^2 sentence φ' can be converted into its equivalent sentence ψ in normal form as follows.

- First, we rewrite every subformula of the form $\exists^k y \theta(x, y)$ with one free variable x into the following form:

$$\begin{aligned} & \theta(x, x) \wedge \exists^{k-1} y \left((x \neq y) \wedge \theta(x, y) \right) \\ & \quad \vee \\ & \exists^k y \left((x \neq y) \wedge \theta(x, y) \right) \end{aligned}$$

After such rewriting, we can assume that every quantifier in φ is of the form $\exists^k y \left((x \neq y) \wedge \theta(x, y) \right)$.

- Second, every quantification $\exists^k y \left((x \neq y) \wedge \theta(x, y) \right)$, in which $\theta(x, y)$ contains a subformula $\alpha(x)$ depending only on x , can be rewritten into the form:

$$\begin{aligned} & \neg \alpha(x) \wedge \exists^k y \left((x \neq y) \wedge \theta_0(x, y) \right) \\ & \quad \vee \\ & \alpha(x) \wedge \exists^k y \left((x \neq y) \wedge \theta_1(x, y) \right) \end{aligned}$$

where $\theta_0(x, y)$ and $\theta_1(x, y)$ are obtained from θ by replacing $\alpha(x)$ with false and true, respectively. We can repeat this until $\theta(x, y)$ no longer has a subformula depending only on x .

After such rewriting we can assume that every quantifier in φ is of the form

$$\exists^k y \left((x \neq y) \wedge \theta(x, y) \right),$$

where $\theta(x, y)$ does not contain any subformula depending only on x .

- Third, every quantification $\exists^k y \left((x \neq y) \wedge \theta(x, y) \right)$ can be rewritten into the form:

$$\bigvee_{f \in \Delta_{\mathcal{R}}^k} \bigwedge_{R \in \mathcal{R}} \exists^{f(R)} y \left(R(x, y) \wedge \theta_R(y) \right)$$

where $\Delta_{\mathcal{R}}^k$ is the set of all functions $f : \mathcal{R} \rightarrow \mathbb{N}$ such that $\sum_{R \in \mathcal{R}} f(R) = k$, and $\theta_R(y)$ is obtained from $\theta(x, y)$ by replacing each $R'(x, y)$ with true if $R = R'$, and false otherwise.

^{||}We would like to remark that the normal form here is different from the standard Scott’s normal form.

By performing these three steps, we get the C^2 sentence ψ in the normal form. Particularly, for every complete structure \mathfrak{A} , we have $\mathfrak{A} \models \varphi'$ if and only if $\mathfrak{A} \models \psi$.

Now from this C^2 sentence ψ in normal form, the construction of its QMLC sentence $\phi = F(\psi)$ can be done inductively as follows. There are two cases.

1. ϑ has no free variable.
 - If ϑ is $\neg\vartheta_1$, then $F(\vartheta) = \neg F(\vartheta_1)$.
 - If ϑ is $\vartheta_1 \wedge \vartheta_2$, then $F(\vartheta) = F(\vartheta_1) \wedge F(\vartheta_2)$.
 - If ϑ is $\exists^k x \vartheta_1(x)$, then $F(\vartheta) = \exists^k F(\vartheta_1(x))$.
2. ϑ has one free variable x .
 - If $\vartheta(x)$ is $P(x)$, then $F(\vartheta(x)) = P$.
 - If $\vartheta(x)$ is $\vartheta_1(x) \wedge \vartheta_2(x)$, then $F(\vartheta(x)) = F(\vartheta_1(x)) \wedge F(\vartheta_2(x))$.
 - If $\vartheta(x)$ is $\neg\vartheta_1(x)$, then $F(\vartheta(x)) = \neg F(\vartheta_1(x))$.
 - If $\vartheta(x)$ is $\exists^k y R(x, y) \wedge \vartheta_1(x, y)$, then $F(\vartheta(x)) = \diamond_R^k F(\vartheta_1(x, y))$.

The case when ϑ has one free variable y can be handled in a symmetrical way. By a straightforward induction, we can show that for every complete structure \mathfrak{A} , $\mathfrak{A} \models \vartheta$ if and only if $\mathfrak{A} \models F(\vartheta)$. In particular, from the equivalences between φ and φ' , between φ' and ψ as well as between ψ and $\phi = F(\psi)$, we obtain that

- for every structure $\mathfrak{A} \models \varphi$, there is a complete structure $\mathfrak{B} \models \phi$ such that $|A| = |B|$;
- for every complete structure $\mathfrak{B} \models \phi$, there is a structure $\mathfrak{A} \models \varphi$ such that $|A| = |B|$.

This concludes our proof of Theorem 4.1. \square

5. Regular graphs. In this section we are going to introduce two types of regular graphs: biregular graphs (bipartite regular graphs) and regular digraphs. The main results in this section are Theorems 5.2 and 5.4. For the sake of readability, we postpone their proofs until Sections 7 and 8.

5.1. Biregular graphs. An ℓ -type bipartite graph is $G = (U, V, E_1, \dots, E_\ell)$, where E_1, \dots, E_ℓ are *pairwise disjoint* subsets of $U \times V$. Elements in E_i are called E_i -edges. It helps to think of G as a bipartite graph in which the edges are coloured with ℓ colours.

For a vertex $u \in U \cup V$, $\deg_{E_i}(u)$ denotes the number of E_i -edges adjacent to it, and $\deg(u) = \sum_{i=1}^{\ell} \deg_{E_i}(u)$. We write $\deg(G) = \max\{\deg(u) \mid u \text{ is a vertex in } G\}$. For an integer $d \in \mathbb{N}$, we write $\deg_{E_i}(u) = \blacktriangleright d$, to denote $\deg_{E_i}(u) \geq d$.

Let \mathbb{N} denote the set of natural numbers $\{0, 1, 2, \dots\}$ and $\blacktriangleright\mathbb{N} = \{\blacktriangleright 0, \blacktriangleright 1, \blacktriangleright 2, \dots\}$ and $\mathbb{B} = \mathbb{N} \cup \blacktriangleright\mathbb{N}$. We write $\mathbb{B}^{\ell \times m}$ to denote the set of $\ell \times m$ matrices whose entries are elements from \mathbb{B} . The entry in row i and column j of a matrix $D \in \mathbb{B}^{\ell \times m}$ is denoted by $D_{i,j}$.

Let $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$. An ℓ -type bipartite graph $G = (U, V, E_1, \dots, E_\ell)$ is (C, D) -biregular, if there is a partition $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ such that the following holds.

- For every $i = 1, \dots, \ell$, for every $j = 1, \dots, m$, for every vertex $u \in U_j$, $\deg_{E_i}(u) = C_{i,j}$.
- For every $i = 1, \dots, \ell$, for every $j = 1, \dots, n$, for every vertex $v \in V_j$, $\deg_{E_i}(v) = D_{i,j}$.

We call the partitions $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ the witness of the (C, D) -biregularity of G . We say that the (C, D) -biregular graph G is of size (\bar{M}, \bar{N}) , if $\bar{M} = (|U_1|, \dots, |U_m|)$ and $\bar{N} = (|V_1|, \dots, |V_n|)$.

THEOREM 5.1. *For every two matrices $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$, there is a*

Presburger formula $\text{BiREG}_{C,D}(\bar{X}, \bar{Y})$, where $\bar{X} = (X_1, \dots, X_m)$ and $\bar{Y} = (Y_1, \dots, Y_n)$ such that the following holds. There exists an ℓ -type (C, D) -biregular graph of size (\bar{M}, \bar{N}) if and only if $\text{BiREG}_{C,D}(\bar{M}, \bar{N})$ holds.

Theorem 5.1 is then generalised to the case of complete bipartite graphs. An ℓ -type bipartite graph $G = (U, V, E_1, \dots, E_\ell)$ is *complete*, if $U \times V = E_1 \cup \dots \cup E_\ell$. If G is also a (C, D) -biregular graph, then we call it a (C, D) -complete-biregular graph.

The following theorem is the main result in this subsection that will be used in the proof in Section 6.

THEOREM 5.2. *For every two matrices $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$, there is a Presburger formula $\text{COMP-BiREG}_{C,D}(\bar{X}, \bar{Y})$, where $\bar{X} = (X_1, \dots, X_m)$ and $\bar{Y} = (Y_1, \dots, Y_n)$ such that the following holds. There exists a (C, D) -complete-biregular graph of size (\bar{M}, \bar{N}) if and only if $\text{COMP-BiREG}_{C,D}(\bar{M}, \bar{N})$ holds.*

5.2. Regular digraphs. An ℓ -type directed graph (or, digraph for short) is a tuple $G = (V, E_1, \dots, E_\ell)$, where E_1, \dots, E_ℓ are *pairwise disjoint* irreflexive relations on V and for every $u, v \in V$, if $(u, v) \in E_1 \cup \dots \cup E_\ell$, then the inverse direction $(v, u) \notin E_1 \cup \dots \cup E_\ell$. Edges in E_i are called E_i -edges.

We will write $\text{in-deg}_{E_i}(u)$ to denote the number of incoming E_i -edges toward the vertex u , and $\text{out-deg}_{E_i}(u)$ the number of outgoing E_i -edges from the vertex u . As before, for an integer $d \in \mathbb{N}$, we write $\text{in-deg}_{E_i}(u) = \blacktriangleright d$ and $\text{out-deg}_{E_i}(u) = \blacktriangleright d$, to indicate that $\text{in-deg}_{E_i}(u) \geq d$ and $\text{out-deg}_{E_i}(u) \geq d$, respectively.

Let $C, D \in \mathbb{B}^{\ell \times m}$. An ℓ -type digraph $G = (V, E_1, \dots, E_\ell)$ is (C, D) -regular-digraph, if there exists a partition $V = V_1 \cup \dots \cup V_m$ such that for each $i = 1, \dots, \ell$, for each $j = 1, \dots, m$, for each vertex $v \in V_j$,

$$\text{in-deg}_{E_i}(v) = C_{i,j} \quad \text{and} \quad \text{out-deg}_{E_i}(v) = D_{i,j}.$$

We call $V_1 \cup \dots \cup V_m$ a witness of the (C, D) -regularity of G and the graph G is of size \bar{N} , if $\bar{N} = (|V_1|, \dots, |V_m|)$.

THEOREM 5.3. *For every $C, D \in \mathbb{B}^{\ell \times m}$, there exists a Presburger formula $\text{REG}_{C,D}(\bar{X})$, where $\bar{X} = (X_1, \dots, X_m)$ such that the following holds. There exists a (C, D) -regular-digraph of size \bar{N} if and only if $\text{REG}_{C,D}(\bar{N})$ holds.*

Similar to Section 5.1, Theorem 5.3 will be generalised to the case of complete regular digraph. An ℓ -type graph $G = (V, E_1, \dots, E_\ell)$ is a *complete* digraph, if for every two different vertices u, v , either (u, v) or (v, u) is in $E_1 \cup \dots \cup E_\ell$. Note that here we still insist that if $(u, v) \in E_1 \cup \dots \cup E_\ell$, the inverse $(v, u) \notin E_1 \cup \dots \cup E_\ell$. If G is also a (C, D) -regular, then we call G a (C, D) -complete-regular digraph.

The following theorem is the main result in this subsection that will be used in the proof in Section 6.

THEOREM 5.4. *For every $C, D \in \mathbb{B}^{\ell \times m}$, there exists a Presburger formula $\text{COMP-REG}_{C,D}(\bar{X})$, where $\bar{X} = (X_1, \dots, X_m)$ such that the following holds. There exists a (C, D) -complete-regular digraph of size \bar{N} if and only if $\text{COMP-REG}_{C,D}(\bar{N})$ holds.*

6. Proof of Theorem 2.1. Now we are ready to prove Theorem 2.1. Let ϕ be a QMLC sentence. Recall that a *basic* QMLC formula is of the form $\exists^k \varphi$, where $\varphi \in \text{MLC}$. We also assume that in ϕ we have “pushed” all the negations inside so that they are applied only to basic QMLC. We are going to construct a Presburger formula $\text{PREB}_\phi(x)$ such that $\text{SPEC}(\phi) = \{n \mid \text{PREB}_\phi(n) \text{ holds}\}$.

Before we proceed, we need a few auxiliary notations. Let \mathcal{P} be the set of unary predicates used in ϕ and $\mathcal{R} = \{R_1, \dots, R_\ell, \bar{R}_1, \dots, \bar{R}_\ell\}$ the set of binary relations

used in ϕ , where \overleftarrow{R}_i is the inverse relation of R_i . Let K be the integer such that for all subformulae $\diamond_R^l \psi$ in ϕ , we have $l \leq K$.

We denote by \mathcal{M}_ϕ the set of all MLC subformulae of ϕ and their negations. A *type* in ϕ is a subset $T \subseteq \mathcal{M}_\phi$ such that

- if $\varphi_1 \wedge \varphi_2 \in T$, then both $\varphi_1, \varphi_2 \in T$;
- $\varphi \in T$ if and only if $\neg\varphi \notin T$;
- if $\neg(\varphi_1 \wedge \varphi_2) \in T$, then at least one of $\neg\varphi_1, \neg\varphi_2 \in T$.

For a structure \mathfrak{A} (not necessarily a model of ϕ) and an element $a \in A$, we define *the type of a in \mathfrak{A}* , denoted by $\text{type}_{\mathfrak{A}}(a) \subseteq \mathcal{M}_\phi$, where $\varphi \in \text{type}_{\mathfrak{A}}(a)$ if and only if $\mathfrak{A}, a \models \varphi$. For a type T , we write $T(\mathfrak{A})$ to denote the set of elements in A with type T . Note that the sets $T(\mathfrak{A})$'s are pairwise disjoint. We let \mathcal{T}_ϕ to be the set of all types in ϕ .

We say that a function $f : \mathcal{T}_\phi \times \mathcal{R} \times \mathcal{T}_\phi \rightarrow \{0, 1, \dots, K\} \cup \{\blacktriangleright K\}$ is *consistent*, if for every $T \in \mathcal{T}_\phi$ the following holds.

- If $\diamond_R^l \mu \in T$, then $\sum_{T' \text{ s.t. } T' \ni \mu} f(T, R, T') \geq l$.
- If $\neg(\diamond_R^l \mu) \in T$, then $\sum_{T' \text{ s.t. } T' \ni \mu} f(T, R, T') \leq l - 1$, and $f(T, R, T') \in \mathbb{N}$, for every $R \in \mathcal{R}$ and for every type $T' \ni \mu$.

In the following we enumerate the set of all consistent functions $\mathcal{F} = \{f_1, \dots, f_m\}$, the set of all types in $\mathcal{T}_\phi = \{T_1, \dots, T_n\}$, and the set $\mathcal{T}_\phi \times \mathcal{F} = \{(T_1, f_1), \dots, (T_n, f_m)\}$.

The desired Presburger formula $\text{PREB}_\phi(x)$ is defined as the formula:

$$\exists X_{(T_1, f_1)} \cdots \exists X_{(T_n, f_m)} \left(x = \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq m} X_{T_i, f_j} \right) \wedge \text{PREB-Atom}_\phi(\bar{X}) \wedge \text{CON}(\bar{X})$$

where $\bar{X} = (X_{(T_1, f_1)}, \dots, X_{(T_n, f_m)})$ is the vector of all the variables $X_{(T, f)}$'s.

We will formally define the formulae $\text{PREB-Atom}_\phi(\bar{X})$ and $\text{CON}(\bar{X})$ in the following paragraphs. The intended meaning of the variable X_{T_i, f_j} and the formulae $\text{PREB-Atom}_\phi(x)$ and $\text{CON}(\bar{X})$ is as follows. The variable X_{T_i, f_j} is to represent the number of elements of type T_i and for each binary relation $R \in \mathcal{R}$ and a type $S \in \mathcal{T}_\phi$, there is $f(T_i, R, S)$ number of outgoing R -edges towards the elements of type S .

Naturally, the total number of all elements in the universe will be the sum of all X_{T_i, f_j} 's, hence, the sum:

$$x = \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq m} X_{T_i, f_j}$$

The formula $\text{PREB-Atom}_\phi(x)$ is to make sure that the satisfiability of the QMLC sentence is preserved. Intuitively, if ϕ is $\exists^k \varphi$, the formula $\text{PREB-Atom}_\phi(x)$ states that there are at least k elements in which φ holds. Formally, it is defined inductively as follows. (Recall that all the negations have been “pushed” inside so that they are applied only to basic QMLC.)

- If $\phi := \exists^k \varphi$, then $\text{PREB-Atom}_\phi := \sum_{(T, f) \text{ s.t. } \varphi \in T} X_{(T, f)} \geq k$.
- If $\phi := \neg \exists^k \varphi$, then $\text{PREB-Atom}_\phi := \sum_{(T, f) \text{ s.t. } \varphi \in T} X_{(T, f)} \leq k - 1$.
- If $\phi := \phi_1 \wedge \phi_2$, then $\text{PREB-Atom}_\phi := \text{PREB-Atom}_{\phi_1} \wedge \text{PREB-Atom}_{\phi_2}$.
- If $\phi := \phi_1 \vee \phi_2$, then $\text{PREB-Atom}_\phi := \text{PREB-Atom}_{\phi_1} \vee \text{PREB-Atom}_{\phi_2}$.

Finally, the formula $\text{CON}(\bar{X})$ is to make sure that the solution to each variable X_{T_i, f_j} is “consistent” to the intended meaning of the type T_i and function f_j . That is, for every types $S, T \in \mathcal{T}_\phi$ the following holds.

- Every solution \bar{M}_T to the variables $\bar{X}_T = (X_{T,f_1}, \dots, X_{T,f_m})$ in the formula $\text{CON}(\bar{X})$ corresponds to a $(D_T, \overleftarrow{D}_T)$ -complete-regular digraph of size \bar{M}_T , where the matrices $D_T, \overleftarrow{D}_T \in \mathbb{B}^{\ell \times m}$ are as follows.

$$D_T := \begin{pmatrix} f_1(T, R_1, T) & f_2(T, R_1, T) & \cdots & f_m(T, R_1, T) \\ f_1(T, R_2, T) & f_2(T, R_2, T) & \cdots & f_m(T, R_2, T) \\ \vdots & \vdots & \ddots & \vdots \\ f_1(T, R_\ell, T) & f_2(T, R_\ell, T) & \cdots & f_m(T, R_\ell, T) \end{pmatrix}$$

and

$$\overleftarrow{D}_T := \begin{pmatrix} f_1(T, \overleftarrow{R}_1, T) & f_2(T, \overleftarrow{R}_1, T) & \cdots & f_m(T, \overleftarrow{R}_1, T) \\ f_1(T, \overleftarrow{R}_2, T) & f_2(T, \overleftarrow{R}_2, T) & \cdots & f_m(T, \overleftarrow{R}_2, T) \\ \vdots & \vdots & \ddots & \vdots \\ f_1(T, \overleftarrow{R}_\ell, T) & f_2(T, \overleftarrow{R}_\ell, T) & \cdots & f_m(T, \overleftarrow{R}_\ell, T) \end{pmatrix}$$

Notice that the matrix D_T contains only the information on the degree of R_1, \dots, R_ℓ , while \overleftarrow{D}_T the information on the degree of $\overleftarrow{R}_1, \dots, \overleftarrow{R}_\ell$. This is because the incoming R_i edges to an element v are precisely the outgoing \overleftarrow{R}_i edges from v , and vice versa, the incoming \overleftarrow{R}_i edges from an element v are precisely the outgoing R_i edges to v .

- Every solution \bar{M}_S, \bar{M}_T to the variables $\bar{X}_S = (X_{S,f_1}, \dots, X_{S,f_m})$ and $\bar{X}_T = (X_{T,f_1}, \dots, X_{T,f_m})$ in the formula $\text{CON}(\bar{X})$ corresponds to a $(D_{S \rightarrow T}, \overleftarrow{D}_{S \rightarrow T})$ -complete-biregular digraph of size (\bar{M}_S, \bar{M}_T) , where $D_{S \rightarrow T}, \overleftarrow{D}_{S \rightarrow T} \in \mathbb{B}^{\ell \times m}$ are the following matrices.

$$D_{S \rightarrow T} := \begin{pmatrix} f_1(S, R_1, T) & f_2(S, R_1, T) & \cdots & f_m(S, R_1, T) \\ f_1(S, R_2, T) & f_2(S, R_2, T) & \cdots & f_m(S, R_2, T) \\ \vdots & \vdots & \ddots & \vdots \\ f_1(S, R_\ell, T) & f_2(S, R_\ell, T) & \cdots & f_m(S, R_\ell, T) \\ f_1(S, \overleftarrow{R}_1, T) & f_2(S, \overleftarrow{R}_1, T) & \cdots & f_m(S, \overleftarrow{R}_1, T) \\ f_1(S, \overleftarrow{R}_2, T) & f_2(S, \overleftarrow{R}_2, T) & \cdots & f_m(S, \overleftarrow{R}_2, T) \\ \vdots & \vdots & \ddots & \vdots \\ f_1(S, \overleftarrow{R}_\ell, T) & f_2(S, \overleftarrow{R}_\ell, T) & \cdots & f_m(S, \overleftarrow{R}_\ell, T) \end{pmatrix}$$

$$\overleftarrow{D}_{S \rightarrow T} := \begin{pmatrix} f_1(T, \overleftarrow{R}_1, S) & f_2(T, \overleftarrow{R}_1, S) & \cdots & f_m(T, \overleftarrow{R}_1, S) \\ f_1(T, \overleftarrow{R}_2, S) & f_2(T, \overleftarrow{R}_2, S) & \cdots & f_m(T, \overleftarrow{R}_2, S) \\ \vdots & \vdots & \ddots & \vdots \\ f_1(T, \overleftarrow{R}_\ell, S) & f_2(T, \overleftarrow{R}_\ell, S) & \cdots & f_m(T, \overleftarrow{R}_\ell, S) \\ f_1(T, R_1, S) & f_2(T, R_1, S) & \cdots & f_m(T, R_1, S) \\ f_1(T, R_2, S) & f_2(T, R_2, S) & \cdots & f_m(T, R_2, S) \\ \vdots & \vdots & \ddots & \vdots \\ f_1(T, R_\ell, S) & f_2(T, R_\ell, S) & \cdots & f_m(T, R_\ell, S) \end{pmatrix}$$

Notice that in the matrix $D_{S \rightarrow T}$ the first ℓ rows contains the information on the degree of R_1, \dots, R_ℓ , and the last ℓ rows the information on the degree

of $\overleftarrow{R}_1, \dots, \overleftarrow{R}_\ell$ from the type S to the type T ; while in the matrix $\overleftarrow{D}_{S \rightarrow T}$ it is the opposite and the direction is from the type T to the type S . Similar as in the D_T, \overleftarrow{D}_t case above, this is because the incoming R_i edges to an element v are precisely the outgoing \overleftarrow{R}_i edges from v , and vice versa, the outgoing R_i edges from an element v are precisely the incoming \overleftarrow{R}_i edges to v .

Now the formula $\text{CON}(\overline{X})$ is simply the conjunction:

$$\begin{aligned} \text{CON}(\overline{X}) := & \bigwedge_{1 \leq i \leq n} \text{COMP-REG}_{D_{T_i}, \overleftarrow{D}_{T_i}}(\overline{X}_T) \\ & \wedge \bigwedge_{1 \leq j < i \leq n} \text{COMP-BiREG}_{D_{T_i \rightarrow T_j}, \overleftarrow{D}_{T_i \rightarrow T_j}}(\overline{X}_{T_i}, \overline{X}_{T_j}) \end{aligned}$$

where $\overline{X}_{T_i} = (X_{(T_i, f_1)}, \dots, X_{(T_i, f_m)})$ is the vector of variables associated with the type T_i , and the formulae $\text{COMP-REG}_{D_{T_i}, \overleftarrow{D}_{T_i}}$ and $\text{COMP-BiREG}_{D_{T_i \rightarrow T_j}, \overleftarrow{D}_{T_i \rightarrow T_j}}$ are the formulae given by Theorems 5.2 and 5.4, respectively.

We are going to show that the set of solutions to $\text{PREB}_\phi(x)$ is precisely the spectrum of ϕ , as stated in the claim below. Abusing the notation, we let PREB_ϕ itself to denote the set $\{n \mid \text{PREB}_\phi(n) \text{ holds}\}$. Recall also that as defined in Section 4, the spectrum of a QMLC sentence ϕ is restricted to the complete structures.

CLAIM 1. *For every QMLC sentence ϕ , $\text{PREB}_\phi = \text{SPEC}(\phi)$, where $\text{PREB}_\phi(x)$ is the formula*

$$\exists X_{(T_1, f_1)} \cdots \exists X_{(T_n, f_m)} \left(x = \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq m} X_{T_i, f_j} \right) \wedge \text{PREB-Atom}_\phi(\overline{X}) \wedge \text{CON}(\overline{X})$$

and $\overline{X} = (X_{(T_1, f_1)}, \dots, X_{(T_n, f_m)})$ is the vector of all the variables $X_{(T, f)}$'s.

The proof is by induction on ϕ . The base case is when ϕ is a basic QMLC sentence or the negation of a basic QMLC sentence. We consider first the case when ϕ is a basic QMLC formula of the form $\exists^k \varphi$, where $\varphi \in \text{MLC}$. In this case $\text{PREB-Atom}_\phi(\overline{X})$ is $\sum_{(T, f) \text{ s.t. } \varphi \in T} X_{(T, f)} \geq k$.

We first show the direction $\text{PREB}_\phi \subseteq \text{SPEC}(\phi)$. Let $N \in \text{PREB}_\phi$. Let $\overline{M} = (M_{T_1, f_1}, \dots, M_{T_n, f_m})$ be an assignment to \overline{X} such that $\text{PREB}_\phi(N)$ holds. That is, the following holds.

$$\left(N = \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq m} M_{T_i, f_j} \right) \wedge \text{PREB-Atom}_\phi(\overline{M}) \wedge \text{CON}(\overline{M})$$

In the following we are going to write \overline{M}_T to denote $(M_{T, f_1}, \dots, M_{T, f_m})$ for every type $T \in \mathcal{T}_\phi$.

Since $x = \sum_{(T, f)} X_{T, f}$, we have $N = \sum_{(T, f)} M_{(T, f)}$. We take a set V of N vertices and we partition $V = V_{(T_1, f_1)} \cup \dots \cup V_{(T_n, f_m)}$ such that $|V_{(T, f)}| = M_{(T, f)}$ for each $T \in \mathcal{T}_\phi$ and $f \in \mathcal{F}$. We denote by $V_T = V_{(T, f_1)} \cup \dots \cup V_{(T, f_m)}$ for each $T \in \mathcal{T}_\phi$.

Since $\text{CON}(\overline{M})$ holds, by Theorems 5.4, for each $T \in \mathcal{T}_\phi$, there exists a $(D_T, \overleftarrow{D}_T)$ -complete-regular digraph $G_T = (V_T, R_{T,1}, \dots, R_{T,\ell})$ of size \overline{M}_T , with $V_T = V_{(T, f_1)} \cup \dots \cup V_{(T, f_m)}$ being the witness of the $(D_T, \overleftarrow{D}_T)$ -regularity. This means that for every vertex $v \in V_{T, f_i}$, for every $R \in \{R_1, \dots, R_\ell\}$,

- out- $\text{deg}_R(v)$ in the graph G_T is $f_i(T, R, T)$;
- in- $\text{deg}_R(v)$ in the graph G_T is $f_i(T, \overleftarrow{R}, T)$.

Now let $\tilde{G}_T = (V_T, R_{T,1}, \dots, R_{T,\ell}, \overleftarrow{R}_{T,1}, \dots, \overleftarrow{R}_{T,\ell})$ be the graph obtained from G_T by taking \overleftarrow{R}_i as the inverse of R_i . Then for each vertex $v \in V_T$,

- $\text{out-deg}_R(v) = \text{in-deg}_{\overleftarrow{R}}(v)$ in the graph \tilde{G}_T ;
- $\text{in-deg}_R(v) = \text{out-deg}_{\overleftarrow{R}}(v)$ in the graph \tilde{G}_T .

Similarly, by Theorem 5.2 for each $T_i, T_j \in \mathcal{T}_\phi$, where $j \leq i - 1$, there exists a $(D_{T_i \rightarrow T_j}, \overleftarrow{D}_{T_i \rightarrow T_j})$ -biregular-complete graph

$$G_{T_i, T_j} = (V_{T_i}, V_{T_j}, R_{T_i, T_j, 1}, \dots, R_{T_i, T_j, \ell}, \overleftarrow{R}_{T_i, T_j, 1}, \dots, \overleftarrow{R}_{T_i, T_j, \ell})$$

of size $(\bar{M}_{T_i}, \bar{M}_{T_j})$, with $V_{T_i} = V_{(T_i, f_1)} \cup \dots \cup V_{(T_i, f_m)}$ and $V_{T_j} = V_{(T_j, f_1)} \cup \dots \cup V_{(T_j, f_m)}$ being the witness of the $(D_{T_i \rightarrow T_j}, \overleftarrow{D}_{T_i \rightarrow T_j})$ -biregularity. This means that for every $R \in \{R_1, \dots, R_\ell, \overleftarrow{R}_1, \dots, \overleftarrow{R}_\ell\}$,

- for every vertex $v \in V_{T_i, f}$, $\text{out-deg}_R(v)$ in the graph G_{T_i, T_j} is $f(T_i, R, T_j)$;
- for every vertex $v \in V_{T_j, f}$, $\text{out-deg}_R(v)$ in the graph G_{T_i, T_j} is $f(T_i, \overleftarrow{R}, T_j)$.

We put the orientation in every the edges in the graph G_{T_i, T_j} going from V_{T_i} to V_{T_j} .

Now let \tilde{G}_{T_i, T_j} be the graph obtained by adding (u, v) into \overleftarrow{R} in the graph G_{T_i, T_j} whenever (v, u) is an R -edge in G_{T_i, T_j} .

Hence, we have for each vertex $v \in V_{T_i} \cup V_{T_j}$, for each $R \in \mathcal{R}$

- $\text{out-deg}_R(v) = \text{in-deg}_{\overleftarrow{R}}(v)$ in the graph \tilde{G}_{T_i, T_j} ;
- $\text{in-deg}_R(v) = \text{out-deg}_{\overleftarrow{R}}(v)$ in the graph \tilde{G}_{T_i, T_j} .

Let $G = (V, R_1, \dots, R_\ell, \overleftarrow{R}_1, \dots, \overleftarrow{R}_\ell)$ be the combination of all the graphs \tilde{G}_{T_i} 's and \tilde{G}_{T_i, T_j} 's. Formally,

$$V = \bigcup_T V(\tilde{G}_T)$$

$$R = \bigcup_T R(\tilde{G}_T) \cup \bigcup_{T_i, T_j} R(\tilde{G}_{T_i, T_j}) \quad \text{for each } R \in \{R_1, \dots, R_\ell, \overleftarrow{R}_1, \dots, \overleftarrow{R}_\ell\},$$

where $V(\tilde{G}_T)$ is the set of vertices in \tilde{G}_T and $R(\tilde{G}_T)$ is the set of R -edges in \tilde{G}_T .

Moreover, we also label each vertex $v \in V$ with a subset of \mathcal{P} as follows. For each $T \in \mathcal{T}_\phi$, for each $v \in V_T$, we “declare” that v is labeled with a unary predicate $P \in \mathcal{P}$ if and only if $P \in T$.

We claim that $G \models \phi$. For that, it is sufficient to show that for each $T \in \mathcal{T}_\phi$, for each $v \in V_T$, $\text{type}_G(v) = T$. The proof is divided into three cases.

- For each unary predicate $P \in \mathcal{P}$, it is by our labelling of the vertices of G that $P(v)$ holds in G if and only if $P \in T$.
- For each $\diamond_R^l \mu \in T$, we have

$$\sum_{T' \text{ s.t. } T' \ni \mu} f(T, R, T') \geq l$$

number of outgoing R -edges from v . Since every function $f \in \mathcal{F}$ is consistent, $\diamond_R^l \mu \in \text{type}_G(v)$.

- Similarly, for each $\diamond_R^l \mu \notin T$, and hence $\neg(\diamond_R^l \mu) \in T$, we have

$$\sum_{T' \text{ s.t. } T' \ni \mu} f(T, R, T') \leq l - 1$$

number of outgoing R -edges from v . Since every function $f \in \mathcal{F}$ is consistent, $\diamond_R^l \mu \notin \text{type}_G(v)$.

Therefore the graph $G \models \phi$, and hence $N \in \text{SPEC}(\phi)$.

Now we prove the direction $\text{PREB}_\phi \supseteq \text{SPEC}(\phi)$. Suppose $\mathfrak{A} \models \phi$ and $|A| = N$. Let $\vec{M} = (M_{(T_1, f_1)}, \dots, M_{(T_n, f_m)})$ where $M_{(T, f)}$ be the number of elements of type T from which there exist $f(T, R, S)$ outgoing R -edges towards the elements of type S . Assign the variable $X_{(T, f)}$ with $M_{(T, f)}$, for each $T \in \mathcal{T}_\phi$ and $f \in \mathcal{F}$. It immediately follows from Theorems 5.2 and 5.4 that $\text{CON}(\vec{M})$ holds. Moreover, $\text{PREB-Atom}_\phi(\vec{M})$ holds, since $\mathfrak{A} \models \phi$. This completes the proof that $\text{PREB}_\phi = \text{SPEC}(\phi)$, when ϕ is a basic QMLC sentence.

When $\phi \in \text{QMLC}$ is the negation of a basic QMLC sentence, say $\neg \exists^k \varphi$, the formula PREB-Atom_ϕ is

$$\sum_{(T, f) \text{ s.t. } \varphi \in T} X_{(T, f)} \leq k - 1$$

which is the negation of

$$\sum_{(T, f) \text{ s.t. } \varphi \in T} X_{(T, f)} \geq k$$

Then $\text{PREB}_\phi = \text{SPEC}(\phi)$ follows immediately from above.

The correctness for the case when ϕ is $\phi_1 \wedge \phi_2$ or $\phi_1 \vee \phi_2$, can be established via straightforward inductive argument. This completes our proof that $\text{PREB}_\phi = \text{SPEC}(\phi)$, and hence, Theorem 2.1.


7. Proof of Theorem 5.2. The proof of Theorem 5.2 is rather long. As a warm-up, we prove the following easy Proposition 7.1 first.

PROPOSITION 7.1. *Let $c, d \geq 0$. For every $M, N \in \mathbb{N}$, the following holds.*

- (a) *There exists a (c, d) -biregular graph of size (M, N) if and only if $N \geq c$, $M \geq d$ and $Mc = Nd$.*
- (b) *There exists a $(c, \blacktriangleright d)$ -biregular graph of size (M, N) if and only if $M \geq d$, $N \geq c$ and $Mc \geq Nd$.*
- (c) *There exists a $(\blacktriangleright c, \blacktriangleright d)$ -biregular graph of size (M, N) if and only if $M \geq d$, $N \geq c$.*

Proof. Let $c, d \geq 0$, and let $M, N \in \mathbb{N}$. We first prove part (a). The ‘‘only if’’ direction follows from the fact that in (c, d) -biregular graph the number of edges is precisely $Mc = Nd$. That $M \geq d$ and $N \geq c$ is straightforward.

The ‘‘if’’ direction is as follows. Let $K = Mc = Nd$. Suppose also $M \geq d$ and $N \geq c$. First, we construct the following graph.



On the left side, we have M vertices, and each has degree c . On the right side, we have $K = Nd$ vertices, and each has degree 1. We are going to merge every d vertices on the right side into one vertex of degree d . The merging is as follows. We

merge every d vertices $v_i, v_{i+N}, \dots, v_{i+(d-1)N}$ into one vertex for every $i = 1, \dots, N$. Since $K = Nd$, it is possible to do such merging. Moreover, $N \geq c$, hence we do not have multiple edges between two vertices. Thus, we obtain the desired (c, d) -biregular graph of size (M, N) .**

Now we consider part (b). The “only if” direction follows from the fact that in $(c, \blacktriangleright d)$ -biregular graph the number of edges is precisely Mc , which should be greater than Nd . That $M \geq d$ and $N \geq c$ is straightforward.

For the “if” direction, the proof is almost the same as above. Suppose $M \geq d$, $N \geq c$ and $Mc \geq Nd$. Let $K = Mc$. We first construct the bipartite graph, in which on the left side, we have M vertices, each of which has degree c ; and on the right side we have $K = Mc$ vertices, each of which has degree 1.

For every $i \in \{1, \dots, N\}$, we set the set $I_i \subseteq \{1, \dots, K\}$ as follows.

$$I_i := \{i + kN \mid 1 \leq i + kN \leq K\}.$$

Since $K = Mc \geq Nd$, the cardinality $|I_i| \geq d$ for every $1 \leq i \leq N$. Now for every $1 \leq i \leq N$, we merge the vertices $\{v_j \mid j \in I_i\}$ into one vertex. Hence, we obtain the desired $(c, \blacktriangleright d)$ -biregular graph of size (M, N) .

Now we prove part (c). The “only if” direction is straightforward. The “if” direction is as follows. Suppose $M \geq d$ and $N \geq c$. There are two cases: either $Mc \geq Nd$, or $Mc < Nd$. In the former case, we construct a $(c, \blacktriangleright d)$ -biregular graph of size (M, N) , while in the latter case, we construct a $(\blacktriangleright c, d)$ -biregular graph of size (M, N) . In either case, we obtain a $(\blacktriangleright c, \blacktriangleright d)$ -biregular graph of size (M, N) . This completes our proof of Proposition 7.1. \square

The proof of Theorem 5.2 is a generalisation of Proposition 7.1 above. It is divided into five successive steps presented in Subsections 7.1– 7.6.

- Subsection 7.1.

It contains the generalisation of Proposition 7.1 to the case of (\bar{c}, \bar{d}) -biregular, where \bar{c} and \bar{d} are vectors over \mathbb{N} . (That is, we consider the 1-type biregular graphs.)

- Subsection 7.2.

In this subsection we use the Presburger characterisation for (\bar{c}, \bar{d}) -biregular graphs to obtain a similar characterisation for ℓ -type (C, D) -biregular graphs, where C, D are matrices over \mathbb{N} .

- Subsection 7.3.

In this subsection we obtain the characterisation for (C, D) -biregular graphs when C, D contain elements from $\blacktriangleright \mathbb{N}$ assuming that the number of vertices whose degrees specified with $\blacktriangleright d$ is “big enough.” It is obtained by using the characterisation obtained in Subsection 7.2.

- Subsection 7.4.

This subsection is the generalisation of Subsection 7.3, where the graphs may contain a “small” number of vertices whose degree is specified with $\blacktriangleright d$. The idea is to encode directly those vertices in the Presburger formula. This is presented formally by our notion of *partial graphs*.

- Subsection 7.5.

**Note that $Mc = Nd$, hence, $N \geq c$ already implies $M \geq d$. So it is not necessary to use the fact that $M \geq d$. Of course, by symmetry, we can first build a bipartite graph in which the left side has N vertices of degree d and the right side has $K = Mc$ vertices of degree 1. Then to build the desired (c, d) -biregular graph, we make use of the fact $M \geq d$.

In this subsection we present the construction of the formula required in Theorem 5.1. It is built from the formula obtained in Subsection 7.4.

- Subsection 7.6.

Finally in this subsection we present the construction of the formula required in Theorem 5.2, which is built from the formula obtained in Subsection 7.5.

In the following we write $\bar{1}$ to denote the vector $(1, \dots, 1) \in \mathbb{N}^m$, for an appropriate $m \geq 1$. That is, $\bar{1}$ is a vector whose components are all one. For two vectors $\bar{c} = (c_1, \dots, c_m) \in \mathbb{N}^m$ and $\bar{d} = (d_1, \dots, d_m) \in \mathbb{N}^m$, the dot product between \bar{c} and \bar{d} is $\bar{c} \cdot \bar{d} = c_1 d_1 + \dots + c_m d_m$.

7.1. When $C = \bar{c} \in \mathbb{N}^{1 \times m}$ and $D = \bar{d} \in \mathbb{N}^{1 \times n}$. In this subsection we consider the case when C and D consist of only one vector each. In this case, we are going to write (\bar{c}, \bar{d}) -biregular graph, where \bar{c} and \bar{d} are the only vectors in C and D , respectively.

LEMMA 7.2. *Let $\bar{c} \in \mathbb{N}^m$ and $\bar{d} \in \mathbb{N}^n$ and both do not contain zero entry. For each $\bar{M} \in \mathbb{N}^m$ and $\bar{N} \in \mathbb{N}^n$ such that $\bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} \geq 2(\bar{c} \cdot \bar{1})(\bar{d} \cdot \bar{1}) + 3$, the following holds. There exists a (\bar{c}, \bar{d}) -biregular graph of size (\bar{M}, \bar{N}) if and only if $\bar{M} \cdot \bar{c} = \bar{N} \cdot \bar{d}$.*

Proof. Let $\bar{c} \in \mathbb{N}^m$ and $\bar{d} \in \mathbb{N}^n$ and both do not contain zero entry. Let $\bar{M} \in \mathbb{N}^m$, $\bar{N} \in \mathbb{N}^n$ such that $\bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} \geq 2(\bar{c} \cdot \bar{1})(\bar{d} \cdot \bar{1}) + 3$.

The “only if” direction is straightforward. If G is a (\bar{c}, \bar{d}) -biregular graph of size (\bar{M}, \bar{N}) , then the number of edges in G is precisely $\bar{M} \cdot \bar{c} = \bar{N} \cdot \bar{d}$.

Now we prove the “if” part. Suppose $\bar{M} \in \mathbb{N}^m$, $\bar{N} \in \mathbb{N}^n$ such that $\bar{M} \cdot \bar{c} = \bar{N} \cdot \bar{d}$. Let $\bar{c} = (c_1, \dots, c_m)$ and $\bar{d} = (d_1, \dots, d_n)$, and $\bar{M} = (M_1, \dots, M_m)$ and $\bar{N} = (N_1, \dots, N_n)$.

We are going to construct a (\bar{c}, \bar{d}) -biregular graph of size (\bar{M}, \bar{N}) . We first construct a preliminary bipartite graph G pictured in Figure 7.1. That is, the left side has $\bar{M} \cdot \bar{1}$ vertices, and there are M_1 vertices of degree c_1 , M_2 nodes of degree c_2 , etc. The right side has $\bar{M} \cdot \bar{c}$ number of vertices, each of degree one.

We are going to do some merging of the vertices on the right side so that there are exactly N_1 vertices of degree d_1 , N_2 vertices of degree d_2 , etc. We do the following. We “group” the vertices on the right side into V_1, \dots, V_n where V_1 has $N_1 d_1$ vertices, V_2 has $N_2 d_2$ vertices, etc. Such grouping is possible because $\bar{M} \cdot \bar{c} = \bar{N} \cdot \bar{d}$.

For each $i \in \{1, \dots, n\}$, we do the following. We merge d_i vertices in V_i into one vertex, so that each vertex in V_i has degree d_i as follows. Let $V_i = \{v_{i,1}, \dots, v_{i,K_i}\}$ where $K_i = N_i d_i$. We merge the vertices $v_1, v_{N_i+1}, v_{2N_i+1}, \dots, v_{(d_i-1)N_i+1}$ into one vertex; the vertices $v_2, v_{N_i+2}, v_{2N_i+2}, \dots, v_{(d_i-1)N_i+2}$ into one vertex; and so on.

After such merging, each vertex in V_i has degree d_i . However, it is possible that after we do the merging, we have “parallel” edges, i.e. more than one edges between two vertices. We are going to “remove” such parallel edges one by one until there are no more parallel edges.

Suppose we have parallel edges between the vertices u and v . We pick an edge (u', v') such that u' is not adjacent to v and v' is not adjacent to u . See the left side of the illustration in Figure 7.2.

Such an edge (u', v') exists since the number of vertices reachable in distance 2 from the vertices u and v is $\leq 2(\bar{c} \cdot \bar{1})(\bar{d} \cdot \bar{1}) + 2$ and the number of vertices is $\bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} \geq 2(\bar{c} \cdot \bar{1})(\bar{d} \cdot \bar{1}) + 3$ and the fact that none of the vertices are of zero degree. (Here we make use of the fact that neither \bar{c} nor \bar{d} contain zero entry.)

Now we delete the edges (u', v') and one of the parallel edge (u, v) , replace it with the edges (u, v') and (u', v) , as illustrated on the right side of the illustration in Figure 7.2. We perform such operation until there are no more parallel edges. This completes the proof of Lemma 7.2. \square




FIG. 7.1. The preliminary graph constructed in the proof of Lemma 7.2.




FIG. 7.2. Eliminating the parallel edges in the proof of Lemma 7.2.

The following theorem is a straightforward application of Lemma 7.2.

THEOREM 7.3. *For every $\bar{c} \in \mathbb{N}^m$ and $\bar{d} \in \mathbb{N}^n$, there exists a Presburger formula $\text{BiREG}_{(\bar{c}, \bar{d})}(\bar{X}, \bar{Y})$, where $\bar{X} = (X_1, \dots, X_m)$ and $\bar{Y} = (Y_1, \dots, Y_n)$ such that the following holds. There exists a (\bar{c}, \bar{d}) -biregular graph of size (\bar{M}, \bar{N}) if and only if the sentence $\text{BiREG}_{(\bar{c}, \bar{d})}(\bar{M}, \bar{N})$ holds.*

Proof. The proof is a direct application of Lemma 7.2. We assume that all the entries in \bar{c} and \bar{d} are not zero. Otherwise, we do the following. Suppose $\bar{c} =$

(c_1, \dots, c_m) , $\bar{d} = (d_1, \dots, d_n)$ and let $I = \{i \mid c_i = 0\}$ and $J = \{j \mid d_j = 0\}$. We define

$$\text{BiREG}_{\bar{c}, \bar{d}}(\bar{X}, \bar{Y}) := \bigwedge_{i \in I} X_i \geq 0 \wedge \bigwedge_{j \in J} Y_j \geq 0 \wedge \text{BiREG}_{\bar{c}', \bar{d}'}(\bar{X}', \bar{Y}'),$$

where \bar{c}' and \bar{X}' are the vectors \bar{c} and \bar{X} without the entries in I , respectively, and \bar{d}' and \bar{Y}' are the vectors \bar{d} and \bar{Y} without the entries in J , respectively.

For $\bar{c} \in \mathbb{N}^m$ and $\bar{d} \in \mathbb{N}^n$ which do not contain zero entry, we define the following set $H_{\bar{c}, \bar{d}}$.

$$H_{\bar{c}, \bar{d}} := \left\{ (\bar{M}, \bar{N}) \mid \begin{array}{l} \bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} \leq 2(\bar{c} \cdot \bar{1})(\bar{d} \cdot \bar{1}) + 2 \text{ and} \\ \text{there exists a } (\bar{c}, \bar{d})\text{-biregular graph of size } (\bar{M}, \bar{N}) \end{array} \right\}$$

Such set can be computed greedily since the number of (\bar{M}, \bar{N}) such that $\bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} \leq 2(\bar{c} \cdot \bar{1})(\bar{d} \cdot \bar{1}) + 2$ is bounded.

Now we define the formula $\text{BiREG}_{\bar{c}, \bar{d}}(\bar{X}, \bar{Y})$ as follows.

$$\left(\bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} \geq 2(\bar{c} \cdot \bar{1})(\bar{d} \cdot \bar{1}) + 3 \wedge (\bar{X} \cdot \bar{c} = \bar{Y} \cdot \bar{d}) \right) \vee \bigvee_{(\bar{M}, \bar{N}) \in H_{\bar{c}, \bar{d}}} \bar{X} = \bar{M} \wedge \bar{Y} = \bar{N}$$

The formula is a Presburger formula since \bar{c} and \bar{d} are constants. Since \bar{c}, \bar{d} do not contain zero entry, we can apply Lemma 7.2 to obtain the correctness of $\text{BiREG}_{\bar{c}, \bar{d}}(\bar{X}, \bar{Y})$. This completes our proof of Theorem 7.3. \square

7.2. When $C \in \mathbb{N}^{\ell \times m}$ and $D \in \mathbb{N}^{\ell \times n}$. Theorem 7.4 below is the generalisation of Theorem 7.3 to the case where $\ell \geq 1$. In the following, for a matrix $C \in \mathbb{N}^{\ell \times m}$, we write $C \cdot \bar{1}$ to denote the sum of all the entries in C .

THEOREM 7.4. *For every $C \in \mathbb{N}^{\ell \times m}$ and $D \in \mathbb{N}^{\ell \times n}$, there exists a Presburger formula $\text{BiREG}_{C, D}(\bar{X}, \bar{Y})$, where $\bar{X} = (X_1, \dots, X_m)$ and $\bar{Y} = (Y_1, \dots, Y_n)$ such that the following holds. There exists a (C, D) -biregular ℓ -type graph of size (\bar{M}, \bar{N}) if and only if the sentence $\text{BiREG}_{C, D}(\bar{M}, \bar{N})$ holds.*

Proof. Let $C \in \mathbb{N}^{\ell \times m}$ and $D \in \mathbb{N}^{\ell \times n}$ be the given matrices. For simplicity, we assume that both C and D do not contain zero column. If column i in matrix C (or, D , respectively) is zero column, then we add the constraint $X_i \geq 0$ (or, $Y_i \geq 0$, respectively) and ignore that column.

Let $\bar{c}_1, \dots, \bar{c}_\ell$ and $\bar{d}_1, \dots, \bar{d}_\ell$ be the row vectors of C and D , respectively. For a vector $\bar{t} = (t_1, \dots, t_m) \in \mathbb{N}^m$, we define the characteristic vector of \bar{t} as $\chi(\bar{t}) := (b_1, \dots, b_m) \in \{0, 1\}^m$ where $b_i = 0$ if $t_i = 0$, and $b_i = 1$ if $t_i \neq 0$.

We first define the following set.

$$H_{C, D} := \left\{ (\bar{M}, \bar{N}) \mid \begin{array}{l} \bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} < 2\ell(C \cdot \bar{1})(D \cdot \bar{1}) + 3\ell \text{ and} \\ \text{there exists a } (C, D)\text{-biregular graph of size } (\bar{M}, \bar{N}) \end{array} \right\}$$

Again, such set can be computed greedily since the number of (\bar{M}, \bar{N}) such that $\bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} < 2\ell(C \cdot \bar{1})(D \cdot \bar{1}) + 3\ell$ is bounded.

Then, the formula $\text{BiREG}_{C, D}(\bar{X}, \bar{Y})$ can be defined inductively as follows. When $\ell = 1$,

$$\text{BiREG}_{C, D}(\bar{X}, \bar{Y}) := \text{BiREG}_{\bar{c}_1, \bar{d}_1}(\bar{X}, \bar{Y})$$

When $\ell \geq 2$,

$$\begin{aligned} \text{BiREG}_{C,D}(\bar{X}, \bar{Y}) := & \bigvee_{(\bar{M}, \bar{N}) \in H_{C,D}} \bar{X} = \bar{M} \wedge \bar{Y} = \bar{N} \\ & \vee \bigvee_{1 \leq j \leq \ell} \left(\bar{X} \cdot \chi(\bar{c}_j) + \bar{Y} \cdot \chi(\bar{d}_j) \geq 2(C \cdot \bar{1})(D \cdot \bar{1}) + 3 \right. \\ & \quad \wedge \text{BiREG}_{C-\bar{c}_j, D-\bar{d}_j}(\bar{X}, \bar{Y}) \\ & \quad \left. \wedge \text{BiREG}_{\bar{c}_j, \bar{d}_j}(\bar{X}, \bar{Y}) \right) \end{aligned}$$

where $C - \bar{c}_j$, $D - \bar{d}_j$ denote the matrices C and D without row j , respectively.

We are going to prove that there exists a (C, D) -biregular graph of size (\bar{M}, \bar{N}) if and only if the statement $\text{BiREG}_{C,D}(\bar{M}, \bar{N})$ holds. The proof is by induction on ℓ . The basis $\ell = 1$ has been established in Theorem 7.3. For the induction step, we assume that it holds for the case of $\ell - 1$ and we are going to prove the case ℓ .

We first prove the “only if” direction. Suppose $G = (U, V, E_1, \dots, E_\ell)$ is (C, D) -biregular of size (\bar{M}, \bar{N}) . If $(\bar{M}, \bar{N}) \in H_{C,D}$, then $\text{BiREG}_{C,D}(\bar{M}, \bar{N})$ holds. So suppose $(\bar{M}, \bar{N}) \notin H_{C,D}$ and $\bar{M} \cdot \bar{1} + \bar{N} \cdot \bar{1} \geq 2\ell(C \cdot \bar{1})(D \cdot \bar{1}) + 3\ell$.

It is not difficult to see that since C and D do not contain zero column, there exists $j \in \{1, \dots, \ell\}$ such that $\bar{M} \cdot \chi(\bar{c}_j) + \bar{N} \cdot \chi(\bar{d}_j) \geq 2(C \cdot \bar{1})(D \cdot \bar{1}) + 3$. Moreover, if $G = (U, V, E_1, \dots, E_\ell)$ is (C, D) -biregular of size (\bar{M}, \bar{N}) , then G is also $(C - \bar{c}_j, D - \bar{d}_j)$ -biregular and (\bar{c}_j, \bar{d}_j) -biregular. By the induction hypothesis, both $\text{BiREG}_{C-\bar{c}_j, D-\bar{d}_j}(\bar{M}, \bar{N})$ and $\text{BiREG}_{\bar{c}_j, \bar{d}_j}(\bar{M}, \bar{N})$ hold.

We now prove the “if” direction. Suppose $\text{BiREG}_{C,D}(\bar{M}, \bar{N})$ holds. If $(\bar{M}, \bar{N}) \in H_{C,D}$, then there exists a (C, D) -biregular graph of size (\bar{M}, \bar{N}) and we are done. So suppose $(\bar{M}, \bar{N}) \notin H_{C,D}$. Hence there exists $j \in \{1, \dots, \ell\}$ such that

$$\begin{aligned} & \bar{M} \cdot \chi(\bar{c}_j) + \bar{N} \cdot \chi(\bar{d}_j) \geq 2(C \cdot \bar{1})(D \cdot \bar{1}) + 3 \\ & \wedge \text{BiREG}_{C-\bar{c}_j, D-\bar{d}_j}(\bar{M}, \bar{N}) \quad \wedge \quad \text{BiREG}_{\bar{c}_j, \bar{d}_j}(\bar{M}, \bar{N}) \end{aligned}$$

For simplicity, we assume that $j = \ell$. By the induction hypothesis, there exists a $(C - \bar{c}_\ell, D - \bar{d}_\ell)$ -biregular graph $G_1 = (U_1, V_1, E_1, \dots, E_{\ell-1})$ of size (\bar{M}, \bar{N}) , and by definition, $E_1, \dots, E_{\ell-1}$ are pairwise disjoint. By Theorem 7.3, there exists a $(\bar{c}_\ell, \bar{d}_\ell)$ -biregular graph $G_2 = (U_2, V_2, E_\ell)$ of size (\bar{M}, \bar{N}) . We can assume that $U_1 = U_2 = U$ and $V_1 = V_2 = V$ since G_1 and G_2 are of the same size (\bar{M}, \bar{N}) .


We are going to combine G_1 and G_2 into one graph to get an ℓ -type (C, D) -biregular graph $G = (U, V, E_1, \dots, E_\ell)$ of size (\bar{M}, \bar{N}) . If $E_\ell \cap (E_1 \cup \dots \cup E_{\ell-1}) = \emptyset$, then the graph $G = (U, V, E_1, \dots, E_\ell)$ is the desired (C, D) -biregular ℓ -type graph of size (\bar{M}, \bar{N}) , and we are done.

Now suppose $E_\ell \cap (E_1 \cup \dots \cup E_{\ell-1}) \neq \emptyset$. We are going to construct another graph $G'_2 = (U, V, E'_\ell)$ such that

$$|E'_\ell \cap (E_1 \cup \dots \cup E_{\ell-1})| < |E_\ell \cap (E_1 \cup \dots \cup E_{\ell-1})|$$

We do this repeatedly until at the end we obtain a graph $G''_2 = (V, E''_\ell)$ such that $E''_\ell \cap (E_1 \cup \dots \cup E_{\ell-1}) = \emptyset$.

Let $(u, v) \in E_\ell \cap (E_1 \cup \dots \cup E_{\ell-1})$. The number of vertices reachable in from u and v within distance 2 (by any of edges in E_1, \dots, E_ℓ) is $\leq 2(C \cdot \bar{1})(D \cdot \bar{1}) + 2$. Since $\bar{M} \cdot \chi(\bar{c}_\ell) + \bar{N} \cdot \chi(\bar{d}_\ell) \geq 2(C \cdot \bar{1})(D \cdot \bar{1}) + 3$, there exists $(u', v') \in E_\ell$ such that $(u, u'), (v, v') \notin E_1 \cup \dots \cup E_{\ell-1}$. See the left side of the illustration below.



Now we define E'_ℓ by deleting the edges $(u, v), (u', v')$ from E_ℓ , while adding the edges $(u, v'), (u', v)$ into E_ℓ . Formally,

$$E'_\ell := (E_\ell - \{(u, v), (u', v')\}) \cup \{(u, v'), (u', v)\}$$

See the right side of the illustration above.

Now it is straightforward that $G' = (U, V, E'_\ell)$ is still a \bar{d}_ℓ -regular graph of size \bar{N} , while

$$|E'_\ell \cap (E_1 \cup \dots \cup E_{\ell-1})| < |E_\ell \cap (E_1 \cup \dots \cup E_{\ell-1})|$$

We perform this operation until $E_{\ell+1} \cap (E_1 \cup \dots \cup E_\ell) = \emptyset$. This completes the proof of Theorem 7.4. \square

7.3. For $C \in \mathbb{B}^{\ell \times n}$ and $D \in \mathbb{B}^{\ell \times m}$ when the number of vertices is “big enough”. Let $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$, where $\bar{c}_1, \dots, \bar{c}_\ell$ and $\bar{d}_1, \dots, \bar{d}_\ell$ are the row vectors of C and D , respectively.

We define the following. Two vectors $\bar{M} = (M_1, \dots, M_m) \in \mathbb{N}^m$ and $\bar{N} = (N_1, \dots, N_n) \in \mathbb{N}^n$ are “big enough” with respect to C, D , if the following inequalities hold.

$$(7.1) \quad \bigwedge_{1 \leq i \leq \ell} \left(\bar{M} \cdot \chi(\bar{c}_i) + \bar{N} \cdot \chi(\bar{d}_i) \geq 2(C \cdot \bar{1})(D \cdot \bar{1}) + 3 \right)$$

$$(7.2) \quad \wedge \bigwedge_{1 \leq i \leq \ell} \left(\left(\sum_{j \text{ such that } D_{i,j} \in \blacktriangleright \mathbb{N}} N_j \right) \geq \max_{1 \leq j \leq m} (C_{i,j}) \right)$$

$$(7.3) \quad \wedge \bigwedge_{1 \leq i \leq \ell} \left(\left(\sum_{j \text{ such that } C_{i,j} \in \blacktriangleright \mathbb{N}} M_j \right) \geq \max_{1 \leq j \leq m} (D_{i,j}) \right)$$

We also need a few auxiliary notations. In the following for a positive integer ℓ , \mathbf{I}_ℓ denotes the $(\ell \times \ell)$ identity matrix. For $\blacktriangleright d \in \blacktriangleright \mathbb{N}$, we write $\lceil \blacktriangleright d \rceil$ to denote the number d . By default, we set $\lceil d \rceil = d$. We define the $+$ operations on \mathbb{B} as follows.

$$\begin{aligned} d_1 + d_2 &= (d_1 + d_2) \\ \blacktriangleright d_1 + d_2 &= d_1 + \blacktriangleright d_2 = \blacktriangleright d_1 + \blacktriangleright d_2 = \blacktriangleright (d_1 + d_2) \end{aligned}$$

We extend $\lceil \cdot \rceil$ and $+$ to vectors and matrices over \mathbb{B} in the natural way, where they are applied componentwise. For two vectors $\bar{t}_1, \bar{t}_2 \in \mathbb{B}^m$, we define the dot product $\bar{t}_1 \cdot \bar{t}_2$ as $\lceil \bar{t}_1 \rceil \cdot \lceil \bar{t}_2 \rceil$. For a matrix $D \in \mathbb{B}^{\ell \times m}$, we write $D \cdot \bar{1}$ to denote the sum $\sum_{i,j} \lceil D_{i,j} \rceil$.

The lemma below characterises the existence of (C, D) -biregular graph of size (\bar{M}, \bar{N}) , where \bar{M}, \bar{N} are big enough with respect to (C, D) and that for every row i , either the row- i of C or of D contains only elements from \mathbb{N} .

LEMMA 7.5. Let $C \in \begin{pmatrix} C^{(1)} \\ C^{(2)} \\ C^{(3)} \end{pmatrix} \in \mathbb{B}^{\ell \times m}$ and $D \in \begin{pmatrix} D^{(1)} \\ D^{(2)} \\ D^{(3)} \end{pmatrix} \in \mathbb{B}^{\ell \times n}$ where

$\ell = \ell_1 + \ell_2 + \ell_3$ and

- $C^{(1)} \in \mathbb{N}^{\ell_1 \times m}$ and $D^{(1)} \in \mathbb{N}^{\ell_1 \times n}$;
- $C^{(2)} \in \mathbb{N}^{\ell_2 \times m}$ and $D^{(2)} \in \mathbb{B}^{\ell_2 \times n}$ and every row in $D^{(2)}$ contains an element of $\blacktriangleright \mathbb{N}$;
- $C^{(3)} \in \mathbb{B}^{\ell_3 \times m}$ and $D^{(3)} \in \mathbb{N}^{\ell_3 \times n}$ and every row in $C^{(3)}$ contains an element of $\blacktriangleright \mathbb{N}$.

Let $\bar{M} \in \mathbb{N}^m$ and $\bar{N} \in \mathbb{N}^n$ be big enough with respect to C and D . Then, the following holds. There exists a (C, D) -biregular graph of size (\bar{M}, \bar{N}) if and only if the statement $\text{BiREG}_{C', D'}((\bar{M}, K_1, \dots, K_{\ell_3}), (\bar{N}, L_1, \dots, L_{\ell_2}))$ holds, where

$$\bullet C' = \left(\begin{array}{c|c} C^{(1)} & 0 \\ \hline C^{(2)} & 0 \\ \hline [C^{(3)}] & \mathbf{I}_{\ell_3} \end{array} \right) \in \mathbb{N}^{\ell \times (m + \ell_3)} \text{ and } D' = \left(\begin{array}{c|c} D^{(1)} & 0 \\ \hline [D^{(2)}] & \mathbf{I}_{\ell_2} \\ \hline D^{(3)} & 0 \end{array} \right) \in \mathbb{N}^{\ell \times (n + \ell_2)}$$

- each $K_i = \bar{d}_{\ell_1 + \ell_2 + i} \cdot \bar{N} - \lfloor \bar{c}_{\ell_1 + \ell_2 + i} \rfloor \cdot \bar{M}$,
- each $L_i = \bar{c}_{\ell_1 + i} \cdot \bar{M} - \lfloor \bar{d}_{\ell_1 + i} \rfloor \cdot \bar{N}$.

Proof. Let $C \in \begin{pmatrix} C^{(1)} \\ C^{(2)} \\ C^{(3)} \end{pmatrix} \in \mathbb{B}^{\ell \times m}$ and $D \in \begin{pmatrix} D^{(1)} \\ D^{(2)} \\ D^{(3)} \end{pmatrix} \in \mathbb{B}^{\ell \times n}$ and ℓ_1, ℓ_2, ℓ_3 ,

$\bar{M} \in \mathbb{N}^m$ and $\bar{N} \in \mathbb{N}^n$ be as in the premises. We also assume that $\bar{c}_1, \dots, \bar{c}_\ell$ and $\bar{d}_1, \dots, \bar{d}_\ell$ are the row vectors of C and D , respectively.

Before we present our proof, we have to remark here that we do not need the condition that \bar{M} and \bar{N} are big enough to establish the “only if” direction. For the “if” direction, we only need Inequalities 7.2 and 7.3. Inequality 7.1 is needed only to establish Theorem 7.6.

We start with the “only if” direction. Suppose $G = (U, V, E_1, \dots, E_\ell)$ is a (C, D) -biregular graph of size (\bar{M}, \bar{N}) . This means there exist a partition $U = U_1 \cup \dots \cup U_m$ and a partition $V = V_1 \cup \dots \cup V_n$ such that for each $i = 1, \dots, \ell$,

- for each $j = 1, \dots, m$, for each $u \in U_j$, $\deg_{E_i}(u) = C_{i,j}$;
- for each $j = 1, \dots, n$, for each $v \in V_j$, $\deg_{E_i}(v) = D_{i,j}$.

Now, the following holds.

- For each $i = \ell_1 + 1, \dots, \ell_1 + \ell_2$, the number of E_i -edges in G is $\bar{c}_i \cdot \bar{M}$, which should be greater than $\lfloor \bar{d}_i \rfloor \cdot \bar{N}$. We set $L_{i - \ell_1} = \bar{c}_i \cdot \bar{M} - \lfloor \bar{d}_i \rfloor \cdot \bar{N}$.
- For each $i = \ell_1 + \ell_2 + 1, \dots, \ell_1 + \ell_2 + \ell_3$, the number of E_i -edges in G is $\bar{d}_i \cdot \bar{N}$, which should be greater than $\lfloor \bar{c}_i \rfloor \cdot \bar{M}$. We set $K_{i - \ell_1 - \ell_2} = \bar{d}_i \cdot \bar{N} - \lfloor \bar{c}_i \rfloor \cdot \bar{M}$.

Let C', D' be as defined in the lemma, and $\bar{K} = (K_1, \dots, K_{\ell_3})$ and $\bar{L} = (L_1, \dots, L_{\ell_2})$. We construct a (C', D') -biregular graph of size $((\bar{M}, \bar{K}), (\bar{N}, \bar{L}))$ as follows.

- For each $i = \ell_1 + 1, \dots, \ell_1 + \ell_2$, for each $j = 1, \dots, n$, for each vertex $v \in V_j$, if $\deg_{E_i}(v) > \lfloor D_{i,j} \rfloor$, then we “split” v into v_0, v_1, \dots, v_k vertices, where
 - $k = \deg_{E_i}(v) - \lfloor D_{i,j} \rfloor$,
 - $\deg_{E_i}(v_0) = \lfloor D_{i,j} \rfloor$, and for each $i' \neq i$, $\deg_{E_{i'}}(v_0) = \deg_{E_{i'}}(v)$,
 - $\deg_{E_i}(v_1) = \deg_{E_i}(v_2) = \dots = \deg_{E_i}(v_k) = 1$, and for each $i' \neq i$, $\deg_{E_{i'}}(v_1) = \deg_{E_{i'}}(v_2) = \dots = \deg_{E_{i'}}(v_k) = 0$.
- Similarly, for each $i = \ell_1 + \ell_2 + 1, \dots, \ell_1 + \ell_2 + \ell_3$, for each $j = 1, \dots, m$, for each vertex $u \in U_j$, if $\deg_{E_i}(u) > \lfloor C_{i,j} \rfloor$, then we “split” u into u_0, u_1, \dots, u_k

vertices, where

- $k = \deg_{E_i}(u) - \lfloor D_{i,j} \rfloor$,
- $\deg_{E_i}(u_0) = \lfloor D_{i,j} \rfloor$, and for each $i' \neq i$, $\deg_{E_{i'}}(u_0) = \deg_{E_{i'}}(u)$,
- $\deg_{E_i}(u_1) = \deg_{E_i}(u_2) = \dots = \deg_{E_i}(u_k) = 1$, and for each $i' \neq i$,
 $\deg_{E_{i'}}(u_1) = \deg_{E_{i'}}(u_2) = \dots = \deg_{E_{i'}}(u_k) = 0$.

It should be obvious that the resulting graph is a (C', D') -biregular graph of size $((\bar{M}, \bar{K}), (\bar{N}, \bar{L}))$.

Now we prove the “if” direction. Suppose $\text{BiREG}_{C', D'}((\bar{M}, \bar{K}), (\bar{N}, \bar{L}))$ holds, where each $L_i = \bar{c}_i \cdot \bar{M} - \lfloor \bar{d}_i \rfloor \cdot \bar{N}$ and $K_i = \bar{d}_i \cdot \bar{N} - \lfloor \bar{c}_i \rfloor \cdot \bar{M}$.

By Theorem 7.4, there exists a (C', D') -biregular graph G of size $((\bar{M}, \bar{K}), (\bar{N}, \bar{L}))$. Let $G = (U \cup A, V \cup B, E_1, \dots, E_\ell)$, where $U \cup A = U_1 \cup \dots \cup U_m \cup A_1 \cup \dots \cup A_{\ell_3}$ and $V \cup B = V_1 \cup \dots \cup V_n \cup B_1 \cup \dots \cup B_{\ell_2}$ are the witness of the (C', D') -biregularity.

To construct a (C, D) -biregular graph of size (\bar{M}, \bar{N}) , we do the following. For each vertex $u \in U$ adjacent by E_i -edges to, say, s vertices in B , we pick s vertices v_1, \dots, v_s from the set

$$\bigcup_{j \text{ such that } d_{i,j} \in \blacktriangleright \mathbb{N}} V_j$$

Such s vertices exist since by Inequality 7.2, $\sum_{j \text{ such that } d_{i,j} \in \blacktriangleright \mathbb{N}} N_j$ is $\geq \max(\bar{c}_i) \geq \deg(u)$. We delete those s vertices in B , and connect u to each of v_1, \dots, v_s by E_i -edges. We do this until the set B is empty. Similarly, by Inequality 7.3, we can perform similar operations until the set A is empty. The resulting graph is a (C, D) -biregular graph of size (\bar{M}, \bar{N}) . This completes the proof of Lemma 7.5. \square

Now Lemma 7.5 tells us the Presburger formula $\text{BiREG}_{C, D}$ for a pair of matrices satisfying the assumption given in Lemma 7.5. More formally, let $C \in \begin{pmatrix} C^{(1)} \\ C^{(2)} \\ C^{(3)} \end{pmatrix} \in$

$\mathbb{B}^{\ell \times m}$ and $D \in \begin{pmatrix} D^{(1)} \\ D^{(2)} \\ D^{(3)} \end{pmatrix} \in \mathbb{B}^{\ell \times n}$ where $\ell = \ell_1 + \ell_2 + \ell_3$ and

- $C^{(1)} \in \mathbb{N}^{\ell_1 \times m}$ and $D^{(1)} \in \mathbb{N}^{\ell_1 \times n}$;
- $C^{(2)} \in \mathbb{N}^{\ell_2 \times m}$ and $D^{(2)} \in \mathbb{B}^{\ell_2 \times n}$ and every row in $D^{(2)}$ contains an element of $\blacktriangleright \mathbb{N}$;
- $C^{(3)} \in \mathbb{B}^{\ell_3 \times m}$ and $D^{(3)} \in \mathbb{N}^{\ell_3 \times n}$ and every row in $C^{(3)}$ contains an element of $\blacktriangleright \mathbb{N}$.

That is, for such C, D , we define $\text{BiREG}_{C, D}(\bar{X}, \bar{Y})$ as follows.

$$(7.4) \quad \text{BiREG}_{C, D}(\bar{X}, \bar{Y}) := \exists Z_1 \dots \exists Z_{\ell_3} \exists Z'_1 \dots \exists Z'_{\ell_2} \\ \text{BiREG}_{C', D'}(\bar{X}, Z_1, \dots, Z_{\ell_3}, \bar{Y}, Z'_1, \dots, Z'_{\ell_2})$$

where $C' = \left(\begin{array}{c|c} C^{(1)} & 0 \\ \hline C^{(2)} & 0 \\ \hline \lfloor C^{(3)} \rfloor & I_{\ell_3} \end{array} \right) \in \mathbb{N}^{\ell \times (m + \ell_3)}$ and $D' = \left(\begin{array}{c|c} D^{(1)} & 0 \\ \hline \lfloor D^{(2)} \rfloor & I_{\ell_2} \\ \hline D^{(3)} & 0 \end{array} \right) \in \mathbb{N}^{\ell \times (n + \ell_2)}$.

Since C', D' consist of entirely \mathbb{N} entries, $\text{BiREG}_{C', D'}$ is defined as in Theorem 7.4. Intuitively, the variables Z_1, \dots, Z_{ℓ_3} are to capture the values K_1, \dots, K_{ℓ_3} and Z'_1, \dots, Z'_{ℓ_2} the values L_1, \dots, L_{ℓ_2} , as stated in Lemma 7.5.

Using this, we can prove the following theorem.

THEOREM 7.6. *For every $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$, there exists a Presburger formula $\widetilde{\text{BiREG}}_{C, D}(\bar{X}, \bar{Y})$ such that for every \bar{M}, \bar{N} big enough with respect to C, D ,*

the following holds. There exists a (C, D) -biregular graph of size (\bar{M}, \bar{N}) if and only if the statement $\widetilde{\text{BiREG}}_{C,D}(\bar{M}, \bar{N})$ holds.

Proof. Let $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$, where $\bar{c}_1, \dots, \bar{c}_\ell$ and $\bar{d}_1, \dots, \bar{d}_\ell$ are the row vectors of C and D , respectively.

We need an additional notation. For a set $I \subseteq \{1, \dots, \ell\}$, we write $C(I)$ be the matrix C' , in which each row vector \bar{c}'_i is defined as $\bar{c}'_i = [\bar{c}_i]$, if $i \in I$, and $\bar{c}'_i = \bar{c}_i$, if $i \notin I$. We can define $D(I)$ similarly.

We define the formula $\text{BiREG}_{C,D}(\bar{X}, \bar{Y})$ as follows.

$$\widetilde{\text{BiREG}}_{C,D}(\bar{X}, \bar{Y}) := \bigvee_{I_0, I_1, I_2} \left(\begin{array}{l} \bigwedge_{i \in I_0} \bar{X} \cdot [\bar{c}_i] = \bar{Y} \cdot [\bar{d}_i] \quad \wedge \\ \bigwedge_{i \in I_1} \bar{X} \cdot [\bar{c}_i] > \bar{Y} \cdot [\bar{d}_i] \quad \wedge \\ \bigwedge_{i \in I_2} \bar{X} \cdot [\bar{c}_i] < \bar{Y} \cdot [\bar{d}_i] \quad \wedge \\ \text{BiREG}_{C(I_0 \cup I_2), D(I_0 \cup I_1)}(\bar{X}, \bar{Y}) \end{array} \right)$$

where each $\text{BiREG}_{C(I_0 \cup I_2), D(I_0 \cup I_1)}(\bar{X}, \bar{Y})$ is as defined in Equation 7.4 and I_0, I_1, I_2 range over the partition $I_0 \cup I_1 \cup I_2 = \{1, 2, \dots, \ell\}$. Obviously, for every partition $I_0 \cup I_1 \cup I_2 = \{1, \dots, \ell\}$, on each row $i = 1, \dots, \ell$, either row- i from $C(I_0 \cup I_2)$, or row- i from $D(I_0 \cup I_1)$, or row- i from both consists entirely of \mathbb{N} . (Our intention is the application of Lemma 7.5 later on.)

We are going to prove that the formula $\widetilde{\text{BiREG}}_{C,D}(\bar{X}, \bar{Y})$ is the desired formula. The “if” direction follows from Lemma 7.5 and that every $C(I_0 \cup I_2), D(I_0 \cup I_1)$ -biregular graph is obviously also a (C, D) -biregular graph.

Now we prove the “only if” direction. Suppose \bar{M}, \bar{N} are big enough for C, D . Let $G = (U, V, E_1, \dots, E_\ell)$ be a (C, D) -biregular graph of size (\bar{M}, \bar{N}) , where $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ are the witnesses of the C, D -biregularity.

We pick the following partition $I_0 \cup I_1 \cup I_2 = \{1, \dots, \ell\}$.

$$\begin{aligned} I_0 &= \{i \mid \bar{M} \cdot [\bar{c}_i] = \bar{M} \cdot [\bar{d}_i]\} \\ I_1 &= \{i \mid \bar{M} \cdot [\bar{c}_i] > \bar{M} \cdot [\bar{d}_i]\} \\ I_2 &= \{i \mid \bar{M} \cdot [\bar{c}_i] < \bar{M} \cdot [\bar{d}_i]\} \end{aligned}$$

We are going to convert the graph G into $(C(I_0 \cup I_1), D(I_0, I_2))$ -biregular graph in which $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ are also the witness of the $(C(I_0 \cup I_1), D(I_0, I_2))$ -biregularity. This, together with Lemma 7.5, implies that $\text{BiREG}_{C(I_0 \cup I_1), D(I_0, I_2)}(\bar{M}, \bar{N})$ holds, and hence, our theorem.

If G is already a $(C(I_0 \cup I_1), D(I_0, I_2))$ -biregular graph, then we are done. Suppose that G is not. We do the following three stages.

Stage 1. We assume that the following holds for the graph G . For every edge $(u, v) \in E_i$ in G , either

$$(7.5) \quad \deg_{E_i}(u) = \lfloor C_{i,j} \rfloor \quad \text{or} \quad \deg_{E_i}(v) = \lfloor D_{i,k} \rfloor.$$

This can be achieved by doing the following. Suppose there is an edge $(u, v) \in E_i$ such that $\deg_{E_i}(u) > \lfloor C_{i,j} \rfloor$ and $\deg_{E_i}(v) > \lfloor D_{i,k} \rfloor$. Since G is (C, D) -biregular, this means that $C_{i,j}, D_{i,k} \in \blacktriangleright \mathbb{N}$.

Deleting the edge (u, v) , we still have $\deg_{E_i}(u) \geq \lfloor C_{i,j} \rfloor$ and $\deg_{E_i}(v) \geq \lfloor D_{i,k} \rfloor$, and hence G is still (C, D) -biregular with $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ being the witness of the (C, D) -biregularity. We repeatedly do this until the graph G satisfies condition (7.5).

Stage 2. We construct a graph $G' = (U \cup S, V \cup T, E'_1, \dots, E'_\ell)$, where for every $i = 1, \dots, \ell$,

- for every $j = 1, \dots, m$, for each $u \in U_j$, $\deg_{E'_i}(u) = \lfloor C_{i,j} \rfloor$,
- for every $s \in S$, $\deg(s) = 1$,^{††}
- for every $k = 1, \dots, n$, for each $v \in V_j$, $\deg_{E'_i}(v) = \lfloor D_{i,j} \rfloor$,
- for every $t \in T$, $\deg(t) = 1$.

The graph G' can be obtained by doing the same trick as in the proof of Lemma 7.5. For every vertex $u \in U_j$, if $\deg_{E_i}(u) - \lfloor C_{i,j} \rfloor = z > 0$, then we “split” u into $z + 1$ vertices u', s_1, \dots, s_z , where

- $\deg_{E_i}(u') = \lfloor C_{i,j} \rfloor$, for all other $h \neq i$, $\deg_{E_h}(u') = \deg_{E_h}(u)$;
- $\deg_{E_i}(s_1) = \dots = \deg_{E_i}(s_z) = 1$, and for all other $h \neq i$, $\deg_{E_h}(s_1) = \dots = \deg_{E_h}(s_z) = 0$.

We can do similar operation to the vertices in V_k . Since G satisfies condition 7.5, there is no edge between vertices in S and T . We also further partition $S = S_1 \cup \dots \cup S_\ell$ and $T = T_1 \cup \dots \cup T_\ell$, where each S_i and T_i contains the vertices whose $\deg_{E_i} = 1$.

Stage 3. Stage 3 is as follows. For each $i = 1, \dots, \ell$, if there are an edge $(s, v) \in E_i$ and an edge $(u, t) \in E_i$, for some $s \in S_i$, $v \in V_k$, $u \in U_j$, $t \in T_i$, we do the following.

- We delete the two edges (s, v) and (u, t) from E_i , as well as the vertices s and t .
- We add an edge (u, v) into E_i .
- If there is already an existing edge $(u, v) \in E_1 \cup \dots \cup E_\ell$, adding another (u, v) may result in “parallel” edges. However, since \bar{M}, \bar{N} is big enough with respect to C, D , and in particular, Inequality 7.1 holds, we can apply the same trick as in the proof of Theorem 7.4 to get rid of the parallel edge, while preserving the degree of the vertices.

We repeatedly do this until for each $i = 1, \dots, \ell$ either $S_i = \emptyset$, or $T_i = \emptyset$. In particular, the following holds.

- If $i \in I_0$, then $S_i = T_i = \emptyset$.
Recall that $i \in I_0$ means that $\bar{M} \cdot \lfloor \bar{c}_i \rfloor = \bar{N} \cdot \lfloor \bar{d}_i \rfloor$, which implies that the initial sets S_i, T_i have the same cardinality. Since we always delete a pair of vertices s, t from S_i, T_i , respectively, we have at the end $S_i = T_i = \emptyset$.
- Likewise, if $i \in I_1$, then $S_i = \emptyset$.
This is because $\bar{M} \cdot \lfloor \bar{c}_i \rfloor > \bar{N} \cdot \lfloor \bar{d}_i \rfloor$, implies that initially $|T_i| > |S_i|$, which further implies that at the end $S_i = \emptyset$.

By symmetrical reasoning, if $i \in I_2$, then $T_i = \emptyset$.

From here, we will “merge” back the vertices in T with vertices in V . This is done as follows. For each vertex $u \in U$ adjacent by E_i -edges to, say, z vertices in T , we pick z vertices v_1, \dots, v_z from the set

$$\bigcup_{j \text{ such that } d_{i,j} \in \mathbb{N}} V_j$$

Such z vertices exist by Inequality 7.2 (because \bar{M}, \bar{N} are big enough w.r.t. C, D). We delete those z vertices in T , and connect u to each of v_1, \dots, v_z by E_i -edges. We do this until the set T is empty. In a similar manner, we can merge back the vertices in S with vertices in U , where the existence of the vertices v_1, \dots, v_z is guaranteed by Inequality 7.3.

The resulting graph is $(C(I_0 \cup I_1), D(I_0, I_2))$ -biregular graph, which by Lemma 7.5 the formula implies that $\text{BiREG}_{C(I_0 \cup I_2), D(I_0 \cup I_1)}(\bar{M}, \bar{N})$ holds. This completes our

^{††}Recall that $\deg(s) = \deg_{E'_1}(s) + \dots + \deg_{E'_\ell}(s)$. Hence, $\deg(s) = 1$ means that there is only one edge adjacent to s .

proof of Theorem 7.6. \square

7.4. The notion of partial bipartite graphs. In this subsection we are going to generalise Theorem 7.6 to the case when it is possible that one of the inequalities 7.1 and 7.2 does not hold. The idea is that those numbers (for which the inequalities do not hold) are encoded into the Presburger formula. For this, we introduce the notion of *partial graph*.

An ℓ -type *partial bipartite graph* is a tuple $\mathcal{P} = (C, D, S, T, f, g)$, where

- $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$,
- S is a finite set of vertices (possibly empty),
- T is a finite set of vertices (possibly empty),
- $f : S \times \{E_1, \dots, E_\ell\} \rightarrow \mathbb{B}$,
- $g : T \times \{E_1, \dots, E_\ell\} \rightarrow \mathbb{B}$.

Obviously, if S or T is empty, then f or g , respectively, is also an “empty” function. In the following the term *partial graph* always means partial bipartite graph.

A completion of the partial graph $\mathcal{P} = (C, D, S, T, f, g)$ is a bipartite graph $G = (U \cup S, V \cup T, E_1, \dots, E_\ell)$ such that there is a partition $U_1 \cup \dots \cup U_m$ of U and a partition $V_1 \cup \dots \cup V_n$ of V such that

- for every $u \in U_j$, $\deg_{E_i}(u) = C_{i,j}$,
- for every $v \in V_j$, $\deg_{E_i}(v) = D_{i,j}$,
- for every $s \in S$, $\deg_{E_i}(s) = f(s, E_i)$,
- for every $t \in T$, $\deg_{E_i}(t) = g(t, E_i)$.

When it is clear from the context, we also call $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ the witness of the (C, D) -biregularity. Note that when both S and T are empty, then the completions of the partial graph \mathcal{P} are simply (C, D) -biregular graphs.

We need a few additional notations.

$$\blacktriangleright c - d = \begin{cases} \blacktriangleright(c - d) & \text{if } c \geq d \\ \blacktriangleright 0 & \text{otherwise} \end{cases}$$

Let $C \in \mathbb{B}^{\ell \times m}$. We define a matrix $\xi(C) \in \mathbb{B}^{\ell \times (\ell+1)m}$ as follows.

$$\xi(C) := \left(C \mid M_1 \mid \dots \mid M_m \right)$$

where each M_i is the matrix obtained by repeating the i th column vector of C for ℓ number of times, and subtracting the identity matrix \mathbf{I}_ℓ . Formally,

$$M_i := \begin{pmatrix} C_{1,i} & C_{1,i} & \cdots & C_{1,i} \\ C_{2,i} & C_{2,i} & \cdots & C_{2,i} \\ \vdots & \vdots & \ddots & \vdots \\ C_{\ell,i} & C_{\ell,i} & \cdots & C_{\ell,i} \end{pmatrix} - \mathbf{I}_\ell$$

Lemma 7.7 below essentially states that every partial graph can be reduced into a “smaller” partial graph with the addition of some linear equalities.

LEMMA 7.7. *Let $\mathcal{P} = (C, D, S, T, f, g)$ be a partial graph, where $T \neq \emptyset$. Let $t \in T$. Then the following holds.*

- (1) *For every completion graph $G = (U \cup S, V \cup T, E_1, \dots, E_\ell)$ of the partial graph \mathcal{P} with $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ being the witness of the (C, D) -biregularity, there exists a completion graph $G' = (U \cup S, (V \cup T) \setminus \{t\}, E'_1, \dots, E'_\ell)$*

of the partial graph $\mathcal{P}' = (\xi(C), D, S, T \setminus \{t\}, f, g')$, with the witness of the $(\xi(C), D)$ -biregularity being

$$\begin{aligned} U &= U'_1 \cup \dots \cup U'_m \cup \\ &\quad (U'_{1,1} \cup \dots \cup U'_{\ell,1}) \cup (U'_{1,2} \cup \dots \cup U'_{\ell,2}) \cup \dots \cup (U'_{1,m} \cup \dots \cup U'_{\ell,m}) \\ V &= V_1 \cup \dots \cup V_n \end{aligned}$$

and

$$\begin{aligned} |U_j| &= |U'_j| + |U'_{1,j}| + \dots + |U'_{\ell,j}| && \text{for each } j = 1, \dots, m \\ \sum_{1 \leq j \leq m} |U'_{i,j}| &= g(t, E_i) && \text{for each } i = 1, \dots, \ell. \end{aligned}$$

(2) Vice versa, for every completion graph $G' = (U \cup S, (V \cup T) \setminus \{t\}, E'_1, \dots, E'_\ell)$ of the partial graph $\mathcal{P}' = (\xi(C), D, S, T \setminus \{t\}, f, g')$, with the witness of the $(\xi(C), D)$ -biregularity being

$$\begin{aligned} U &= U'_1 \cup \dots \cup U'_m \cup \\ &\quad (U'_{1,1} \cup \dots \cup U'_{\ell,1}) \cup (U'_{1,2} \cup \dots \cup U'_{\ell,2}) \cup \dots \cup (U'_{1,m} \cup \dots \cup U'_{\ell,m}) \\ V &= V_1 \cup \dots \cup V_n \end{aligned}$$

and for each $i = 1, \dots, \ell$, $\sum_{1 \leq j \leq m} |U'_{i,j}| = g(t, E_i)$, there exists a completion graph $G = (U \cup S, V \cup T, E_1, \dots, E_\ell)$ of the partial graph \mathcal{P} with $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ being the witness of the (C, D) -biregularity, and

$$|U_j| = |U'_j| + |U'_{1,j}| + \dots + |U'_{\ell,j}| \quad \text{for each } j = 1, \dots, m.$$

Proof. Let $\mathcal{P} = (C, D, S, T, f, g)$ be a partial graph, where $T \neq \emptyset$ and $t \in T$. First, we prove part (1). Let $G = (U \cup S, V \cup T, E_1, \dots, E_\ell)$ be a completion graph of \mathcal{P} with $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ being the witness of the (C, D) -biregularity.

For each $j = 1, \dots, m$, we partition U_j into

$$U_j = U'_j \cup (U'_{1,j} \cup \dots \cup U'_{\ell,j}),$$

where

- U'_j be the set of vertices in U_j that are not adjacent to the vertex t ,
- for each $i = 1, \dots, \ell$, $U'_{i,j}$ is the set of vertices in U_j adjacent to t via E_i -edges.

Now deleting the vertex t and all its adjacent edges, we obtain the desired completion graph $G' = (U \cup S, (V \cup T) \setminus \{t\}, E'_1, \dots, E'_\ell)$ of $\mathcal{P}' = (\xi(C), D, S, T \setminus \{t\}, f, g')$.

Now we prove part (2). Let $G' = (U \cup S, (V \cup T) \setminus \{t\}, E'_1, \dots, E'_\ell)$ be a completion of the partial graph $\mathcal{P}' = (\xi(C), D, S, T \setminus \{t\}, f, g')$, with the witness of the $(\xi(C), D)$ -biregularity being

$$\begin{aligned} U &= U'_1 \cup \dots \cup U'_m \cup \\ &\quad (U'_{1,1} \cup \dots \cup U'_{\ell,1}) \cup (U'_{1,2} \cup \dots \cup U'_{\ell,2}) \cup \dots \cup (U'_{1,m} \cup \dots \cup U'_{\ell,m}) \\ V &= V_1 \cup \dots \cup V_n \end{aligned}$$

and for each $i = 1, \dots, \ell$, $\sum_{1 \leq j \leq m} |U'_{i,j}| = g(t, E_i)$.

The desired completion graph $G = (U \cup S, V \cup T, E_1, \dots, E_\ell)$ of the partial graph \mathcal{P} can be obtained as follows. We put the vertex t back inside T . Then, for each

$i = 1, \dots, \ell$ and for each $j = 1, \dots, m$, we connect t with every vertex $u \in U_{i,j}'$ with E_i -edge. This way we obtain the completion graph G with $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ being the witness of the (C, D) -biregularity, and

$$|U_i| = |U_i'| + |U_{1,i}'| + \dots + |U_{\ell,i}'| \quad \text{for each } i = 1, \dots, m.$$

This completes our proof of Lemma 7.7. \square

Following Lemma 7.7 above, we show that every partial graph can be translated into a Presburger formula that captures any of its completion, as stated in the following theorem.

THEOREM 7.8. *For every partial graph $\mathcal{P} = (C, D, S, T, f, g)$, we can construct a Presburger formula $\Psi_{\mathcal{P}}(\bar{X}, \bar{Y})$ such that for every \bar{M} and \bar{N} big enough w.r.t. C, D , the following holds. There exists a completion graph $G = (U \cup S, V \cup T, E_1, \dots, E_{\ell})$, such that $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ and $\bar{M} = (|U_1|, \dots, |U_m|)$ and $\bar{N} = (|V_1|, \dots, |V_n|)$ if and only if $\Psi_{\mathcal{P}}(\bar{M}, \bar{N})$ holds.*

Proof. Let $\mathcal{P} = (C, D, S, T, f, g)$ be a partial graph. If the matrix C is empty, there are only finitely many completion of \mathcal{P} . In this case $\Psi_{\mathcal{P}}$ simply contains the enumeration of the sizes of all possible completions of \mathcal{P} . We can define $\Psi_{\mathcal{P}}$ in a similar manner when D is empty.

Now suppose both the matrices C and D are not empty. The construction of $\Psi_{\mathcal{P}}$ is done inductively as follows. The base case is $S \cup T = \emptyset$, in which case $\Psi_{\mathcal{P}}$ is defined as follows.

$$\Psi_{\mathcal{P}}(\bar{X}, \bar{Y}) := \widetilde{\text{BiREG}}_{C,D}(\bar{X}, \bar{Y}),$$

where $\widetilde{\text{BiREG}}_{C,D}(\bar{X}, \bar{Y})$ is as defined in Theorem 7.6.

Towards the induction step, let $S \cup T \neq \emptyset$. Suppose $T \neq \emptyset$ and $t \in T$. (The case when $S \neq \emptyset$ can be handled in a symmetrical manner.)

We define $\Psi_{\mathcal{P}}(\bar{X}, \bar{Y})$ as follows.

$$\begin{aligned} \Psi_{\mathcal{P}}(\bar{X}, \bar{Y}) &:= \exists Z_1 \dots \exists Z_m \exists Z_{1,1} \dots \exists Z_{\ell,1} \exists Z_{1,2} \dots \exists Z_{\ell,2} \dots \exists Z_{1,m} \dots \exists Z_{\ell,m} \\ &\wedge \bigwedge_{1 \leq i \leq m} X_i = Z_i + Z_{1,i} + \dots + Z_{\ell,i} \\ &\wedge \bigwedge_{1 \leq i \leq \ell} \sum_{1 \leq j \leq m} Z_{i,j} = g(t, E_i) \\ &\wedge \Psi_{\mathcal{P}'}((Z_1, \dots, Z_m, Z_{1,1}, \dots, Z_{\ell,1}, \dots, Z_{1,m}, \dots, Z_{\ell,m}), \bar{Y}) \end{aligned}$$

where $\mathcal{P}' = (\xi(C), D, S, T \setminus \{t\}, f, g')$ and g' is the function g restricted to $T \setminus \{t\}$.

By Theorem 7.6 in the previous section, the correctness of the base case is established. The induction step follows from Lemma 7.7, and hence, shows that the formula $\Psi_{\mathcal{P}}$ is the desired formula. This completes our proof of Theorem 7.8. \square

7.5. Constructing the formula $\text{BiREG}_{C,D}(\bar{X}, \bar{Y})$ for Theorem 5.1. We need the following notions. Let $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$. We say that a partial graph $\mathcal{P} = (C', D', S, T, f, g)$ is compatible with (C, D) with respect to a subset $I \subseteq \{1, \dots, m\}$ and a subset $J \subseteq \{1, \dots, n\}$, and the partitions $S = S_1 \cup \dots \cup S_{m''}$ and $T = T_1 \cup \dots \cup T_{n''}$, if the following four conditions hold.

- $C' \in \mathbb{B}^{\ell \times m'}$ is obtained by deleting the columns I in C .
- $D' \in \mathbb{B}^{\ell \times n'}$ is obtained by deleting the columns J in D .

- Let $C'' \in \mathbb{B}^{\ell \times m''}$ be the matrix whose columns are the columns I in C where $m'' = m - m'$. The matrix C'' is simply the matrix form of the function f . That is, for each $k = 1, \dots, m''$, for every vertex $s \in S_k$, for every $i = 1, \dots, \ell$, $f(s, E_i) = C''_{i,k}$.
- Let $D'' \in \mathbb{B}^{\ell \times n''}$ be the matrix whose columns are the columns J in D where $n'' = n - n'$. The matrix D'' is simply the matrix form of the function g . That is, for each $k = 1, \dots, n''$, for every vertex $t \in T_k$, for every $i = 1, \dots, \ell$, $g(t, E_i) = D''_{i,k}$.

For a subset $I \subseteq \{1, \dots, m\}$, and a variable vector $\bar{X} = (X_1, \dots, X_\ell)$, we write \bar{X}_I to denote the variables obtained by deleting X_i whenever $i \in I$. We can define \bar{Y}_J similarly when $J \subseteq \{1, \dots, n\}$ and $\bar{Y} = (Y_1, \dots, Y_n)$.

The formula $\text{BiREG}_{C,D}(\bar{X}, \bar{Y})$ as required in Theorem 5.1 is as follows.

$$\text{BiREG}_{C,D}(\bar{X}, \bar{Y}) := \bigvee_{\mathcal{P}} \left(\begin{array}{l} \Psi_{\mathcal{P}}(\bar{X}_I, \bar{Y}_J) \quad \wedge \quad \varphi \\ \wedge X_{i_1} = |S_1| \quad \wedge X_{i_2} = |S_2| \quad \wedge \cdots \quad \wedge X_{i_{m''}} = |S_{m''}| \\ \wedge Y_{j_1} = |T_1| \quad \wedge Y_{j_2} = |T_2| \quad \wedge \cdots \quad \wedge Y_{j_{n''}} = |T_{n''}| \end{array} \right)$$

where

- the disjunction ranges over all partial graphs $\mathcal{P} = (C', D', S, T, f, g)$ compatible with (C, D) w.r.t. $I = \{i_1, \dots, i_{m''}\}$ and $J = \{j_1, \dots, j_{n''}\}$, as well as the partitions $S = S_1 \cup \dots \cup S_{m''}$ and $T = T_1 \cup \dots \cup T_{n''}$,
- the formula $\psi_{\mathcal{P}}$ is as defined in Theorem 7.8,
- φ states that \bar{X}_I, \bar{Y}_J are big enough w.r.t. (C', D') , as defined in the Inequalities (7.1), (7.2) and (7.3).

The correctness of the formula $\text{BiREG}_{C,D}$ follows immediately from the correctness of the formula $\Psi_{\mathcal{P}}$ in Theorem 7.8. This completes our proof of Theorem 5.1.

7.6. Constructing the formula $\text{COMP-BiREG}_{C,D}(\bar{X}, \bar{Y})$ for Theorem 5.2.

We start with Lemma 7.9 which essentially states that if there exists a (C, D) -biregular-complete graph of “big enough” size, then for every column i in C and every column j in D , there is a row l such that both $C_{l,i}, D_{l,j} \in \blacktriangleright\mathbb{N}$. This means that a (C, D) -biregular-complete graph $G = (U, V, E_1, \dots, E_\ell)$ of “big enough” size (\bar{M}, \bar{N}) , then we can connect every pair of vertices $u \in U$ and $v \in V$ with one of the edges without violating the (C, D) -biregularity.

LEMMA 7.9. *Let $G = (U, V, E_1, \dots, E_\ell)$ be an ℓ -type (C, D) -biregular graph and $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ be the witness of the (C, D) -biregularity. Suppose that for each i, j , we have*

$$|U_i|, |V_j| \geq \lfloor C \rfloor \cdot \bar{1} + \lfloor D \rfloor \cdot \bar{1} + 1.$$

If G is a complete bipartite graph, then for every $i \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$, there exists $l \in \{1, \dots, \ell\}$ such that both $C_{l,i}, D_{l,j} \in \blacktriangleright\mathbb{N}$.

Proof. Let $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$, and $G = (U, V, E_1, \dots, E_\ell)$ be a (C, D) -biregular-complete graph, where $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ are the witness of the (C, D) -biregularity. Suppose each $|U_i|$ and $|V_j|$ satisfy the inequality above.

For the sake of contradiction, we assume that there exist $i, j \in \{1, \dots, m\}$ such that for all $l \in \{1, \dots, \ell\}$, either $C_{l,i} \in \mathbb{N}$ or $D_{l,j} \in \mathbb{N}$. This means that for each $l \in \{1, \dots, \ell\}$, the number of E_l -edges between U_i and V_j is $|U_i|C_{l,i}$, if $C_{l,i} \in \mathbb{N}$, or $|V_j|D_{l,j}$, if $D_{l,j} \in \mathbb{N}$. For each $l = 1, \dots, \ell$,

$$K_l = \begin{cases} |U_i|C_{l,i} & \text{if } C_{l,i} \in \mathbb{N}, \\ |V_j|D_{l,j} & \text{if } D_{l,j} \in \mathbb{N}. \end{cases}$$

Now the total number of edges between U_i and V_j must be $\sum_{1 \leq l \leq \ell} K_l$, which must be equal to $|U_i| \times |V_j|$ since G is a complete bipartite graph.

However, from the inequality

$$|U_i|, |V_j| \geq \lfloor C \rfloor \cdot \bar{1} + \lfloor D \rfloor \cdot \bar{1} + 1,$$

a straightforward calculation shows that K is strictly less than $|U_i| \times |V_j|$, a contradiction. Therefore, for every $i \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$, there exists $l \in \{1, \dots, \ell\}$ such that both $C_{l,i}, D_{l,j} \in \blacktriangleright \mathbb{N}$. This completes the proof of our lemma. \square

We say that a pair of matrices $(C, D) \in \mathbb{B}^{\ell \times m} \times \mathbb{B}^{\ell \times n}$ is an *easy pair of matrices*, if for every $i \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$ there exists $l \in \{1, \dots, \ell\}$ such that both $C_{l,i}, D_{l,j} \in \blacktriangleright \mathbb{N}$.

Lemma 7.10 says that if (C, D) is an easy pair of matrices, then the formula $\text{BiREG}_{C,D}(\bar{X}, \bar{Y})$ as defined in Subsection 7.5 is sufficient as the required formula $\text{COMP-BiREG}_{C,D}(\bar{X}, \bar{Y})$ in Theorem 5.2.

LEMMA 7.10. *Let (C, D) be an easy pair of matrices, where $C \in \mathbb{B}^{\ell \times m}$ and $D \in \mathbb{B}^{\ell \times n}$. Then the following holds. There exists a (C, D) -biregular-complete graph of size (\bar{M}, \bar{N}) if and only if $\text{BiREG}_{C,D}(\bar{M}, \bar{N})$ holds.*

Proof. The “only if” direction follows directly from Theorem 5.1. Now we prove the “if” direction. Suppose $\text{BiREG}_{C,D}(\bar{M}, \bar{N})$ holds. By Theorem 5.1, there exists a (C, D) -biregular graph $G = (U, V, E_1, \dots, E_\ell)$ of size (\bar{M}, \bar{N}) . This graph G is not necessarily complete. So suppose $U = U_1 \cup \dots \cup U_m$ and $V = V_1 \cup \dots \cup V_n$ is the witness of the (C, D) -biregularity. If G is not complete, then we perform the following. For every $u \in U$ and $v \in V$ such that $(u, v) \notin E_1 \cup \dots \cup E_\ell$, we do the following.


- Let $u \in U_i$ and $v \in V_j$.
- Pick an index $l \in \{1, \dots, \ell\}$ such that $C_{l,i}, D_{l,j} \in \blacktriangleright \mathbb{N}$. (Such an index l exists since (C, D) is an easy pair.)
- Connect u and v with an E_l -edge.

The resulting graph is now complete and still (C, D) -biregular. This completes our proof of Lemma 7.10. \square

If (C, D) is not an easy pair, then by Lemma 7.9, the values in the entries (in \bar{X} and \bar{Y}) corresponding to the columns in C and D that make them not an easy pair must be bounded. These values can be encoded as partial graphs as described in the previous section. This completes our proof of Theorem 5.2.

8. Proof of Theorem 5.4. The proof is by observing that the existence of a (C, D) -directed-regular graph of size \bar{N} is equivalent to the existence of a (C, D) -biregular graph of size (\bar{N}, \bar{N}) . We explain it more precisely below.

- Suppose $G = (V, E_1, \dots, E_\ell)$ is a (C, D) -directed-regular graph of size \bar{N} . Then, for every vertex $v \in V$, we “split” it into two vertices u and w such that u is only adjacent to the incoming edges of v and w to the outgoing edges of v . See the illustration below. The left-hand side shows the vertex v before the splitting, and the right-hand side shows the vertices u and w after the splitting.




Let U be the set of vertices u 's and W the set of vertices w 's after splitting all the vertices in V . Ignoring the orientation of the edges, the resulting graph is a bipartite graph with vertices $U \cup W$ and it is a (C, D) -biregular graph of size (\bar{N}, \bar{N}) .


- Suppose $G = (U, W, E_1, \dots, E_\ell)$ is a (C, D) -biregular graph of size (\bar{N}, \bar{N}) . Let $U = U_1 \cup \dots \cup U_m$ and $W = W_1 \cup \dots \cup W_m$ be the witness of the (C, D) -biregularity. We denote by $U_i = \{u_{i,1}, \dots, u_{i,K_i}\}$ and $W_i = \{w_{i,1}, \dots, w_{i,K_i}\}$ for each $i = 1, \dots, m$.

Now we put the orientation on all the edges from U to W . Then we merge every two vertices $u_{i,j}$ and $w_{i,j}$ into one vertex $v_{i,j}$. This way, we obtain a (C, D) -directed-regular graph $G = (V, E_1, \dots, E_\ell)$ with $V = V_1 \cup \dots \cup V_m$ being the witness of the (C, D) -regularity where $V_i = \{v_{i,1}, \dots, v_{i,K_i}\}$ for each $i = 1, \dots, m$.

However, with such merging it is possible that there is a self-loop (v, v) in G or a pair of edges $(v, v'), (v', v) \in E_1 \cup \dots \cup E_\ell$. We can get rid of the self-loop (v, v) without violating the (C, D) -directed-regularity as follows. The trick is similar to the one used before. Assuming that the size of each $|V_1|, \dots, |V_m|$ is big enough and there are enough edges in each E_1, \dots, E_ℓ , there is an edge (v', v'') of the same type such that both v', v'' are not adjacent to v . Deleting the edge (v, v) and (v', v'') , and adding the edges (v', v) and (v, v'') , we obtain a (C, D) -directed-regular graph with one less self-loop. We do this repeatedly until there is no more self-loop. See the illustration below.



Similarly, we can get rid of a pair of edges $(v, v'), (v', v) \in E_1 \cup \dots \cup E_\ell$ without violating the (C, D) -directed-regularity as follows. Again, the trick is similar to the one used before. Assuming that the size of each $|V_1|, \dots, |V_m|$ is big enough and there are enough edges in each E_1, \dots, E_ℓ , there is an edge (w, w') of the same type such that both w, w' are not adjacent to either v or v' . Deleting the edge (v, v') and (w, w') , and adding the edges (v, w') and (w, v') , we obtain a (C, D) -directed-regular graph with one less parallel edges. We do this repeatedly until there are no more parallel edges. See the illustration below.



If some sets V_i or $|E_i|$ are of a fixed size, they can be encoded in a partial graph in the same manner as discussed in Subsection 7.4.

We omit the technical details since we essentially run through the same arguments used in the previous section.

9. Concluding remarks. In this paper we have shown that the spectra of \mathcal{C}^2 formulae are semilinear sets. The proof is by constructing the Presburger formulae that express precisely the spectra. As far as our knowledge is concerned, the logic \mathcal{C}^2 is the first logic whose spectra is closed under complement without any restriction on the vocabulary nor in the interpretation.

Furthermore, from our proof we can easily deduce a few easy corollaries. The first is that the family of models of a QMLC formula can be viewed as a collection of biregular graphs and regular digraphs in the following sense. Let ϕ be a QMLC formula and $\mathcal{R} = \{R_1, \dots, R_\ell, \overleftarrow{R}_1, \dots, \overleftarrow{R}_\ell\}$ be the set of binary relations used in ϕ and that \overleftarrow{R}_i is the inversed relation of R_i . Let K be the integer such that for all subformulae $\diamond_R^l \psi$ in ϕ , we have $l \leq K$.

Let \mathcal{T}_ϕ be the set of all types in ϕ . For a model $\mathfrak{A} \models \phi$, we partition $A = \bigcup_{T \in \mathcal{T}_\phi} A_T$, where $A_T = \{a \in A \mid a \text{ is of type } T\}$. Then the model \mathfrak{A} can be seen as a collection of regular graphs in the following sense. Recall the matrices $D_T, \overleftarrow{D}_T, D_{S \rightarrow T}$ and $D_{T \rightarrow S}$ as defined in Section 6. For a type T , by restricting the relations R_1, \dots, R_ℓ on the elements in A_T , we obtain a $(D_T, \overleftarrow{D}_T)$ -regular digraph $G_T = (A_T, E_1, \dots, E_\ell)$, where for each E_i ,

$$E_i = R_i \cap (A_T \times A_T)$$

For two different types S, T , by restricting the relations on $A_S \times A_T$, we obtain a $(D_{S \rightarrow T}, \overleftarrow{D}_{S \rightarrow T})$ -biregular graph $G_{S,T} = (A_S, A_T, E_1, \dots, E_\ell, E'_1, \dots, E'_\ell)$, where each E_i, E'_i are

$$E_i = R_i \cap (A_S \times A_T) \quad \text{and} \quad E'_i = \overleftarrow{R}_i \cap (A_S \times A_T)$$

Theorem 2.1 can be further generalised as follows. Let $\mathcal{P} = (P_1, \dots, P_l)$, where P_1, P_2, \dots, P_l are unary predicates. Define the *image* of a structure \mathfrak{A} as $\text{Image}_{\mathcal{P}}(\mathfrak{A}) = (|P_1^{\mathfrak{A}}|, \dots, |P_l^{\mathfrak{A}}|)$. We also define the image of a formula φ with predicates from \mathcal{P} as $\text{Image}_{\mathcal{P}}(\varphi) = \{\text{Image}_{\mathcal{P}}(\mathfrak{A}) \mid \mathfrak{A} \models \varphi\}$. It must be noted here that $P_1^{\mathfrak{A}}, \dots, P_l^{\mathfrak{A}}$ are not necessarily disjoint, and that they may not cover the whole domain A . For this reason, the notion of image is more general than the notion of many-sorted spectrum which requires the unary predicates to partition the whole domain. With a slight adjustment in our proof in Section 6, we can obtain the following two corollaries.

COROLLARY 9.1. *Let $\phi \in \mathcal{C}^2$ and $\mathcal{P} = (P_1, \dots, P_l)$, where P_1, \dots, P_l be a set of unary predicates in ϕ . The set $\{\text{Image}_{\mathcal{P}}(\mathfrak{A}) \mid \mathfrak{A} \models \phi\}$ is semilinear.*

COROLLARY 9.2. *Let $\mathcal{P} = (P_1, \dots, P_l)$. The following problem is decidable. Given a \mathcal{C}^2 formula ϕ and a Presburger formula $\Psi(x_1, \dots, x_l)$, determine whether there exists a structure $\mathfrak{A} \models \phi$ such that $\Psi(\text{Image}_{\mathcal{P}}(\mathfrak{A}))$ holds.*

There are still a few more questions that we would like to investigate for future work. The first natural question is: how can \mathcal{C}^2 be extended while keeping decidability for the satisfiability and image membership problems? Using three variables (FO^3) one can easily encode a grid; therefore, the satisfiability problem is no longer decidable (and thus also the image membership problem). However, we could extend \mathcal{C}^2 by giving access to a relation having a property which is undefinable in \mathcal{C}^2 , such as transitivity. In particular, $\mathcal{C}^2(<)$, that is, the logic \mathcal{C}^2 with access to a total order on the universe, seems powerful: Petri net reachability [30, 24, 25] reduces to image membership for $\mathcal{C}^2(<)$ formulae. We do not know whether a reduction exists in the other direction. Another possible extension is to add an equivalence relation to \mathcal{C}^2 .

Acknowledgment. We thank the anonymous referees for their comments and suggestions which greatly improve our paper. The second author also acknowledges the generous financial support of FWO, under the scheme FWO Pegasus Marie Curie fellowship.

REFERENCES

- [1] G. Asser. Das Repräsentantenproblem in Prädikatenkalkül der ersten Stufe mit Identität. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, 1: 252–263 (1955).
- [2] G. van den Broeck, W. Meert, A. Darwiche. Skolemization for Weighted First-Order Model Counting. In KR 2014.
- [3] A. Durand, N. D. Jones, J. A. Makowsky, M. More. Fifty years of the spectrum problem: survey and new results. *Bulletin of Symbolic Logic* 18(4): 505–553 (2012).
- [4] H.-D. Ebbinghaus, J. Flum. *Finite model theory*. Perspectives in Mathematical Logic, Springer 1995.
- [5] R. Fagin. *Contributions to the model theory of finite structures*. Ph.D. thesis, University of California, Berkeley, 1973.
- [6] R. Fagin. Finite model theory—a personal perspective. *Theoretical Computer Science* 116: 3–31 (1993).
- [7] E. Fischer, J. A. Makowsky. On spectra of sentences of monadic second order logic with counting. *Journal of Symbolic Logic* 69(3):617–640 (2004).
- [8] S. Ginsburg, E. H. Spanier. Semigroups, Presburger Formulas, and Languages. *Pacific Journal of Mathematics* 16:285–296 (1966).
- [9] É. Grandjean. The spectra of first-order sentences and computational complexity. *SIAM Journal on Computing* 13(2): 356–373 (1984).
- [10] É. Grandjean. Universal quantifiers and time complexity of random access machines. *Mathematical Systems Theory* 18(2): 171–187 (1985).
- [11] É. Grandjean. First-order spectra with one variable. *Journal of Computer and System Sciences* 40(2): 136–153 (1990).
- [12] E. Grädel. Why are Modal Logics so Robustly Decidable? *Current Trends in Theoretical Computer Science* 2001.
- [13] E. Grädel, P. G. Kolaitis, M. Y. Vardi. On the Decision Problem for Two-Variable First-Order Logic. *The Bulletin of Symbolic Logic*, 3(1): 53–69 (1997).
- [14] E. Grädel, M. Otto. On Logics with Two Variables. *Theoretical Computer Science* 224(1-2): 73–113 (1999).
- [15] E. Grädel, M. Otto, E. Rosen. Two-Variable Logic with Counting is Decidable. In LICS 1997.
- [16] M. Grohe, J. A. Makowsky, Editors. *Model Theoretic Methods in Finite Combinatorics*. Contemporary Mathematics 558, American Mathematical Society, 2011.
- [17] Y. Gurevich and S. Shelah. Spectra of monadic second-order formulas with one unary function. In LICS 2003.

- [18] S. Hakimi. On realizability of a set of integers as degrees of the vertices of a linear graph. I. *Journal of the Soc. Indust. Appl. Math.* 10(3): 496–506 (1962).
- [19] I. Pratt-Hartmann. Complexity of the Two-Variable Fragment with Counting Quantifiers. *Journal of Logic, Language and Information* 14(3): 369–395 (2005).
- [20] A. Hunter. Spectrum hierarchies and subdiagonal functions. In LICS 2003.
- [21] N. Immerman. *Descriptive complexity*. Graduate texts in computer science, Springer 1999.
- [22] N. Jones, A. Selman. Turing machines and the spectra of first-order formulas. *The Journal of Symbolic Logic*, 39: 139–150 (1974). (Also in STOC 1972.)
- [23] H. Landau. On dominance relations and the structure of animal societies. III. The condition for a score structure. *Bulletin of Mathematical Biophysics* 15(2): 143–148 (1953).
- [24] S. Kosaraju. Decability of reachability in vector addition systems. In STOC 1982.
- [25] J. Leroux. The General Vector Addition System Reachability Problem by Presburger Inductive Invariants. In LICS 2009.
- [26] E. Kieronski, J. Michaliszyn, I. Pratt-Hartmann, L. Tendera. Two-Variable First-Order Logic with Equivalence Closure. In LICS 2012
- [27] L. Libkin. *Elements of Finite Model Theory*. Springer 2004.
- [28] J. Lynch. Complexity classes and theories of finite models. *Mathematical Systems Theory* 15(2): 127–144 (1982).
- [29] C. Lutz, U. Sattler, F. Wolter. Modal logic and the two-variable fragment. In CSL 2001.
- [30] E. Mayr. An algorithm for the general petri net reachability problem. In STOC 1981.
- [31] M. Mortimer. On languages with two variables. *Mathematical Logic Quarterly*, 21(1): 135–140 (1975).
- [32] M. Otto. *Bounded Variable Logics and Counting – A Study in Finite Models* Lecture Notes in Logic volume 9, Springer 1997.
- [33] L. Pacholski, W. Szwast, L. Tendera. Complexity Results for First-Order Two-Variable Logic with Counting. *SIAM Journal on Computing* 29(4): 1083–1117 (2000).
- [34] I. Pratt-Hartman. Logics with counting and equivalence. In CSL-LICS 2014.
- [35] D. Rödding, H. Schwichtenberg. Bemerkungen zum Spektralproblem. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, 18: 1–12 (1972).
- [36] H. Scholz. Ein ungelöstes Problem in der symbolischen Logik. *The journal of Symbolic Logic*, 17: 160 (1952).
- [37] W. Szwast, L. Tendera. FO^2 with one transitive relation is decidable. In STACS 2013.
- [38] K. N. Verma, H. Seidl, T. Schwentick. On the Complexity of Equational Horn Clauses. In CADE 2005.
- [39] M. Vardi. Why is Modal Logic So Robustly Decidable? In *Descriptive Complexity and Finite Models* 1996.
- [40] A. Woods. *Some Problems in Logic and Number Theory and their Connections*. Ph.D. thesis, University of Manchester, 1981.
- [41] C. Wrathall. Rudimentary Predicates and Relative Computation. In *SIAM Journal on Computing* 7(2): 194–209 (1978).