

Analyzing pace frequencies in bipedal primates and primate
"predecessors" reveals mechanisms that regulate foot inversion and
thus ensure foot stability at touchdown

Peer-reviewed author version

VAN ZWIETEN, Koos Jaap; NARAIN, Faridi; De Munter, Stephanie; Kosten, Lauren;
Lamur, Kenneth S.; SCHMIDT, Klaus; LIPPENS, Peter; ZOUBOVA, Irina; Piskun,
Oleg E. & Varzin, Sergey A. (2015) Analyzing pace frequencies in bipedal primates
and primate "predecessors" reveals mechanisms that regulate foot inversion and
thus ensure foot stability at touchdown. In: Xth Annual All-Russian Research and
Practical Conference with International Participation "Health - The Base of Human
Potential: Problems and Ways to Solve Them": Proceedings of the conference, p. 820-822.

DOI: 10.13140/RG.2.1.3688.8400

Handle: <http://hdl.handle.net/1942/20087>

UDK 611.738.3

***K.J. van Zwieten ^a, F.H.M. Narain ^a, S. De Munter ^a,
L. Kosten ^a, K.S. Lamur ^b, K.P. Schmidt ^a, P.L. Lippens ^a,
I.A. Zubova ^c, O.E. Piskun ^c, S.A. Varzin ^c***

**ANALYZING PACE FREQUENCIES IN BIPEDAL PRIMATES AND PRIMATE
“PREDECESSORS” REVEALS MECHANISMS THAT REGULATE FOOT INVERSION
AND THUS ENSURE FOOT STABILITY AT TOUCHDOWN**

^a University of Hasselt, Diepenbeek, Belgium, ^b Anton-de-Kom University of Suriname, Paramaribo, Suriname, ^c Peter-the-Great St. Petersburg Polytechnic University, St. Petersburg, Russia koosjaap.vanzwieten@uhasselt.be


Introduction and hypothesis: High-pitched sounds of unintentional lateral foot-shuffling along the ground during the swing phase of gait have been recognized as almost pathognomonic signs of fatigue in lower leg and foot evertor muscles in the early stages of Multiple Sclerosis (MS) [1]. MS patients diagnosed in the early stages of the disease encounter several ominous events like tripping and falling. Recovery from foot inversion at toe-off and during the swing phase by lower leg internal rotation as well as stabilization of foot inversion after touchdown by subsequent foot eversion are affected. Inversion is defined as turning the sole of the foot inward, while eversion is defined as turning the sole outward. In order to explore the key features of this phenomenon extensive investigation of normal functional morphology of the lower leg is worthwhile. Studying normal functional morphology of the lower leg, particularly in primates and predecessors, is an alternative approach that may enlighten the evolutionary path leading to this phenomenon. Recent literature shows that in walking bipedally, various arboreal New World primate species use a “forefoot first” strategy, after which heel contact occurs [2]. A similar walking scenario is seen in arboreal New World marsupials like the opossum, a quadrupedal primate “predecessor”. In opossum walking the swing phase ends with the foot in the inverted position to the next touchdown with the forefoot [3]. We therefore hypothesized that those quadrupedal marsupials which are mainly or exclusively terrestrial like e.g., wombats, will also display a “forefoot first” strategy at touchdown. However, while lower leg rotational mechanisms in wombats are well-described [4], currently “no detailed analyses of wombat locomotion have been carried out. (...) Wombats are plantigrade, walking on the soles of their feet with three distinct gaits, walk, trot and bound” [5].

Material and methods: Various semi-domesticated fully terrestrial *Vombatus ursinus* (common wombats) were videotaped (November 2013) before dusk, walking in their natural environments at Cradle Mountain Road (C132), Cradle Mountain, Tasmania. Following current guidelines, we took utmost care to leave all animals undisturbed [6]. Posterior and lateral views were analyzed frame by frame.

Observational results: Average wombat walking speed is about 2,5 km/h, with hind-limb pace frequencies of about 1 Hz. In man, this is : 0,5 Hz (walking) - 2,5 Hz (running) [3]. Wombat toe-off occurs with the foot in the everted position (Fig. 1). Halfway recovery there is inversion of the foot, coupled to shank external rotation (Fig. 2). At touchdown a “forefoot first” position is observed. The wombat recovery phase namely ends with a first ground-contact by the (presumably metatarsophalangeal) plantar pads [7, 8] (Fig. 3, red arrowhead), with the foot still in inversion. This is immediately followed by a rapid full foot placement, foot in eversion.

Conclusion: Marsupials such as wombats, though being exclusively quadrupedal and non-arboreal, show basically the same strategies of foot eversion to ensure safe touchdown as bipedally walking arboreal New World primate species [2], with morphologically guided adjustments. Analyzing this hopefully helps to understand gait training problems in patients too, with emphasis on foot inversion and eversion.

Acknowledgements: The authors gratefully acknowledge Jikke van Osnabrugge, PT for making the videos.



References.

1. van Zwieten KJ, Narain FHM, Kosten L, Wens I, Eijnde BO, Vandersteen M, Schmidt KP (2013) Reappraisal of gait patterns in minimally impaired Multiple Sclerosis patients reveals characteristic foot shuffling sounds. International Sound and Vibration Digest, 14, 1, 5-6.

2. Schmitt D, Larson SG (1995) Heel contact as a function of substrate type and speed in primates. *American Journal of Physical Anthropology*, 96, 1, 39-50.
3. Narain FHM (2013) Foot inversion and eversion in stance and swing of the gait cycle of the hind limb of the opossum (*Didelphis marsupialis*) - some comparative-anatomical and functional-morphological aspects. PhD Thesis, University of Hasselt, Belgium.
4. Home E (1808) An account of some peculiarities in the anatomical structure of the wombat. *Philosophical Transactions of the Royal Society*, 98, 2, 304-312.
5. Wells RT (1989) Vombatidae. In *Fauna of Australia*. Vol. 1B, Mammalia, 32, AGPS, Canberra.
6. Dennis L, Rolls K, Fowler A, Dineen A-M (2010) A Guide to the Care of Bare-Nosed Wombats (*Vombatus ursinus*) Version 3. Fourth Crossing Wildlife, www.fourthcrossingwildlife.com
7. Boardman W (1943) On the external characters of the pouch young of some Australian Marsupials. *Australian Zoologist*, 10, 2, 138-160.
8. Vogelnest L, Allan G (2015) Radiology of Australian Mammals. CSIRO Publishing, Melbourne.