# Made available by Hasselt University Library in https://documentserver.uhasselt.be

Analysis and upscaling of a reactive transport model in fractured porous media with nonlinear transmission condition Non Peer-reviewed author version

POP, Sorin; Bogers, J. & Kumar, K. (2017) Analysis and upscaling of a reactive transport model in fractured porous media with nonlinear transmission condition. In: Vietnam Journal of Mathematics, 45 (1-2), pag. 77-102.

DOI: 10.1007/s10013-016-0198-7 Handle: http://hdl.handle.net/1942/21498



# Analysis and upscaling of a reactive transport model in fractured porous media involving a nonlinear transmission condition

Pop, I. U. Bogers, J. Kumar, K.

GRK 1398/2 (DFG) DN 81 -754 (NWO) 215627 (RCN)

- Geschäftsstelle -Pfaffenwaldring 61 70569 Stuttgart Telefon: 0711/685-60399 Telefax: 0711/685-60430 E-Mail: nupus@iws.uni-stuttgart.de http://www.nupus.uni-stuttgart.de

# Analysis and upscaling of a reactive transport model in fractured porous media involving nonlinear a transmission condition

Iuliu Sorin Pop $^{1,3}$   $\cdot$  Jeroen Bogers $^2$   $\cdot$  Kundan Kumar $^3$ 

Received: date / Accepted: date

Abstract We consider a reactive transport model in a fractured porous medium. The particularity appears in the conditions imposed at the interface separating the block and the fracture, which involves a nonlinear transmission condition. Assuming that the fracture has thickness  $\varepsilon$ , we analyze the resulting problem and prove the convergence towards a *reduced* model in the limit  $\varepsilon \searrow 0$ . The resulting is a model defined on an interface (the reduced fracture) and acting as a boundary condition for the equations defined in the block. Using both formal and rigorous arguments, we obtain the reduced models for different flow regimes, expressed through a moderate, or a high Peclét number.

Keywords Fractured porous media  $\cdot$  Upscaling  $\cdot$  Reactive transport  $\cdot$  Nonlinear transmission conditions

## 1 Introduction

Fractures are ubiquitous in porous media and have strong influence on the flow and transport. Several energy and environmental applications including carbon sequestration, geothermal energy, hydraulic fracturing, petroleum extraction, or ground water contamination, are involving flow and reactive transport in fractured porous media. Typically, fractures are thin and long formations along which medium properties like permeability, or porosity, are different from the adjacent formations (the blocks). This leads to media with high contrasting properties appearing in anisotropic regions, and involving jump-type discontinuities across interfaces between the fracture and the block. This makes the

 $<sup>^1\</sup>rm Eindhoven$  University of Technology, Mathematics and Computer Science, PO Box 513, 5600 MB Eindhoven, The Netherlands, E-mail: i.pop@tue.nl

<sup>· &</sup>lt;sup>2</sup>ASML Netherlands B.V., De Run 6501, 5504DR Veldhoven, The Netherlands, E-mail: j.j.p.bogers@gmail.com

 $<sup>\</sup>cdot$   $^3$  University of Bergen, Department of Mathematics, Postboks 7803, 5020 Bergen, Norway, E-mail: kundan.kumar@math.uib.no

numerical simulation of mathematical models in such domains a very challenging task. Discretization schemes resolving the full thin fracture regions explicitly is prohibitively expensive. Moreover, often the main interest is in the transport or flow along fractures and not in the transverse details. This suggests considering the so called *reduced models* for the processes inside fractures, which are transversally averaged along such formations, and reduce the fracture itself to an interface separating two blocks. The resulting problem is defined in a bulk domain (the union of the porous blocks), as well as on surfaces (the lower dimensional approximation of the fractures) embedded in the bulk domain. Nevertheless, the surface should incorporate the averaged processes in the fracture, including the coupling conditions between the fracture and the adjacent blocks. This requires a consistent procedure for developing such reduced models.

In this work, both formal and rigorous procedures are used for deriving reduced models that describe reactive transport in fractured media. The flow is assumed to be known in the thin fractures, and the transport is modelled by a convection-diffusion equation. Inside blocks, which are typically less permeable than fractures, no flow is encountered, and species undergo only diffusive transport. The particularity of the model lies in the conditions coupling the models inside fractures and blocks at the common interface. Next to the normal flux continuity, a nonlinear transmission condition is imposed. For the ease of presentation, we consider a simple geometry, where a large rectangular domain (the porous block) is coupled to a rectangular fracture with small thickness  $\varepsilon$ . After transversal averaging in the fracture and letting  $\varepsilon$  go to zero, the fracture model becomes a boundary condition for the partial differential equation in the bulk domain. This procedure is carried out formally in two situations, when the flow and transport processes in the fracture are in balance (moderate Peclét regime), and when flow is dominating (high Peclét regime). Then, convergence is obtained rigorously for the moderate Peclét regime.

Reactive transport in heterogeneous media lead naturally to nonlinear transmission conditions. Examples in this sense appear in bubble columns or extraction processes involving multiphase systems. In ionic fluids, which are strongly non-ideal, nonlinear reactions are encountered at interfaces separating two domains [33]. Well-known examples in this sense are Langmuir, Freundlich or Monod type reactions. Such models are also the outcome of the upscaling procedure in [20,26,27,28,29]. Nonlinear transmission conditions also appear in multiphase flow models in porous media [10], where e.g. pressure continuity at interface separating a fine and a coarse porous block induces a nonlinear relation between the oil saturations at the two sides of the interface.

Due to their high permeability in the fractures one encounters both convection and diffusion processes. Related, at least two different time scales can be identified: a convective time scale  $T_C$ , and a diffusive one  $T_D$ . Their ratio defines the non-dimensional Peclét number,  $Pe = \frac{T_D}{T_C}$ . The observed transport behaviour depends strongly on this number, in particular, when the convection dominates (high Peclét), the net diffusion is enhanced by the convective

strength itself exhibiting the well-known Taylor dispersion [34, 19, 25]. The upscaling procedure needs to take these complexities into account.

The results in this paper are the following. For a fixed  $\varepsilon$ , the existence of solutions for the system involving two parabolic models posed in adjacent domains, and involving a nonlinear transmission conditions is proved. This is obtained by Rothe's method [16] and essentially uses the existence results for elliptic problems obtained in [14]. We also mention that the full model (i.e. when  $\varepsilon > 0$  considered here is similar to the one in [15]. Here the existence results are obtained directly for the original unknowns, and without employing a nonlinear transformation of these in order to obtain the continuity across the interface. This is closer to the approaches in [6,7,23] and allows extending the present results to the case of heterogeneous (e.g. spatially dependent) transmission conditions. Next, we derive the reduced models by averaging in the transversal direction of the fracture, and using formal asymptotic methods. This is achieved for two regimes in the fracture, moderate Péclet,  $Pe = \mathcal{O}(1)$ , and transport dominated regime  $Pe = \mathcal{O}(\varepsilon^{-1})$ . Finally, for the case when  $Pe = \mathcal{O}(1)$ , we rigorously prove convergence of the full model, including the nonlinear transmission condition, to the simple, upscaled model obtained by formal asymptotics.

The paper is structured as follows. In Section 2, we introduce the mathematical model and in Section 3 we prove the existence of a solution for a fixed  $\varepsilon$ with the necessary a priori estimates derived in Section 3.4. Next, we perform a formal upscaling for both moderate and high Peclét number in sections 4 and 4.2, respectively. Section 5 deals with rigorous derivation of the upscaled equations obtained in Section 4. Full details can be found in [4], Chapters 4, 5 and 9. The paper concludes with discussions and outlook in Section 6.

#### 2 Model and assumptions

The problem is stated in a dimensionless framework and we refer e.g. to [9,11, 19] for a non-dimensionalization step. Let T > 0 be a maximal time and  $\Omega_p$  (the porous block) and  $\Omega_f^{\varepsilon}$  (the fracture) be two adjacent domains separated by the interface  $\Gamma$ :

$$\begin{split} \Omega_p &:= \{(x,y) \in \mathbb{R}^2 | 0 < x < 1, -1 < y < 0\},\\ \Omega_f^{\varepsilon} &:= \{(x,y) \in \mathbb{R}^2 | 0 < x < 1, 0 < y < \varepsilon\},\\ \Gamma &:= \{(x,y) \in \mathbb{R}^2 | 0 < x < 1, y = 0\},\\ \Gamma_{f,1}^{\varepsilon} &:= \{(x,y) \in \mathbb{R}^2 | x = 0, 0 < y < \varepsilon\},\\ \Gamma_{f,3}^{\varepsilon} &:= \{(x,y) \in \mathbb{R}^2 | x = 1, 0 < y < \varepsilon\},\\ \Gamma_{p,1}^{\varepsilon} &:= \{(x,y) \in \mathbb{R}^2 | x = 0, -1 < y < 0\},\\ \Gamma_{p,3}^{\varepsilon} &:= \{(x,y) \in \mathbb{R}^2 | x = 1, -1 < y < 0\},\\ \Gamma_{p,2}^{\varepsilon} &:= \{(x,y) \in \mathbb{R}^2 | 0 < x < 1, y = -1\},\\ \Gamma_{f,2}^{\varepsilon} &:= \{(x,y) \in \mathbb{R}^2 | 0 < x < 1, y = \varepsilon\}. \end{split}$$
(1)



Fig. 1 Geometry depicting a fracture coupled to an adjacent porous medium. The fracture is of thickness  $2\varepsilon$ , and due to symmetry it is sufficient to consider only half of the fracture.  $\Gamma_{f,2}^{\varepsilon}$  is this line of symmetry.

The remaining boundaries of  $\Omega_p$  and  $\Omega_f$  are  $\Gamma_{p,1}$ , respectively  $\Gamma_{f,1}$  (1 = 1, 2, 3), see Figure 1. Here  $\varepsilon > 0$  is the half-fracture aperture (because of symmetry) and is assumed small compared to the fracture length.

## 2.1 Notation

Before stating the model, we mention that standard notations from the functional analysis are being used below. By  $(\cdot, \cdot)$  we mean  $L^2$  inner product or the duality pairing between  $W^{1,2}$  and its dual. By  $Tr(\cdot)$  we denote the trace operator. Further, with X being a Banach space, C(0,T;X) stands for the Xvalued functions that are continuous over [0,T], and  $L^p(0,T;X)$  is the usual Bochner space. Next, for the weak solutions we will use the spaces

$$V_p := \{ u \in W^{1,2}(\Omega_p) | u = 0 \text{ on } \Gamma_{p,1} \cup \Gamma_{p,3} \},$$
(2)

$$V_f^{\varepsilon} := \{ u \in W^{1,2}(\Omega_f^{\varepsilon}) | u = 0 \text{ on } \Gamma_{f,1}^{\varepsilon} \cup \Gamma_{f,3}^{\varepsilon} \},$$
(3)

with  $\Gamma_{p,1},\Gamma_{p,3}$ ,  $\Gamma_{f,1}^{\varepsilon}$  and  $\Gamma_{f,3}^{\varepsilon}$  defined in (1). Since  $V_p$  is a closed subspace of the Hilbert space  $W^{1,2}(\Omega_p)$ , it is again a Hilbert space, equipped with the  $W^{1,2}(\Omega_p)$  norm and inner product. Similarly,  $V_f^{\varepsilon}$  is a Hilbert space equipped with the  $W^{1,2}(\Omega_f^{\varepsilon})$  norm and inner product. For  $u \in L^2(0,T;V_p^*)$  and  $v \in L^2(0,T;V_p)$ , we define

$$(u,v)_{\varOmega_p^T} := \int_0^T \langle u(t), v(t) \rangle_{\varOmega_p} dt, \text{ where } \langle u(t), v(t) \rangle_{\varOmega_p} := \langle u(t), v(t) \rangle_{V_p^*, V_p}.$$

The counterparts for the fracture domains are defined similarly. Finally, let

$$\mathcal{W}_{p} := \{ u \in L^{2}(0, T; V_{p}) | \partial_{t} u \in L^{2}(0, T; V_{p}^{*}) \},$$
(4)

$$\mathcal{W}_f^{\varepsilon} := \{ u \in L^2(0, T; V_f^{\varepsilon}) | \partial_t u \in L^2(0, T; (V_f^{\varepsilon})^*) \}.$$
(5)

We observe that, by Assumption  $A_r$  below, if  $u \in W^{1,2}(\Omega)$ , then  $r(u) \in W^{1,2}(\Omega)$  as well, and hence Tr(r(u)) is well defined, belonging to  $L^2(\partial \Omega)$ .

#### 2.2 Mathematical model

The unknown quantities are the concentrations  $u_p^{\varepsilon}, u_f^{\varepsilon}$  of two chemical species defined in  $\Omega_p$  and  $\Omega_f^{\varepsilon}$ , respectively. The transporting mechanisms are diffusion in  $\Omega_p$  and convection and diffusion in  $\Omega_f^{\varepsilon}$ , respectively. The models in the two domains are coupled at  $\Gamma$  through two conditions: the continuity of the normal fluxes, and the nonlinear reactions (the nonlinear transmission condition). We assume symmetry along  $\Gamma_{f,2}^{\varepsilon}$ . This is summarized in:

$$\begin{cases} \partial_t u_f^{\varepsilon} - \Delta u_f^{\varepsilon} = f_p, & \text{in } (0, T] \times \Omega_p, \\ \partial_t u_f^{\varepsilon} + \mathbf{q}^{\varepsilon} \cdot \nabla u_f^{\varepsilon} - \Delta u_f^{\varepsilon} = f_f, & \text{in } (0, T] \times \Omega_f^{\varepsilon}, \\ u_f^{\varepsilon} = r(u_p^{\varepsilon}), & \text{at } (0, T] \times \Gamma, \\ \partial_{\boldsymbol{\nu}} u_p^{\varepsilon} = \partial_{\boldsymbol{\nu}} u_f^{\varepsilon}, & \text{at } (0, T] \times \Gamma. \end{cases}$$

$$(6)$$

Boundary and initial conditions are specified below. Here the diffusion coefficients are taken exactly 1, but extending the presentation to more general coefficients  $D_p$  and  $D_f$ , or positive definite tensors, is immediate. In Section 4, where formal upscaling will be carried out, we consider an  $\varepsilon$  order diffusion in the fracture as well to include the convection dominated regime.  $f_p$  and  $f_f^{\varepsilon}$  are source terms, and  $\mathbf{q}^{\varepsilon}$  is a given fluid velocity, all satisfying assumptions mentioned below.

#### 2.3 Assumptions

The particularity is in the nonlinear transmission conditions at  $\Gamma$ , involving the function r. For this we assume

 $(A_r)$  The function  $r \in C^1(\mathbb{R})$  satisfies r(0) = 0, and there exist m, M > 0 such that for all  $u \in \mathbb{R}, 0 < m \le r'(u) \le M < \infty$ .

The concentrations vanish on the vertical boundaries,

$$u_f^{\varepsilon} = 0, \text{ on } \Gamma_{f,1}^{\varepsilon} \cup \Gamma_{f,3}^{\varepsilon}, \text{ and } u_p^{\varepsilon} = 0, \text{ on } \Gamma_{p,1} \cup \Gamma_{p,3}.$$
 (7)

**Remark 1** By Assumption  $A_r$ , r has a  $C^1$  inverse  $r^{-1}$  satisfying

$$0 < \frac{1}{M} \le (r^{-1})'(u) \le \frac{1}{m} < \infty, \quad \text{for all } u \in \mathbb{R}.$$

This gives  $0 \le (r^{-1})(u) \le \frac{1}{m}u$  for all  $u \ge 0$ .

Below we will use the antiderivative of r, defined as

$$R: \mathbb{R} \to \mathbb{R}, \qquad R(u) := \int_0^u r(v) dv. \tag{8}$$

Using Assumption  $A_r$  and the Mean Value Theorem, the following can be proved straightforwardly

**Proposition 2** The function R has the following elementary properties: (i)  $R(x) \ge 0$ , for all  $x \in \mathbb{R}$ . (ii)  $R(y) - R(x) \le r(y)(y-x)$ , for all  $x, y \in \mathbb{R}$ . (iii)  $R(x) \ge \frac{m}{2}x^2$ , for all  $x \in \mathbb{R}$ .

Further, we assume

 $\begin{array}{l} (A_f) \mbox{ The functions } f_p \in C(0,T;L^2(\varOmega_p)) \mbox{ and } f_f \in C(0,T;L^2(\varOmega_f^\varepsilon)) \mbox{ are bounded} \\ \mbox{ and positive: there exist } M_f > 0 \mbox{ such that for all } t \in [0,T], \end{array}$ 

 $0 \leq f_f(t, \cdot) \leq M_f$  a.e. in  $\Omega_f^{\varepsilon}$ , and  $0 \leq f_p(t, \cdot) \leq M_f$  a.e. in  $\Omega_p$ .

 $({\cal A}_I)$  The initial conditions are positive and essentially bounded: there exist  $M_I>0$  such that

 $0 \le u_{I,f} \le M_I$  a.e. in  $\Omega_f^{\varepsilon}$ , and  $0 \le u_{I,p} \le M_I$  a.e. in  $\Omega_p$ .

 $(A_q)$  The velocity field  $\mathbf{q}^{\varepsilon} = (q^{\varepsilon,1}, q^{\varepsilon,2}) \in [W^{1,2}(\Omega_f^{\varepsilon})]^2$  isatisfies

 $\nabla \cdot \mathbf{q}^{\varepsilon} = 0 \text{ in } \Omega_{f}^{\varepsilon}, \text{ and } \mathbf{q}^{\varepsilon} = \mathbf{0} \text{ on } \Gamma.$ 

Further,  $q^{\varepsilon,2} = 0$  at  $\Gamma_{f,2}^{\varepsilon}$  (symmetry). Finally,  $\mathbf{q}^{\varepsilon}$  is essentially bounded: there exists  $M_q > 0$  s.t.  $||\mathbf{q}^{\varepsilon}|| \leq M_q$  a.e. in  $\Omega_f^{\varepsilon}$ .

Concerning  $A_q$ , this holds true if e.g., **q** solves the Stokes model with homogeneous Dirichlet boundary conditions, see [17,21]. To simplify the calculations, the initial data are assumed  $\varepsilon$ -independent, i.e.  $u_p^{\varepsilon}(0, x, y) = u_{I,p}(x, y)$ and  $u_f^{\varepsilon}(0, x, y) = u_{I,f}(x)$ . Similarly, the source terms are  $\varepsilon$ -independent too.

#### 3 Existence of a weak solution

=

We first prove the existence of a weak solution for the original model in (6), with the initial and boundary conditions stated above. This is defined in

**Definition 3** A pair  $(u_f, u_p) \in L^2(0, T; V_f^{\varepsilon}) \times L^2(0, T; V_p)$  is called a weak solution of (6) if  $u_f = r(u_p)$  on  $\Gamma$  (in the sense of traces) and

$$- (u_f, \partial_t \phi_f)_{\Omega_f^{\varepsilon,T}} - (u_p, \partial_t \phi_p)_{\Omega_p^T} + (\mathbf{q} \cdot \nabla u_f, \phi_f)_{\Omega_f^{\varepsilon,T}} + (\nabla u_f, \nabla \phi_f)_{\Omega_f^{\varepsilon,T}} + (\nabla u_p, \nabla \phi_p)_{\Omega_p^T}$$
(9)  
$$(f_f, \phi_f)_{\Omega_t^T} + (f_p, \phi_p)_{\Omega_p^T} + (u_{I,f}, \phi_f(0))_{\Omega_t^{\varepsilon}} + (u_{I,p}, \phi_p(0))_{\Omega_p},$$

for all  $(\phi_f, \phi_p) \in W^{1,2}(0, T; V_f^{\varepsilon}) \times W^{1,2}(0, T; V_p)$  such that  $\phi_f = \phi_p$  on  $\Gamma$  and  $\phi_f(T) = \phi_p(T) = 0$ .

The existence of a weak solution is obtained by the Rothe's method [16]. In doing so, the  $\varepsilon$  dependence in the a-priori estimates is stated explicitly.

## 3.1 Discretization in time

Letting  $\Delta t$  be a fixed time step, taking  $u_i^0 = u_{I,i}$   $(i \in \{f, p\})$ , we construct the Euler implicit approximations  $\{u_i^k\}_{k \in \mathbb{N}}$  of  $u_i$  at  $t_k = k\Delta t$ . This leads to a sequence of elliptic problems involving again a nonlinear transmission condition. We omit the strong form here and provide directly the definition of a weak solution, which is stated in

**Definition 4** Let k > 0 and let  $(u_f^{k-1}, u_p^{k-1}) \in V_f^{\varepsilon} \times V_p$  be given. A weak solution to the time discrete problem at  $t_k$  is a pair  $(u_f^k, u_p^k) \in V_f^{\varepsilon} \times V_p$  satisfying  $u_f^k = r(u_p^k)$  on  $\Gamma$  (in the sense of traces) and

$$\left(\frac{u_f^k - u_f^{k-1}}{\Delta t}, \phi_f\right)_{\Omega_f^\varepsilon} + \left(\frac{u_p^k - u_p^{k-1}}{\Delta t}, \phi_p\right)_{\Omega_p} + (\nabla u_f^k, \nabla \phi_f)_{\Omega_f^\varepsilon} + (\nabla u_p^k, \nabla \phi_p)_{\Omega_p} + (\mathbf{q} \cdot \nabla u_f^k, \phi_f)_{\Omega_f^\varepsilon} = (f_f(t_k), \phi_f)_{\Omega_f^\varepsilon} + (f_p(t_k), \phi_p)_{\Omega_p} \tag{10}$$

for all  $(\phi_f, \phi_p) \in V_f^{\varepsilon} \times V_p$  such that  $\phi_f = \phi_p$  on  $\Gamma$ .

Such elliptic problems are studied in [14], where the existence and uniqueness of a weak solution is proved.

#### 3.2 A priori estimates

We start by observing that for any  $u \in V_p$ , since  $r \in C^1$  we have  $r(u) \in V_f$  as well. This will be used below to prove

**Lemma 1** For the sequence of time discrete weak solutions in Definition 4, a C > 0 not depending on  $\Delta t$  exists s.t.

$$\max_{j \in \{1,...,N\}} \left\| \left\| u_{f}^{j} \right\|_{\Omega_{f}^{\varepsilon}}^{2} + \max_{j \in \{1,...,N\}} \int_{\Omega_{p}} R(u_{p}^{j}) dx + \sum_{k=1}^{N} \left\| \left\| u_{f}^{k} - u_{f}^{k-1} \right\|_{\Omega_{f}^{\varepsilon}}^{2} + \Delta t \sum_{k=1}^{N} \left\{ \left\| \nabla u_{f}^{k} \right\|_{\Omega_{f}^{\varepsilon}}^{2} + \left\| \nabla u_{p}^{k} \right\|_{\Omega_{p}}^{2} \right\} \leq C.$$
(11)

*Proof* We test with  $\phi_f := u_f^k$  and  $\phi_p := r(u_p^k)$  in (10) and denote the resulting terms by  $I_1, I_2, ..., I_7$ . Clearly,

$$I_{1} = \frac{1}{2} \left| \left| u_{f}^{k} \right| \right|_{\Omega_{f}^{\varepsilon}}^{2} + \frac{1}{2} \left| \left| u_{f}^{k} - u_{f}^{k-1} \right| \right|_{\Omega_{f}^{\varepsilon}}^{2} - \frac{1}{2} \left| \left| u_{f}^{k-1} \right| \right|_{\Omega_{f}^{\varepsilon}}^{2}.$$

To treat  $I_2$ , we use Proposition 2 to obtain

$$I_2 \ge \int_{\Omega_p} R(u_p^k) dx - \int_{\Omega_p} R(u_p^{k-1}) dx.$$

Note that, since R is positive,  $\int_{\Omega_p} R(u_p^k) dx \ge 0$  for all  $k \in \{0, ..., N\}$ .

For  $I_3$ , using the Poincaré inequality gives

$$I_3 \geq \frac{1}{2} \left| \left| \nabla u_f^k \right| \right|_{\Omega_f^\varepsilon}^2 + \frac{1}{2C_f} \left| \left| u_f^k \right| \right|_{\Omega_f^\varepsilon}^2,$$

for a  $C_f$  depending only on the geometry of  $\Omega_f^{\varepsilon}$ . Actually, in view of the boundary conditions on  $\Gamma_{\varepsilon,1}$  and  $\Gamma_{\varepsilon,3}$ , the constant  $C_f$  is  $\varepsilon$  independent.

For  $I_4$  we proceed similarly, to obtain

$$I_4 \ge m \left| \left| \nabla u_p^k \right| \right|_{\Omega_p}^2 \ge \frac{m}{2} \left| \left| \nabla u_p^k \right| \right|_{\Omega_p}^2 + \frac{m}{2C_p} \left| \left| u_p^k \right| \right|_{\Omega_p}^2$$

where  $C_p$  depends only on the geometry of  $\Omega_p$ .

The convection term  $I_5$  vanishes. Indeed,

$$\begin{aligned} (\mathbf{q} \cdot \nabla u_f^k, u_f^k)_{\Omega_f^{\varepsilon}} &= (\mathbf{q} \cdot \nabla u_f^k, u_f^k)_{\Omega_f^{\varepsilon}} = \frac{1}{2} (\mathbf{q}, (\nabla u_f^k)^2)_{\Omega_f^{\varepsilon}} \\ &= \frac{1}{2} ((u_f^k)^2 \mathbf{q}, \boldsymbol{\nu})_{\partial \Omega_f^{\varepsilon}} - \frac{1}{2} ((u_f^k)^2, \nabla \cdot \mathbf{q})_{\Omega_f^{\varepsilon}} = 0, \end{aligned}$$

by the boundary conditions on  $\partial \varOmega_f^\varepsilon$  and the properties of  ${\bf q}$ 

Using the Cauchy-Schwarz inequality and Young's inequality gives for  $I_6$  and  $I_7$ 

$$|I_{6}| \leq ||f_{f}(t_{k})||_{\Omega_{f}^{\varepsilon}} ||u_{f}^{k}||_{\Omega_{f}^{\varepsilon}} \leq \frac{1}{2C_{f}} ||u_{f}^{k}||_{\Omega_{f}^{\varepsilon}}^{2} + \frac{C_{f}}{2} ||f_{f}(t_{k})||_{\Omega_{f}^{\varepsilon}}^{2}, \text{ and} |I_{7}| \leq M ||f_{p}(t_{k})||_{\Omega_{p}} ||u_{p}^{k}||_{\Omega_{p}} \leq M^{2} \frac{C_{p}}{2m} ||f_{p}(t_{k})||_{\Omega_{p}}^{2} + \frac{m}{2C_{p}} ||u_{p}^{k}||_{\Omega_{p}}^{2}$$

Using the estimates above into (10) and summing for k = 1, ..., j (where  $j \le N$  is arbitrary) gives

$$\begin{split} &\frac{1}{2} \left| \left| u_{f}^{j} \right| \right|_{\Omega_{f}}^{2} + \frac{1}{2} \sum_{k=1}^{j} \left| \left| u_{f}^{k} - u_{f}^{k-1} \right| \right|_{\Omega_{f}^{\varepsilon}}^{2} + \int_{\Omega_{p}} R(u_{p}^{j}) dx \\ &+ \frac{\Delta t}{2} \sum_{k=1}^{j} \left| \left| \nabla u_{f}^{k} \right| \right|_{\Omega_{f}^{\varepsilon}}^{2} + \frac{m \Delta t}{2} \sum_{k=1}^{j} \left| \left| \nabla u_{p}^{k} \right| \right|_{\Omega_{p}}^{2} \\ &\leq \frac{1}{2} \left| \left| u_{f}^{0} \right| \right|_{\Omega_{f}^{\varepsilon}}^{2} + \int_{\Omega_{p}} R(u_{p}^{0}) dx + \Delta t \frac{C_{p}}{2} \sum_{k=1}^{j} \left| \left| f_{f}(t_{k}) \right| \right|_{\Omega_{f}^{\varepsilon}}^{2} + \Delta t \frac{C_{p} M^{2}}{2m} \sum_{k=1}^{j} \left| \left| f_{p}(t_{k}) \right| \right|_{\Omega_{p}}^{2}. \end{split}$$

In view of assumptions  $A_I$  and  $A_f$ , the sums on the right are bounded uniformly in  $\Delta t$  and j. This proves the estimates.

In a similar fashion one obtains

**Lemma 2** For the sequence of time discrete weak solutions in Definition 4, a C > 0 not depending on  $\Delta t$  exists s.t.

$$\max_{j \in \{1,\dots,N\}} \left| \left| u_p^j \right| \right|_{\Omega_p}^2 + \sum_{k=1}^N \left| \left| u_p^k - u_p^{k-1} \right| \right|_{\Omega_p}^2 \le C.$$
(12)

Proof As before,  $\phi_f := r^{-1}(u_f^k) \in V_f^{\varepsilon}$ , and it can be used together with  $\phi_p := u_p^k$  as test functions in (10). The proof follows as above, but involves the antiderivative  $R^*$  of  $r^{-1}$  for showing that the convection term vanishes:

$$(\mathbf{q} \cdot \nabla u_f^k, r^{-1}(u_f^k))_{\Omega_f^{\varepsilon}} = (\mathbf{q}, \nabla R^*(u_{\varepsilon,f}^k))_{\Omega_f^{\varepsilon}},$$

and the rest follows as before.

#### 3.3 Interpolation in time and convergence

Having obtained the a priori estimates, the time discrete pairs are used to construct piecewise linear and piecewise constant interpolations in time. More precisely, for almost every  $t \in (t_{k-1}, t_k]$ , we define

$$\bar{U}_{\Delta t}^{f}(t) := u_{f}^{k}, \quad \hat{U}_{\Delta t}^{f}(t) := u_{f}^{k-1} + \frac{t - t_{k-1}}{\Delta t} (u_{f}^{k} - u_{f}^{k-1}), \quad (13)$$

$$\bar{U}^{p}_{\Delta t}(t) := u^{k}_{p}, \quad \hat{U}^{p}_{\Delta t}(t) := u^{k-1}_{p} + \frac{t - t_{k-1}}{\Delta t} (u^{k}_{p} - u^{k-1}_{p}). \tag{14}$$

Further, the piecewise constant interpolation of the source terms will be used

$$\bar{f}^i_{\Delta t}(t) := f_i(t_k).$$

For the ease of writing, we take  $i \in \{p, f\}$ , and omit the superscript  $\varepsilon$  for the quantities, domains, or spaces involving the fracture. Using the a priori estimates in Lemmata 1 and 2, one gets for  $\overline{U}_{\Delta t}^{i}$ :

**Lemma 3**  $\{\overline{U}_{\Delta t}^i\}_{\Delta t>0}$  is bounded uniformly w.r.t  $\Delta t$  in  $L^{\infty}(0,T;L^2(\Omega_i)) \cap L^2(0,T;V_i)$ .

Due to the a priori bounds in Lemma 3, there exists a subsequence (along  $\Delta t \searrow 0$ ) of the time interpolations in (13)–(14) that converges weakly in  $L^2(0,T;V_f) \times L^2(0,T;V_p)$ . Here we show that the weak limit is a solution in the sense of Definition 3. We start with a strong convergence result, which is needed for the convergence on the boundary  $\Gamma$ . In doing so, we use the following result (Lemma 3.2 in [23]).

**Lemma 4** Let X be a Hilbert space. The strong convergence

$$\hat{U}^i_{\Delta t} \to U_i \qquad \text{in } L^2(0,T;X)$$

implies the strong convergence

$$\bar{U}^i_{\Delta t} \to U_i \qquad \text{in } L^2(0,T;X). \tag{15}$$

Based on this, we obtain the strong convergence in  $L^2(0,T;L^2(\Omega_i))$ .

**Lemma 5** Along a sequence  $\Delta t \searrow 0$ , the piecewise constant in time approximations  $(\bar{U}_{\Delta t}^f, \bar{U}_{\Delta t}^p)$  converge strongly in  $L^2(0,T; L^2(\Omega_f)) \times L^2(0,T; L^2(\Omega_p))$ .

Proof The proof is given for the fracture (i = f), the arguments for i = pbeing exactly the same. For  $t \in (t_{k-1}, t_k]$  we regard  $\partial_t \hat{U}_{\Delta t} = \frac{u_f^{k} - u_f^{k-1}}{\Delta t}$  as an element in  $L^2(0, T; W^{-1,2}(\Omega_f))$ , where  $W^{-1,2}(\Omega_f)$  is the dual of  $W_0^{1,2}(\Omega_f)$  (the  $W^{1,2}(\Omega_f)$  functions having a vanishing trace over the entire  $\partial \Omega_f$ ) identified by the duality pairing

$$\langle \partial_t \hat{U}^f_{\Delta t}(t), \phi_f \rangle_{W^{-1,2}(\Omega_f), W^{1,2}_0(\Omega_f)} = \frac{1}{\Delta t} (u_f^k - u_f^{k-1}, \phi_f)_{\Omega_f},$$

for all  $\phi_f \in W_0^{1,2}(\Omega_f)$ . Testing in (10) with  $\phi_p \equiv 0$  and an arbitrary  $\phi_f \in W_0^{1,2}(\Omega_f)$  gives

$$\begin{aligned} \langle \partial_t \hat{U}^f_{\Delta t}(t), \phi_f \rangle_{(W^{-1,2}(\Omega_f)), W^{1,2}_0(\Omega_f)} + (\nabla u^k_f, \nabla \phi_f)_{\Omega_f} \\ + (\mathbf{q} \cdot \nabla u^k_f, \phi_f)_{\Omega_f} &= (f_f(t_k), \phi_f)_{\Omega_f}, \end{aligned}$$

providing for  $t \in (t_{k-1}, t_k]$ 

$$\left|\left|\partial_t U_{\Delta t}^f(t)\right|\right|_{W^{-1,2}(\Omega_f)} \le (1+M_q) \left|\left|\nabla u_f^k\right|\right|_{\Omega_f} + \left|\left|f_f(t_k)\right|\right|_{\Omega_f} + \left|f_f(t_k)\right|_{\Omega_f} + \left|$$

Here  $M_q$  is the bound on the velocity profile, as introduced in Assumption  $A_q$ . Using now the estimates in Lemma 1 and the assumptions of  $f_f$  one gets that  $\partial_t U^f_{\Delta t}$  is bounded in  $L^2(0,T;W^{-1,2}(\Omega_f))$  uniformly w.r.t.  $\Delta t$ .

By [32], the above boundedness together with the uniform boundedness of  $\hat{U}_{\Delta t}^{f}$  in  $L^{2}(0,T;V_{f})$  provide the existence of a limit  $U_{f}$  s.t. along a sequence  $\Delta t \searrow 0$ ,

$$\hat{U}_{\Delta t}^f \to U_f, \qquad \text{strongly in } L^2(0,T;L^2(\Omega_f))$$

Further, Lemma 4 gives the strong convergence for  $\bar{U}_{\Delta t}^{f}$  to the same limit  $U_{f}$ , and the proof is finished.

By Lemmata 3 and 5, a sequence  $\Delta t$  exists s.t.

$$\bar{U}_{\Delta t}^{f} \to U_{f}, \text{ weakly in } L^{2}(0,T;V_{f}) \text{ and strongly in } L^{2}(0,T;L^{2}(\Omega_{f})),$$
 $\bar{U}_{\Delta t}^{p} \to U_{p}, \text{ weakly in } L^{2}(0,T;V_{p}) \text{ and strongly in } L^{2}(0,T;L^{2}(\Omega_{p})).$ 
(16)

We show that the limit pair is a weak solution

**Theorem 5** The pair  $(U_f, U_p)$  is a solution in the sense of Definition 3.

Proof We start by summing (10) from j = 1, ..., k. Recalling (13) and (14), for any  $(\phi_f, \phi_p) \in V_f \times V_p$  such that  $\phi_f = \phi_p$  on  $\Gamma$  one gets for every  $t \in (t_{k-1}, t_k)$ 

$$\begin{split} (\bar{U}_{\Delta t}^{f}(t),\phi_{f})_{\Omega_{f}} + (\bar{U}_{\Delta t}^{p}(t),\phi_{p})_{\Omega_{p}} + \int_{0}^{t} (\nabla \bar{U}_{\Delta t}^{f}(\tau),\nabla\phi_{f})_{\Omega_{f}}d\tau \\ &+ \int_{0}^{t} (\nabla \bar{U}_{\Delta t}^{p}(\tau),\nabla\phi_{p})_{\Omega_{p}}d\tau + \int_{0}^{t} (\mathbf{q}\cdot\nabla \bar{U}_{\Delta t}^{f}(\tau),\phi_{f})_{\Omega_{f}}d\tau \\ &- \int_{0}^{t} (\bar{f}_{f}(\tau),\phi_{f})_{\Omega_{f}}d\tau - \int_{0}^{t} (\bar{f}_{p}(\tau),\phi_{p})_{\Omega_{p}}d\tau - (u_{I,f},\phi_{f})_{\Omega_{f}} - (u_{I,p},\phi_{p})_{\Omega_{f}}d\tau \\ &= \int_{t}^{t_{k}} (\bar{f}_{f}(\tau),\phi_{f})_{\Omega_{f}}d\tau + \int_{t}^{t_{k}} (\bar{f}_{p}(\tau),\phi_{p})_{\Omega_{p}}d\tau - \int_{t}^{t_{k}} (\nabla \bar{U}_{\Delta t}^{f}(\tau),\nabla\phi_{f})_{\Omega_{f}}d\tau \\ &- \int_{t}^{t_{k}} (\nabla \bar{U}_{\Delta t}^{p}(\tau),\nabla\phi_{p})_{\Omega_{p}}d\tau - \int_{t}^{t_{k}} (\mathbf{q}\cdot\nabla \bar{U}_{\Delta t}^{f}(\tau),\phi_{f})_{\Omega_{f}}d\tau, \end{split}$$

The terms on the right are accounting for the fact that, actually, the upper limit in the time integrals on the left should be  $t_k$ . Next, take  $\phi_i \in L^2(0,T;V_i)$ , such that  $\phi_f = \phi_p$  on  $\Gamma$ , and integrate over [0,T], to obtain

$$\int_{0}^{T} (\bar{U}_{\Delta t}^{f}(t), \phi_{f}(t))_{\Omega_{f}} dt + \int_{0}^{T} (\bar{U}_{\Delta t}^{p}(t), \phi_{p}(t))_{\Omega_{p}} dt \\
+ \int_{0}^{T} \int_{0}^{t} (\nabla \bar{U}_{\Delta t}^{f}(\tau), \nabla \phi_{f}(t))_{\Omega_{f}} d\tau dt + \int_{0}^{T} \int_{0}^{t} (\nabla \bar{U}_{\Delta t}^{p}(\tau), \nabla \phi_{p}(t))_{\Omega_{p}} d\tau dt \\
+ \int_{0}^{T} \int_{0}^{t} (\mathbf{q} \cdot \nabla \bar{U}_{\Delta t}^{f}(\tau), \phi_{f}(t))_{\Omega_{f}} d\tau dt - \int_{0}^{T} \int_{0}^{t} (\bar{f}_{f}(\tau), \phi_{f}(t))_{\Omega_{f}} d\tau dt \\
- \int_{0}^{T} \int_{0}^{t} (\bar{f}_{p}(\tau), \phi_{p}(t))_{\Omega_{p}} d\tau dt - \int_{0}^{T} (u_{I,f}, \phi_{f}(t))_{\Omega_{f}} dt - \int_{0}^{T} (u_{I,p}, \phi_{p}(t))_{\Omega_{p}} dt \\
= \sum_{k=1}^{N} \int_{t_{k-1}}^{t_{k}} \int_{t}^{t_{k}} (\bar{f}_{f}(\tau), \phi_{f}(t))_{\Omega_{f}} d\tau dt + \sum_{k=1}^{N} \int_{t_{k-1}}^{t_{k}} \int_{t}^{t_{k}} (\bar{f}_{p}(\tau), \phi_{p}(t))_{\Omega_{p}} d\tau dt \\
- \sum_{k=1}^{N} \int_{t_{k-1}}^{t_{k}} \int_{t}^{t_{k}} (\nabla \bar{U}_{\Delta t}^{f}(\tau), \nabla \phi_{f}(t))_{\Omega_{f}} d\tau dt - \sum_{k=1}^{N} \int_{t_{k-1}}^{t_{k}} \int_{t}^{t_{k}} (\nabla \bar{U}_{\Delta t}^{p}(\tau), \nabla \phi_{p}(t))_{\Omega_{p}} d\tau dt \\
- \sum_{k=1}^{N} \int_{t_{k-1}}^{t_{k}} \int_{t}^{t_{k}} (\mathbf{q} \cdot \nabla \bar{U}_{\Delta t}^{f}(\tau), \phi_{f}(t))_{\Omega_{f}} d\tau dt. \tag{17}$$

By (16), for  $\Delta t \searrow 0$  one has

$$\int_0^T (\bar{U}^i_{\Delta t}(t), \phi_i(t))_{\Omega_i} dt \to \int_0^T (U_i(t), \phi_i(t))_{\Omega_i} dt, \tag{18}$$

$$\int_{0}^{T} \int_{0}^{t} (\nabla \bar{U}_{\Delta t}^{i}(\tau), \nabla \phi_{i}(t))_{\Omega_{i}} d\tau dt \rightarrow \int_{0}^{T} \int_{0}^{t} (\nabla U_{i}(\tau), \nabla \phi_{i}(t))_{\Omega_{i}} d\tau dt, \quad (19)$$
$$\int_{0}^{T} \int_{0}^{t} (\mathbf{q} \cdot \nabla \bar{U}_{\Delta t}^{f}(\tau), \phi_{f}(t))_{\Omega_{f}} d\tau dt \rightarrow \int_{0}^{T} \int_{0}^{t} (\mathbf{q} \cdot \nabla U_{f}(\tau), \phi_{f}(t))_{\Omega_{f}} d\tau dt. \quad (20)$$

Furthermore, since  $f_i \in C(0,T;L^2(\varOmega_i))$  we also have

$$\int_0^T \int_0^t (\bar{f}_i(\tau), \phi_i(t))_{\Omega_i} d\tau dt \to \int_0^T \int_0^t (f_i(\tau), \phi_i(t))_{\Omega_i} d\tau dt.$$
(21)

We denote the terms on the right hand side of (17) by  $T_1, ..., T_5$ . Since  $f_f \in C(0,T; L^2(\Omega_f))$ , for  $T_1$  one has

$$|T_1| \le (\Delta t)^2 \sum_{k=1}^N ||f_f(t_k)||_{\Omega_f}^2 + \frac{\Delta t}{4} ||\phi_f||_{\Omega_f}^2 \le C\Delta t,$$

with C independent of  $\Delta t$ . Therefore  $T_1$  is vanishing as  $\Delta t \searrow 0$ .

For  $T_3$  we use the priori estimate (12) to obtain

$$\begin{aligned} |T_f| &\leq \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \int_t^{t_k} \left| (\nabla \bar{U}_{\Delta t}^f(\tau), \nabla \phi_f(t))_{\Omega_f} \right| d\tau dt \\ &\leq \Delta t \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \left| |\nabla u_f^k| \right|_{\Omega_f} \left| |\nabla \phi_f(t)| \right|_{\Omega_i} dt \\ &\leq \Delta t \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \left| |\nabla u_f^k| \right|_{\Omega_f}^2 dt + \frac{\Delta t}{4} \sum_{k=1}^N \left| |\nabla \phi_f(t)| \right|_{\Omega_f}^2 dt \\ &\leq (\Delta t)^2 \sum_{k=1}^N \left| |\nabla u_f^k| \right|_{\Omega_f}^2 + \frac{\Delta t}{4} \int_0^T \left| |\nabla \phi_f(t)| \right|_{\Omega_f}^2 dt \leq C \Delta t, \end{aligned}$$

with C independent of  $\Delta t$ . This shows that  $T_3$  vanishes as well as  $\Delta t \searrow 0$ .

The proofs for  $T_2$ ,  $T_4$  and  $T_5$  are similar and we omit them here. This shows that for all  $(\phi_f, \phi_p) \in L^2(0, T; V_f) \times L^2(0, T; V_p)$  s.t.  $\phi_f = \phi_p$  on  $\Gamma$ , we have

$$\int_{0}^{T} (U_{f}(t), \phi_{f}(t))_{\Omega_{f}} dt + \int_{0}^{T} (U_{p}(t), \phi_{p}(t))_{\Omega_{p}} dt + \int_{0}^{T} \int_{0}^{t} (\nabla U_{f}(\tau), \nabla \phi_{f}(t))_{\Omega_{f}} d\tau dt \\
+ \int_{0}^{T} \int_{0}^{t} (\nabla U_{p}(\tau), \nabla \phi_{p}(t))_{\Omega_{p}} d\tau dt + \int_{0}^{T} \int_{0}^{t} (\mathbf{q} \cdot \nabla U(\tau), \phi_{f}(t))_{\Omega_{f}} d\tau dt \\
= \int_{0}^{T} \int_{0}^{t} (f_{f}(\tau), \phi_{f}(t))_{\Omega_{f}} d\tau dt + \int_{0}^{T} \int_{0}^{t} (f_{p}(\tau), \phi_{p}(t))_{\Omega_{p}} d\tau dt \qquad (22) \\
+ \int_{0}^{T} (u_{I,f}, \phi_{f}(t))_{\Omega_{f}} dt + \int_{0}^{T} (u_{I,p}, \phi_{p}(t))_{\Omega_{p}} dt.$$

Next, let  $\psi_i \in W^{1,2}(0,T;V_i)$  (i = f, p) s.t.  $\psi_f = \psi_p$  on  $\Gamma$  and  $\psi_f(T) = \psi_p(T) = 0$ . We observe that

$$\int_0^T \int_0^t (f_f(\tau), \partial_t \psi_f(t))_{\Omega_f} d\tau dt$$
  
=  $-\int_0^T \int_{\Omega_f} f_f(t, x) \psi_f(t, x) dx dt + \int_{\Omega_f} \left[ \int_0^t f_f(\tau, x) d\tau \ \psi_f(t, x) \right]_{t=0}^{t=T} dx$   
=  $-\int_0^T (f_f(t), \psi_f(t))_{\Omega_f} dt.$ 

Analogously, one gets similar results for the integrals involving  $f_p,$  as well as

$$\int_0^T \int_0^t (\nabla U_i(\tau), \nabla \partial_t \psi_i(t))_{\Omega_i} d\tau dt = -\int_0^T (\nabla U_i(t), \nabla \psi_i(t))_{\Omega_i} dt \ (i = f, p)$$
$$\int_0^T \int_0^t (\mathbf{q} \cdot \nabla U_f(\tau), \partial_t \psi_f(t))_{\Omega_f} d\tau dt = -\int_0^T (\mathbf{q} \cdot \nabla U_f(t), \psi_f(t))_{\Omega_f} dt.$$

With  $\phi_i := \partial_t \psi_i$  as test functions in (22) gives

$$\begin{split} &\int_0^T (U_f(t), \partial_t \psi_f(t))_{\Omega_f} dt + \int_0^T (U_p(t), \partial_t \psi_p(t))_{\Omega_p} dt - \int_0^T (\nabla U_f(t), \nabla \psi_f(t))_{\Omega_f} dt \\ &- \int_0^T (\nabla U_p(t), \nabla \psi_p(t))_{\Omega_p} dt - \int_0^T (\mathbf{q} \cdot \nabla U_f(t), \psi_f(t))_{\Omega_f} dt \\ &= - \int_0^T (f_f(t), \psi_f(t))_{\Omega_f} dt - \int_0^T (f_p(t), \psi_p(t))_{\Omega_p} dt \\ &- (u_{I,f}, \psi_f(0))_{\Omega_f} - (u_{I,p}, \psi_p(0))_{\Omega_p}. \end{split}$$

Therefore, (9) holds true.

It only remains to show that  $U_f = r(U_p)$  on  $\Gamma$ . In doing so, we estimate

$$\begin{aligned} \left| \left| U_{f} - r(U_{p}) \right| \right|_{\Gamma^{T}} &\leq \left| \left| U_{f} - \bar{U}_{\Delta t}^{f} \right| \right|_{\Gamma^{T}} \\ &+ \left| \left| r(U_{p}) - r(\bar{U}_{\Delta t}^{p}) \right| \right|_{\Gamma^{T}} + \left| \left| \bar{U}_{\Delta t}^{f} - r(\bar{U}_{\Delta t}^{p}) \right| \right|_{\Gamma^{T}}. \end{aligned}$$

$$(23)$$

The third term on the right vanishes since, by definition,  $\bar{U}_{\Delta t}^f = r(\bar{U}_{\Delta t}^p)$  on  $\Gamma$ . For the second term, by the trace inequality one has

$$\begin{aligned} \left| \left| r(U_p) - r(\bar{U}_{\Delta t}^p) \right| \right|_{\Gamma^T}^2 &\leq M^2 \int_0^T \left| \left| U_p(t) - \bar{U}_{\Delta t}^p(t) \right| \right|_{\Gamma}^2 dt \\ &\leq C \left| \left| U_p - \bar{U}_{\Delta t}^p \right| \right|_{\Omega_p^T} \left( \left| \left| \nabla U_p - \nabla \bar{U}_{\Delta t}^p \right| \right|_{\Omega_p^T} + \left| \left| U_p - \bar{U}_{\Delta t}^p \right| \right|_{\Omega_p^T} \right), \quad (24) \end{aligned}$$

with C only depending on the geometry of  $\Omega_p$ . As  $\Delta t \searrow 0$ ,  $||U_p - \bar{U}_{\Delta t}^p||_{\Omega_p^T} \to 0$ by the strong convergence in Lemma 5. The weak convergence (16) implies that  $||\nabla U_p - \nabla \bar{U}_{\Delta t}^p||_{\Omega_p^T}$  is bounded uniformly, and by (24) the second term on the right in (23) vanishes. A similar argument can be applied to the first term on the right, showing that the traces of  $U_f$  and  $r(U_p)$  are equal a.e. on  $\Gamma$ . This concludes the proof.

#### 3.4 Positivity and $L^{\infty}$ bounds

As stated in Assumption  $A_I$ , the initial conditions are positive and essentially bounded. Since the Dirichlet boundary conditions are homogeneous one expects that the solution is positive and essentially bounded as well. This is proved below. We start with the proof for the time discrete concentrations  $u_f^k$ and  $u_p^k$ , which immediately give similar results for  $u_f$  and  $u_p$ . The procedure is quite standard and makes use of the nonpositive, respectively nonnegative cuts,  $[\cdot]_+$  and  $[\cdot]_-$ :

$$[u]_{+} := \begin{cases} 0, & \text{if } u \leq 0, \\ u, & \text{if } u > 0, \end{cases} \qquad [u]_{-} := \begin{cases} u, & \text{if } u < 0, \\ 0, & \text{if } u \geq 0. \end{cases}$$
(25)

We start with the lower bounds, which are proved in

)

**Lemma 6** For any  $k \in \{0, \ldots, N\}$ ,  $u_f^k \ge 0$  and  $u_p^k \ge a.e.$ .

Proof The proof is by mathematical induction. By Assumption  $A_I$ , the statement holds for k = 0. Next, assuming  $u_f^{k-1} \ge 0$  and  $u_p^{k-1} \ge 0$  a.e., the same is obtained for  $u_f^k$  and  $u_p^k$  when testing in (10) with  $\phi_f := [u_f^k]_-$  and  $\phi_p := [r(u_p^k)]_-$ . We omit the detailed arguments here, as these are standard.

For the upper bounds, with M,  $M_f$ ,  $M_I$  introduced in the assumptions, we let  $M_u := \max\{M_I, r(M_I), M_f, MM_f\}$  and prove

**Lemma 7** For all  $k \in \{0, \ldots, N\}$  one has

 $u_f^k \le M_u(k\Delta t+1)$  and  $u_p^k \le r^{-1} (M_u(k\Delta t+1)).$ 

Proof Here we use again mathematical induction and follow ideas in [8]. The statement holds trivially for k = 0. Next, assume  $u_f^k \leq M_u((k-1)\Delta t+1)$  and  $u_p^k \leq r^{-1} \left( M_u((k-1)\Delta t+1) \right)$ . Since  $u_f^k = r(u_p^k)$  on  $\Gamma$ ,  $\phi_f := [u_f^k - M_u(k\Delta t+1)]_+$  and  $\phi_p := [r(u_p^k) - M_u(k\Delta t+1)]_+$  can be used as test functions in (10), giving

$$(u_{f}^{k} - M_{u}(k\Delta t + 1), [u_{f}^{k} - M_{u}(k\Delta t + 1)]_{+})_{\Omega_{f}} + (u_{p}^{k} - r^{-1}(M_{u}(k\Delta t + 1)), [r(u_{p}^{k}) - M_{u}(k\Delta t + 1)]_{+})_{\Omega_{p}} + \Delta t (\nabla (u_{f}^{k} - M_{u}k(\Delta t + 1)), \nabla [u_{f}^{k} - M_{u}k(\Delta t + 1)]_{+})_{\Omega_{f}} + \Delta t (\nabla (u_{p}^{k} - M_{u}k(\Delta t + 1)), \nabla [r(u_{p}^{k}) - M_{u}k(\Delta t + 1)]_{+})_{\Omega_{f}} + \Delta t (\mathbf{q} \cdot \nabla (u_{f}^{k} - M_{u}(k\Delta t + 1)), [u_{f}^{k} - M_{u}(k\Delta t + 1)]_{+})_{\Omega_{f}} = (u_{f}^{k-1} - M_{u}((k-1)\Delta t + 1), [u_{f}^{k} - M_{u}(k\Delta t + 1)]_{+})_{\Omega_{f}} + (u_{p}^{k-1} - r^{-1}(M_{u}((k-1)\Delta t + 1)), [u_{p}^{k} - M_{u}(k\Delta t + 1)]_{+})_{\Omega_{p}} + (\Delta t f_{p} - r^{-1}(M_{u}(k\Delta t + 1)) + r^{-1}(M_{u}((k-1)\Delta t + 1)), [u_{p}^{k} - M_{u}(k\Delta t + 1)]_{+})_{\Omega_{p}} + \Delta t (f_{f} - M_{u}, [u_{f}^{k} - M_{u}(k\Delta t + 1)]_{+})_{\Omega_{f}}.$$

$$(26)$$

The first three terms in (26) are positive, while for the fourth one uses the properties of r to obtain

$$\left(\nabla(u_p^k - M_u k(\varDelta t+1)), \nabla[r(u_p^k) - M_u k(\varDelta t+1)]_+\right)_{\varOmega_p} \ge m \|\nabla[u_p^k - M_u k(\varDelta t+1)]_+)\|_{\varOmega_p}^2$$

As in the proof of Lemma 1, the last term on the left vanishes.

Also, by the induction hypothesis, the first two terms on the right are nonpositive, and the same holds for the third by Assumption  $A_f$  and since  $M_u \ge M_f$ . Finally, for a.e.  $x \in \Omega_p$ , a  $\xi_x \in (M_u((k-1)\Delta t+1), M_u(k\Delta t+1))$ exists s.t.

$$\Delta t f_p - \left(r^{-1}(M_u(k\Delta t+1)) - r^{-1}(M_u((k-1)\Delta t+1))\right)$$
$$= \Delta t f_p - (r^{-1})'(\xi_x) M_u \Delta t \le \Delta t M_{f_p} - \frac{1}{M} M_u \Delta t \le 0.$$

Therefore the last term is nonpositive as well, showing that  $u_f^k \leq M_u(k\Delta t+1)$ and  $u_p^k \leq r^{-1}(M_u(k\Delta t+1))$  a.e..

Lemmata 6 and 7 provide similar bounds for the solution pair  $(u_f, u_p)$ .

**Lemma 8** The solution pair in Definition 3 is essentially bounded. For a.e.  $t \in [0,T]$  one has

$$0 \le u_f(t) \le M_u(t+1)$$
 almost everywhere in  $\Omega_f$ , (27)

$$0 \le u_p(t) \le \frac{M_u}{m}(t+1)$$
 almost everywhere in  $\Omega_p$ . (28)

*Proof* The proof uses the convergence of the interpolations in (13) and (14). In view of the results above, these are nonnegative a.e.. Further, the upper bounds follow straightforwardly for the interpolations in the fracture subdomain, and by using Remark 1 for the interpolation in the porous block.

#### 4 Formal upscaling

We use the formal asymptotic expansions for the variables and use the transverse averaging to obtain the upscaled equations. Specifically, we let the fracture thickness  $\varepsilon$  go to zero and reduce the fracture model to a boundary condition. We refer to [9,25] for a general procedure applied to convection dominated regimes, and to [19,30,31] for more specific applications related to precipitation-dissolution models, or to biofilm growth in porous media. However, these papers refer strictly to the fracture region and do not consider the coupling with a porous block. For simplicity, a Poiseuille flow is considered, but the procedure can be applied to more general situations straightforwardly:

$$\mathbf{q} = (q^{\varepsilon}(y), 0), \qquad \text{with } q^{\varepsilon}(y) = \frac{3}{2}Q\frac{y}{\varepsilon}\left(2 - \frac{y}{\varepsilon}\right).$$
 (29)

Here Q is the average of the fluid velocity in the longitudinal direction,  $Q = \frac{1}{2\varepsilon} \int_0^{2\varepsilon} q_{if}^{\varepsilon}(y) dy$ , with Q > 0. By this choice, the given velocity field satisfied the Stokes equation, and its transversal average does not vanish as  $\varepsilon \searrow 0$ .

For the upscaling we use the transformation  $z := y/\varepsilon$  to rescale the fracture domain to  $\Omega_f = (0, 1) \times (0, 1)$ . However, the porous block  $\Omega_p$  remains unchanged. For consistency, we change the name of the transversal variable yinto z there as well. In this context, the upper part of the boundary becomes  $\Gamma_f^2 = (0, 1) \times \{1\}$ , and (6) transforms into

$$\begin{cases} \partial_t u_p^{\varepsilon} - \Delta u_p^{\varepsilon} = f_p, & \text{in } (0, T] \times \Omega_p, \\ \partial_t u_f^{\varepsilon} + q^{\varepsilon} \partial_x u_f^{\varepsilon} - (\partial_{xx} u_f^{\varepsilon} + \frac{1}{\varepsilon^2} \partial_{zz} u_f^{\varepsilon}) = f_f, & \text{in } (0, T] \times \Omega_f, \\ u_f^{\varepsilon} = r(u_p^{\varepsilon}), & \text{at } (0, T] \times \Gamma, \\ \partial_z u_p^{\varepsilon} = \frac{1}{\varepsilon} \partial_z u_f^{\varepsilon}, & \text{at } (0, T] \times \Gamma, \\ \partial_z u_f^{\varepsilon} = 0, & \text{at } (0, T] \times \Gamma_f^2. \end{cases}$$
(30)

Observe that the partial derivatives  $\frac{\partial}{\partial y}$  become now  $\frac{1}{\varepsilon}\frac{\partial}{\partial z}$ , and that  $q^{\varepsilon} = q^{\varepsilon}(z) = \frac{3}{2}Qz(2-z)$ .

In the fracture domain we redefine  $\tilde{u}_f^{\varepsilon}$  in terms of the new variable z,  $\tilde{u}_f^{\varepsilon}(t, x, z) = u_f^{\varepsilon}(t, x, \varepsilon z)$ . By an abuse of notation allowing to avoid an excess of symbols, we give up the  $\tilde{}$ . Further, we assume the following asymptotic expansions for the pair  $(u_f^{\varepsilon}, u_p^{\varepsilon})$ :

$$u_f^{\varepsilon}(t, x, z) = u_0^f(t, x, z) + \varepsilon u_1^f(t, x, z) + \mathcal{O}(\varepsilon^2),$$
(31)

$$u_p^{\varepsilon}(t, x, z) = u_0^p(t, x, z) + \varepsilon u_1^p(t, x, y) + \mathcal{O}(\varepsilon^2).$$
(32)

Substituting the expansions (31) and (32) in (30) gives

$$\partial_t (u_0^p + \varepsilon u_1^p) - \Delta (u_0^p + \varepsilon u_1^p) = f_p + \mathcal{O}(\varepsilon^2), \quad \text{in } (0, T] \times \Omega_p, \qquad (33)$$
$$\partial_t u_0^f + \frac{3}{2} Q z(2-z) \partial_x u_0^f - \partial_{xx} u_0^f$$

$$-\frac{1}{\varepsilon^2}\partial_{zz}(u_0^f + \varepsilon u_1^f + \varepsilon^2 u_2^f) = f_f + \mathcal{O}(\varepsilon), \quad \text{in } (0,T] \times \Omega_f, \quad (34)$$

$$u_0^f + \varepsilon u_1^f = r(u_0^p) + \varepsilon r'(u_0^p) u_1^p + \mathcal{O}(\varepsilon^2), \quad \text{at } (0,T] \times \Gamma,$$
(35)

$$\partial_z (u_0^p + \varepsilon u_1^p) = \frac{1}{\varepsilon} \partial_z (u_0^f + \varepsilon u_1^f + \varepsilon^2 u_2^f) + \mathcal{O}(\varepsilon), \quad \text{at } (0, T] \times \Gamma, \tag{36}$$

$$\partial_z (u_0^f + \varepsilon u_1^f + \varepsilon^2 u_2^f) = \mathcal{O}(\varepsilon^3), \quad \text{at } (0,T] \times \Gamma_f^2.$$
 (37)

The  $\varepsilon^{-2}$  term in (34), and the lowes order terms in (36) and (37) imply

$$\partial_{zz} u_0^f = 0, \quad \text{in } \Omega_f, \qquad \text{and} \qquad \partial_z u_0^f = 0, \quad \text{at } \Gamma_f^2 \cup \Gamma.$$

This gives

$$u_0^f(x,z,t) = u_0^f(x,t).$$

In a similar fashion one gets

$$u_1^f(x, z, t) = u_1^f(x, t).$$

Hence, from equation (36), we obtain

$$\partial_z u_0^p = 0, \qquad \text{at } (0,T] \times \Gamma.$$
 (38)

The  $\varepsilon^0$  term of (35) gives

$$u_0^f = r(u_0^p), \quad \text{at } (0, T] \times \Gamma.$$
 (39)

Furthermore, the  $\varepsilon^0$  term of (33) implies

$$\partial_t u_0^p - \Delta u_0^p = f_p, \qquad \text{in } (0, T] \times \Omega_p.$$
(40)

From the above, one obtains the effective model

$$(P_{\mathrm{Up}_{0}}) \begin{cases} \partial_{t} u_{0}^{p} - \Delta u_{0}^{p} = f_{p}, & \text{in } (0, T] \times \Omega_{p}, \\ \partial_{z} u_{0}^{p} = 0, & \text{at } (0, T] \times \Gamma, \\ u_{0}^{f} = r(u_{0}^{p}), & \text{at } (0, T] \times \Gamma. \end{cases}$$
(41)

The model above is completed by the initial conditions and the (homogeneous) Dirichlet boundary conditions as stated in Section 2. Note that  $u_0^p$  can be solved independently of  $u_0^f$ . The last boundary condition in 41 provides the effective concentration in the fracture,  $u_0^f$ .

#### 4.1 Higher order approximation

The procedure above can be continued to find effective equations having better approximation properties than (41). To this end, we consider the  $\varepsilon^0$  term of equation (34),

$$\partial_t u_0^f - Q \frac{3}{2} z(z-2) \partial_x u_0^f - \partial_{xx} u_0^f - \partial_{zz} u_2^f = f_f \qquad \text{in } \Omega_f.$$
(42)

Integrating (42) from z = 0 to z = 1, since  $u_0^f$  does not depend on z and using (37) one gets

$$\partial_t u_0^f + Q \partial_x u_0^f - \partial_{xx} u_0^f - \bar{f}_f = \partial_z u_2^f \Big|_{z=0}^{z=1}$$

for  $(t, x) \in (0, T] \times (0, 1)$ , where  $\bar{f}_f := \int_{-1}^0 f_f dz$ . By (36) and (37), this becomes

$$-\partial_z u_1^p \big|_{z=0} = \partial_t u_0^f + Q \partial_x u_0^f - \partial_{xx} u_0^f - \bar{f}_f.$$

$$\tag{43}$$

The  $\varepsilon^1$  term of (35) gives

$$u_1^f = r'(u_0^p)u_1^p, \qquad \text{at } (0,T] \times \Gamma,$$
 (44)

and the  $\varepsilon^1$  term of (33) implies

$$\partial_t u_1^p - \Delta u_1^p = 0, \qquad \text{in } (0, T] \times \Omega_p.$$
 (45)

Defining the effective quantities

$$u_e^p := u_0^p + \varepsilon u_1^p, \quad \text{and} \quad u_e^f := u_0^f + \varepsilon u_1^f, \tag{46}$$

adding (38) to (43), (39) to (44), and (40) to (45) gives

$$u_e^f = r(u_0^p) + \varepsilon r'(u_0^p) u_1^p, \qquad \text{at } (0,T] \times \Gamma, \qquad (48)$$

$$-\partial_z u_e^p = \varepsilon \Big( \partial_t u_0^f + Q \partial_x u_0^f - \partial_{xx} u_0 - \bar{f}_f \Big), \qquad \text{at } (0, T] \times \Gamma.$$
(49)

Up to  $\varepsilon^2$  terms, (47)-(49) becomes

$$(P_{\mathrm{Up}_{1}}) \begin{cases} \partial_{t} u_{e}^{p} = \Delta u_{e}^{p} + f_{p}, & \text{in } (0,T] \times \Omega_{p}, \\ -\partial_{z} u_{e}^{p} = \varepsilon (\partial_{t} u_{e}^{f} + Q \partial_{x} u_{e}^{f} - \partial_{xx} u_{e}^{f} - \bar{f}_{f}), & \text{at } (0,T] \times \Gamma, \\ u_{e}^{f} = r(u_{e}^{p}), & \text{at } (0,T] \times \Gamma. \end{cases}$$
(50)

As before, initial and boundary conditions complete the model. Furthermore, substituting the third equation in the second one leads to a (nonlinear) model only in terms of  $u_e^p$ , and again can be solved independently. Then  $u_e^f$  can be determined from the third equation.

# 4.2 Formal upscaling of drift dominated model

As for the original model, we now determine upscaled (effective) equations for the model with dominating drift, where  $Pe = \mathcal{O}(\varepsilon^{-1})$ . Our approach is inspired from similar exercises carried out for precipitation-dissolution type models in [19,25], but uses the drift model discussed e.g. in [1,2,3], and assumes that the velocity is now very high, i.e. of order  $\varepsilon^{-1}$ . The system of equations is given by

$$\begin{cases} \partial_t u_p^{\varepsilon} - \Delta u_p^{\varepsilon} = f_p^{\varepsilon}, & \text{in } (0, T] \times \Omega_p, \\ \partial_t u_f^{\varepsilon} + \frac{1}{\varepsilon} \mathbf{q}^{\varepsilon} \cdot \nabla u_f^{\varepsilon} - \Delta u_f^{\varepsilon} = f_f^{\varepsilon}, & \text{in } (0, T] \times \Omega_f^{\varepsilon}, \\ u_f^{\varepsilon} = r(u_p^{\varepsilon}), & \text{at } (0, T] \times \Gamma, \\ \partial_{\boldsymbol{\nu}} u_p^{\varepsilon} = \partial_{\boldsymbol{\nu}} u_f^{\varepsilon}, & \text{at } (0, T] \times \Gamma. \end{cases}$$

$$(51)$$

We proceed agian by rescaling the fractured domain. Using the asymptotic expansions (31) - (32), (51) becomes

$$\partial_t (u_0^p + \varepsilon u_1^p) - \Delta (u_0^p + \varepsilon u_1^p) = f_p + \mathcal{O}(\varepsilon^2), \quad \text{in } (0, T] \times \Omega_p, \tag{52}$$
$$\partial_t u_0^f + \frac{3}{2} Q_2 (2-z) \partial_\tau u_0^f - \partial_{\tau \tau} u_0^f$$

$$-\frac{1}{\varepsilon^2}\partial_{zz}(u_0^f + \varepsilon u_1^f + \varepsilon^2 u_2^f) = f_f + \mathcal{O}(\varepsilon), \quad \text{in } (0, T] \times \Omega_f, \quad (53)$$

$$u_0^f + \varepsilon u_1^f = r(u_0^p) + \varepsilon r'(u_0^p) u_1^p + \mathcal{O}(\varepsilon^2), \quad \text{at } (0,T] \times \Gamma, \tag{54}$$

$$\partial_{z}(u_{0}^{p} + \varepsilon u_{1}^{p}) = \frac{1}{\varepsilon} \partial_{z}(u_{0}^{f} + \varepsilon u_{1}^{f} + \varepsilon^{2} u_{2}^{f}) + \mathcal{O}(\varepsilon), \quad \text{at } (0, T] \times \Gamma, \quad (55)$$
$$\partial_{z}(u_{0}^{f} + \varepsilon u_{1}^{f} + \varepsilon^{2} u_{2}^{f}) = \mathcal{O}(\varepsilon^{3}), \quad \text{at } (0, T] \times \Gamma_{f}^{2}. \quad (56)$$

As before, we obtain  $u_0^f(x, z, t) = u_0^f(x, t)$ . Further, the  $\varepsilon^{-1}$  terms in (53) give

$$-\frac{3}{2}Qz(z-2)\partial_x u_0^f - \partial_{zz}u_1^f = 0, \qquad \text{in } \Omega_f.$$
(57)

Integrating (57) from z = 0 to z = 1, using that  $u_0$  is independent of z and the boundary condition (56) yields

$$-Q\partial_x u_0^f = \partial_z u_1^f \big|_{z=0}.$$

Using the  $\varepsilon^0$  terms in (55), this gives

$$-\partial_z u_0^p = Q \partial_x u_0^f, \qquad \text{at } (0,T] \times \Gamma.$$
(58)

Furthermore, the  $\varepsilon^0$  term from (54) implies

$$u_0^f = r(u_0^p), \qquad \text{on } \Gamma$$

and the  $\varepsilon^0$  term of (52) gives

$$\partial_t u_0^p - \Delta u_0^p = f_p, \qquad \text{in } \Omega_p$$

Summarizing, the leading order terms solve the model

$$(P_{\mathrm{Dr}_{0}}) \begin{cases} \partial_{t} u_{0}^{p} - \Delta u_{0}^{p} = f_{p}, & \text{in } (0, T] \times \Omega_{p}, \\ -\partial_{z} u_{0}^{p} = Q \partial_{x} u_{0}^{f}, & \text{at } (0, T] \times \Gamma, \\ u_{0}^{f} = r(u_{0}^{p}), & \text{at } (0, T] \times \Gamma. \end{cases}$$
(59)

Boundary and initial conditions are needed to complete the model, which, as before can be reduced to the a (nonlinear) model in  $u_0^p$ .

Further, a more accurate model, is obtained when including higher order terms. The  $\varepsilon^0$  term in (53) is

$$\partial_t u_0^f + \frac{3}{2} Qz(2-z) \partial_x u_1^f - \partial_{xx} u_0^f - \partial_{zz} u_2^f = f_f.$$

$$\tag{60}$$

Integrating (60) from z = 0 to z = 1 gives for  $(t, x) \in (0, T] \times (0, 1)$ 

$$\partial_t u_0^f - \int_0^1 \frac{3}{2} Qz(z-2) \partial_x u_1^f dz - \partial_{xx} u_0^f - \bar{f}_f = -\partial_z u_2^f \Big|_{z=0}^{z=1}$$

where  $\bar{f}_f = \int_0^1 f_f dz$ . Hence, by (55) and (56), we have

$$-\partial_z u_1^p \Big|_{z=0} = \partial_t u_0^f - \int_0^1 \frac{3}{2} Q z(z-2) \partial_x u_1^f dz - \partial_{xx} u_0^f - \bar{f}_f.$$
(61)

With the effective quantities

$$u_e^p := u_0^p + \varepsilon u_1^p$$
 and  $u_e^f := u_0^f + \varepsilon \overline{u}_1^f$ , where  $\overline{u}_1^f := \int_0^1 u_1^f dz$ ,

adding (58) and (61) gives

$$-\partial_z u_e^p = Q \partial_x u_0^f + \varepsilon \Big( \partial_t u_0^f - \int_0^1 \frac{3}{2} Q z(z-2) \partial_x u_1^f dz - \partial_{xx} u_0^f - \bar{f}_f \Big).$$
(62)

By (57), in  $(0,T] \times \Omega_f$  one has

$$\partial_z u_1^f = \frac{3}{2} Q \partial_x u_0^f \left(\frac{z^2 - z^3}{3}\right) + c_1(x, t).$$

Since  $\partial_z u_1 = 0$  at  $\Gamma_f^2$ , we get  $c_1(x,t) = -Q \partial_x u_0^f$ . This gives

$$u_1^f = -\frac{3}{2}Q(\frac{z^4}{12} - \frac{z^3}{3})\partial_x u_0^f - Qz\partial_x u_0^f + c_2(x,t).$$

Integrating the above from z = 0 to z = 1, and after some elementary calculations, one has  $c_2(x,t) = \bar{u}_1^f + \frac{2}{5}Q\partial_x u_0^f$ , hence

$$u_1^f(x,z,t) = -\frac{3}{2}Q(\frac{z^4}{12} - \frac{z^3}{3})\partial_x u_0^f - Qz\partial_x u_0^f + \bar{u}_1^f + \frac{2}{5}Q\partial_x u_0^f.$$
 (63)

This can be used to compute

$$\int_0^1 \frac{3}{2} Qz(z-2)\partial_x u_1^f dz = \frac{3}{35} Q^2 \partial_{xx} u_0^f - Q \partial_x \bar{u}_1^f.$$

Inserting this in (62) gives for  $(t, x) \in (0, T] \times (0, 1)$ 

$$\begin{aligned} -\partial_z u_e^p &= Q \partial_x u_0^f + \varepsilon \Big( \partial_t u_0^f - \frac{3}{35} Q^2 \partial_{xx} u_0^f + Q \partial_x \bar{u}_1^f - \partial_{xx} u_0^f - \bar{f}_f \Big) \\ &= Q \partial_x u_e^f + \varepsilon \Big( \partial_t u_0^f - \big(1 + \frac{3}{35} Q^2\big) \partial_{xx} u_0^f - \partial_{xx} u_0^f - \bar{f}_f \Big). \end{aligned}$$

Up to an  $O(\varepsilon^2)$  error, this rewrites

$$-\partial_z u_e^p = Q \partial_x u_e^f + \varepsilon \Big( \partial_t u_e^f - \Big(1 + \frac{3}{35}Q^2\Big) \partial_{xx} u_e^f - \bar{f}_f \Big).$$

Similarly, the nonlinear transmission condition becomes

$$r(u_p^{\varepsilon}|_{z=0}) = r(u_e^p|_{z=0}) + O(\varepsilon^2).$$

Furthermore, using the expression for  $u_1^f$  obtained in (63), we have

$$\begin{split} u_f^{\varepsilon}|_{z=0} &= u_e^f + \varepsilon (u_1^f|_{z=0} - \bar{u}_1^f) + O(\varepsilon^2) \\ &= u_e^f + \varepsilon \frac{2Q}{5} \partial_x u_0^f + O(\varepsilon^2) = u_e^f + \varepsilon \frac{2Q}{5} \partial_x u_e^f + O(\varepsilon^2). \end{split}$$

Adding the  $\varepsilon^0$  and  $\varepsilon^1$  terms of (52), gives

$$\partial_t u_e^p - \Delta u_e^p = f_p, \qquad \text{in } \Omega_p.$$

Summarizing, we have the following set of effective equations

$$(P_{\mathrm{Dr}_{1}}) \begin{cases} \partial_{t} u_{e}^{p} = \Delta u_{e}^{p} + f_{p}, & \text{in } (0, T] \times \Omega_{p}, \\ -\partial_{z} u_{e}^{p} = Q \partial_{x} u_{e}^{f} \\ + \varepsilon \Big( \partial_{t} u_{e}^{f} - \big(1 + \frac{3Q^{2}}{35}\big) \partial_{xx} u_{e}^{f} - \bar{f}_{f} \Big), & \text{at } (0, T] \times \Gamma, \end{cases}$$

$$(64)$$

$$r(u_{e}^{p}) = u_{e}^{f} + \frac{2Q}{5} \partial_{x} u_{e}^{f}, & \text{at } (0, T] \times \Gamma. \end{cases}$$

In this particular situation, the third equation gives

$$u_e^f = e^{-\frac{5}{2Q}x} \int_0^x \frac{5}{2Q} e^{\frac{5}{2Q}\xi} r(u_e^p) d\xi,$$
(65)

allowing again to decouple the problem for  $u_e^p$ .

# 5 Rigorous upscaling for moderate Peclét (Pe = O(1))

In this Section we give a rigorous convergence proof for the upscaled model obtained in Section 4, when diffusion and transport are in balance. We follow ideas from [11], pp.200-208 (a detailed presentation can be found in [4], Chapters 4 and 9; see also [18] for an extension of the results to periodic media, and [12] for a similar dimensionality reduction approach). As in the formal upscaling, to simplify the exposition we assume the parabolic velocity profile in (29). For the proofs we derive  $\varepsilon$ -independent a priori estimates, and use compactness arguments to pass to the limit. Note that most of these a priori estimates immediately follow from the ones already obtained when proving existence.

5.1 Weak formulation

As in Section 4, the fractured part is rescaled vertically by  $z = y/\varepsilon$ . Then 0 < z < 1 and the domain  $\Omega_f^{\varepsilon}$  transforms to  $\Omega_f = (0, 1) \times (0, 1)$ . Furthermore,  $q^{\varepsilon}(z) = q(z) := -\frac{3}{2}Qz(z-2)$ . We define,

$$\tilde{u}_f^\varepsilon(t,x,z):=u_f^\varepsilon(t,x,z\varepsilon), \ \hat{f}_f(t,x,z):=f_f(t,x,z\varepsilon), \ \tilde{u}_{I,f}(x,z):=u_{I,f}(x,z\varepsilon).$$

We have  $\tilde{u}_f^{\varepsilon} \in L^2(0,T;V_f)$ ,  $\tilde{u}_f^{\varepsilon} = r(u_p^{\varepsilon})$  on  $\Gamma$  and (3) becomes

$$-(u_{p}^{\varepsilon},\partial_{t}\phi_{p})_{\Omega_{p}^{T}}-\varepsilon(\tilde{u}_{f}^{\varepsilon},\partial_{t}\phi_{f})_{\Omega_{f}^{T}}+D_{p}(\nabla u_{p}^{\varepsilon},\nabla\phi_{p})_{\Omega_{p}^{T}}+\varepsilon D_{f}(\partial_{x}\tilde{u}_{f}^{\varepsilon},\partial_{x}\phi_{f})_{\Omega_{f}^{T}}$$
$$+\frac{D_{f}}{\varepsilon}(\partial_{z}\tilde{u}_{f}^{\varepsilon},\partial_{z}\phi_{f})_{\Omega_{f}^{T}}+\varepsilon(q\partial_{x}\tilde{u}_{f}^{\varepsilon},\phi_{f})_{\Omega_{f}^{T}}$$
$$=(f_{p}^{\varepsilon},\phi_{p})_{\Omega_{p}^{T}}+\varepsilon(\tilde{f}_{f}^{\varepsilon},\phi_{f})_{\Omega_{f}^{T}}+(u_{I,p},\phi_{p}(0))_{\Omega_{p}}+\varepsilon(\tilde{u}_{I,f},\phi_{f}(0))_{\Omega_{f}}$$
(66)

for all  $(\phi_f, \phi_p) \in W^{1,2}(0, T, V_f) \times W^{1,2}(0, T, V_p)$  such that  $\phi_p = \phi_f$  on  $\Gamma$ , and  $\phi_f(T) = \phi_p(T) = 0$ .

In the fracture, we define the (vertical) average

$$U_f^{\varepsilon}(t,x) := \int_0^1 \tilde{u}_f^{\varepsilon}(t,x,z) dz.$$

Furthermore, we let  $\Omega_{AV} := (0, 1)$ . Since  $\tilde{u}_f^{\varepsilon} \in L^2(0, T; V_f^{\varepsilon}), U_f^{\varepsilon} \in L^2(0, T; W_0^{1,2}(\Omega_{AV}))$ . For consistency of notation, we let  $U_p(t, x, z) := u_p(t, x, z)$ . We also define  $\bar{f}_f(t, x) := \int_0^1 \tilde{f}_f(t, x, z) dz$  and  $\bar{u}_{L_f}(x) := \int_0^1 \tilde{u}_{L_f}(x, z) dz$ .

$$\begin{split} \bar{f}_f(t,x) &:= \int_0^1 \tilde{f}_f(t,x,z) dz \text{ and } \bar{u}_{I,f}(x) := \int_0^1 \tilde{u}_{I,f}(x,z) dz. \\ \text{Below we show that as } \varepsilon \searrow 0, \text{ the pair } (U_p^\varepsilon, U_f^\varepsilon) \text{ converges towards } (U_p, U_f), \\ \text{the weak solution of the upscaled Problem } P_{Up_0} \text{ in } (41). \text{ This is defined below} \end{split}$$

**Definition 6** A pair  $(U_f, U_p) \in L^2(0, T; L^2(\Omega_{AV})) \times L^2(0, T; V_p)$  is a weak solution to Problem  $P_{U_{p_0}}$  if for the trace of  $U_p$  on  $\Gamma$  one has  $U_f = r(U_p)$ , and

$$-(U_p,\partial_t\phi_p)_{\Omega_p^T} + (\nabla U_p,\nabla\phi_p)_{\Omega_p^T} = (f_p,\phi_p)_{\Omega_p^T} + (u_{I,p},\phi_p(0))_{\Omega_p},$$

for all  $\phi_p \in W^{1,2}(0,T;V_p)$  such that  $\phi_p(T) = 0$ .

In the convergence proof the transversal average of the fracture model will be used. For estimating this we test in (66) with z-independent functions in the fracture, that is,  $\phi_f(t, x, z) = \phi_f(t, x)$ . This gives

$$-\varepsilon (U_{f}^{\varepsilon},\partial_{t}\phi_{f})_{(\Omega_{AV})^{T}} - (U_{p}^{\varepsilon},\partial_{t}\phi_{p})_{\Omega_{p}^{T}} + (\nabla U_{p}^{\varepsilon},\nabla\phi_{p})_{\Omega_{p}^{T}} + \varepsilon (\partial_{x}U_{f}^{\varepsilon},\partial_{x}\phi_{f})_{\Omega_{AV}^{T}} + \varepsilon Q(\partial_{x}U_{f}^{\varepsilon},\phi_{f})_{\Omega_{AV}^{T}} = \varepsilon (\bar{f}_{f},\phi_{f})_{\Omega_{AV}^{T}} + (f_{p},\phi_{p})_{\Omega_{p}^{T}} + \varepsilon (\int_{0}^{1} (Q-q)\partial_{x}u_{f}^{\varepsilon},\phi_{f})_{\Omega_{AV}^{T}} + \varepsilon (\bar{u}_{I,f},\phi_{f}(0))_{\Omega_{AV}} + (u_{I,p},\phi_{p}(0))_{\Omega_{AV}},$$
(67)

for all  $(\phi_f, \phi_p) \in L^2(0, T, W_0^{1,2}(\Omega_{AV})) \times L^2(0, T, V_p)$  such that  $\phi_p = \phi_f$  on  $\Gamma$ , and  $\phi_f(T) = \phi_p(T) = 0$ . We now pass to the limit  $\varepsilon \searrow 0$  in (67). To do so, we adapt the proofs in Section 3.2 to obtain the a priori estimates

**Lemma 9** There exists C > 0 independent of  $\varepsilon$  such that

$$\left|\left|u_{p}^{\varepsilon}\right|\right|_{L^{2}(0,T;V_{p})}^{2}+\varepsilon\left|\left|\partial_{x}\tilde{u}_{f}^{\varepsilon}\right|\right|_{\Omega_{f}^{T}}^{2}+\frac{1}{\varepsilon}\left|\left|\partial_{z}\tilde{u}_{f}^{\varepsilon}\right|\right|_{\Omega_{f}^{T}}^{2}\leq C.$$

Similar estimates can be obtained for  $U_p^{\varepsilon}$  and  $U_f^{\varepsilon}$ :

**Lemma 10** There exists a constant C > 0 not depending on  $\varepsilon$  such that

$$\left|\left|U_{p}^{\varepsilon}\right|\right|_{L^{2}(0,T;V_{p})}^{2}+\left|\left|U_{f}^{\varepsilon}\right|\right|_{\Omega_{AV}^{T}}^{2}+\varepsilon\left|\left|\partial_{x}U_{f}^{\varepsilon}\right|\right|_{\Omega_{AV}^{T}}^{2}\leq C$$

Moreover, the essential bounds in Lemma 8 remain valid for  $U_p^\varepsilon$  and  $U_f^\varepsilon.$  For all  $t\in(0,T],$ 

$$0 \le U_f^{\varepsilon}(t) \le M_u(t+1)$$
 almost everywhere in  $\Omega_{AV}$ , (68)

$$0 \le U_p^{\varepsilon}(t) \le \frac{M_u}{m}(t+1)$$
 almost everywhere in  $\Omega_p$ . (69)

This implies the weak convergence

$$U_p^{\varepsilon} \rightharpoonup U_p$$
, weakly in  $L^2(0,T;V_p)$ , (70)

$$U_f^{\varepsilon} \rightharpoonup U_f$$
, weakly in  $L^2(0,T; L^2(\Omega_{AV}))$ . (71)

Next, as for Lemma 5 one obtains the strong convergence of  $U_p^{\varepsilon}$  to  $U_p$  in  $L^2(0,T; L^2(\Omega_p))$ , providing the strong convergence for the traces on  $\Gamma$ .

**Lemma 11** Along a sequence  $\varepsilon \searrow 0$ , one has

$$U_p^{\varepsilon} \to U_p$$
 strongly in  $L^2(0,T;L^2(\Omega_p))$ .

Finally, combining Lemma 9 and Proposition 4.3 in [11] gives

**Lemma 12** There exists a C > 0 not depending on  $\varepsilon$  s.t. for any  $z_0 \in [0, 1]$  one has

$$\left|\left|\tilde{u}_{f}^{\varepsilon}(\cdot,\cdot,z_{0})-U_{f}^{\varepsilon}\right|\right|_{\Omega_{AV}^{T}}^{2}\leq\varepsilon C.$$

Using this, one can follow the steps in the proof of Lemma 4.5 in [11] to obtain

**Lemma 13** A constant C > 0 not depending on  $\varepsilon$  exists such that for all  $\phi_f \in L^2(0,T; W_0^{1,2}(\Omega_{AV}))$ 

$$\left| \Big( \int_0^1 (Q-q) dz \partial_x \tilde{u}_f^\varepsilon, \phi_f \Big)_{\Omega_{AV}^T} \right| \le Q \sqrt{\frac{53C}{40}} \, ||\partial_x \phi_f||_{L^2(\Omega_{AV}^T)}$$

We now have sufficient estimates to let  $\varepsilon \searrow 0$ , and show that the limiting pair  $(U_p, U_f)$  is a weak solution introduced in Definition 6. The result is contained in the following theorem.

**Theorem 7** The pair  $(U_p, U_f)$  is a weak solution introduced in Definition 6.

*Proof* For arbitrary test functions, we denote the terms of (67) by  $I_1, ..., I_{10}$ and analyze their limit as  $\varepsilon \searrow 0$ . The weak convergence in (70) gives

$$I_2 \to -(U_p, \partial_t \phi_p)_{\Omega_p^T},$$
  
$$I_3 \to (\nabla U_p, \nabla \phi_p)_{\Omega^T}.$$

Further, by Lemma 10 one has

$$|I_1| \le \varepsilon |(U_f^{\varepsilon}, \partial_t \phi_f)_{\Omega_{AV}^T}| \le \varepsilon \left| \left| U_f^{\varepsilon} \right| \right|_{\Omega_{AV}^T} ||\partial_t \phi_f||_{\Omega_{AV}^T} \le C\varepsilon \, ||\partial_t \phi_f||_{\Omega_{AV}^T} \to 0$$

The argument can be repeated for  $I_4$ ,  $I_5$ ,  $I_6$  and  $I_9$  to conclude that all have 0 limit as well. Moreover, by Lemma 13, the same holds for  $I_8$ .

The terms  $I_7$  and  $I_{10}$  do not change in the limit, since they do not depend on  $\varepsilon$ . Hence, for all  $\phi_p \in W^{1,2}(0,T;V_p)$  such that  $\phi_p(T) = 0$ ,  $U_p$  satisfies

$$-(U_p,\partial_t\phi_p)_{\Omega_p^T} + (\nabla U_p,\nabla\phi_p)_{\Omega_p^T} = (f_p,\phi_p)_{\Omega_p^T} + (u_{I,p},\phi_p(0))_{\Omega_p}.$$

It remains to show that  $r(U_p) = U_f$  on  $\Gamma$ . To this end, for arbitrary  $\phi \in L^2(0,T; L^2(\Gamma))$  we use the triangle inequality to estimate

$$|(r(U_p) - U_f, \phi)_{\Gamma^T}| \leq |(r(U_p^{\varepsilon}) - r(U_p), \phi)_{\Gamma^T}| + |(U_f^{\varepsilon} - U_f, \phi)_{\Gamma^T}| + |(r(u_p^{\varepsilon}) - \tilde{u}_f^{\varepsilon}, \phi)_{\Gamma^T}| + |(U_f^{\varepsilon} - \tilde{u}_f^{\varepsilon}, \phi)_{\Gamma^T}|.$$

We denote the terms on the right hand side by  $\tilde{I}_1, ..., \tilde{I}_4$  and let  $\varepsilon \searrow 0$ . For  $\tilde{I}_1$  we follow the steps concluding the proof of Theorem 5, namely use the trace inequality, (70) and the strong convergence in Lemma 11 to conclude that  $||r(U_p^{\varepsilon}) - r(U_p)||_{\Gamma^T}$  vanishes, implying that  $\tilde{I}_1 \to 0$ . Next,  $\tilde{I}_2$  goes to zero by the weak convergence (71).  $\tilde{I}_3 = 0$  trivially, by the nonlinear transmission condition, and  $\tilde{I}_4$  goes to zero by Lemma 12. Finally, since  $\phi \in L^2(0,T; L^2(\Gamma))$  was chosen arbitrarily, we conclude that  $U_f = r(U_p)$  on  $\Gamma$ , finishing the proof.

## 6 Discussion and outlook

We have derived upscaled equations for a time dependent reactive transport process in a fractured porous domain. The particularity lies in the nonlinear transmission conditions at the fracture interfaces. Both formal and rigorous upscaling procedures have been employed. The key role in this work is played by the existence results obtained in [14] for similar kind of elliptic problems. Related to this, efficient solution strategies for such kind of coupled problems need to be developed, including domain decomposition approaches as discussed in [5].

Acknowledgement ISP and KK are members of the International Research Training Group NUPUS funded by the German Research Foundation DFG (GRK 1398), the Netherlands Organization for Scientific Research NWO (DN 81-754) and by the Research Council of Norway (215627). They also acknowledge the Akademia grant supporting this work. Also, the work of ISP is supported by the Shell-NWO/FOM CSER programme (project 14CSER016). We thank Prof. M. van Sint Annaland (Eindhoven) for the discussions about the nonlinear transmission conditions in the context of reactive flows. Finally, we thank Prof. W. Jäger (Heidelberg) for stimulating discussions, guidance and continuous support.

#### References

- Allaire, G., Brizzi, R., Mikelić, A., Piatnitski, A., Two-scale expansion with drift approach to the Taylor dispersion for reactive transport through porous media, Chem. Eng. Sci. 65 (2010): 2292--2300.
- Allaire, G. and Hutridurga, H., Homogenization of reactive flows in porous media and competition between bulk and surface diffusion, IMA J. Appl. Math. 77 (2012): 788–815
- 3. Allaire, G., Mikelić, A. and Piatnitski, A., Homogenization approach to the dispersion theory for reactive transport through porous media, SIAM J. Math. Anal. 42 (2010): 125–144.
- Bogers, J.J.P., Transport in Fractured Media: Well-posedness, upscaling and numerical schemes for problems involving a nonlinear transmission condition, MSc thesis, Eindhoven University of Technology, 2011.
- Bogers, J.J.P., Kumar, K., Notten, P.H.L., Oudenhoven, J.F.M. and Pop, I.S., A multiscale domain decomposition approach for chemical vapor deposition. J. Comput. Appl. Math. 246 (2013), 65-73.
- Cances, C., Finite volume scheme for two-phase flows in heterogeneous porous media involving capillary pressure discontinuities. ESAIM: Math. Model. Num. 43 (2009): 973–1001.
- Cances, C. and Pierre, M., An existence result for multidimensional immiscible twophase flows with discontinuous capillary pressure field. SIAM J. Math. Anal. 44 (2012): 966–992.
- Devigne, V. M., Pop, I. S., van Duijn, C. J. and Clopeau, T., A numerical scheme for the pore-scale simulation of crystal dissolution and precipitation in porous media. SIAM J. Numer Anal 46 (2008): 895–919.
- van Duijn, C. J., Mikelić, A., Pop, I. S. and Rosier, C., Effective dispersion equations for reactive flows with dominant Peclét and Damköhler numbers. Advances in Chemical Engineering, 34 (2008): 1–45.

- van Duijn, C. J., Molenaar, J. and de Neef, M.J., The Effect of Capillary Forces on Immiscible Two-phase Flow in Heterogeneous Porous Media. Transport Porous Med, 21 (1995): 71–93.
- 11. van Duijn, C. J. and Pop, I. S., Crystal dissolution and precipitation in porous media: pore scale analysis. J. Reine. Angew. Math. 577 (2004): 171–211.
- Fatima, T., Ijioma, E.R., Ogawa, T. and Muntean, A., Homogenization and dimension reduction of filtration combustion in heterogeneous thin layers. Netw. Heterog. Media 9 (2014), 709-737.
- 13. Gilbarg, D. and Trudinger, N. S., Elliptic partial differential equations of second order. Vol. 224. Springer Science & Business Media, 2001.
- Jäger, W. and Kutev, N., Discontinuous solutions of the nonlinear transmission problem for quasilinear elliptic equations. In: Preprint IWR der Universität Heidelberg (1998): 1–37.
- Jäger, W. and Simon, L., On transmission problems for nonlinear parabolic differential equations. Ann. Univ. Sci. Budap. 45 (2002): 143–158.
- Kačur, J., Method of Rothe in evolution equations. Springer Berlin Heidelberg, 1986.
   Kellogg, R. B. and J. E. Osborn., A regularity result for the Stokes problem in a convex polygon. J. of Funct. Anal. 21 (1976): 397–431.
- Kumar, K., Neuss-Radu, M. and Pop, I. S., Homogenization of a pore scale model for precipitation and dissolution in porous media. arXiv preprint arXiv:1401.7025 (2014).
- Kumar, K., van Noorden, T. L. and Pop, I. S. Effective dispersion equations for reactive flows involving free boundaries at the microscale. Multiscale Model. Sim. 9 (2011): 29–58.
- Kumar, K., van Noorden, T. and Pop, I. S. Upscaling of reactive flows in domains with moving oscillating boundaries. Discrete Contin. Dyn. Sys. Ser. S 7 (2014): 95–111.
- Kweon, J. R. and Kellogg, R. B., Compressible Stokes problem on nonconvex polygonal domains. J. Differ. Equations 176 (2001): 290–314.
- 22. Evans, L. C., Partial Differential Equations. American Mathematics Society, 2010.
- Lenzinger, M. and Schweizer, B., Two-phase flow equations with outflow boundary conditions in the hydrophobic-hydrophilic case. Nonlinear Anal.-Theor. 73 (2010): 840–853.
- Ma, T. F. and Munoz Rivera, J. E., Positive solutions for a nonlinear nonlocal elliptic transmission problem. Appl. Math. Lett. 16 (2003): 243–248.
- 25. Mikelić, A., Devigne, V. and van Duijn C. J., Rigorous upscaling of the reactive flow through a pore, under dominant Peclet and Damkohler numbers. SIAM journal on mathematical analysis 38 (2006): 1262–1287.
- Muntean, A., Neuss-Radu, M., A multiscale Galerkin approach for a class of nonlinear coupled reaction diffusion systems in complex media. J. Math. anal. Appl. 371 (2010): 705–718.
- Musuuza, J.L., Radu, F.A and Attinger, S., The effect of dispersion on the stability of density-driven flows in saturated homogeneous porous media. Adv. Water Resour. 34 (2011): 417–432.
- Neuss-Radu, M., Ludwig, S. and Jäger, W., Multiscale analysis and simulation of a reaction-diffusion problem with transmission conditions. Nonlinear Anal. Real World Appl. 11 (2010): 4572–4585.
- Neuss-Radu, M. and Jäger, W., Effective transmission conditions for reactiondiffusion processes in domains separated by an interface. SIAM J. Math. Anal. 39 (2007): 687–720.
- van Noorden, T. L., Crystal precipitation and dissolution in a thin strip. Eur. J. Appl. Math. 20 (2009): 69–91.
- van Noorden, T. L., Pop, I. S., Ebigbo, A. and Helmig, R., An upscaled model for biofilm growth in a thin strip. Water Resour. Res. 46 (2010): W06505.
- 32. Simon, J., Compact sets in the space  $L^p(0,T;B)$ . Ann. Mat. Pura Appl. 146 (1986): 65–96.
- Lyu, Z., Zhou, T., Chen, L., Ye, Y., Sundmacher, K., Qi, Z., Simulation based ionic liquid screening for benzenecyclohexane extractive separation. Chem. Eng. Sci. 113 (2014): 45-53.

 Taylor, G., Dispersion of soluble matter in solvent flowing slowly through a tube. Proc. Math. Phys. Eng. Sci. 219. 1137. The Royal Society, 1953.

Zeidler, E., Nonlinear functional analysis and its applications. Vol. 2. Springer Science & Business Media, 1989.

# **Published Preprints**

http://www.nupus.uni-stuttgart.de

2007/1	Cao, Y. / Eikemo, B. / Helmig, R.: Fractional flow formulation for two-phase flow in porous media
2008/1	Helmig, R. / Weiss, A. / Wohlmuth, B.: Variational inequalities for modeling flow in heterogeneous porous media with entry pressure
2008/2	Cao, Y. / Helmig, R. / Wohlmuth, B.: Convergence study and comparison of the multipoint flux approximation L-method
2008/3	van Duijn, C.J. / Pop, I.S. / Niessner, J. / Hassanizadeh, S.M.: Philip's redistribution problem revisited: the role of fluid-fluid interfacial areas
2008/4	Niessner, J. / Hassanizadeh, S.M.: Modeling kinetic interphase mass transfer for two-phase flow in porous media including fluid-fluid interfacial area
2008/5	Niessner, J. / Hassanizadeh, S.M.: A model for two-phase flow in porous media including fluid-fluid interfacial area
2008/6	Cao, Y. / Helmig, R. / Wohlmuth, B.: Geometrical interpretation of the multipoint flux approximation L-method
2008/7	Vervoort, R.W. / van der Zee, S.E.A.T.M.: Simulating the effect of capillary flux on the soil water balance in a stochastic ecohydrological framework
2008/8	Niessner, J. / Hassanizadeh, S.M.: Two-phase flow and transport in porous media including fluid-fluid interfacial area
2008/9	Haslauer, C.P. / Bárdossy, A, / Sudicky, E.A.: Geostatistical analysis of hydraulic conductivity fields using copulas
2008/10	Wolff, M.: Comparison of mathematical and numerical models for twophase flow in porous media
2008/11	Darcis, M.: Implementation of a numerical model for the convection-enhanced delivery of therapeutic agents into brain tumors
2008/12	Cao, Y. / Helmig, R. / Wohlmuth, B.: Convergence of the multipoint flux approximation L-method for homogeneous media on uniform grids
2008/13	Ochs, S.O.: Development of a multiphase multicomponent model for PEMFC
2008/14	Walter, L.: Towards a model concept for coupling porous gas diffusion layer and gas distributor in PEM fuel cells
2009/1	Hægland, H. / Assteerawatt, A. / Helmig, R. / Dahle, H.K.: Streamline approach for a discrete fracture-matrix system
2009/2	Assteerawatt, A. / Hægland, H. / Helmig, R. / Bárdossy, A. / Dahle, H.K.: Simulation of flow and transport in a geostatistical fracture-matrix system
2009/3	Heimann, F.: An unfitted discontinuous Galerkin method for two-phase flow
2009/4	Hilfer, R. / Doster, F.: Percolation as a basic concept for macroscopic capillarity

2009/5	van Noorden, T.L. / Pop, I.S. / Ebigbo, A. / Helmig, R.: An effective model for biofilm growth in a thin strip
2009/6	Baber, K.: Modeling the transfer of therapeutic agents from the vascular space to the tissue compartment (a continuum approach)
2009/7	Faigle, B.: Two-phase flow modeling in porous media with kinetic interphase mass transfer processes in fractures
2009/8	Fritz, J. / Flemisch, B. / Helmig, R.: Multiphysics modeling of advection-dominated two-phase compositional flow in porous media
2009/9	Støverud, K.: Modeling convection-enhanced delivery into brain tissue using information from magnetic resonance imaging
2009/10	Doster, F. / Zegeling, P. / Hilfer, R.: Numerical solutions of a generalized theory for macroscopic capillarity
2009/11	Kissling, F. / Rohde, C.: The computation of non-classical shock waves with a heterogeneous multiscale method
2010/1	Rosenbrand, E.: Modelling biofilm distribution and its effect on two-phase flow in porous media
2010/2	Schöniger, A.: Parameter estimation by ensemble Kalman filters with transformed data
2010/3	Ebigbo, A. / Helmig, R. / Cunningham, A.B. / Class, H. / Gerlach, R.: Modelling biofilm growth in the presence of carbon dioxide and water flow in the subsurface
2010/4	Lauser, A. / Hager, C. / Helmig, R. / Wohlmuth, B.: A new approach for phase transitions in miscible multi-phase flow in porous media
2011/1	Linders, B.: Experimental investigations on horizontal redistribution
2011/2	Rau, M.T.: Geostatistical analysis of three-dimensional hydraulic conductivity fields by means of maximum Gauss copula
2011/3	Kraus, D.: Two phase flow in homogeneous porous media - The role of dynamic capillary pressure in modeling gravity driven fingering
2011/4	Brugman, R.: Dimensionless analysis of convection enhanced drug delivery to brain tissues
2011/5	Sinsbeck, M.: Adaptive grid refinement for two-phase flow in porous media
2011/6	Kissling, F. / Helmig, R. / Rohde, C.: A multi-scale approach for the modelling of infiltration processes in the unsaturated zone
2012/1	Köppl, T. / Wohlmuth, B. / Helmig, R.: Reduced one-dimensional modelling and numerical simulation for mass transport in fluids
2012/2	Kumar, K. / Pop, I.S. / Radu, F.A.: Convergence analysis for a conformal discretization of a model for precipitation and dissolution in porous media
2012/3	Hommel, J.: Modelling biofilm induced calcite precipitation and its effect on two phase flow in porous media

2012/4	Estrella, D.: Experimental and numerical approximation methods for zero-valent iron transport around injection wells
2012/5	Heimhuber, R.: Efficient history matching for reduced reservoir models with PCE-based bootstrap filters
2012/6	Kissling, F. / Karlsen, K.H.: On the singular limit of a two-phase flow equation with heterogeneities and dynamic capillary pressure
2012/7	Fritz, S.: Experimental investigations of water infiltration into unsaturated soil - Analysis of dynamic capillarity effects
2012/8	Strohmer, V.: Numerische Analysis von nahezu parallelen Strömungen in porösen Medien
2012/9	Kissling, F. / Rohde, C.: The computation of nonclassical shock waves in porous media with a heterogeneous multiscale method: The multidimensional case
2012/10	Fetzer, T.: Numerical analysis of the influence of turbulence on exchange processes between porous-medium and free flow
2012/11	Schröder, P.: A response surface bootstrap filter to calibrate CO2 injection models
2013/1	Brunner, F. / Radu, F.A. / Knabner, P.: Analysis of an upwind-mixed hybrid finite element method for transport problems
2013/2	Köppel, M.: Flow modelling of coupled fracture-matrix porous media systems with a two mesh concept
2013/3	van Helvoort, M.: Upscaling of processes involving rough boundaries
2013/4	Redeker, M. / Haasdonk, B.: A POD-EIM reduced two-scale model for crystal growth
2013/5	Vogler, D.: A comparison of different model reduction techniques for model calibration and risk assessment
2014/1	Song, N.: Investigation of a decoupling scheme for the modeling of reactive transport
2014/2	Aydogdu, A.B.: Phase field modelling of critical shear band evolution in granular media on the basis of a micropolar porous medium theory
2014/3	Becker, B.: Investigation of error estimates for cell centered finite volume schemes: Analysis and improvement of grid adaptation strategies in DuMux
2014/4	Moghaddam, N.D.: Sorption of methane and ethane on Belgian black shale using a manometric setup
2014/5	Schwenck, N. / Flemisch, B. / Helmig, R. / Wohlmuth, B.: Dimensionally reduced flow models in fractured porous media: crossings and boundaries
2014/6	Redeker, M. / Pop, S. / Rohde, C.: Upscaling of a tri-phase phase-field model for precipitation in porous media
2014/7	Faigle, B. / Ahmed Elfeel, M. / Helmig, R. / Becker, B. / Flemisch, B. / Geiger, S.: Multi-physics modeling of non-isothermal compositional flow on adaptive grids

2014/8	Radu, F.A. / Nordbotten, J.M. / Pop, I.S. / Kumar, K.: A robust linearization scheme for finite volume based discretizations for simulation of two-phase flow in porous media
2015/1	Jambhekar, V.A. / Helmig, R. / Schröder, N. / Shokri, N.: Free-flow-porous-media coupling for evaporation-driven transport and precipitation of salt
2015/2	Bause, M. / Radu, F. A./ Köcher, U.: Error analysis for discretizations of parabolic problems using continuous finite elements in time and mixed finite elements in space
2015/3	van Duijn, C. J. / Cao, X. / Pop, I. S.: Two-phase flow in porous media: dynamic capillarity and heterogeneous media
2015/4	Pop, I. U. / Bogers, J. / Kumar, K.: Analysis and upscaling of a reactive transport model in fractured porous media involving a nonlinear transmission condition