Made available by Hasselt University Library in https://documentserver.uhasselt.be

The Assessment of Motor Fatigability in Persons With Multiple Sclerosis Peer-reviewed author version

SEVERIJNS, Deborah; Zijdewind, Inge; Dalgas, Ulrik; LAMERS, Ilse; Lismont, Caroline & FEYS, Peter (2017) The Assessment of Motor Fatigability in Persons With Multiple Sclerosis. In: Neurorehabilitation and neural repair,.

DOI: 10.1177/1545968317690831 Handle: http://hdl.handle.net/1942/23380

The assessment of motor fatigability in persons with multiple sclerosis: a systematic review

Deborah Severijns, MSc^{1*}; Inge Zijdewind, PhD,²; Ulrik Dalgas, PhD³; Ilse Lamers, PhD^{1;} Caroline Lismont, BSc^{1;} Peter Feys, PhD¹ ¹ REVAL Rehabilitation research center - BIOMED, Faculty of Medicine and Life Sciences, Hasselt University, Hasselt, Belgium ² Department of Neuroscience, University Medical Center Groningen, University of Groningen, Groningen, Netherlands ³ Section of Sport Science, Department of Public Health, Aarhus, Aarhus University, Denmark.

*Corresponding author: Deborah Severijns Agoralaan, Building A, B-3590 Diepenbeek Belgium <u>deborah.severijns@uhasselt.be</u> 0032 11 29 21 29

Wordcount abstract: 246 Wordcount main text: 5337 Number of figures: 2 Number of tables: 3

Abstract

Background

Persons with Multiple Sclerosis (PwMS) are often characterized by increased motor fatigability, which is a performance change on an objectively measured criterion after any type of voluntary muscle contractions. This review summarizes the existing literature, to determine which protocols and outcome measures are best to detect or study motor fatigability and the underlying mechanisms in MS.

Methods

Two electronic databases, Pubmed and Web of Science, were searched for relevant articles published until august 2016 with a combination of 'Multiple Sclerosis', 'fatigability', 'muscle fatigue' and 'motor fatigue'.

Results

A total of 48 papers were retained for data-extraction. A variety of fatigability protocols were reported, protocols showed differences in type (isometric vs. concentric) duration (15 to 180 seconds) and number of contractions (fixed or until exhaustion). Twelve papers reported motor fatigability during functional movements, predominantly assessed by changes in walking speed. Eleven studies evaluated the mechanisms underlying motor fatigability, using additional electrical nerve or transcranial magnetic stimulation. Three papers reported psychometrics of the outcomes.

Conclusions

The disparity of protocols and outcome measures to study different aspects of motor fatigability in PwMS impedes direct comparison between data. Most protocols use maximal single joint isometric contractions, with the advantage of high standardization. As there is no head to head comparison of the different protocols,

and only limited information on psychometric properties of outcomes, there is currently no gold standard to assess motor fatigability. The disability level, disease phenotype and studied limb may influence the assessment of motor fatigability in PwMS.

Key words: muscle fatigue, fatigability, motor fatigue, multiple sclerosis

Introduction

Multiple Sclerosis (MS) is a progressive, chronic disease of the central nervous system, with symptoms such as weakness and fatigue.¹ Fatigue is often defined as a "subjective lack of physical and/or mental energy that is perceived by the individual or caregiver to interfere with usual and desired activities".² Fatigue is thus a subjective symptom that interferes with task performance in daily life. Kluger et al. proposed a unified taxonomy, where the symptom of fatigue is affected by perceptions of fatigue and performance fatigability.³ Performance fatigability can be studied within different domains. When studying the motor domain, the following definition of motor fatigability (adapted from Kluger et al.³) can be used: "Motor fatigability is the magnitude or rate of change of motor performance on an objectively measured reference criterion after any type of voluntary activity or exercise". Several reviews on the assessment and treatment of fatigue in persons with MS (PwMS) or other neurological diseases exists.⁴⁻⁶ In contrast, no overview has been reported of protocols and outcomes that can identify pathological motor fatigability in PwMS. Motor fatigability is, however, important to consider in PwMS, as it affects the ability to perform sustained activities of daily living.⁷ Further, it will have an impact on the exercise capacity of PwMS, limiting the possibility to exercise regularly. The assessment of muscle weakness is routinely performed, but muscle weakness and motor fatigability are different concepts.⁸ PwMS might not show muscle weakness on a single assessment of muscle strength, but they might show increased motor fatigability with a fast decline in muscle strength during longer or repeated test protocols.

Importantly, self-reported measures for fatigue do not provide a valid indicator of motor fatigability after physical activities or vice versa,⁹ since the two concepts are

most often unrelated.^{10,11} Consequently, it is important to quantify (pathological) motor fatigability applying objective, valid, reliable and responsive outcomes.¹⁰ To unravel the relation between motor fatigability and activities of daily living, it is furthermore important to include outcome measures on both the body function level and the activity level of the International Classification of Functioning and Health (ICF), when assessing fatigability.¹² It is not known if PwMS show motor fatigability after repeated and sustained muscle contractions of isolated joints, reflecting the "ICF body function level" as well as after functional activities such as walking reflecting the "ICF activity level". From studies in healthy persons, it is known that motor fatigability is task specific and that protocol specifications affect the findings and the underlying mechanisms of motor fatigability.¹³ In healthy subjects, different protocols are applied to assess different aspects of motor fatigability, where submaximal protocols most likely challenge the central nervous system, and high intensity exercises the peripheral neuromuscular system.^{14,15} It is, however, not known if PwMS express pathological motor fatigability during all tasks or whether the results of different protocols can be compared.

Therefore, the aim of this systematic review was 1) to provide an extensive overview of the applied motor fatigability assessment protocols and outcome measures within the ICF framework to detect the most optimal assessment to quantify motor fatigability in PwMS, 2) to summarize the available information on psychometric properties for the located protocols, including the discriminant ability and the effect of exercise interventions.

<u>Methods</u>

Data sources and study selection

To review the literature, the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement were adopted.¹⁶ The current review was registered on PROSPERO (CRD42016032320). Two databases (PubMed and Web of Science (WOS)), were searched, using a combination of Mesh terms/keywords: Multiple sclerosis AND (muscle fatigue OR fatigability OR motor fatigue). Literature searching started in September 2015, ended in November 2015 and was updated on August 23rd, 2016.

Studies were selected when written in English, published within the last 20 years, having undergone peer review and enrolled ≥10 PwMS. Furthermore, studies had to generate/assess an objective parameter quantifying motor fatigability based on voluntary contractions. Interventional studies were only included when the intervention consisted of physical exercises. Descriptive, exploratory or experimental full length studies were included; conference papers, meeting reports, letters and reviews were excluded.

All results were screened based on title and abstract by 2 independent reviewers (DS and CL). The entire full text was read when the abstract did not provide sufficient information. In cases of disagreement between the two reviewers or if it was unclear whether a study should be included, a third reviewer (IL) was consulted. The reference lists of included articles were checked for further relevant articles.

Quality assessment

The methodological quality was determined by the Downs and Black checklist.¹⁷ Questions 6, 8, 14, 15, 19, 23, 24, and 26 were removed from the original scale when assessing observational studies as these questions were not applicable.

A maximal score of 18 could therefore be assigned to observational studies as compared to 26 for randomized controlled trials. The score was converted to a percentage score for each study, with a score above 65% indicating sufficient methodological quality.¹⁸ Since the main goal of the review was to evaluate details of fatigability protocols, no studies were excluded, based on the quality assessment.

Data extraction

The following data were extracted from the selected articles: 1) study aim, ICF assessment level, 2) sample characteristics 3) fatigability protocol specifications and outcome parameters, 4) information on psychometrics of outcomes and 5) the main findings on motor fatigability in studies comparing healthy controls (HC) to PwMS. The papers reporting protocols of a single muscle group or a single limb, during non-functional activities, were classified at the ICF body function level (category b 730-muscle power functions and b740-muscle endurance functions). Papers examining motor fatigability during functional activities, were classified at the ICF activity level (d450-469-walking and moving). Details of the intervention and the effects on motor fatigability were documented from interventional studies.

<u>Results</u>

As presented in detail in Figure 1, 48 papers were included from the initial 494 hits (353 from WOS and 141 from Pubmed). The data for quality assessment of the included studies is shown in supplementary Table 1 and 2. The results of the quality assessment for observational studies, show that the scores range from 35%-82% and for the interventional studies between 41%-74%. The main problems of the observational studies were the lack of reporting: the difference between asked and

recruited participants (Q 11 and 12), the recruitment period (Q22), possible confounders (Q25) and the power of the study (Q27). For the interventional studies all studies scored 0 on the representation of the study population (Q12), blinding of subjects and assessors (Q14-15) and concealment of allocation (Q24). Further, also the reporting was limited on: the confounders (Q25), losses to follow up (Q26), and the power of the study (Q27).

insert Figure 1

Study objectives and patient characteristics

Study objectives were heterogeneous (supplementary Table 3). In 23 papers the objective was to assess motor fatigability in PwMS, while 11 papers aimed to detect underlying mechanisms. Eight papers aimed to study the correlation between self-reported fatigue and motor fatigability, while seven papers reported the influence of motor fatigability on other outcomes such as dual tasking, balance, motor cortex excitability and brain activation patterns. Five papers reported the effect of an intervention on motor fatigability. Three papers reported on the reliability of the applied outcomes for fatigability.

Most papers (n=36) investigated the ICF body function level. Two papers reported both at ICF body function level and ICF activity level. Of the protocols at body function level, 20 examined the upper, 16 the lower limbs and two papers both upper and lower limbs. Twelve papers reported motor fatigability protocols at the ICF activity level. One of these combined outcomes at body function level with an exercise on activity level.

Table 1 summarizes sample characteristics. The sample size ranged from 10 to 208. Thirteen papers included all types of MS, eleven papers only included relapsing remitting MS, and three papers only secondary progressive MS. Three

papers by the same author, reported on PwMS with clinically isolated syndrome.¹⁹⁻²¹ The average Expanded Disability Status Scale (EDSS) ranged from 1.3 to 6. The studies that reported on self-reported subjective fatigue enrolled both fatigued and non-fatigued PwMS.

insert Table 1

Fatigability protocols

A schematic overview of type of protocols, and frequency of application in clinical research in MS, is provided in Figure 2. Table 2 documents the protocols specifications.

insert Figure 2

Protocols assessing motor fatigability at the ICF body function level

In 30/38 protocols, isometric protocols were used (ICF category b7300-7301, 7400-7401). Ten studies examined the lower limb, 18 the upper limb and two studies assessed both the upper and lower limbs. Most studies applied maximal voluntary contractions (MVCs) to assess motor fatigability. One study reported both maximal and submaximal protocols.²² Generally, the duration of an MVC was predetermined (15s²³, 30s^{8,11,19-21,24-26}, 45s^{27,28}, 60s²⁹, 120s³⁰⁻³⁶, 180s³⁷⁻³⁹). In one study the endpoint was not clearly defined ("until strength declined").⁴⁰ For protocols applying intermittent MVCs, the number of contractions ranged from 8 repetitions of 4 s²³, to 11 or 15 repetitive contractions without a specified duration^{24,40}, the number of contractions that can be performed in 30 seconds⁸, 18 repetitions of 5 seconds⁴¹, up to 2 minutes at 2 Hz (240 repetitions)⁴².

Six papers reported submaximal isometric tasks of 10 to 50% of the MVC. One study evaluated a 60s sustained contraction at 25, 50, 75 and 100%MVC,²⁹ while one study applied 50%MVC until failure.⁴³ The three other protocols used repetitive contractions until inability to produce target force (e.g. 30s vs. 30s rest, or blocks of 44s with 5s rest)^{35,44,45} or with a specified duration (6 minutes of exercises).²²

Eight papers reported non-isometric protocols (ICF categories b7401 or 7402). Two studies asked as many contractions as possible in a certain time frame.^{46,47} Three other studies used an isokinetic protocol in the lower limbs, with 20 or 30 repetitions at 180°/s^{48,49} or 50 repetitions at 60°/s.⁵⁰ One study reported squeezing a rubber ball maximally for 10 times.⁵¹ Motor fatigability during multi-joint movements of the lower limb was assessed with leg presses until exhaustion.^{52,53}

Ten studies applied additional electrical stimulation of the nerve innervating the target muscle to determine underlying mechanisms.^{23,27-33,36,42} Two studies interspersed the voluntary exercises with transcranial magnetic stimulation (TMS) to gain further insight into corticospinal excitability,^{42,46} six studies used TMS before and after a fatigability protocol.^{28,37,39,43-45}

Protocols assessing ICF activity level

In ten studies, walking for a fixed time (6 or 12 min)⁵⁴⁻⁵⁷, a fixed distance (500m)^{8,25,58} or until voluntary exhaustion,⁵⁹⁻⁶¹ at either comfortable or fastest walking speed was applied. Three walking studies used a treadmill⁵⁹⁻⁶¹, seven studies tested free overground walking.^{8,25,54-58} One study examined motor fatigability during 5 minutes wheelchair driving at a speed of 1m/sec in a manual wheelchair,⁶² while another used a functional steering task with a joystick driven wheelchair, mimicking daily life activities.⁶³

Outcome measures: the quantification of motor fatigability and underlying mechanisms

Table 2 summarizes the identified outcomes to document motor fatigability. Overall the outcomes could be categorized as 1) strength-based outcomes, 2) indirect outcomes and 3) neurophysiological outcomes.

Insert Table 2

Strength-based outcomes

The majority of the isometric protocols (29/48) used strength-based outcomes, to assess motor fatigability. Despite the different names, (e.g. "percent decrement of force", "static fatigue index", "rate of decline in strength") most outcomes directly measure the strength decline. Indices of motor fatigability are mostly based on the ratio between the initial and the final strength during sustained (23/48) or repetitive contractions (8/48). For the sustained contractions, an index was also based on the area under the curve or the slope of the strength decline.^{8,11,19-21,24-26} Some papers only report a statistical comparison of the maximal strength, assessed before and after a specific task, rather than an index.^{22,29,47,64} This approach is also used in one study after walking.⁵⁴ During isokinetic protocols, the ratio between the work done during the first contractions vs. the last contractions,^{48,49} or the slope of torque decline⁵⁰ was used.

Indirect outcomes

Four papers reported indirect outcomes based on endurance time. For example, three studies reported the time until MVC declined to 50% of the initial MVC.³⁷⁻³⁹ Five studies evaluated the "number of repetitions": e.g. the number of contractions until the subject was unable to maintain 50%MVC,⁴⁵ the number of repetitions performed in a fixed amount of time,⁴⁷ or until inability to maintain a target force ^{35,52,53} were reported. The performance such as the change in tapping speed⁴⁶

and the trajectory of movements⁴⁷ during single joint movements was described as an indirect measure of motor fatigability.

After walking, indices were based on walking velocity or distance, such as the deceleration index,⁵⁸ distance walked index⁵⁵ and ambulatory fatigue index⁸. Four studies calculated an index based on gait kinematics.^{57,59-61} The performance outcomes after manual wheelchair driving were velocity and propulsion frequency.⁶² Electromyography (EMG) was used in nine studies.^{22,31,32,35,36,44,47,63,64} Four studies reported the median or mean frequency of the EMG signal,^{22,47,63,64} while the remaining 6 studies reported a measure of the EMG amplitude (e.g. root mean square).

Neurophysiological outcomes

Besides the assessment of motor fatigability as such, neurophysiological outcomes were reported in 17 studies to explore underlying mechanisms of motor fatigability (Table 2). The twitch interpolation technique⁶⁵ was frequently applied to determine the changes in central and the peripheral activation of the muscles under investigation^{23,27-33,36,42,43} and used as an indication of a loss of central drive (central fatigue) or peripheral muscle fatigue.⁶⁶ .

Psychometric properties

The only information on psychometrics in PwMS that was detected with the current literature search was related to reliability of the outcome measures. No information was detected on the validity and responsiveness of motor fatigability measures. The discriminant ability, however, might partially represent construct validity.

Reliability

Three studies investigated the reliability of motor fatigability outcomes. Schwid et al.⁸ and Surakka et al.²⁶ reported on the reliability of static fatigue indices, based

on a 30s sustained isometric contraction. The static fatigue index with the best reliability in PwMS (intraclass correlation coefficients (ICC) range 0.71-0.96) that best discriminated between HC and PwMS, was calculated by dividing the total area under the strength curve by the hypothetical area under the curve, when no strength decline would occur (i.e. the multiplication of maximal strength times contraction duration from 5 sec after the start of the contraction). Other fatigue indices showed lower ICCs.⁸ A fatigue index based on a ratio between the initial strength and the final strength had ICCs of 0.46-0.77. A fatigue index, based on the area under the first 5 sec vs; the last 5 sec. had ICC's of 0.5-0.73. A fatigue index, based on the area under the curve of the entire contraction period, showed ICC's of 0.64-0.93. A slightly modified static fatigue index also showed good reliability (ICC of 0.68-0.86).²⁶ Lambert et al.⁴⁹ reported poor to moderate reliability of an index, based on the work performed during the first 15 vs. the last 15 contractions of an isokinetic protocol consisting of 30 maximal contractions at 180°/s (ICC's from 0.36 to 0.51). In accordance, poor reliability of a dynamic fatigue index was reported, with ICC's from 0.20 to 0.44.8 One report showed poor reliability of an ambulatory fatigue index, determined during a 500 meter walk (ICC: 0.21-0.36).8

Discriminant ability between (subgroups of) PwMS and healthy controls

Table 2 shows that 34/48 papers enrolled PwMS and HC. Not all studies compared groups statistically. Twenty-four papers reported a statistically significant difference between HC and PwMS, and 17 of these used isometric (repetitive or sustained) MVCs. Of these 17, strength based outcomes showed a group difference for motor fatigability in the upper limb^{22,24,28,36,40,42,51}, the lower limb,^{19,20,29} and in both the upper and lower limb.^{8,11} The number of times that an outcome showed a difference between PwMS and HC, is shown in Table 3. Three protocols with

sustained MVCs investigated the difference between subgroups of PwMS and found that motor fatigability was related to disability level (EDSS score)²⁴, the presence of pyramidal signs¹¹ and the involvement of more than one functional system.²⁰ Wolkorte et al. further found that secondary progressive PwMS show more motor fatigability, compared to relapsing remitting MS.³⁶ Seven papers reported nonsignificantly different strength based outcomes. Six of these studied the upper limb (hand grip or intrinsic hand muscles), with a sustained contraction of 2 minutes^{31,32} or 3 minutes ^{37,38}, or with intermittent isometric hand grip contractions^{24,41}. One paper studied knee extension with a sustained contraction of 2 minutes.³⁰ Four out of six studies with non-significant results included PwMS with RRMS or mild MS (average EDSS<3.5). Five of the latter studies, using a sustained MVC, did, however, report significantly different neurophysiological outcomes. Intermittent concentric maximal contractions elicited a greater decline in strength in the upper limb compared to healthy controls.⁵¹ One study, using maximal concentric knee contractions, showed only a difference in total work, not for the work index.⁴⁹ One study reported different motor fatigability between mild and moderate MS, assessed with the decline in peak torque after 50 isokinetic knee extensions.⁵⁰

For the submaximal protocols contractions, only one strength outcome (force variability) differentiated HC from PwMS for the upper limb,³⁵ while no difference in decline in strength, the number of contractions or exercise duration was observed. One submaximal foot tapping protocol did detect a difference in strength decline between PwMS and HC.⁴⁶ Significant differences were reported for motor fatigability during walking^{8,54,56,61} and after 5 minutes of wheelchair driving.⁶² Three walking studies showed that PwMS with higher EDSS levels experienced more motor fatigability.^{55,56,58}

The effect of exercise interventions on motor fatigability

Although there is no study investigating responsiveness of outcomes for motor fatigability, a limited number of exercise intervention studies revealed that motor fatigability can be improved. Five studies evaluated the effects of different exercise interventions on motor fatigability (Supplementary Table 4). Studies aimed to improve motor fatigability of the lower limb.^{25,48,52,53,64} Interventions included progressive resistance training,^{52,53} inpatient rehabilitation followed by home exercises,²⁵ electrical stimulation⁶⁴ and whole body vibration.⁴⁸ Three intervention studies showed improvements of motor fatigability, based on the number of repetitions,⁵² the knee flexion fatigue index,²⁵ and the quadriceps peak torque after a 30s sustained contraction.⁶⁴

insert Table 3

Discussion

The main findings of this systematic review are that 1) most assessments of motor fatigability at the ICF body function level are based on isometric maximal contractions; 2) the assessments of motor fatigability at the ICF activity level are predominantly based on changes in performance during a functional activity; 3) the majority of the studies at both the body function level and activity level were able to discriminate between PwMS and HC; and 4) information on the psychometrics of the applied fatigability protocols is limited.

Selecting the right protocol and outcomes when assessing motor fatigability

The included study protocols differed substantially on protocol components such as the type, intensity and duration of the fatiguing task, type of movement, and the assessed limb or body part. As motor fatigability is task specific,⁶⁷ these components will influence the assessment of motor fatigability.

Body function level

Most protocols assessed isometric MVCs, which is probably explained by the physiological definition of fatigability that most of the studies adopted.^{14,68} Furthermore, isometric contractions have the advantage that they can be highly standardized. In addition, subjects quickly fatigue when they perform MVCs, making strength deficits easier to detect.⁶⁹ The duration of most protocols is generally fixed, which is preferable, as protocols performed until volitional exhaustion show poor reliability.⁷⁰ The maximal contraction duration was 3 minutes,³⁷⁻³⁹ but most applied 30 seconds. Most studies assessed motor fatigability with strength-based measures. These are easy to interpret, but results of different contraction times are difficult to compare, as other mechanisms are likely responsible for the quick decline in strength within 30 seconds compared to over a longer period.^{34,71} Nonetheless, the reliability of fatigue indices does not seem to depend on protocol length in HC.⁷² Numerous calculation methods exist for fatigability indices. Using a ratio of the initial vs. the final force level might cause problems as the force variability increases during the final stage of a sustained contraction.⁷³ This problem may, however, be overcome by looking at the total area under the curve after sustained contractions.²⁶ Studies evaluating the underlying mechanisms of motor fatigability, require standardized (high intensity) isometric contractions, often supplemented by TMS ^{39,42-46} or peripheral nerve stimulation^{23,27,28,30-33,36,42} before and after the fatiguing task. An indepth discussion on the underlying mechanisms of motor fatigability, is provided in a recently published review.⁷⁴ Motor fatigability is most often assessed during sustained contractions, however, intermittent contractions seem more relevant for daily life. Although isokinetic protocols are sparsely used,⁴⁸⁻⁵⁰ these might be interesting, since they are even more energetically demanding.¹⁴ One should take

into account that also here, results on motor fatigability of sustained contractions cannot be interchanged with intermittent contractions, as the physiological reaction in the muscle tissue is different,¹⁴ and possibly also the brain activation patterns.⁷⁵ Some results might indicate that intermittent contractions are less suitable to determine pathological motor fatigability, as these do not differentiate PwMS from healthy controls.²⁴

The majority of the protocols used maximal muscle contractions. Only six studies applied a submaximal isometric protocol.^{22,29,35,43-45} One could argue that intermittent submaximal contractions more closely simulate "real-world" situations.^{76,77} Submaximal protocols might be difficult to standardize, as these might be influenced more by problems with selective motor control requiring more concentration. To evaluate changes in muscle output due to submaximal exercises, a submaximal fatiguing exercise interrupted with brief maximal contractions might be applied ⁷⁸ as has been done in a few studies.^{35,46,47} Further, PwMS with muscle weakness may already perform near their maximal strength during ADL, thus when asking a submaximal percentage of their strength, this might be underestimating the problem of fatigability in daily life. A further challenge to this approach relates to the higher absolute strength usually exerted by HC. This causes a reduced blood perfusion in HC and thus increases the peripheral fatigue, although the proportional strength is identical, which contributes to sex-related differences in fatigability.⁷⁹ Indirect outcomes such as performance measures⁴⁷ have the advantage that they are directly related to the task. However, the interpretation is more difficult, as they cannot be attributed to one muscle group.

Most protocols on body function level assess the upper limbs, which contrasts the fact that, clinically, motor fatigability is mostly reported in the lower limbs, e.g.. during walking.

Clinically, it could be hypothesized that fatigability in the lower limbs is influencing activities of daily living more profoundly, and thus is reported more frequently, as fatigued legs require rest by sitting down or lying. This fatigue effect might only be reported in the upper limbs after very specific tasks, such as sewing, which require continuous activation. One could further argue that investigating themotor fatigability in the upper limbs is less relevant, as the upper limbs are normally less impaired by the disease process.⁸ The reason that most of the research is ,however, on the upper limbs, is probably caused by the fact that many studies included neurophysiological investigations to investigate the mechanisms of fatigability. These investigations are technically more straightforward to perform accurately in the upper limbs than in the lower limbs as relevant muscles can more easily be accessed and stimulated. Further, when investigating the lower limbs, a more profound effect of deconditioning (besides the disease effect of MS) can be expected. Until now, only one study compared motor fatigability in the upper limb and the lower limb,⁸ stating that these are related in PwMS. This information is contrasted by a study in healthy controls, stating that there is no relation between hand grip fatigability and fatigability of the knee extensors.⁸⁰ In shorth, the generalizability of motor fatigability across upper and lower limb muscles is unknown More can probably be learned in future studies comparing upper and lower limb fatigability. Activity level

Fatigability determined on the body function level might be more precise and has several advantages, such as the ability to determine underlying mechanisms

within the same tasks (strength decline combined with neurophysiological measurements). However, the investigation at the ICF activity level, might be more relevant for daily life as these activities are most often closer linked to quality of life. Most studies attempting to evaluate motor fatigability at ICF activity level have applied walking paradigms.^{8,25,56,58-61} These protocols seem relevant for clinical practice, but the interpretation is complicated, as outcomes based on physical performance during functional activities, such as the deceleration in walking speed, give no information on underlying causes. Observing changes in gait dynamics seems an attractive approach and has been applied during walking on a treadmill at maximal speed.^{60,61} However, it is not straightforward to interpret changes in kinematics as an increase in motor fatigability (decreased neural drive) versus an increased manifestation of underlying impairments of spasticity or ataxia. To overcome this, some studies use EMG parameters as an additional indirect measure of motor fatigability. During sustained submaximal static contractions notable changes occur as the signal amplitude (RMS) increases, and shifts in the EMG spectrum towards lower frequencies take place. Both parameters are used as markers of muscle fatigue. However, EMG variables could not discriminate HC from PwMS in several studies.^{31,47} Further, it has been argued that amplitude characteristics are not reliable^{14,81,82} and that the use of EMG outcomes during dynamic contractions is often difficult to interpret. Mcloughlin et al.⁵⁴ tried to integrate the ICF body function level and the ICF activity level, by assessing strength and balance before and after a 6 minutes walking paradigm and showed that PwMS showed greater loss of strength and balance after 6 minutes walking than HC. Is motor fatigability increased in multiple sclerosis?

Overall, literature shows that motor fatigability is increased in PwMS. As discussed above, the ability to detect differences between HC and PwMS seems to be related to the type of tasks (sustained or intermittent), contraction intensity (maximal or submaximal) and the limbs examined (upper or lower limb). Further, it is likely that not all PwMS show abnormal motor fatigability, as reported earlier.^{24,55} Strength-based outcomes were not always sufficiently sensitive to detect differences in mildly impaired PwMS, despite different neurophysiological outcomes.³² Seven studies found no significant difference between HC and PwMS for strength-based parameters. Maximal protocols, which could not discriminate HC and PwMS, used a low number of repetitive hand grip contractions^{24,41} or a long isometric contraction for 2/3 minutes^{30,31}, where subjects are in the plateau phase of strength decline. Further explanations for the lack of discrimination might be the small samples, MS phenotype³⁶ or disability level of the participants.²⁴

Of note, none of the submaximal protocols, assessing the upper limb with strength outcomes, discriminated HC from PwMS, whereas one lower limb study did show a more profound force decline after a foot tapping exercise.⁴⁶ This may again suggest a different response between upper and lower limbs, caused by earlier and more substantial impairments in the lower limbs of PwMS.⁸

Limitations in the current literature review

The aim of this review was to detect the best protocol to assess and study motor fatigability. Therefore, we narrowed the literature search to included studies that specifically mentioned motor fatigue, muscle fatigue or fatigability. This was done to avoid non-relevant articles, but might have excluded studies where motor fatigability was a secondary outcome. The quality checklist of the included studies showed that a substantial part of the included research, suffered from methodological

issues. Previously, Meyer et al used a cut-off score of 65% for inclusion.¹⁸ However, if we used this value as a cut-off, only 22 studies would have been retained, leading to exclusion of several relevant protocols and outcomes.

Clinical assessment of motor fatigability

In 1977, it was already stated that it was necessary to investigate muscle endurance as an additional index of paresis.⁸³ Later, it was stated that some PwMS do not show strength deficits acutely, but experience abnormal motor fatigability over time.³⁷ To be able to use a protocol in a clinical setting, the protocol should be feasible, quick, easy to interpret and reliable. Based on the current review, no single protocol can be recommended for clinical use, due to the lack of comparison of protocols head to head and the lack of knowledge on psychometrics. To detect the PwMS prone to a slight increase in motor fatigability, a strenuous approach is needed, by determining the change in strength during sustained MVCs. Currently, the only outcome that has documented sufficient reliability in PwMS is a fatigue index based on the area under the curve, of a sustained MVC of 30 s.⁸ Previously, it was stated, that documenting the difference in strength before and after an exercise protocol is clinically feasible.⁶⁸ This is a feasible approach for physical exercises or daily life activities. With this approach, one should take into account that submaximal exercises might not elicit increased motor fatigability, when they are executed at the subjects own relative strength. Further, PwMS are likely to use a higher percentage of their maximal strength in daily life, compared to healthy controls.⁸⁴ Unfortunately, the current clinical tools to assess muscle strength, do not include standard protocols and outcomes measures for motor fatigability. Furthermore, the lack of cut-off values for this kind of protocols hinders clinical interpretation.

Recommendations for future research

This literature review revealed a number of limitations that needs to be kept in mind when interpreting the existing studies and designing future studies. To begin with, one should clearly define the concept that is assessed.^{3,7} Second, the existing psychometrics information only included limited information on the reliability of a few protocols.^{8,26} Related to this, no cut-off value exists to determine if motor fatigability is pathological. Future studies should identify norm values for motor fatigability. obtained from a large, well described population-based group of HC. Third, before collecting norm data, a consensus has to be reached on the best protocol. This is not possible at the moment due the almost complete lack of studies comparing different protocols head to head. Fourth, future intervention studies should evaluate patients with established motor fatigability at study entry, based on well-defined cut-off values, and perform comparisons of different interventions. Some research suggests that submaximal exercises are better to target the central nervous system.⁸⁵ As pathological motor fatigability in PwMS seems more related to central mechanisms, submaximal exercises with a high number of repetitions or a long duration might attenuate motor fatigability. There is already (limited) information on the influence on white matter structure of for example task specific training in PwMS.⁸⁶ However, for the improvement of strength, high intensity exercises are advised. It was suggested that motor fatigability can be improved by increasing the muscle mass with strength training,^{87,88} which might also increase neural drive in PwMS⁸⁹ and thus improve motor fatigability indirectly. Fifth, there is limited information on the relation between motor fatigability in different muscle groups⁸ and on the relation between motor fatigability on the different levels of the ICF. Although there seems to be a relation between motor fatigability on body function level and gait parameters,¹⁹ there are no reports explicitly examining the relation between motor fatigability on body function

level and activity level. It might be expected that motor fatigability affects functional tasks when sustained or high intensity muscle activity is required. The clinical meaning of protocols and outcomes for motor fatigability (i.e. the relation to quality of life or the participation in society) was not studied so far. It is, however, important that the influence of motor fatigability on daily functioning is elucidated, as the daily life functioning has a large impact on the quality of life of the patient.⁹⁰ Finally, the difference between PwMS and HC in underlying mechanisms of motor fatigability have predominantly been studied in the upper limbs, with only two reports evaluating the lower limbs.

Conclusion

A variety of protocols and outcome measures are applied to study different aspects of motor fatigability in PwMS, thus challenging comparability across studies. Most protocols use maximal single joint isometric contractions, with the advantage of high standardization. Protocols determining motor fatigability during submaximal or functional activities might, however, be more relevant for the PwMS. Due to the lack of head to head comparisons of the different protocols and a lack of information on psychometric properties, no gold standard is currently available to determine increased motor fatigability in PwMS.



Figure 1. Schematic representation of the literature search.



Figure 2. A schematic overview of the type of experimental protocols, used to assess motor fatigability in multiple sclerosis. UL: upper limb; LL: Lower limb; SB: strength based outcomes; IO: indirect outcomes for motor fatigability; NP: neurophysiological outcomes

References

- 1. Klenner FR. Fatigue, normal and pathological, with special consideration of myasthenia gravis and multiple sclerosis. *South Med Surg.* 1949;111(9):273-277.
- 2. Multiple Sclerosis Council for Clinical Practice Guidelines. *Fatigue and Multiple Sclerosis: evidence-based management strategies for fatigue in Multiple Sclerosis.* Paralyzed Veterans of America; 1998.
- 3. Kluger BM, Krupp LB, Enoka RM. Fatigue and fatigability in neurologic illnesses: proposal for a unified taxonomy. *Neurology*. 2013;80(4):409-416.
- 4. Induruwa I, Constantinescu CS, Gran B. Fatigue in multiple sclerosis a brief review. *J Neurol Sci.* 2012;323(1-2):9-15.
- 5. Patejdl R, Penner IK, Noack TK, Zettl UK. Multiple sclerosis and fatigue: A review on the contribution of inflammation and immune-mediated neurodegeneration. *Autoimmunity reviews*. 2016;15(3):210-220.
- 6. Chalah MA, Riachi N, Ahdab R, Creange A, Lefaucheur JP, Ayache SS. Fatigue in Multiple Sclerosis: Neural Correlates and the Role of Non-Invasive Brain Stimulation. *Front Cell Neurosci.* 2015;9:460.
- 7. Rudroff T, Kindred JH, Ketelhut NB. Fatigue in Multiple Sclerosis: Misconceptions and Future Research Directions. *Front Neurol.* 2016;7:122.
- 8. Schwid SR, Thornton CA, Pandya S, et al. Quantitative assessment of motor fatigue and strength in MS. *Neurology*. 1999;53(4):743-750.
- Feys P, Gijbels D, Romberg A, et al. Effect of time of day on walking capacity and self-reported fatigue in persons with multiple sclerosis: a multi-center trial. *Mult Scler*. 2012;18(3):351-357.
- 10. Schwid SR, Covington M, Segal BM, Goodman AD. Fatigue in multiple sclerosis: current understanding and future directions. *J Rehabil Res Dev.* 2002;39(2):211-224.
- 11. Djaldetti R, Ziv I, Achiron A, Melamed E. Fatigue in multiple sclerosis compared with chronic fatigue syndrome: A quantitative assessment. *Neurology.* 1996;46(3):632-635.
- 12. World Health Organisation. *International Classification of Functioning, Disability and Health: ICF.* Geneva: WHO; 2001.
- 13. Enoka RM. Mechanisms of muscle fatigue: Central factors and task dependency. *J Electromyogr Kinesiol.* 1995;5(3):141-149.
- 14. Vollestad NK. Measurement of human muscle fatigue. *J Neurosci Methods.* 1997;74(2):219-227.
- 15. Taylor JL, Gandevia SC. A comparison of central aspects of fatigue in submaximal and maximal voluntary contractions. *J Appl Physiol (1985).* 2008;104(2):542-550.
- 16. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *PLoS Med.* 2009;6(7):e1000100.
- 17. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health.* 1998;52(6):377-384.
- 18. Meyer S, Karttunen AH, Thijs V, Feys H, Verheyden G. How do somatosensory deficits in the arm and hand relate to upper limb impairment, activity, and participation problems after stroke? A systematic review. *Phys Ther.* 2014;94(9):1220-1231.
- 19. Kalron A, Achiron A, Dvir Z. Muscular and gait abnormalities in persons with early onset multiple sclerosis. *J Neurol Phys Ther.* 2011;35(4):164-169.
- 20. Kalron A, Achiron A, Dvir Z. Motor impairments at presentation of clinically isolated syndrome suggestive of multiple sclerosis: Characterization of different disease subtypes. *NeuroRehabilitation*. 2012;31(2):147-155.

- Kalron A, Dvir Z, Gurevich M, Achiron A. Do motor impairments detected on onset of multiple sclerosis suggest an early second attack? A prospective study. *NeuroRehabilitation*. 2013;33(3):423-430.
- 22. Severijns D, Lemmens M, Thoelen R, Feys P. Motor fatigability after low-intensity hand grip exercises in persons with multiple sclerosis. *Multiple Sclerosis and Related Disorders*.10:7-13.
- 23. Andreasen AK, Jakobsen J, Petersen T, Andersen H. Fatigued patients with multiple sclerosis have impaired central muscle activation. *Mult Scler.* 2009;15(7):818-827.
- 24. Severijns D, Lamers I, Kerkhofs L, Feys P. Hand grip fatigability in persons with multiple sclerosis according to hand dominance and disease progression. *J Rehabil Med.* 2015;47(2):154-160.
- 25. Surakka J, Romberg A, Ruutiainen J, et al. Effects of aerobic and strength exercise on motor fatigue in men and women with multiple sclerosis: a randomized controlled trial. *Clin Rehabil.* 2004;18(7):737-746.
- 26. Surakka J, Romberg A, Ruutiainen J, Virtanen A, Aunola S, Maentaka K. Assessment of muscle strength and motor fatigue with a knee dynamometer in subjects with multiple sclerosis: a new fatigue index. *Clin Rehabil.* 2004;18(6):652-659.
- 27. Romani A, Bergamaschi R, Candeloro E, Alfonsi E, Callieco R, Cosi V. Fatigue in multiple sclerosis: multidimensional assessment and response to symptomatic treatment. *Mult Scler*. 2004;10(4):462-468.
- 28. Sheean GL, Murray NMF, Rothwell JC, Miller DH, Thompson AJ. An electrophysiological study of the mechanism of fatigue in multiple sclerosis. *Brain.* 1997;120:299-315.
- 29. Latash M, Kalugina E, Nicholas J, Orpett C, Stefoski D, Davis F. Myogenic and central neurogenic factors in fatigue in multiple sclerosis. *Mult Scler.* 1996;1(4):236-241.
- 30. Skurvydas A, Brazaitis M, Andrejeva J, Mickeviciene D, Streckis V. The effect of multiple sclerosis and gender on central and peripheral fatigue during 2-min MVC. *Clin Neurophysiol*. 2011;122(4):767-776.
- 31. Steens A, de Vries A, Hemmen J, et al. Fatigue perceived by multiple sclerosis patients is associated with muscle fatigue. *Neurorehabil Neural Repair*. 2012;26(1):48-57.
- 32. Steens A, Heersema DJ, Maurits NM, Renken RJ, Zijdewind I. Mechanisms underlying muscle fatigue differ between multiple sclerosis patients and controls: a combined electrophysiological and neuroimaging study. *Neuroimage*. 2012;59(4):3110-3118.
- 33. Streckis V, Skurvydas A, Mamkus G. Effect of the time of day on central and peripheral fatigue during 2-min maximal voluntary contractions in persons with multiple sclerosis: gender differences. *J Electromyogr Kinesiol.* 2014;24(5):601-606.
- 34. Wolkorte R, Heersema DJ, Zijdewind I. Muscle Fatigability During a Sustained Index Finger Abduction and Depression Scores Are Associated With Perceived Fatigue in Patients With Relapsing-Remitting Multiple Sclerosis. *Neurorehabil Neural Repair.* 2015;29(8):796-802.
- 35. Wolkorte R, Heersema DJ, Zijdewind I. Reduced Dual-Task Performance in MS Patients Is Further Decreased by Muscle Fatigue. *Neurorehabil Neural Repair.* 2015;29(5):424-435.
- 36. Wolkorte R, Heersema DJ, Zijdewind I. Reduced Voluntary Activation During Brief and Sustained Contractions of a Hand Muscle in Secondary-Progressive Multiple Sclerosis Patients. *Neurorehabil Neural Repair.* 2016;30(4):307-316.
- 37. Petajan JH, White AT. Motor-evoked potentials in response to fatiguing grip exercise in multiple sclerosis patients. *Clin Neurophysiol.* 2000;111(12):2188-2195.
- 38. White AT, Lee JN, Light AR, Light KC. Brain activation in multiple sclerosis: a BOLD fMRI study of the effects of fatiguing hand exercise. *Mult Scler.* 2009;15(5):580-586.
- 39. White AT, Vanhaitsma TA, Vener J, Davis SL. Effect of passive whole body heating on central conduction and cortical excitability in multiple sclerosis patients and healthy controls. *J Appl Physiol (1985).* 2013;114(12):1697-1704.
- 40. Iriarte J, de Castro P. Correlation between symptom fatigue and muscular fatigue in multiple sclerosis. *Eur J Neurol.* 1998;5(6):579-585.

- 41. Ickmans K, Simoens F, Nijs J, et al. Recovery of peripheral muscle function from fatiguing exercise and daily physical activity level in patients with multiple sclerosis: a case-control study. *Clin Neurol Neurosurg.* 2014;122:97-105.
- 42. Scheidegger O, Kamm CP, Humpert SJ, Rosler KM. Corticospinal output during muscular fatigue differs in multiple sclerosis patients compared to healthy controls. *Mult Scler*. 2012;18(10):1500-1506.
- 43. Liepert J, Mingers D, Heesen C, Baumer T, Weiller C. Motor cortex excitability and fatigue in multiple sclerosis: a transcranial magnetic stimulation study. *Mult Scler.* 2005;11(3):316-321.
- 44. Thickbroom GW, Sacco P, Kermode AG, et al. Central motor drive and perception of effort during fatigue in multiple sclerosis. *J Neurol.* 2006;253(8):1048-1053.
- 45. Perretti A, Balbi P, Orefice G, et al. Post-exercise facilitation and depression of motor evoked potentials to transcranial magnetic stimulation: a study in multiple sclerosis. *Clin Neurophysiol.* 2004;115(9):2128-2133.
- 46. Thickbroom GW, Sacco P, Faulkner DL, Kermode AG, Mastaglia FL. Enhanced corticomotor excitability with dynamic fatiguing exercise of the lower limb in multiple sclerosis. *J Neurol.* 2008;255(7):1001-1005.
- 47. Severijns D, Octavia JR, Kerkhofs L, Coninx K, Lamers I, Feys P. Investigation of Fatigability during Repetitive Robot-Mediated Arm Training in People with Multiple Sclerosis. *PLoS One.* 2015;10(7):e0133729.
- 48. Broekmans T, Roelants M, Alders G, Feys P, Thijs H, Eijnde BO. Exploring the effects of a 20week whole-body vibration training programme on leg muscle performance and function in persons with multiple sclerosis. *J Rehabil Med.* 2010;42(9):866-872.
- 49. Lambert CP, Archer RL, Evans WJ. Muscle strength and fatigue during isokinetic exercise in individuals with multiple sclerosis. *Med Sci Sports Exerc.* 2001;33(10):1613-1619.
- 50. Hameau S, Zory R, Latrille C, Roche N, Bensmail D. Relationship between neuromuscular and perceived fatigue and locomotor performance in patients with multiple sclerosis. *Eur J Phys Rehabil Med.* 2016.
- 51. Greim B, Benecke R, Zettl UK. Qualitative and quantitative assessment of fatigue in multiple sclerosis (MS). *J Neurol*. 2007;254 Suppl 2:II58-64.
- 52. de Souza-Teixeira F, Castilla S, Ayan C, Garcia-Lopez D, Gonzalez-Gallego J, de Paz JA. Effects of Resistance Training in Multiple Sclerosis. *Int J Sports Med.* 2009;30(4):245-250.
- 53. Dodd KJ, Taylor NF, Shields N, Prasad D, McDonald E, Gillon A. Progressive resistance training did not improve walking but can improve muscle performance, quality of life and fatigue in adults with multiple sclerosis: a randomized controlled trial. *Mult Scler.* 2011;17(11):1362-1374.
- 54. McLoughlin JV, Barr CJ, Crotty M, Sturnieks DL, Lord SR. Six minutes of walking leads to reduced lower limb strength and increased postural sway in people with Multiple Sclerosis. *NeuroRehabilitation.* 2014;35(3):503-508.
- 55. Leone C, Severijns D, Dolezalova V, et al. Prevalence of Walking-Related Motor Fatigue in Persons With Multiple Sclerosis: Decline in Walking Distance Induced by the 6-Minute Walk Test. *Neurorehabil Neural Repair.* 2016;30(4):373-383.
- 56. Burschka JM, Keune PM, Menge U, Hofstadt-van Oy U, Oschmann P, Hoos O. An exploration of impaired walking dynamics and fatigue in multiple sclerosis. *BMC Neurol.* 2012;12:161.
- 57. Engelhard MM, Dandu SR, Patek SD, Lach JC, Goldman MD. Quantifying six-minute walk induced gait deterioration with inertial sensors in multiple sclerosis subjects. *Gait Posture*. 2016;49:340-345.
- 58. Phan-Ba R, Calay P, Grodent P, et al. Motor fatigue measurement by distance-induced slow down of walking speed in multiple sclerosis. *PLoS One.* 2012;7(4):e34744.
- 59. Sehle A, Mundermann A, Starrost K, et al. Objective assessment of motor fatigue in Multiple Sclerosis using kinematic gait analysis: a pilot study. *J Neuroeng Rehabil.* 2011;8:59.

- 60. Sehle A, Vieten M, Mundermann A, Dettmers C. Difference in Motor Fatigue between Patients with Stroke and Patients with Multiple Sclerosis: A Pilot Study. *Front Neurol.* 2014;5:279.
- 61. Sehle A, Vieten M, Sailer S, Mundermann A, Dettmers C. Objective assessment of motor fatigue in multiple sclerosis: the Fatigue index Kliniken Schmieder (FKS). *J Neurol.* 2014;261(9):1752-1762.
- 62. Fay BT, Boninger ML, Fitzgerald SG, Souza AL, Cooper RA, Koontz AM. Manual wheelchair pushrim dynamics in people with multiple sclerosis. *Arch Phys Med Rehabil.* 2004;85(6):935-942.
- 63. Jonkers I, Nuyens G, Seghers J, Nuttin M, Spaepen A. Muscular effort in multiple sclerosis patients during powered wheelchair manoeuvres. *Clin Biomech (Bristol, Avon).* 2004;19(9):929-938.
- 64. Korkmaz NC, Kirdi N, Temucin CM, Armutlu K, Yakut Y, Karabudak R. Improvement of muscle strength and fatigue with high voltage pulsed galvanic stimulation in multiple sclerosis patients--a non-randomized controlled trial. *J Pak Med Assoc.* 2011;61(8):736-743.
- 65. Allen GM, Gandevia SC, McKenzie DK. Reliability of measurements of muscle strength and voluntary activation using twitch interpolation. *Muscle Nerve.* 1995;18(6):593-600.
- 66. Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev.* 2001;81(4):1725-1789.
- 67. Enoka RM. Muscle fatigue--from motor units to clinical symptoms. *J Biomech.* 2012;45(3):427-433.
- 68. Dobkin BH. Fatigue versus activity-dependent fatigability in patients with central or peripheral motor impairments. *Neurorehabil Neural Repair.* 2008;22(2):105-110.
- 69. Sogaard K, Gandevia SC, Todd G, Petersen NT, Taylor JL. The effect of sustained low-intensity contractions on supraspinal fatigue in human elbow flexor muscles. *J Physiol.* 2006;573(Pt 2):511-523.
- 70. Reuter SE, Massy-Westropp N, Evans AM. Reliability and validity of indices of hand-grip strength and endurance. *Aust Occup Ther J.* 2011;58(2):82-87.
- 71. Larsson B, Karlsson S, Eriksson M, Gerdle B. Test-retest reliability of EMG and peak torque during repetitive maximum concentric knee extensions. *J Electromyogr Kinesiol.* 2003;13(3):281-287.
- 72. Bosquet L, Maquet D, Forthomme B, Nowak N, Lehance C, Croisier JL. Effect of the lengthening of the protocol on the reliability of muscle fatigue indicators. *Int J Sports Med.* 2010;31(2):82-88.
- 73. Contessa P, Adam A, De Luca CJ. Motor unit control and force fluctuation during fatigue. *J Appl Physiol (1985).* 2009;107(1):235-243.
- 74. Zijdewind I, Prak R, Wolkorte R. Fatigue and Fatigability in Persons with Multiple Sclerosis. *Exerc Sport Sci Rev.* 2016.
- 75. Liu JZ, Zhang L, Yao B, Sahgal V, Yue GH. Fatigue induced by intermittent maximal voluntary contractions is associated with significant losses in muscle output but limited reductions in functional MRI-measured brain activation level. *Brain Res.* 2005;1040(1-2):44-54.
- 76. Lloyd AR, Gandevia SC, Hales JP. Muscle performance, voluntary activation, twitch properties and perceived effort in normal subjects and patients with the chronic fatigue syndrome. *Brain.* 1991;114 (Pt 1A):85-98.
- 77. Sheean GL, Murray NM, Rothwell JC, Miller DH, Thompson AJ. An electrophysiological study of the mechanism of fatigue in multiple sclerosis. *Brain.* 1997;120 299-315.
- 78. Zijdewind I, Zwarts MJ, Kernell D. Influence of a voluntary fatigue test on the contralateral homologous muscle in humans? *Neurosci Lett.* 1998;253(1):41-44.
- 79. Gonzales JU, Scheuermann BW. Absence of gender differences in the fatigability of the forearm muscles during intermittent isometric handgrip exercise. *J Sports Sci Med.* 2007;6(1):98-105.

- 80. White C, Dixon K, Samuel D, Stokes M. Handgrip and quadriceps muscle endurance testing in young adults. *Springerplus.* 2013;2:451.
- 81. Gerdle B, Larsson B, Karlsson S. Criterion validation of surface EMG variables as fatigue indicators using peak torque: a study of repetitive maximum isokinetic knee extensions. *J Electromyogr Kinesiol.* 2000;10(4):225-232.
- 82. Dideriksen JL, Farina D, Enoka RM. Influence of fatigue on the simulated relation between the amplitude of the surface electromyogram and muscle force. *Philos Trans A Math Phys Eng Sci.* 2010;368(1920):2765-2781.
- 83. Pedersen L, Raun NE. Endurance test as index of paresis in patients with multiple sclerosis. *Acta Neurol Scand.* 1977;55(2):163-168.
- 84. Iyengar V, Santos MJ, Ko M, Aruin AS. Grip force control in individuals with multiple sclerosis. *Neurorehabil Neural Repair.* 2009;23(8):855-861.
- 85. Yoon T, Schlinder Delap B, Griffith EE, Hunter SK. Mechanisms of fatigue differ after low- and high-force fatiguing contractions in men and women. *Muscle Nerve.* 2007;36(4):515-524.
- 86. Bonzano L, Tacchino A, Brichetto G, et al. Upper limb motor rehabilitation impacts white matter microstructure in multiple sclerosis. *Neuroimage*. 2014;90:107-116.
- 87. de Haan A, de Ruiter CJ, van Der Woude LH, Jongen PJ. Contractile properties and fatigue of quadriceps muscles in multiple sclerosis. *Muscle Nerve.* 2000;23(10):1534-1541.
- 88. Kent-Braun JA, Sharma KR, Miller RG, Weiner MW. Postexercise phosphocreatine resynthesis is slowed in multiple sclerosis. *Muscle Nerve.* 1994;17(8):835-841.
- 89. Fimland MS, Helgerud J, Gruber M, Leivseth G, Hoff J. Enhanced neural drive after maximal strength training in multiple sclerosis patients. *Eur J Appl Physiol.* 2010;110(2):435-443.
- 90. Enoka RM, Duchateau J. Translating Fatigue to Human Performance. *Med Sci Sports Exerc.* 2016.