

On the realisability of double-cross matrices by polylines in the plane

Peer-reviewed author version

KUIJPERS, Bart & MOELANS, Bart (2017) On the realisability of double-cross matrices by polylines in the plane. In: JOURNAL OF COMPUTER AND SYSTEM SCIENCES, 86, p. 117-135.

DOI: 10.1016/j.jcss.2016.12.001

Handle: <http://hdl.handle.net/1942/23452>

# On the realisability of double-cross matrices by polylines in the plane

Bart Kuijpers<sup>1,\*</sup>, Bart Moelans<sup>1,2,\*</sup>

*UHasselt – Hasselt University and transnational University Limburg, Belgium*

*Unleashed, Hasselt, Belgium*

---

## Abstract

We study a decision problem, that emerges from the area of spatial reasoning. This decision problem concerns the description of polylines in the plane by means of their *double-cross matrix*. In such a matrix, the relative position of each pair of line segments in a polyline is expressed by means of a 4-tuple over  $\{-, 0, +\}$ . However, not any such matrix of 4-tuples is the double-cross matrix of a polyline. This gives rise to the decision problem: given a matrix of such 4-tuples, decide whether it is the double-cross matrix of a polyline. This problem is decidable, but it is **NP-hard**. In this paper, we give polynomial-time algorithms for the cases where consecutive line segments in a polyline make angles that are multiples of  $90^\circ$  or  $45^\circ$  and for the case where, apart from an input matrix, the successive angles of a polyline are also given as input.

*Keywords:* Spatial reasoning, Double-cross calculus, Qualitative description of polylines, Computational algebraic geometry, Algorithmic complexity

---

---

\*Corresponding author

<sup>1</sup>UHasselt – Hasselt University and transnational University Limburg, Databases and Theoretical Computer Science Research Group, Agoralaan, Gebouw D, 3590 Diepenbeek, Belgium, bart.kuijpers@uhasselt.be

<sup>2</sup>Unleashed, Kempische Steenweg 309, Box 1, 3500 Hasselt, Belgium, bart.moelans@unleashed.be

## 1. Introduction and summary of results

Polylines arise in Geographical Information Science (GIS) in a multitude of ways. One recent example comes from the collection of moving object data, where trajectories of moving persons (or animals), that carry GPS-equipped devices, are collected in the form of time-space points that are registered at certain (ir)regular moments in time. The spatial trace of this movement is a collection of points in two-dimensional geographical space, that form a polyline, when in between the measured sample points, for instance, linear interpolation is applied (Güting and Schneider (2005)). Another example of the use of polylines comes from shape recognition and retrieval, which arises in domains, such as computer vision and image analysis. Here, closed polylines (or polygons) often occur as the boundary of two-dimensional shapes or regions.

In examples, such as the above, there are, roughly speaking, two very distinct approaches to deal with polygonal curves and shapes. On the one hand, there are the *quantitative* approaches and, on the other hand, there are the *qualitative* approaches. Initially, most research efforts have dealt with the quantitative methods (Bookstein (1986); Dryden and Mardia (1998); Kent and Mardia (1986); Mokhtarian and Mackworth (1992)). Only afterwards, the qualitative approaches have gained more attention, mainly supported by research in cognitive science that provides evidence that qualitative models of shape representation are much more expressive than their quantitative counterpart and reflect better the way in which humans reason about their environment (Gero (1999)). The principles behind qualitative approaches to deal with polylines are also related to the field of *spatial reasoning*, which has as one of its main objectives to present geographic information in a qualitative way, to facilitate reasoning about it. For an overview of spatial and for spatio-temporal reasoning, we refer to Chapter 12 in (Giannotti and Pedreschi (2008)). The reason for using a *qualitative representation* is that the available information is often imprecise, partial and subjective (Freksa (1992)).

One of the formalisms to qualitatively describe polylines in the plane is given by the *double-cross calculus*. In this method, a *double-cross matrix* captures the relative position (or orientation) of any two line segments in a polyline by describing it with respect to a double cross based on the starting points of these line segments (Freksa (1992); Zimmermann and Freksa (1996)). For an overview of the use of constraint calculi in qualitative spatial

reasoning, we refer to (Renz and Nebel (2007)). In the  $N \times N$  double-cross matrix of a polyline with  $N$  line segments (or  $N + 1$  vertices), the relative position (or orientation) of two (oriented) line segments in a polyline is encoded by means of a 4-tuple, whose entries come from the set  $\{-, 0, +\}$ .

However, not every  $N \times N$  matrix of 4-tuples from  $\{-, 0, +\}$  is the double-cross matrix of a polyline with  $N + 1$  vertices. This gives rise to the following decision problem: *Given an  $N \times N$  matrix of 4-tuples from  $\{-, 0, +\}$ , decide whether or not it is the double-cross matrix of a polyline (with  $N + 1$  vertices), and if it is, given an example of a polyline that realises the matrix.*

To start with, we give a known collection of polynomial (in)equalities on the coordinates of the vertices of a polyline, that express the information contained in the double-cross matrix of a polyline. Since first-order logic over the reals (or elementary geometry) is decidable (Tarski (1951)), it follows that this decision problem is also decidable. However, we are left with the question of its time complexity.

In computational algebraic geometry, the problem can be viewed as a satisfiability problem of a system of quadratic equations in  $2(N + 1)$  variables. However, the known best algorithms to solve our problem (including the production of sample points) take exponential time. Our decision problem has many particularities: the polynomials are homogeneous of degree 2; they use few monomials and each of them uses only six variables. Nevertheless, the problem is known to be NP-hard (Scivos and Nebel (2001); Renz and Nebel (2007)). Whether or not this problem is in NP is less obvious, since no apriori polynomial bound on the complexity of sample points (to be guessed) is obvious. We discuss this problem in more detail in Section 3.

In this paper, we focus on subclasses of the above decision problem, for which we can give *polynomial time* decision algorithms. A first subclass is obtained by restricting the attention to polylines in which consecutive line segments make angles that are multiples of  $90^\circ$ . For this sub-problem, we give a  $O(N^2)$ -time decision procedure. Next, we turn our attention to polylines in which consecutive line segments make angles that are multiples of  $45^\circ$ . To solve the more complicated case of  $45^\circ$ -polylines, we introduce the polar-coordinate representation of double-cross matrices. We give two-way translations between the Cartesian- and the polar-coordinate representations. Using polar coordinates, our decision problem can be reduced to a linear programming problem (with algebraic coefficients, however). For the particular decision problem of a double-cross matrix  $M$  being realisable (or not) by a  $45^\circ$ -polyline, we can make use of the fact that the entries of  $M$  above its diag-

onal give exact information on the angles that a polyline, that would realise  $M$ , should have. This one-to-one correspondence between the qualitative double-cross information and the angle information implies that our decision problem simplifies to deciding whether or not appropriate segment lengths of a polyline exist. The latter problem is a linear programming problem, that can be solved in polynomial time. In fact, we show that this situation can be generalised and we first show that whenever the consecutive angles of a polyline are given, it can be decided in polynomial time whether a matrix  $M$  can be realised by a polyline (with the given angle sequence). Next, we apply this more general result to the case of  $45^\circ$ -polylines to obtain a polynomial time decision procedure. This result has some implications on the convexity of the solution set consisting of all  $45^\circ$ -polylines that realise a matrix. It is not the intention of this paper to discuss implementations of and experiments with the proposed methods.

*Organisation.* This paper is organised as follows. Section 2 gives the definition of a polyline, the double-cross matrix of a polyline and the known results on the algebraic interpretation of the double-cross matrix. In Section 3, we state our decision problem in a more technical way and discuss some of its general properties. Section 4 gives a  $O(N^2)$ -time decision procedure for the case of  $90^\circ$ -polylines. In Section 5, we introduce the polar-coordinate representation of double-cross matrices. In Section 6, we use the double-cross conditions in polar form to show the existence of a polynomial-time realisability test in the case where, apart from an input matrix, also the successive angles of a polyline are given as input. Section 7 gives a polynomial-time decision procedure for the case of  $45^\circ$ -polylines. The paper ends with concluding remarks that include variants of our decision problem.

## 2. Definition and preliminaries

In this section, we give the definitions of a polyline, an  $\alpha$ -polyline and of the double-cross matrix of a polyline. We also give an algebraic interpretation of the double-cross matrix.

We start with the following notational conventions. Let  $\mathbf{R}$  denote the sets of the real numbers, and let  $\mathbf{R}^2$  denote the two-dimensional real plane. To stress that some real values are constants, we use sans serif characters:  $x, y, x_0, y_0, x_1, y_1, \dots$ . Real variables are denoted in normal characters. For constant points of  $\mathbf{R}^2$ , we use the sans serif characters  $\mathbf{p}, \mathbf{p}_0, \mathbf{p}_1, \dots$ .

## 2.1. Polyline and $\alpha$ -polylines

The following definition specifies what we mean by *polylines*. We define polylines as a finite sequences of points in  $\mathbf{R}^2$  (which is often used as their finite representation). When we add the line segments between consecutive points we obtain what we call the *semantics* of the polyline. We also introduce some terminology about polylines.

**Definition 1.** A *polyline* (in  $\mathbf{R}^2$ ) is an ordered list  $P = \langle (x_0, y_0), (x_1, y_1), \dots, (x_N, y_N) \rangle$  of points in  $\mathbf{R}^2$ . We call the points  $(x_i, y_i)$ ,  $0 \leq i \leq N$ , the *vertices* of the polyline. We assume that no two consecutive vertices are identical, that is:  $(x_i, y_i) \neq (x_{i+1}, y_{i+1})$ , for  $0 \leq i < N$ .

The vertices  $(x_0, y_0)$  and  $(x_N, y_N)$  are respectively called the *start* and *end vertex* of  $P$ . The line segments connecting the points  $(x_i, y_i)$  and  $(x_{i+1}, y_{i+1})$ , for  $0 \leq i < N$ , are called the *(line) segments* of the polyline  $P$ . The *semantics* of  $P$ , denoted  $\text{sem}(P)$ , is the union of the line segments of  $P$ . We call  $N$ , the number of line segments, the *size* of the polyline  $P$ .  $\square$

Figure 1 gives an example of two polylines,  $P_1$  and  $P_2$ , of size 4 and their semantics. Further on, we will loosely use the term polyline also to refer to the semantics of a polyline, although, *stricto sensu*, a polyline is a ordered list of points in  $\mathbf{R}^2$ .

We remark that, by the above definition, two polylines with a different number of vertices, may have the same semantics. We also remark that the line segments, appearing in the semantics, may intersect in points which may or may be not vertices. Finally, we remark that it is reasonable to assume that polylines coming from GIS applications have vertices with rational coordinates (or that are finitely representable in some other way).

We use the following additional notational conventions. As a standard, for vertices of a polyline, we abbreviate  $(x_i, y_i)$  by  $\mathbf{p}_i$ . The *(located) vector*<sup>3</sup> from  $\mathbf{p}_i$  to  $\mathbf{p}_j$  is denoted by  $\overrightarrow{\mathbf{p}_i \mathbf{p}_j}$ . The counter-clockwise angle (expressed in degrees) measured from  $\overrightarrow{\mathbf{p}_i \mathbf{p}_j}$  to  $\overrightarrow{\mathbf{p}_i \mathbf{p}_k}$  is denoted by  $\angle(\overrightarrow{\mathbf{p}_i \mathbf{p}_j}, \overrightarrow{\mathbf{p}_i \mathbf{p}_k})$ , as illustrated in Figure 2.

In this paper, we use 45°- and 90°-polylines, which are special cases of  $\alpha$ -polylines

---

<sup>3</sup>By the *located vector* from  $\mathbf{p}$  to  $\mathbf{q}$ , we mean an ordered pair  $(\mathbf{p}, \mathbf{q})$  of points of  $\mathbf{R}^2$ , which we denote  $\overrightarrow{\mathbf{p} \mathbf{q}}$ . We use this concept to represent the oriented line segment between  $\mathbf{p}$  and  $\mathbf{q}$ .




Figure 1: An example of two polylines,  $P_1$  and  $P_2$ , of size 4 (the dots) and their semantics (the lines).




Figure 2: The counter-clockwise angle  $\angle(\overrightarrow{p_i p_j}, \overrightarrow{p_i p_k})$  from  $\overrightarrow{p_i p_j}$  to  $\overrightarrow{p_i p_k}$ .

**Definition 2.** Let  $\alpha$ ,  $0^\circ < \alpha < 360^\circ$ , be an angle such that  $\frac{360^\circ}{\alpha} = k_\alpha$  is a natural number. Let  $P = \langle p_0, p_1, \dots, p_N \rangle$  be a polyline. We call  $P$  an  $\alpha$ -polyline if all angles  $\angle(\overrightarrow{p_i p_{i-1}}, \overrightarrow{p_i p_{i+1}})$  are multiples of  $\alpha$ , for  $0 < i < N$ , that is, if  $\angle(\overrightarrow{p_i p_{i-1}}, \overrightarrow{p_i p_{i+1}})$  is of the form  $n_i \alpha$ , with  $n_i \in \{0, 1, \dots, k_\alpha\}$ .  $\square$

Figure 3 shows the  $90^\circ$ -polyline  $P_1$  and the  $45^\circ$ -polylines  $P_1$  and  $P_2$ . Indeed, in the polyline  $P_1$ , for instance, the consecutive angles are  $90^\circ, 90^\circ, 270^\circ$  and  $270^\circ$ , assuming that the start vertex is at the left bottom.

## 2.2. The double-cross matrix of a polyline

As mentioned in the Introduction, in the double-cross formalism, the relative position (or orientation) of two (located) vectors of a polyline is encoded by means of a 4-tuple, whose entries come from the set  $\{-, 0, +\}$  (Freksa (1992); Zimmermann and Freksa (1996)). Such a 4-tuple expresses the relative orientation of two vectors with respect to each other.




Figure 3: An example of a 90°-polyline ( $P_1$ ) and two 45°-polylines ( $P_1$  and  $P_2$ ).

In this section, we define the *double-cross matrix* of a polyline. We associate to a polyline  $P = \langle \mathbf{p}_0, \mathbf{p}_1, \dots, \mathbf{p}_N \rangle$  the (located) vectors  $\overrightarrow{\mathbf{p}_0\mathbf{p}_1}, \overrightarrow{\mathbf{p}_1\mathbf{p}_2}, \dots, \overrightarrow{\mathbf{p}_{N-1}\mathbf{p}_N}$ , representing the oriented line segments between the consecutive vertices of  $P$ . Because of the assumption in Definition 1, the vectors  $\overrightarrow{\mathbf{p}_0\mathbf{p}_1}, \overrightarrow{\mathbf{p}_1\mathbf{p}_2}, \dots, \overrightarrow{\mathbf{p}_{N-1}\mathbf{p}_N}$  all have a strictly positive length. In the double-cross formalism, the relative orientation between  $\overrightarrow{\mathbf{p}_i\mathbf{p}_{i+1}}$  and  $\overrightarrow{\mathbf{p}_j\mathbf{p}_{j+1}}$  is given by means of a 4-tuple  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$ . We follow the traditional notation of this 4-tuple *without commas*. To determine  $C_1, C_2, C_3$  and  $C_4$ , for  $\mathbf{p}_i \neq \mathbf{p}_j$ , first of all, a *double cross* is defined for the vectors  $\overrightarrow{\mathbf{p}_i\mathbf{p}_{i+1}}$  and  $\overrightarrow{\mathbf{p}_j\mathbf{p}_{j+1}}$ , determined by the following three lines:

- the line  $L_{ij}$  through  $\mathbf{p}_i$  and  $\mathbf{p}_j$ ;
- the line  $P_{iji}$  through  $\mathbf{p}_i$ , perpendicular on  $L_{ij}$ ; and
- the line  $P_{ijj}$  through  $\mathbf{p}_j$ , perpendicular on  $L_{ij}$ .

These three lines are illustrated in Figure 4. These three lines determine a cross at  $\mathbf{p}_i$  and a cross at  $\mathbf{p}_j$ . Hence the name “double cross.” The entries  $C_1, C_2, C_3$  and  $C_4$  express in which quadrants or on which half lines  $\mathbf{p}_{i+1}$  and  $\mathbf{p}_{j+1}$  are located with respect to the double cross.

We now define this more formally and follow the historical use of the double cross. In this definition, an interval  $(a, b)$  of angles, represents the open interval between  $a$  and  $b$  on the counter-clockwise oriented circle.

**Definition 3.** Let  $P = \langle \mathbf{p}_0, \mathbf{p}_1, \dots, \mathbf{p}_N \rangle$  be a polyline with associated vectors  $\overrightarrow{\mathbf{p}_0\mathbf{p}_1}, \overrightarrow{\mathbf{p}_1\mathbf{p}_2}, \dots, \overrightarrow{\mathbf{p}_{N-1}\mathbf{p}_N}$ . For  $\overrightarrow{\mathbf{p}_i\mathbf{p}_{i+1}}$  and  $\overrightarrow{\mathbf{p}_j\mathbf{p}_{j+1}}$  with  $0 \leq i, j < N$ ,  $i \neq j$  and  $\mathbf{p}_i \neq \mathbf{p}_j$ , we define

$$\text{DC}(\overrightarrow{\mathbf{p}_i\mathbf{p}_{i+1}}, \overrightarrow{\mathbf{p}_j\mathbf{p}_{j+1}}) = (C_1 \ C_2 \ C_3 \ C_4)$$






Figure 4: The double cross (in blue): the lines  $L_{ij}$ ,  $P_{iji}$  and  $P_{ijj}$ .

as follows:

$$C_1 = \begin{cases} - & \text{if } \angle(\overrightarrow{p_i p_j}, \overrightarrow{p_i p_{i+1}}) \in (-90^\circ, 90^\circ) \\ 0 & \text{if } \angle(\overrightarrow{p_i p_j}, \overrightarrow{p_i p_{i+1}}) \in \{-90^\circ, 90^\circ\} \\ + & \text{else} \end{cases}$$

$$C_2 = \begin{cases} - & \text{if } \angle(\overrightarrow{p_j p_i}, \overrightarrow{p_j p_{j+1}}) \in (-90^\circ, 90^\circ) \\ 0 & \text{if } \angle(\overrightarrow{p_j p_i}, \overrightarrow{p_j p_{j+1}}) \in \{-90^\circ, 90^\circ\} \\ + & \text{else} \end{cases}$$

$$C_3 = \begin{cases} - & \text{if } \angle(\overrightarrow{p_i p_j}, \overrightarrow{p_i p_{i+1}}) \in (0^\circ, 180^\circ) \\ 0 & \text{if } \angle(\overrightarrow{p_i p_j}, \overrightarrow{p_i p_{i+1}}) \in \{0^\circ, 180^\circ\} \\ + & \text{else} \end{cases}$$

$$C_4 = \begin{cases} - & \text{if } \angle(\overrightarrow{p_j p_i}, \overrightarrow{p_j p_{j+1}}) \in (0^\circ, 180^\circ) \\ 0 & \text{if } \angle(\overrightarrow{p_j p_i}, \overrightarrow{p_j p_{j+1}}) \in \{0^\circ, 180^\circ\} \\ + & \text{else.} \end{cases}$$

For  $\overrightarrow{p_i p_{i+1}}$  and  $\overrightarrow{p_j p_{j+1}}$ , with  $p_i = p_j$ , we define, for reasons of continuity,<sup>4</sup>

$$\text{DC}(\overrightarrow{p_i p_{i+1}}, \overrightarrow{p_j p_{j+1}}) = (0 \ 0 \ 0 \ 0).$$

The *double-cross matrix* of  $P$ , denoted  $\text{DCM}(P)$ , is the  $N \times N$  matrix with the entries  $\text{DCM}(P)[i, j] = \text{DC}(\overrightarrow{p_i p_{i+1}}, \overrightarrow{p_j p_{j+1}})$ , for  $0 \leq i, j < N$ .  $\square$

<sup>4</sup>This argumentation is given in (Forbus (1990)).

So, in particular, when  $i = j$ , we have  $\text{DC}(\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}, \overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}}) = (0 \ 0 \ 0 \ 0)$ .

We remark that the values  $C_1$  and  $C_3$  describe the location of the point  $\mathbf{p}_{i+1}$  or, equivalently, the orientation of the vector  $\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}$  with respect to the cross at  $\mathbf{p}_i$  (formed by the lines  $L_{ij}$  and  $P_{iji}$ ). We see that each of the four quadrants and four half lines determined by the cross at  $\mathbf{p}_i$  are determined by a unique combination of  $C_1$  and  $C_3$  values. Similarly, the values  $C_2$  and  $C_4$  describe the location of the point  $\mathbf{p}_{j+1}$  or, equivalently, the orientation of the vector  $\overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}}$  with respect to the cross at  $\mathbf{p}_j$  (formed by the lines  $L_{ij}$  and  $P_{ijj}$ ). The quadrants and half lines where  $C_1$ ,  $C_2$ ,  $C_3$  and  $C_4$  take different values are graphically illustrated in Figure 5. For example, the 4-tuple  $\text{DC}(\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}, \overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}})$  for the vectors  $\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}$  and  $\overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}}$ , shown in Figure 4, is  $(+ \ - \ - \ -)$ .




Figure 5: The quadrants and half lines where  $C_1$ ,  $C_2$ ,  $C_3$  and  $C_4$  take different values.

For example, the entries of the double-cross matrix of the polylines  $P_1$  and  $P_2$  of Figure 1 are given in Table 1. Polyline, such as  $P_1$  and  $P_2$  of Figure 1, that have the same double-cross matrix, are called *double-cross similar*.

This first example can be used to illustrate some properties of this matrix (Kuijpers et al. (2006)). First, we observe that on the diagonal always  $(0 \ 0 \ 0 \ 0)$  appears. We also see that there is a certain degree of *symmetry along the diagonal*. If  $\text{DC}(\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}, \overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}}) = (C_1 \ C_2 \ C_3 \ C_4)$ , then we have  $\text{DC}(\overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}}, \overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}) = (C_2 \ C_1 \ C_4 \ C_3)$ . These two observations imply that it suffices to know a double-cross matrix *above its diagonal*.

Input matrices, that do not possess these symmetry properties, are therefore, *a priori*, *not* realisable.

	$\overrightarrow{p_0 p_1}$	$\overrightarrow{p_1 p_2}$	$\overrightarrow{p_2 p_3}$	$\overrightarrow{p_3 p_4}$
$\overrightarrow{p_0 p_1}$	(0 0 0 0)	(- + 0 +)	(- + + +)	(- + + +)
$\overrightarrow{p_1 p_2}$	(+ - + 0)	(0 0 0 0)	(- + 0 +)	(- + + +)
$\overrightarrow{p_2 p_3}$	(+ - + +)	(+ - + 0)	(0 0 0 0)	(- + 0 +)
$\overrightarrow{p_3 p_4}$	(+ - + +)	(+ - + +)	(+ - + 0)	(0 0 0 0)

Table 1: The entries of the double-cross matrix of the polylines  $P_1$  and  $P_2$  of Figure 1.

### 2.3. An algebraic interpretation of the double-cross matrix

In this section, we give an algebraic interpretation of the double-cross matrix. In the following theorem, taken from (Kuijpers et al. (2006)), we use the function

$$\text{sign} : \mathbf{R} \rightarrow \{-, 0, +\} : x \mapsto \text{sign}(x) = \begin{cases} - & \text{if } x < 0; \\ 0 & \text{if } x = 0; \text{ and} \\ + & \text{if } x > 0. \end{cases}$$

**Theorem 1.** *Let  $P = \langle (x_0, y_0), (x_1, y_1), \dots, (x_N, y_N) \rangle$  be a polyline and let  $p_i = (x_i, y_i)$ , for  $0 \leq i \leq N$ . Then,  $\text{DC}(\overrightarrow{p_i p_{i+1}}, \overrightarrow{p_j p_{j+1}}) = (C_1 \ C_2 \ C_3 \ C_4)$  with*

$$\begin{aligned} C_1 &= - \text{sign}((x_j - x_i) \cdot (x_{i+1} - x_i) + (y_j - y_i) \cdot (y_{i+1} - y_i)); \\ C_2 &= \text{sign}((x_j - x_i) \cdot (x_{j+1} - x_j) + (y_j - y_i) \cdot (y_{j+1} - y_j)); \\ C_3 &= - \text{sign}((x_j - x_i) \cdot (y_{i+1} - y_i) - (y_j - y_i) \cdot (x_{i+1} - x_i)); \text{ and} \\ C_4 &= \text{sign}((x_j - x_i) \cdot (y_{j+1} - y_j) - (y_j - y_i) \cdot (x_{j+1} - x_j)). \end{aligned}$$

□

## 3. Problem statement and discussion

In this section, we state the decision problem, already given in the Introduction, more formally and we devote some theoretical discussion to it.

### 3.1. Problem statement

In this papers, we address the following decision problem (relative to some class  $\mathcal{P}$  of polylines in the plane  $\mathbf{R}^2$ ).

**Problem 1 (Realisability).** Given is an  $N \times N$  matrix  $M$  of 4-tuples

$$(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4.$$

- (a) Decide whether  $M$  is the double-cross matrix of some polyline (from a class  $\mathcal{P}$ ) in  $\mathbf{R}^2$  of size  $N$ ; and
- (b) If the answer to question (a) is *yes*, then produce an example of a polyline  $P$  with  $\text{DCM}(P) = M$ .  $\square$

Initially, we take the class of polylines  $\mathcal{P}$  as broad as possible. For instance, it is sufficient to look at polylines that have, as first line segment, the unit interval on the  $x$ -axis of  $\mathbf{R}^2$  and whose vertices have algebraic coordinates.

### 3.2. Discussion

By Theorem 1, the entries of an input matrix  $M$  to Problem 1 can be translated into sign conditions on quadratic polynomial equalities and inequalities. Therefore, Problem 1 is equivalent to deciding the first-order sentence

$$\exists x_0 \exists y_0 \exists x_1 \exists y_1 \cdots \exists x_N \exists y_N \bigwedge_{0 \leq i < j \leq N} \begin{cases} (x_j - x_i) \cdot (x_{i+1} - x_i) + (y_j - y_i) \cdot (y_{i+1} - y_i) & \alpha_{ij} & 0 \\ (x_j - x_i) \cdot (x_{j+1} - x_j) + (y_j - y_i) \cdot (y_{j+1} - y_j) & \beta_{ij} & 0 \\ (x_j - x_i) \cdot (y_{i+1} - y_i) - (y_j - y_i) \cdot (x_{i+1} - x_i) & \gamma_{ij} & 0 \\ (x_j - x_i) \cdot (y_{j+1} - y_j) - (y_j - y_i) \cdot (x_{j+1} - x_j) & \delta_{ij} & 0 \end{cases}$$

where  $\alpha_{ij}, \beta_{ij}, \gamma_{ij}, \delta_{ij} \in \{=, <, >\}$ , for  $0 \leq i < j \leq N$  are determined by the input matrix  $M$ , over the reals. The minus signs before the equations for  $C_1$  and  $C_3$  are assumed to be incorporated in the  $\alpha_{ij}$  and  $\gamma_{ij}$ . We remark that the above sentence expresses the entries of the input matrix strictly above its diagonal (as we can apriori discard non-symmetric input matrices).

The  $4 \frac{N(N-1)}{2}$  equalities and inequalities describe a semi-algebraic subsets of  $\mathbf{R}^{2(N+1)}$  (Bochnak et al. (1998)). We make the following observations about this system:

- there are  $2N(N-1)$  (in)equalities in  $2(N+1)$  variables  $x_0, y_0, \dots, x_N, y_N$ ;
- each polynomial uses 6 variables from  $x_0, y_0, x_1, y_1, \dots, x_N, y_N$  and has at most 8 monomial terms;
- each of the polynomials is homogeneous of degree 2;
- all the coefficients of the polynomials are 0, 1 or  $-1$ .

The first-order theory of the real ordered field is decidable (Tarski (1951)) and various implementations of decision procedures, that are based on Cylindrical Algebraic Decomposition (Collins (1975)) or other techniques, for this theory exist. We refer to QEPCAD (Hong (2000)), REDLOG (Dolzmann and Sturm (1997)) and MATHEMATICA (Wolfram Research (2015)) as a few examples. This type of software could be used, in theory, to answer Problem 1 (a) in practice. If there is a solution, these implementations also provide, as a byproduct of the above decision problem, sample points, thus, also, effectively answering question Problem 1 (b). But it is also known that the above mentioned implementations are slow and fail in practice to produce answers as soon as the number of variables increases. This is due to the intrinsic high time complexity of quantifier elimination in the ordered field of the reals (Heintz et al. (2013)). The theory of computational algebraic geometry gives an upper complexity bound. In particular, Theorem 13.13 in (Basu et al. (2006)) gives an upper bound on determining realisable sign conditions of a collection of polynomials. When applied to our decision problem, we obtain that there exists an algorithm to compute the set of *all* realisable sign conditions of the above system of polynomial (in)equalities with complexity  $(2N(N-1))^{2N+3} \cdot 2^{O(N)}$ . The complexity of deciding the satisfiability of the system is the same, as well as that of generating a sample point in case of non-emptiness. The use of alternative data structures to codify the polynomials can improve the time complexity, but not below exponential time (Giusti and Heintz (2001)). For a more recent discussion on lower bounds of the complexity, we refer to (Heintz et al. (2013)). The general problem of deciding an existential sentence in the first-order theory of the reals is known to be **NP**-hard (and to be in **PSPACE**) (Canny (1988)).

However, we have the following, negative result: Problem 1 (a) is **NP**-hard (Scivos and Nebel (2001); Renz and Nebel (2007)). Whether or not this problem is in **NP** is less obvious. It is known that if there is a solution to the above system of polynomial (in)equalities, there is also an solution with algebraic coordinates (Basu et al. (2006)). We could, for instance, try to guess the coordinates of the vertices of a polyline and then verify whether it satisfies the above system. Guessing algebraic coordinates could be implemented by guessing a polynomial and a root of this polynomial. However, an apriori polynomial bound on the complexity of sample points (to be guessed) is not obvious (Basu et al. (2006)). Above, we have observed that each polynomial uses at most 6 variables from  $x_0, y_0, x_1, y_1, \dots, x_N, y_N$  and has at most 8 monomial terms. This implies our problem is part of the field of

“fewnomials” (Khovanskii (1991)), where problems are notoriously difficult. And our problem and the production of sample points, is not covered by the known solutions there.

On the positive side, we can remark that, from the definition of the double-cross matrix in Section 2.2, it is clear that translations, rotations and scalings of a polyline do not change its double-cross matrix. Double-cross matrices are, in fact, invariant under *similarities* of  $\mathbf{R}^2$ . Thus, we can conclude, that if Problem 1 (a) has a positive answer, we can always find a polyline, to witness this fact, that starts of with the vertices  $(\mathbf{x}_0, \mathbf{y}_0) = (0, 0)$  and  $(\mathbf{x}_1, \mathbf{y}_1) = (1, 0)$  and in which the other vertices have coordinates that are algebraic numbers.

#### 4. A realisability test for $90^\circ$ -polylines

In this section, we give an efficient solution for a special case of Problem 1, for  $\mathcal{P} = \mathcal{P}_{90^\circ}$ , the class of  $90^\circ$ -polylines (again with vertices with algebraic coordinates). As we have remarked, for the problem of realisability, we may assume, without loss of any generality, that the polyline that realises a matrix  $M$ , if it exists, starts with the unit interval on the  $x$ -axis, that is,  $\mathbf{p}_0 = (\mathbf{x}_0, \mathbf{y}_0) = (0, 0)$  and  $\mathbf{p}_1 = (\mathbf{x}_1, \mathbf{y}_1) = (1, 0)$ .

The following, straightforward, property gives a first necessary condition for the input to our decision problem, the matrix  $M$ .

**Property 1.** *Let  $P = \langle \mathbf{p}_0, \mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_N \rangle$  be a polyline. A necessary and sufficient condition for  $P$  to be a  $90^\circ$ -polyline is that for all  $i$ ,  $0 \leq i < N - 1$ ,  $\text{DC}(\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}, \overrightarrow{\mathbf{p}_{i+1} \mathbf{p}_{i+2}}) =$*

- $(- \ - \ 0 \ 0)$  (*reverse turn*);
- $(- \ 0 \ 0 \ -)$  (*right turn*);
- $(- \ + \ 0 \ 0)$  (*straight*); or
- $(- \ 0 \ 0 \ +)$  (*left turn*). □

Since we take  $\mathbf{p}_0 = (\mathbf{x}_0, \mathbf{y}_0) = (0, 0)$  and  $\mathbf{p}_1 = (\mathbf{x}_1, \mathbf{y}_1) = (1, 0)$ , all line segments of the polyline, realising  $M$ , should be parallel to one of the coordinate axis of  $\mathbf{R}^2$ . In fact, we have for each  $i$ ,  $0 \leq i < N$  that  $\mathbf{x}_i = \mathbf{x}_{i+1} \wedge (\mathbf{y}_i < \mathbf{y}_{i+1} \vee \mathbf{y}_i > \mathbf{y}_{i+1})$  or  $\mathbf{y}_i = \mathbf{y}_{i+1} \wedge (\mathbf{x}_i < \mathbf{x}_{i+1} \vee \mathbf{x}_i > \mathbf{x}_{i+1})$ . Here,

for  $0 \leq i < N$ , we are in exactly one of the following four situations (always with  $\ell_{i+1} > 0$ ):

$$\left\{ \begin{array}{l} x_{i+1} = x_i + \ell_{i+1} \\ y_{i+1} = y_i; \end{array} \right. \quad \left\{ \begin{array}{l} x_{i+1} = x_i - \ell_{i+1} \\ y_{i+1} = y_i; \end{array} \right. \quad \left\{ \begin{array}{l} x_{i+1} = x_i \\ y_{i+1} = y_i + \ell_{i+1}; \end{array} \right. \quad \left\{ \begin{array}{l} x_{i+1} = x_i \\ y_{i+1} = y_i - \ell_{i+1}. \end{array} \right.$$

Before we give an efficient solution to Problem 1 for  $\mathcal{P} = \mathcal{P}_{90^\circ}$ , we prove a lemma that explains in which quadrant, determined by a line segment of a polyline, that is parallel to one of the coordinate axis of  $\mathbf{R}^2$ , a vertex of a polyline is located. For clarity, we state and prove the lemma for segments of a polyline, that are parallel to one of the coordinate axis of  $\mathbf{R}^2$  and that coincide with the unit interval on the  $x$ - and  $y$ -axis (or their negatives), but the lemma can be easily extended and applied to any polyline segments that are parallel to one of the coordinate axis of  $\mathbf{R}^2$ , after applying a scaling and translation of  $\mathbf{R}^2$ .

**Lemma 1.** *Let  $P = \langle \mathbf{p}_0, \mathbf{p}_1, \dots, \mathbf{p}_N \rangle$  be a polyline and assume that  $\mathbf{p}_i = (0, 0)$  and  $\mathbf{p}_{i+1} = (\pm 1, 0)$  or  $\mathbf{p}_i = (0, 0)$  and  $\mathbf{p}_{i+1} = (0, \pm 1)$ . Let  $\mathbf{p}_j = (x_j, y_j)$ , for  $0 \leq i \leq N$  and  $i + 1 < j$ . From the first and third component of  $\text{DC}(\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}, \overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}})$ , we can determine  $\text{sign}(x_j)$  and  $\text{sign}(y_j)$ .*

**Proof.** First, let  $\mathbf{p}_i = (0, 0)$  and  $\mathbf{p}_{i+1} = (\pm 1, 0)$ . From Theorem 1 it is clear that  $C_1 = -\text{sign}((x_j - 0) \cdot \pm 1 + y_j \cdot 0) = -\text{sign}(\pm x_j)$  and that  $C_3 = -\text{sign}(x_j \cdot 0 - y_j \cdot \pm 1) = \text{sign}(\pm y_j)$ .

Secondly, let  $\mathbf{p}_i = (0, 0)$  and  $\mathbf{p}_{i+1} = (0, \pm 1)$ . Similarly, Theorem 1 implies  $C_1 = -\text{sign}((x_j - 0) \cdot 0 + y_j \cdot \pm 1) = -\text{sign}(\pm y_j)$  and that  $C_3 = -\text{sign}(x_j \cdot \pm 1 - y_j \cdot 0) = -\text{sign}(\pm x_j)$ .  $\square$

**Theorem 2.** *It can be decided in time  $O(N^2)$  whether a  $N \times N$  matrix  $M$  of 4-tuples  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$  is the double-cross matrix of some  $90^\circ$ -polyline in  $\mathbf{R}^2$  of size  $N$ . If  $M$  this is the case, also witnesses to this can be produced in time  $O(N^2)$ .*

**Proof.** We now describe a decision procedure for Problem 1: in a first step, we determine the relationship ( $<$ ,  $=$ ,  $>$ ) between coordinates of consecutive vertices. In a second step, we do it for all remaining vertices.

Let  $M$  be a  $N \times N$  matrix of 4-tuples  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$ . Since we want to determine the existence of a  $90^\circ$ -polyline  $P = \langle \mathbf{p}_0, \mathbf{p}_1, \dots, \mathbf{p}_N \rangle$ , with  $\mathbf{p}_0 = (0, 0)$  and  $\mathbf{p}_1 = (1, 0)$ , that realises  $M$ , we know that all the segments of  $P$  are parallel to one of the coordinate axis of  $\mathbf{R}^2$ . in the following, we assume that  $\mathbf{p}_i = (x_i, y_i)$ , for  $0 \leq i \leq N$ .

As an apriori step, we check whether  $M$  does not have  $(0 \ 0 \ 0 \ 0)$  entries on its diagonal or doesn't have the "symmetry" properties, discussed in Section 2.2. If  $M$  fails this symmetry-test, we can already answer *no*, else we proceed.

**Step 1.** First, we inspect all entries  $M[i, i + 1]$ ,  $0 \leq i < N$  of  $M$ . They should all be of the form

- $(- \ - \ 0 \ 0)$  (reverse turn);
- $(- \ 0 \ 0 \ -)$  (right turn);
- $(- \ + \ 0 \ 0)$  (straight); or
- $(- \ 0 \ 0 \ +)$  (left turn).

If this is not the case, we can already answer *no*. In the other case, we deduce, from the entries of  $M$  above the diagonal, the arrangement<sup>5</sup> of the consecutive  $x$ -coordinates  $x_i$  and  $x_{i+1}$  and of the consecutive  $y$ -coordinates  $y_i$  and  $y_{i+1}$  of candidate vertices of a polyline. Then we proceed to Step 2.

**Step 2.** Now, we inspect all entries  $M[i, j]$ ,  $1 \leq i + 1 < j < N$  of  $M$ . Per entry, two cases, corresponding to the line segment that connects  $\mathbf{p}_i$  and  $\mathbf{p}_{i+1}$  being *parallel to the  $y$ -axis* or *parallel to the  $x$ -axis*, have to be considered.

*Case 1 (parallel to the  $y$ -axis or  $x_i = x_{i+1}$ ):* For  $y_i$  and  $y_{i+1}$ , we have only two possible arrangements:  $y_i < y_{i+1}$  or  $y_i > y_{i+1}$ . We use the "parallel to the  $y$ -axis" version of Lemma 1, to determine, from the matrix entry  $M[i, j]$ , the quadrant in which  $(x_j, y_j)$  is located compared to the line segment that connects  $\mathbf{p}_i$  and  $\mathbf{p}_{i+1}$ . This gives us the the arrangement of  $x_i$  and  $x_j$  on the one hand and of  $y_i$  and  $y_j$  on the other hand.

*Case 2 (parallel to the  $x$ -axis or  $y_i = y_{i+1}$ ):* This case is analogous to Case 1, but now we use the "parallel to the  $x$ -axis" version of version of Lemma 1.

---

<sup>5</sup>By *arrangement*, we mean which of the cases  $x_i < x_{i+1}$ ,  $x_i = x_{i+1}$  and  $x_i > x_{i+1}$  holds.



At this point, we have now complete information on how the  $x$ -coordinate values  $x_0, x_1, \dots, x_N$  are pairwise arranged (or ordered) and how the  $y$ -coordinate values  $y_0, y_1, \dots, y_N$  are pairwise arranged. We can store this arrangement information in two matrices (similarly to the double-cross matrix). The first matrix can be used to verify whether an ordering of  $x_0, x_1, \dots, x_N$  is possible. To this purpose, we scan this matrix column per column. The first column will allow us to place  $x_0$  and  $x_1$  on the real line (according to their arrangement). This results in at most five locations to place  $x_2$  (before; between; after; or on  $x_0$  and  $x_1$ ). The second column of the matrix tells us where. We repeat this process until all the candidate values  $x_0, x_1, \dots, x_N$  are placed on the real line. Next, we use the second matrix to place the  $y$ -coordinate values  $y_0, y_1, \dots, y_N$  on the  $y$ -axis. If, in this process, we find it impossible to find a location to place one of the  $x_i$  or  $y_i$  (due to a contradiction), we answer *no*. If we have never found a contradiction and all  $x$ - and  $y$ -values can be ordered, we are ready to answer *yes*. This ordering process takes  $O(N^2)$  time.

If we have found  $k_x$  different values  $x_0, x_1, \dots, x_N$  and  $k_y$  different values  $y_0, y_1, \dots, y_N$ , we can draw an example of a polyline that realises  $M$  on the grid  $\{0, 1, \dots, k_x - 1\} \times \mathbf{R} \cup \mathbf{R} \times \{0, 1, \dots, k_y - 1\}$ , with vertices belonging to  $\{0, 1, \dots, k_x - 1\} \times \{0, 1, \dots, k_y - 1\}$ . This drawing serves as a sample point and answers Problem 1 (b).

It is clear that the above inspection of the matrix  $M$  takes  $O(N^2)$  time. The reconstruction of a polyline can be done in the same amount of time. This completes the proof.  $\square$

## 5. The polar coordinate representation of a polyline

In this section, we define the polar coordinate representation of a polyline and we describe how to go from the Cartesian coordinate representation to the polar coordinate representation and vice versa.

**Definition 4.** Let  $P = \langle (x_0, y_0), (x_1, y_1), \dots, (x_N, y_N) \rangle$  be a polyline (in Cartesian coordinate representation) and let  $\mathbf{p}_i = (x_i, y_i)$ ,  $0 \leq i \leq N$ . The *polar coordinate representation* of the polyline  $P$  is the list

$$\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle,$$

where  $\ell_i$  is the length of the line segment  $\mathbf{p}_{i-1}\mathbf{p}_i$  and  $\theta_i$  is the counter-clockwise angel at  $\mathbf{p}_i$  between the line connecting  $\mathbf{p}_i$  and  $\mathbf{p}_{i-1}$  and the line connecting  $\mathbf{p}_i$  and  $\mathbf{p}_{i+1}$ .  $\square$

If at  $\mathbf{p}_i$ , the polyline turns to the left or goes straight,  $\theta_i = 180^\circ - \angle(\overrightarrow{\mathbf{p}_i\mathbf{p}_{i-1}}, \overrightarrow{\mathbf{p}_i\mathbf{p}_{i+1}})$  and if at  $\mathbf{p}_i$ , the polyline turns to the right or returns,  $\theta_i = 180^\circ + \angle(\overrightarrow{\mathbf{p}_i\mathbf{p}_{i-1}}, \overrightarrow{\mathbf{p}_i\mathbf{p}_{i+1}})$ .

So,  $\theta_i$  captures the (counter-clockwise) change in direction when going from the line segment  $\mathbf{p}_{i-1}\mathbf{p}_i$  to the line segment  $\mathbf{p}_i\mathbf{p}_{i+1}$ . This is illustrated in Figure 6.




Figure 6: The polar coordinates  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \ell_3, \theta_3, \ell_4, \theta_4, \ell_5 \rangle$  (in red) of the polyline  $\langle \mathbf{p}_0, \mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4, \mathbf{p}_5 \rangle$  (in black).

### 5.1. From the Cartesian coordinate to the polar coordinate representation

To convert a polyline  $P = \langle (x_0, y_0), (x_1, y_1), \dots, (x_N, y_N) \rangle$  given by the Cartesian coordinates of its vertices to polar coordinate representation is easy. For  $\ell_i$ , we take the length of the line segment  $\mathbf{p}_{i-1}\mathbf{p}_i$ . By definition  $\theta_i = 180^\circ - \angle(\overrightarrow{\mathbf{p}_i\mathbf{p}_{i-1}}, \overrightarrow{\mathbf{p}_i\mathbf{p}_{i+1}})$  if the polyline turns to the left or goes straight and  $\theta_i = 180^\circ + \angle(\overrightarrow{\mathbf{p}_i\mathbf{p}_{i-1}}, \overrightarrow{\mathbf{p}_i\mathbf{p}_{i+1}})$  if the polyline turns to the right or returns. Therefore, the angle  $\theta_i$  is given by the formula

$$\pi - \arccos\left(\frac{(x_{i-1} - x_i, y_{i-1} - y_i) \cdot (x_{i+1} - x_i, y_{i+1} - y_i)}{|(x_{i-1} - x_i, y_{i-1} - y_i)| \cdot |(x_{i+1} - x_i, y_{i+1} - y_i)|}\right)$$

if the polyline turns to the left or goes straight, and by

$$\pi + \arccos\left(\frac{(x_{i-1} - x_i, y_{i-1} - y_i) \cdot (x_{i+1} - x_i, y_{i+1} - y_i)}{|(x_{i-1} - x_i, y_{i-1} - y_i)| \cdot |(x_{i+1} - x_i, y_{i+1} - y_i)|}\right)$$

if the polyline turns to the right or returns.<sup>6</sup>

### 5.2. From the polar coordinate to the Cartesian coordinate representation

Now, we turn to transforming the polar coordinate representation into the classical Cartesian coordinate representation, which is more laborious. Here, we can use some techniques that are also known in the description of robot arms with multiple joints (see, for instance, Chapter 6 of (Cox et al. (1997))).

Hereto, we first need some technical results. Let  $P = \langle (x_0, y_0), (x_1, y_1), \dots, (x_N, y_N) \rangle$  be a polyline and let  $p_i = (x_i, y_i)$ ,  $0 \leq i \leq N$ . In each vertex  $(x_i, y_i)$ , we create a local coordinate system  $(X_i, Y_i)$ . The origin of this coordinate system is  $(x_i, y_i)$  and the positive  $X_i$ -axis is points from  $(x_i, y_i)$  to  $(x_{i+1}, y_{i+1})$ . The  $Y_i$ -axis is perpendicular to the  $X_i$ -axis in  $(x_i, y_i)$  in the usual way. This is illustrated in Figure 7.




Figure 7: The local coordinate systems  $(X_{i-1}, Y_{i-1})$  (in blue) and  $(X_i, Y_i)$  (in green) on the vertices  $p_{i-1}$  and  $p_i$  of a polyline.

The following property is well known from linear algebra and also from the field of multiple joint robot arms (see, Chapter 6, page 262, in Cox et al. (1997)).

---

<sup>6</sup>Here, the  $\cdot$  in the numerator denotes the *inner product* of two vectors and the  $\cdot$  in the denominator is the product of norms.

**Property 2.** Let  $\mathbf{p}_{i-1}$ ,  $\mathbf{p}_i$  and  $\mathbf{p}_{i+1}$  be three consecutive vertices on a polyline  $P = \langle (\mathbf{x}_0, \mathbf{y}_0), (\mathbf{x}_1, \mathbf{y}_1), \dots, (\mathbf{x}_N, \mathbf{y}_N) \rangle$  with  $\mathbf{p}_i = (\mathbf{x}_i, \mathbf{y}_i)$ ,  $0 \leq i \leq N$ . If a point  $\mathbf{q}$  in  $\mathbf{R}^2$  has coordinates  $(a_{i-1}, b_{i-1})$  and  $(a_i, b_i)$ , respectively, in the local coordinate systems  $(X_{i-1}, Y_{i-1})$  and  $(X_i, Y_i)$ , respectively, then

$$\begin{pmatrix} a_{i-1} \\ b_{i-1} \\ 1 \end{pmatrix} = \begin{pmatrix} \cos \theta_i & -\sin \theta_i & \ell_i \\ \sin \theta_i & \cos \theta_i & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} a_i \\ b_i \\ 1 \end{pmatrix}.$$

□

For a polyline  $P$ , given by its polar coordinate representation  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$ , we set

$$P_i = \begin{pmatrix} \cos \theta_i & -\sin \theta_i & \ell_i \\ \sin \theta_i & \cos \theta_i & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

From now on, we only consider polylines with  $(\mathbf{x}_0, \mathbf{y}_0) = (0, 0)$  and  $(\mathbf{x}_1, \mathbf{y}_1) = (1, 0)$ , such that  $(X_0, Y_0)$  is the standard coordinate system.

The following property, based on the previous property, has a straightforward induction proof.

**Property 3.** Let  $P = \langle (\mathbf{x}_0, \mathbf{y}_0), (\mathbf{x}_1, \mathbf{y}_1), \dots, (\mathbf{x}_N, \mathbf{y}_N) \rangle$  be a polyline. If a point  $\mathbf{q}$  in  $\mathbf{R}^2$  has coordinates  $(a_i, b_i)$  in the local coordinate system  $(X_i, Y_i)$ , then it has absolute Cartesian coordinates  $(a_0, b_0)$  in  $(X_0, Y_0)$ , with

$$\begin{pmatrix} a_0 \\ b_0 \\ 1 \end{pmatrix} = P_1 \cdot P_2 \cdots P_i \cdot \begin{pmatrix} a_i \\ b_i \\ 1 \end{pmatrix}.$$

□

The following property tells us what the matrix product  $P_1 \cdot P_2 \cdots P_i$  looks like.

**Property 4.** For  $1 \leq i < N$ , we have

$$P_1 \cdot P_2 \cdots P_i = \begin{pmatrix} \cos \Theta_1^i & -\sin \Theta_1^i & \sum_{j=1}^i \ell_j \cos \Theta_1^{j-1} \\ \sin \Theta_1^i & \cos \Theta_1^i & \sum_{j=1}^i \ell_j \sin \Theta_1^{j-1} \\ 0 & 0 & 1 \end{pmatrix},$$

where  $\Theta_i^j$  abbreviates  $\theta_i + \theta_{i+1} + \cdots + \theta_j$ , for  $i \leq j$ .

**Proof.** We proceed by induction on  $i$ . For  $i = 1$ , we have  $\ell_1 \cos 0 = \ell_1$  and  $\ell_1 \sin 0 = 0$ , which clearly gives  $P_1$ .

Now, we proceed from  $i$  to  $i+1$ . By the induction hypothesis,  $P_1 \cdot P_2 \cdots P_i \cdot P_{i+1}$  equals

$$\begin{pmatrix} \cos \Theta_1^i & -\sin \Theta_1^i & \sum_{j=1}^i \ell_j \cos \Theta_1^{j-1} \\ \sin \Theta_1^i & \cos \Theta_1^i & \sum_{j=1}^i \ell_j \sin \Theta_1^{j-1} \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \cos \theta_{i+1} & -\sin \theta_{i+1} & \ell_{i+1} \\ \sin \theta_{i+1} & \cos \theta_{i+1} & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

which is

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

with

- $a_{11} = \cos \Theta_1^i \cdot \cos \theta_{i+1} - \sin \Theta_1^i \cdot \sin \theta_{i+1} = \cos (\Theta_1^i + \theta_{i+1}) = \cos (\Theta_1^{i+1});$
- $a_{12} = -\cos \Theta_1^i \cdot \sin \theta_{i+1} - \sin \Theta_1^i \cdot \cos \theta_{i+1} = -\sin (\Theta_1^i + \theta_{i+1}) = -\sin (\Theta_1^{i+1});$
- $a_{13} = \ell_{i+1} \cos \Theta_1^i + \sum_{j=1}^i \ell_j \cos \Theta_1^{j-1} = \sum_{j=1}^{i+1} \ell_j \cos \Theta_1^{j-1};$
- $a_{21} = \sin \Theta_1^i \cdot \cos \theta_{i+1} + \cos \Theta_1^i \cdot \sin \theta_{i+1} = \sin (\Theta_1^i + \theta_{i+1}) = \sin (\Theta_1^{i+1});$
- $a_{22} = -\sin \Theta_1^i \cdot \sin \theta_{i+1} + \cos \Theta_1^i \cdot \cos \theta_{i+1} = \cos (\Theta_1^i + \theta_{i+1}) = \cos (\Theta_1^{i+1});$
- $a_{23} = \ell_{i+1} \sin \Theta_1^i + \sum_{j=1}^i \ell_j \sin \Theta_1^{j-1} = \sum_{j=1}^{i+1} \ell_j \sin \Theta_1^{j-1};$
- $a_{31} = 0 + 0 + 0 = 0;$
- $a_{32} = 0 + 0 + 0 = 0;$  and
- $a_{33} = 0 + 0 + 1 = 1;$

where we have used the well-known formulas for cosinus and sinus of the sum of angles. This gives the desired matrix and concludes the proof.  $\square$

The following theorem tells us how to translate from polar coordinates to Cartesian coordinates.

**Theorem 3.** Let  $P = \langle (x_0, y_0), (x_1, y_1), \dots, (x_N, y_N) \rangle$  be a polyline that is given by its polar coordinate representation  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$ . If we assume that  $(x_0, y_0) = (0, 0)$  and  $(x_1, y_1) = (1, 0)$ , then

$$\begin{cases} x_i &= \sum_{j=1}^i \ell_j \cos(\theta_1 + \dots + \theta_{j-1}) \\ y_i &= \sum_{j=1}^i \ell_j \sin(\theta_1 + \dots + \theta_{j-1}) \end{cases}$$

for  $2 \leq i \leq N$ .

We remark that we could also have written

$$\begin{cases} x_i &= 1 + \sum_{j=2}^i \ell_j \cos(\theta_1 + \dots + \theta_{j-1}) \\ y_i &= \sum_{j=2}^i \ell_j \sin(\theta_1 + \dots + \theta_{j-1}) \end{cases}$$

in the statement of this theorem, since  $\ell_1 = 1$ ,  $\cos 0 = 1$  and  $\sin 0 = 0$ . For esthetic reasons, we will stick to the earlier expressions.

**Proof.** In the local coordinate system  $(X_{i-1}, Y_{i-1})$ , the coordinates of  $\mathbf{p}_i = (x_i, y_i)$  are  $(\ell_i, 0)$ . By Property 3, the coordinates of  $\mathbf{p}_i$  in the standard coordinate system  $(X_0, Y_0)$  are given by

$$\begin{pmatrix} x_i \\ y_i \\ 1 \end{pmatrix} = P_1 \cdot P_2 \cdots P_{i-1} \cdot \begin{pmatrix} \ell_i \\ 0 \\ 1 \end{pmatrix}.$$

By Property 4, this means

$$\begin{pmatrix} x_i \\ y_i \\ 1 \end{pmatrix} = \begin{pmatrix} \cos \Theta_1^{i-1} & -\sin \Theta_1^{i-1} & \sum_{j=1}^{i-1} \ell_j \cos \Theta_1^{j-1} \\ \sin \Theta_1^{i-1} & \cos \Theta_1^{i-1} & \sum_{j=1}^{i-1} \ell_j \sin \Theta_1^{j-1} \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \ell_i \\ 0 \\ 1 \end{pmatrix}$$

or

$$\begin{cases} x_i &= \ell_i \cos \Theta_1^{i-1} + \sum_{j=1}^{i-1} \ell_j \cos \Theta_1^{j-1} = \sum_{j=1}^i \ell_j \cos \Theta_1^{j-1} \\ y_i &= \ell_i \sin \Theta_1^{i-1} + \sum_{j=1}^{i-1} \ell_j \sin \Theta_1^{j-1} = \sum_{j=1}^i \ell_j \sin \Theta_1^{j-1} \end{cases}$$

which concludes the proof.  $\square$

### 5.3. The double-cross conditions for polar coordinates

For a polyline  $P = \langle (x_0, y_0), (x_1, y_1), \dots, (x_N, y_N) \rangle$  with  $\mathbf{p}_i = (x_i, y_i)$ ,  $0 \leq i \leq N$ , Theorem 1, gives us sign conditions on polynomials for  $C_1, C_2, C_3$  and  $C_4$  in  $\text{DC}(\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}, \overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}}) = (C_1 \ C_2 \ C_3 \ C_4)$ . Now, if the polyline  $P = \langle (x_0, y_0), (x_1, y_1), \dots, (x_N, y_N) \rangle$  is given by its polar coordinate representation  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$ , Theorem 1 allows us to translate these conditions into polar coordinates.

Where needed, we use the abbreviations

$$\begin{cases} c_i &= \cos \Theta_1^i = \cos (\theta_1 + \dots + \theta_i); \\ s_i &= \sin \Theta_1^i = \sin (\theta_1 + \dots + \theta_i). \end{cases}$$

to control the length of the expressions.

The following theorem gives the double-cross conditions in polar form.

**Theorem 4.** *If now the polyline  $P = \langle \mathbf{p}_0, \mathbf{p}_1, \dots, \mathbf{p}_N \rangle$  is given by its polar coordinate representation  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$ , and if we assume that  $\mathbf{p}_0 = (0, 0)$  and  $\mathbf{p}_1 = (1, 0)$ , then  $\text{DC}(\overrightarrow{\mathbf{p}_i \mathbf{p}_{i+1}}, \overrightarrow{\mathbf{p}_j \mathbf{p}_{j+1}}) = (C_1 \ C_2 \ C_3 \ C_4)$ , for  $0 \leq i < j < N$ , are expressed in polar coordinates as follows:*

$$\begin{aligned} C_1 &= - \text{sign}(\sum_{k=i+1}^j \ell_k \cos (\theta_{i+1} + \dots + \theta_{k-1})); \\ C_2 &= \text{sign}(\sum_{k=i+1}^j \ell_k \cos (\theta_k + \dots + \theta_j)); \\ C_3 &= - \text{sign}(\sum_{k=i+1}^j \ell_k \sin (\theta_{i+1} + \dots + \theta_{k-1})); \text{ and} \\ C_4 &= \text{sign}(\sum_{k=i+1}^j \ell_k \sin (\theta_k + \dots + \theta_j)), \end{aligned}$$

where we agree that the empty sum of angles equals 0.

**Proof.** Let  $P$  be as in the statement of the theorem. From Theorem 3, we get

$$\begin{cases} x_i &= \sum_{k=1}^i \ell_k \cos (\theta_1 + \dots + \theta_{k-1}) \\ y_i &= \sum_{k=1}^i \ell_k \sin (\theta_1 + \dots + \theta_{k-1}), \end{cases}$$

for  $0 \leq i \leq N$ . So, we obtain, for  $0 \leq i < j < N$ ,

$$\left\{ \begin{array}{lcl} x_j - x_i & = & \sum_{k=1}^j \ell_k \cos(\theta_1 + \dots + \theta_{k-1}) - \sum_{k=1}^i \ell_k \cos(\theta_1 + \dots + \theta_{k-1}) \\ & = & \sum_{k=i+1}^j \ell_k \cos(\theta_1 + \dots + \theta_{k-1}) \\ y_j - y_i & = & \sum_{k=1}^j \ell_k \sin(\theta_1 + \dots + \theta_{k-1}) - \sum_{k=1}^i \ell_k \sin(\theta_1 + \dots + \theta_{k-1}) \\ & = & \sum_{k=i+1}^j \ell_k \sin(\theta_1 + \dots + \theta_{k-1}) \\ x_{i+1} - x_i & = & \sum_{k=1}^{i+1} \ell_k \cos(\theta_1 + \dots + \theta_{k-1}) - \sum_{k=1}^i \ell_k \cos(\theta_1 + \dots + \theta_{k-1}) \\ & = & \ell_{i+1} \cos(\theta_1 + \dots + \theta_i) \\ y_{i+1} - y_i & = & \sum_{k=1}^{i+1} \ell_k \sin(\theta_1 + \dots + \theta_{k-1}) - \sum_{k=1}^i \ell_k \sin(\theta_1 + \dots + \theta_{k-1}) \\ & = & \ell_{i+1} \sin(\theta_1 + \dots + \theta_i) \\ x_{j+1} - x_j & = & \sum_{k=1}^{j+1} \ell_k \cos(\theta_1 + \dots + \theta_{k-1}) - \sum_{k=1}^j \ell_k \cos(\theta_1 + \dots + \theta_{k-1}) \\ & = & \ell_{j+1} \cos(\theta_1 + \dots + \theta_j) \\ y_{j+1} - y_j & = & \sum_{k=1}^{j+1} \ell_k \sin(\theta_1 + \dots + \theta_{k-1}) - \sum_{k=1}^j \ell_k \sin(\theta_1 + \dots + \theta_{k-1}) \\ & = & \ell_{j+1} \sin(\theta_1 + \dots + \theta_j). \end{array} \right.$$

If we plug these identities in the equations of Theorem 1, we get

$$\begin{aligned} C_1 &= - \operatorname{sign}(\sum_{k=i+1}^j \ell_k (\mathbf{c}_i \mathbf{c}_{k-1} + \mathbf{s}_i \mathbf{s}_{k-1})) \\ &= - \operatorname{sign}(\sum_{k=i+1}^j \ell_k \cos(\theta_{i+1} + \dots + \theta_{k-1})); \\ C_2 &= \operatorname{sign}(\sum_{k=i+1}^j \ell_k (\mathbf{c}_j \mathbf{c}_{k-1} + \mathbf{s}_j \mathbf{s}_{k-1})) \\ &= \operatorname{sign}(\sum_{k=i+1}^j \ell_k \cos(\theta_k + \dots + \theta_j)); \\ C_3 &= - \operatorname{sign}(\sum_{k=i+1}^j \ell_k (\mathbf{s}_i \mathbf{c}_{k-1} - \mathbf{c}_i \mathbf{s}_{k-1})) \\ &= - \operatorname{sign}(\sum_{k=i+1}^j \ell_k \sin(\theta_{i+1} + \dots + \theta_{k-1})); \text{ and} \\ C_4 &= \operatorname{sign}(\sum_{k=i+1}^j \ell_k (\mathbf{s}_j \mathbf{c}_{k-1} - \mathbf{c}_j \mathbf{s}_{k-1})) \\ &= \operatorname{sign}(\sum_{k=i+1}^j \ell_k \sin(\theta_k + \dots + \theta_j)). \end{aligned}$$

In the last equalities we used the well-known formulas  $\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$  and  $\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$ . This concludes the proof.  $\square$

We remark that all the double-cross conditions in the above theorem are *linear* expressions in the lengths  $\ell_1, \dots, \ell_{N-1}$ . We also remark that an alternative way to write these conditions is

$$\begin{aligned} C_1 &= - \operatorname{sign}(\ell_{i+1} + \sum_{k=i+2}^j \ell_k \cos(\theta_{i+1} + \dots + \theta_{k-1})); \\ C_2 &= \operatorname{sign}(\sum_{k=i+1}^j \ell_k \cos(\theta_k + \dots + \theta_j)); \\ C_3 &= - \operatorname{sign}(\sum_{k=i+2}^j \ell_k \sin(\theta_{i+1} + \dots + \theta_{k-1})); \text{ and} \\ C_4 &= \operatorname{sign}(\sum_{k=i+1}^j \ell_k \sin(\theta_k + \dots + \theta_j)). \end{aligned}$$



We end this section with a remark about the double cross entries for consecutive line segments.

Because of the special location of  $(x_0, y_0) = (0, 0)$  and  $(x_1, y_1) = (1, 0)$ , we look at a special case of this theorem, namely  $i = 0$  and  $j = 1$ . Here, we have  $DC(\overrightarrow{p_0 p_1}, \overrightarrow{p_1 p_2}) = (C_1 \ C_2 \ C_3 \ C_4)$ , with

$$\begin{aligned} C_1 &= - \quad \text{sign}(1) = -; \\ C_2 &= \quad \text{sign}(\ell_1 + \ell_2 c_1 - 1) = \text{sign}(\ell_2 c_1); \\ C_3 &= - \quad \text{sign}(0) = 0; \quad \text{and} \\ C_4 &= \quad \text{sign}(\ell_2 s_1); \end{aligned}$$

Because, by the assumption in Definition 1, two consecutive vertices in a polyline are never identical, we have  $\ell_2 > 0$ , we can simplify conditions  $C_2$  and  $C_4$  and we get

$$\begin{aligned} C_1 &= - \\ C_2 &= \text{sign}(\cos \theta_1) \\ C_3 &= 0 \quad \text{and} \\ C_4 &= \text{sign}(\sin \theta_1). \end{aligned}$$

More generally, we look at the following special case of consecutive line segments of a polyline.

**Corollary 1.** *If now the polyline  $P = \langle p_0, p_1, \dots, p_N \rangle$  is given by its polar coordinate representation  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$ , and if we assume that  $p_0 = (0, 0)$  and  $p_1 = (1, 0)$ , then  $DC(\overrightarrow{p_i p_{i+1}}, \overrightarrow{p_{i+1} p_{i+2}}) = (C_1 \ C_2 \ C_3 \ C_4)$ , for  $0 \leq i < N - 1$ , are expressed in polar coordinates as follows:*

$$\begin{aligned} C_1 &= -; \\ C_2 &= \text{sign}(\cos \theta_{i+1}); \\ C_3 &= 0; \\ C_4 &= \text{sign}(\sin \theta_{i+1}). \end{aligned}$$

□

## 6. A polynomial-time realisability test for the case where the successive angles of a polyline are given

In this section, we use the double-cross conditions in polar form, from Theorem 4, to show the existence of a polynomial-time realisability test in the

case where, apart from an  $N \times N$  input matrix  $M$ , also the successive angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$  of a polyline are given as input. In this setting, the problem is to decide whether  $M$  is the double-cross matrix of some polyline of size  $N$  in  $\mathbf{R}^2$  with polar representation  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$ . Essentially, this decision problem asks whether there exist lengths  $\ell_1, \ell_2, \dots, \ell_{N-1}, \ell_N > 0$  that satisfy the conditions expressed by the matrix  $M$ . For the problem of realisability, here again, we may assume, without loss of generality, that a polyline that realises a matrix  $M$ , if it exists, starts with the unit interval on the  $x$ -axis. This permits us to use the results on the polar representation from Section 5.

However, since the given angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$  are part of the input, we need to impose some restrictions concerning their finite representability. We also need to impose some further technical conditions on these angles and on their finite sums, that appear in Theorem 4. The role of these technical restrictions will become clear in the proof of Theorem 5. To capture these restrictions, we define the notion of a “degree-bounded set of angles.” In the following,  $\mathbf{Q}$  denotes the set of the rational numbers. A set  $\Theta$  of angles is *degree-bounded* if

1.  $\Theta$  is closed under addition;
2. for any  $\theta \in \Theta$ ,  $\sin \theta$  (and thus,  $\cos \theta$ ) is an algebraic number;<sup>7</sup>
3. there exists a natural number  $d$  such that for any finite subset  $\{\theta_1, \dots, \theta_k\}$  of  $\Theta$ , the field extension  $\mathbf{Q}(\sin \theta_1, \dots, \sin \theta_k, \cos \theta_1, \dots, \cos \theta_k)$  has a degree less or equal to  $d$  over  $\mathbf{Q}$ ; and
4. there exists a natural number  $E$  such that any  $\theta \in \Theta$  has an encoding of size less or equal to  $E$ .<sup>8</sup>

---

<sup>7</sup>A real number  $a$  is called *algebraic* if there exists a polynomial  $F$  with rational coefficients such that  $F(a) = 0$ . In this case,  $a$  is also the root of a unique, monic polynomial over the rationals, called the *minimal polynomial of  $a$*  and the *degree of  $a$*  is defined to be the degree of its minimal polynomial (Lang (1986)).

<sup>8</sup>An real algebraic number  $a$  can be encoded by a triple  $(F, q_1, q_2)$ , where  $F$  is the minimal polynomial of  $a$  and  $q_1$  and  $q_2$  are rational numbers such that  $a$  is the unique root of  $F$  in the interval  $[q_1, q_2]$ . Since  $q_1$  and  $q_2$  can be determined within a time complexity that is polynomial in the bit size of the coefficients of  $F$ , the total bit size of the coefficients of  $F$  can then be taken as a measure for the encoding size of  $a$ . For further details, we refer to (Adler and Beling (1994)).

For a degree-bounded set of angles  $\Theta$ , we refer to the above  $d$  and  $E$  as the *extension degree bound* of  $\Theta$  and the *encoding bound* of  $\Theta$ , respectively.

Now, we give examples of degree-bounded sets of angles, that are relevant to this paper. We show that, for any angle  $\alpha$ ,  $0^\circ < \alpha < 360^\circ$ , such that  $\frac{360^\circ}{\alpha} = k_\alpha$  is a natural number (see Definition 2), the set

$$\Theta_{k_\alpha} = \{j \cdot \alpha \mid 0 \leq j < k_\alpha\}$$

is degree-bounded. Indeed, such sets are closed under addition of angles and  $\sin(j \cdot \alpha)$  and  $\cos(j \cdot \alpha)$  are algebraic numbers for  $0 \leq j < k_\alpha$ . The latter follows from the well-known equality  $(\cos \alpha + i \sin \alpha)^{k_\alpha} = 1$  of complex numbers. Expanding the real part of the left side of this equality gives a polynomial equation in  $\cos \alpha$  and  $\sin^2 \alpha = 1 - \cos^2 \alpha$ , which shows that  $\cos \alpha$  is an algebraic number. By looking at the imaginary part, we see that  $\sin \alpha$  is also algebraic. Since the set of algebraic numbers form a field, it follows from the well-known formulas for sines and cosines of sums of angles that also  $\sin(j \cdot \alpha)$  and  $\cos(j \cdot \alpha)$  are algebraic numbers for  $0 \leq j < k_\alpha$ . In fact, this argumentation shows that the sines and cosines of any rational multiple of  $180^\circ$ , the so-called *trigonomometric numbers*, are algebraic numbers.

From the above equation, it also follows that the degrees of these numbers are bounded by  $k_\alpha$ . Thus, for any finite subset  $\{\theta_1, \dots, \theta_k\}$  of  $\Theta_{k_\alpha}$ , the degree of the field extension  $\mathbf{Q}(\sin \theta_1, \dots, \sin \theta_k, \cos \theta_1, \dots, \cos \theta_k)$  over  $\mathbf{Q}$  is bounded by  $k_\alpha^{2k}$ , which, in its turn, is bounded by  $k_\alpha^{2k_\alpha}$  (which can serve as  $d$ ). The algebraic numbers  $\sin(j \cdot \alpha)$  and  $\cos(j \cdot \alpha)$  are related to the cyclotomic integers and for a uniform bound on their encoding size, we refer to Section 4 of (Adler and Beling (1994)).

Thus, the sets of angles, that are of interest to this paper, are degree-bounded and in Section 7, we show, in more detail, that  $\Theta_{90^\circ} = \{0^\circ, 90^\circ, 180^\circ, 270^\circ\}$  and  $\Theta_{45^\circ} = \{0^\circ, 45^\circ, 90^\circ, 135^\circ, 180^\circ, 225^\circ, 270^\circ, 315^\circ\}$  are degree-bounded.

Now, we state and prove the main theorem of this section.

**Theorem 5.** *Let  $\Theta$  be degree-bounded set of angles. There is a polynomial-time decision procedure that determines, on input an  $N \times N$  matrix  $M$  of 4-tuples  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$  and a sequence of angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$  from  $\Theta$ , whether  $M$  is the double-cross matrix of a polyline of size  $N$  with successive angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$  in its polar representation. If the answer is positive, also witnesses to this fact can be produced in polynomial time.*

**Proof.** Let  $\Theta$  be a degree-bounded set of angles with extension degree bound  $d$  and encoding bound  $E$ .

We describe a procedure for the decision problem in the statement of this theorem. Let the input be an  $N \times N$  matrix  $M$  of 4-tuples  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$  and a sequence of angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$  from  $\Theta$ . Our decision procedure determines whether there exist lengths  $\ell_1, \ell_2, \dots, \ell_N > 0$ , such that  $M$  is the double-cross matrix of the polyline (that starts with the unit interval on the  $x$ -axis, as its first line segment) with polar representation  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$ .

As an apriori step, we check whether  $M$  does not have  $(0 \ 0 \ 0 \ 0)$  entries on its diagonal or does not have the “symmetry” properties, discussed in Section 2.2. We also check, whether the given sequence of angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$  satisfies the entries  $M[i, i+1]$ ,  $0 \leq i < N$ , of  $M$ . Hereto, we can use Corollary 1. So,  $C_1$  should be  $-$  and  $C_3$  should be  $0$ . And the entries  $C_2 = \text{sign}(\cos \theta_i)$  and  $C_4 = \text{sign}(\sin \theta_i)$  in all  $M[i, i+1]$  should be consistent with the given angles  $\theta_i$ . If any of these tests fail, we can already answer *no*. In the other case, we proceed to determine if there exist lengths  $\ell_1, \ell_2, \dots, \ell_N > 0$  that satisfy the other conditions expressed by the matrix  $M$ .

We have already remarked that, since the given angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$  belong to  $\Theta$ , all the cosine values  $\cos(\theta_{i+1} + \dots + \theta_{k-1})$  and  $\cos(\theta_k + \dots + \theta_j)$  and all the sine values  $\sin(\theta_{i+1} + \dots + \theta_{k-1})$  and  $\sin(\theta_k + \dots + \theta_j)$ , that appear in the expressions given in Theorem 4, are algebraic numbers. This implies that the double-cross conditions, given by Theorem 4, together with the constraints that the  $\ell_i$  are strictly positive lengths, can be seen as linear constraint conditions in  $\ell_1, \ell_2, \dots, \ell_N$  of the form

$$\left\{ \begin{array}{llll} - & \sum_{k=i+1}^j \ell_k \cos(\theta_{i+1} + \dots + \theta_{k-1}) & \alpha_{ij} & 0 \quad (0 \leq i < j < N) \\ & \sum_{k=i+1}^j \ell_k \cos(\theta_k + \dots + \theta_j) & \beta_{ij} & 0 \quad (0 \leq i < j < N) \\ - & \sum_{k=i+1}^j \ell_k \sin(\theta_{i+1} + \dots + \theta_{k-1}) & \gamma_{ij} & 0 \quad (0 \leq i < j < N) \\ & \sum_{k=i+1}^j \ell_k \sin(\theta_k + \dots + \theta_j) & \delta_{ij} & 0 \quad (0 \leq i < j < N) \\ \ell_i & & & > 0 \quad (0 < i \leq N) \end{array} \right. \quad (*)$$

with  $\alpha_{ij}, \beta_{ij}, \gamma_{ij}, \delta_{ij} \in \{=, <, >\}$ , determined by the entries of the matrix  $M$ . Since all the cosines and sines in these expressions are algebraic constants, all these conditions are linear in  $\ell_1, \ell_2, \dots, \ell_N$ . Therefore,  $(*)$  can be seen as

a *linear programming problem*, or at least almost. Normally, in a linear programming problem, linear polynomial conditions of the form

$$a_1\ell_1 + a_2\ell_2 + \cdots + a_N\ell_N \geq 0,$$

with rational coefficients  $a_i$ , are expected to appear, together with the additional conditions

$$\ell_i \geq 0 \quad (0 \leq i \leq N).$$

So, we are left with three problems to see (\*) as a traditional *linear programming problem*:

- (1) we have  $\ell_i > 0$  for  $0 < i \leq N$  and not the traditional  $\ell_i \geq 0$ ;
- (2) we have  $\alpha_{ij}, \beta_{ij}, \gamma_{ij}, \delta_{ij} \in \{=, <, >\}$  and not the traditional  $\geq$ ; and
- (3) we possibly have irrational coefficients  $a_i$ .

The linear polynomial condition

$$a_1\ell_1 + a_2\ell_2 + \cdots + a_N\ell_N = 0$$

is obviously equivalent to

$$a_1\ell_1 + a_2\ell_2 + \cdots + a_N\ell_N \geq 0 \text{ and } a_1\ell_1 + a_2\ell_2 + \cdots + a_N\ell_N \leq 0.$$

This solves the case of equality. Obviously,

$$a_1\ell_1 + a_2\ell_2 + \cdots + a_N\ell_N < 0$$

is equivalent to

$$-a_1\ell_1 - a_2\ell_2 - \cdots - a_N\ell_N > 0.$$

So, we are left with  $a_1\ell_1 + a_2\ell_2 + \cdots + a_N\ell_N > 0$ . To solve the problem of the strict inequalities in (1) and (2), there is a known trick from the linear programming literature that we can use (see page 22 of Matousek and Gärtner (2007)). We introduce a new variable  $\delta$ , which stands for the “gap” between the left and the right side of each inequality and we try to make this gap as large as possible. We consider the linear program

$$\begin{aligned} & \text{maximize } \delta \\ & \text{subject to } a_1\ell_1 + a_2\ell_2 + \cdots + a_N\ell_N - \delta \geq 0 \\ & \text{and } \delta \geq 0 \end{aligned}$$

and observe that  $a_1\ell_1 + a_2\ell_2 + \dots + a_N\ell_N > 0$  is equivalent to the optimal solution  $\delta$  of this linear program being strictly positive. Furthermore, a single  $\delta$  can be used to deal with several strict inequalities all at once. Indeed, the linear program has now an extra variable  $\delta$  and the optimal  $\delta$  is strictly positive exactly when the original system with strict inequalities has a solution.

We define the sets  $S_=$ ,  $S_<$  and  $S_>$  to consists of all the linear polynomials (in  $\ell_1, \ell_2, \dots, \ell_N$ ), that appear in the first four lines of (\*) and for which  $\alpha_{ij}, \beta_{ij}, \gamma_{ij}, \delta_{ij}$  is, respectively,  $=$ ,  $<$  and  $>$ .

Now, we can see that the existence of  $\ell_1, \ell_2, \dots, \ell_N$  that satisfy (\*) is equivalent to the optimal solution  $\delta$  of the following linear programming problem being strictly positive:

$$\begin{aligned}
& \text{maximize } \delta \\
& \text{subject to} \quad P(\ell_1, \ell_2, \dots, \ell_N) \geq 0, \quad \text{for } P \in S_= \\
& \quad \quad \quad -P(\ell_1, \ell_2, \dots, \ell_N) \geq 0, \quad \text{for } P \in S_= \\
& \quad \quad \quad P(\ell_1, \ell_2, \dots, \ell_N) - \delta \geq 0, \quad \text{for } P \in S_> \\
& \quad \quad \quad -P(\ell_1, \ell_2, \dots, \ell_N) - \delta \geq 0, \quad \text{for } P \in S_< \\
& \text{and} \quad \quad \quad \ell_i - \delta \geq 0, \quad \text{for } 0 < i \leq N.
\end{aligned}$$

What remains is Problem (3), namely that we may have the irrational coefficients in our linear programming problem. However, a result by Adler and Beling (Adler and Beling (1994)) shows that linear programming with algebraic coefficients also has a time complexity that is a polynomial of

- (i) the “rank” of the linear system of inequalities, which, applied to our example, is  $O(N^2)$ ;
- (ii) the degree of the field extension of the rationals in which we work, which, in our case, is bounded by the constant  $d$  (the field extension degree bound of  $\Theta$ ); and
- (iii) the encoding size of  $\Theta$ , which, in our case, is bounded by the constant  $E$  (the encoding bound of  $\Theta$ ).

So, the above linear programming problem has a solution that can be determined in polynomial time in  $N$ ,  $d$  and  $E$ . Since, for given  $\Theta$ ,  $d$  and  $E$  are constants, we conclude that our linear programming problem can be solved

in polynomial time in  $N$ . Solving the above linear program, also produce example lengths  $\ell_1, \ell_2, \dots, \ell_N$  (if they exist). This completes the proof.  $\square$

## 7. A realisability test for $45^\circ$ -polylines and some remarks on convexity

In this section, we describe how it can be decided whether a given  $N \times N$  matrix is realisable in the plane by a  $45^\circ$ -polyline. That is, we look at Problem 1 for  $\mathcal{P} = \mathcal{P}_{45^\circ}$ , the class of  $45^\circ$ -polylines (again, with vertices with algebraic coordinates). At the end of this section, we discuss some implications of our result on the convexity of the solution set, determined by a matrix that is realisable in the plane by a  $45^\circ$ -polyline.

### 7.1. A realisability test for $45^\circ$ -polylines

For the problem of realisability, here again, we may assume, without loss of any generality, that the polyline that realises a matrix  $M$ , if it exists, starts with the unit interval on the  $x$ -axis, that is,  $\mathbf{p}_0 = (\mathbf{x}_0, \mathbf{y}_0) = (0, 0)$  and  $\mathbf{p}_1 = (\mathbf{x}_1, \mathbf{y}_1) = (1, 0)$ . This also permits us, to use the results on the polar representation from Section 5 and the results of the previous section, that depend on it.

The proof of the following theorem relies on the fact that, when we restrict the angles that appear in a polyline to be multiples of  $45^\circ$ , there is a one to one correspondence between the qualitative and the quantitative representation (modulo the lengths of the segments). More specifically, the entries above the diagonal of an input matrix  $M$  uniquely determine the angles of a polyline that can realise  $M$ . This correspondence, in combination with the result of the previous section, implies the main result of the paper: *the realisability problem for  $45^\circ$ -polylines is solvable in polynomial time.*

**Theorem 6.** *It can be decided in polynomial time whether an  $N \times N$  matrix  $M$  of 4-tuples  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$  is the double-cross matrix of some  $45^\circ$ -polyline of size  $N$  in  $\mathbf{R}^2$ . If this is the case, also witnesses to this can be produced in polynomial time.*

**Proof.** We now describe a decision procedure that solves Problem 1 for  $\mathcal{P} = \mathcal{P}_{45^\circ}$ . Let  $M$  be a  $N \times N$  input matrix of 4-tuples  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$ . In a first step, we determine the polar angles of the polyline,

we attempt to construct. Once these angles have been determined, in a second step, we decide, using the method of Theorem 5, whether appropriate lengths of the line segments can be found. As before, as an apriori step, we check whether  $M$  doesn't have  $(0 \ 0 \ 0 \ 0)$  entries on its diagonal or doesn't have the "symmetry" properties, discussed in Section 2.2. If  $M$  fails this symmetry-test, we can already answer *no*, else we proceed.

**Step 1 (Determining the angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$ ).** First, we inspect the entries  $M[i, i+1]$ ,  $0 \leq i < N$  of  $M$ . Hereto, we use Corollary 1. So,  $C_1$  should be  $-$  and  $C_3$  should be  $0$ . If this is not the case, we can already answer *no*. From  $C_2$  and  $C_4$  in all entries  $M[i, i+1]$ , we can determine the angles  $\theta_i$  as is shown in the following table.

$C_2$	$C_4$	$\theta_i$
0	0	answer <i>no</i>
0	+	$270^\circ$
0	-	$90^\circ$
+	0	$180^\circ$
+	+	$225^\circ$
+	-	$135^\circ$
-	0	$0^\circ$
-	+	$315^\circ$
-	-	$45^\circ$

Obviously, if both  $C_2 = \text{sign}(\cos \theta_i)$  and  $C_4 = \text{sign}(\sin \theta_i)$  are 0, we have an impossible situation (see Corollary 1). So, at this point, or we have answered *no*, or we know all the angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$  of a possible realisation of  $M$ . In the latter case, we proceed to Step 2.

**Step 2 (Determining  $\ell_1, \ell_2, \dots, \ell_N$ ).** Once, we have determined the angles  $\theta_1, \theta_2, \dots, \theta_{N-1}$ , we can compute all the values  $\cos(\theta_{i+1} + \dots + \theta_{k-1})$ ,  $\cos(\theta_k + \dots + \theta_j)$ ,  $\sin(\theta_{i+1} + \dots + \theta_{k-1})$  and  $\sin(\theta_k + \dots + \theta_j)$  that appear in the expressions given in Theorem 4. Since all these sums of angles are multiples of  $45^\circ$ , these cosines and sines will take values as shown in the following table.



$\alpha$	$\cos \alpha$	$\sin \alpha$
$0^\circ$	1	0
$45^\circ$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$
$90^\circ$	0	1
$135^\circ$	$-\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$
$180^\circ$	-1	0
$225^\circ$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$
$270^\circ$	0	-1
$315^\circ$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$

This means that the double-cross conditions given by Theorem 4, together with the constraints that the  $\ell_i$  are strictly positive lengths, can be seen as linear constraint conditions in  $\ell_1, \ell_2, \dots, \ell_N$ . To this system of linear polynomial (in)equalities, the method of Theorem 5 can be applied. Indeed, as we have observed in Section 6, the set of angles  $\Theta_{45^\circ} = \{0^\circ, 45^\circ, 90^\circ, 135^\circ, 180^\circ, 225^\circ, 270^\circ, 315^\circ\}$  is degree-bounded.

The degree of the extension of the rationals in which the linear programming problem has its coefficients is, in this case, 2. Indeed, we have  $(\mathbb{Q}(\sqrt{2}) : \mathbb{Q}) = 2$ , since the minimal polynomial of  $\sqrt{2}$  is  $x^2 - 2$ . So, we have extension degree bound  $d = 2$ . As discussed in the previous section, the encoding bound  $E$  depends on the number of bits needed to encode the coefficients  $\frac{\sqrt{2}}{2}$  and  $-\frac{\sqrt{2}}{2}$ . An encoding of  $\frac{\sqrt{2}}{2}$  is given by  $(x^2 - \frac{1}{2}, 0, 1)$ , since  $\frac{\sqrt{2}}{2}$  is the only root of the polynomial  $x^2 - \frac{1}{2}$  in the interval  $[0, 1]$ . Six bits are needed to encode the coefficients of this polynomial and the boundary points of this interval. We have a similar bound for  $-\frac{\sqrt{2}}{2}$ . Depending on the particular way we encode polynomials (for instance, in sparse or dense encoding), this shows that also the encoding bound  $E$  can be taken to be a small constant. For further details, we refer to Section 4 of (Adler and Beling (1994)).

Thus, Theorem 5, applied to this setting, tells us that our linear programming problem can be solved in polynomial time in  $N$ . The linear program, as discussed in the proof of Theorem 5, can also produce example lengths  $\ell_1, \ell_2, \dots, \ell_N$  (if they exist). This completes the proof.  $\square$

## 7.2. Convexity properties of 45°-polylines

From Step 1 of the proof of Theorem 6, it follows that a matrix  $M$  that is realisable by a 45°-polyline determines the angles  $\theta_i$  uniquely for  $0 < i < N$ . This proves the following corollary.

**Corollary 2.** *If an  $N \times N$  matrix  $M$  of 4-tuples  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$  is realisable by two 45°-polylines  $P_1$  and  $P_2$ , that start with the line segment connecting  $(0,0)$  and  $(1,0)$  and have polar-coordinate representations  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$  and  $\langle \ell'_1, \theta'_1, \ell'_2, \theta'_2, \dots, \ell'_{N-1}, \theta'_{N-1}, \ell'_N \rangle$ , respectively, then  $\theta_i = \theta'_i$  for  $0 < i < N$ .  $\square$*

Also from the proof of Theorem 6, the following property follows.

**Corollary 3.** *If an  $N \times N$  matrix  $M$  of 4-tuples  $(C_1 \ C_2 \ C_3 \ C_4) \in \{-, 0, +\}^4$  is realisable by two 45°-polylines  $P_1$  and  $P_2$ , that start with the line segment connecting  $(0,0)$  and  $(1,0)$  and have polar-coordinate representations  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$  and  $\langle \ell'_1, \theta'_1, \ell'_2, \theta'_2, \dots, \ell'_{N-1}, \theta'_{N-1}, \ell'_N \rangle$ , respectively, then for any real numbers  $\alpha_1, \alpha_2 > 0$ , the 45°-polyline given by the polar coordinate representation  $\langle \alpha_1 \cdot \ell_1 + \alpha_2 \cdot \ell'_1, \theta_1, \alpha_1 \cdot \ell_2 + \alpha_2 \cdot \ell'_2, \theta_2, \dots, \alpha_1 \cdot \ell_{N-1} + \alpha_2 \cdot \ell'_{N-1}, \theta_{N-1}, \alpha_1 \cdot \ell_N + \alpha_2 \cdot \ell'_N \rangle$  also realises  $M$ .*

**Proof.** Corollary 2 takes care of the angles. From Step 2 of the proof of the previous theorem it follows that if  $P_1$  and  $P_2$  are realisations of a matrix  $M$  their lengths satisfy the same set of linear conditions of the form  $a_1 \ell_1 + a_2 \ell_2 + \dots + a_N \ell_N \alpha$ , with  $\alpha \in \{<, =, >\}$ . Suppose that we have

$$\begin{cases} a_1 \ell_1 + a_2 \ell_2 + \dots + a_N \ell_N > 0 \text{ and} \\ a_1 \ell'_1 + a_2 \ell'_2 + \dots + a_N \ell'_N > 0 \end{cases}$$

for  $P_1$  and  $P_2$ , for any of these linear conditions. Since both  $\alpha_1 > 0$  and  $\alpha_2 > 0$ , we also have

$$\begin{cases} \alpha_1 \cdot (a_1 \ell_1 + a_2 \ell_2 + \dots + a_N \ell_N) > 0 \text{ and} \\ \alpha_2 \cdot (a_1 \ell'_1 + a_2 \ell'_2 + \dots + a_N \ell'_N) > 0. \end{cases}$$

So, also the sum of the two left hand sides,

$$\alpha_1 \cdot (a_1 \ell_1 + a_2 \ell_2 + \cdots + a_N \ell_N) + \alpha_2 \cdot (a_1 \ell'_1 + a_2 \ell'_2 + \cdots + a_N \ell'_N)$$

will be strictly larger than 0. The same argument hold when  $\alpha$  is  $=$  or  $<$ . This completes the proof.  $\square$

We end this chapter with the following *convexity property* for  $45^\circ$ -polylines.

**Corollary 4.** *The set of  $45^\circ$ -polylines, that start with the line segment connecting  $(0, 0)$  and  $(1, 0)$ , and that realise an  $N \times N$  matrix  $M$  of 4-tuples  $(C_1 C_2 C_3 C_4) \in \{-, 0, +\}^4$ , is a convex set.*

**Proof.** Let  $M$  be a  $N \times N$  matrix of 4-tuples  $(C_1 C_2 C_3 C_4) \in \{-, 0, +\}^4$ . If  $M$  cannot be realised by a  $45^\circ$ -polyline (that start with the line segment connecting  $(0, 0)$  and  $(1, 0)$ ), or by exactly one such polyline, then the statement is trivially true.

On the other hand, let  $P_1$  and  $P_2$  be two  $45^\circ$ -polylines that realise  $M$ . We have to show that if  $P_1$  and  $P_2$  have polar-coordinate representations  $\langle \ell_1, \theta_1, \ell_2, \theta_2, \dots, \ell_{N-1}, \theta_{N-1}, \ell_N \rangle$  and  $\langle \ell'_1, \theta'_1, \ell'_2, \theta'_2, \dots, \ell'_{N-1}, \theta'_{N-1}, \ell'_N \rangle$ , respectively, then for any  $\lambda$ , with  $0 \leq \lambda \leq 1$ , the  $45^\circ$ -polyline given by the polar coordinate representation

$$\begin{aligned} &\langle \lambda \cdot \ell_1 + (1 - \lambda) \cdot \ell'_1, \lambda \cdot \theta_1 + (1 - \lambda) \cdot \theta'_1, \\ &\lambda \cdot \ell_2 + (1 - \lambda) \cdot \ell'_2, \lambda \cdot \theta_2 + (1 - \lambda) \cdot \theta'_2, \dots, \lambda \cdot \ell_{N-1} + (1 - \lambda) \cdot \ell'_{N-1}, \\ &\lambda \cdot \theta_{N-1} + (1 - \lambda) \cdot \theta'_{N-1}, \lambda \cdot \ell_N + (1 - \lambda) \cdot \ell'_N \rangle \end{aligned}$$

also realises  $M$ .

From Corollary 2, it is clear that  $\lambda \cdot \theta_i + (1 - \lambda) \cdot \theta'_i = \theta_i = \theta'_i$  for  $0 < i < N$ . For  $\lambda$  with  $0 \leq \lambda \leq 1$ , we observe that if we take  $\lambda = 0$ , we get  $P_2$  and if we take  $\lambda = 1$ , we get  $P_1$ . This leaves us with the case  $0 < \lambda < 1$ . But here, both  $\lambda$  and  $1 - \lambda$  are strictly larger than 0 and Corollary 3 applies with  $\alpha_1 = \lambda$  and  $\alpha_2 = 1 - \lambda$ . This completes the proof.  $\square$

## 8. Conclusion and discussion

We have studied the decision problem that asks whether a  $N \times N$  matrix of 4-tuples from  $\{-, 0, +\}$  is the double-cross matrix of a polyline with  $N$  line

segments. This problem is, in general, NP-hard. In this paper, we have given a conceptually easy  $O(N^2)$ -time algorithms for the case where the attention is restricted to polylines in which consecutive line segments make angles that are multiples of  $90^\circ$ . Next, we have given a more complicated algorithm that solves the question for  $45^\circ$ -polylines. For this more complicated case of  $45^\circ$ -polylines, we have introduced the polar-coordinate representation of double-cross matrices.

We emphasise that in both examples, the key to avoid the exponential blow-up, relies on the fact that the entries of the input matrix above its diagonal completely determine the angles between two consecutive segments (in linear time). This one-to-one correspondence between the qualitative double-cross information and the angle information implies that our decision problem simplifies to deciding whether or not appropriate segment lengths of a polyline exist.

If we would be interested, for instance, in the realisability question for  $30^\circ$ -polylines, we would, no longer, be able to rely on such a one-to-one correspondence between a part of the matrix and the angles. In fact, in this case, we would have to consider two possibilities per quadrant of the double cross (essentially corresponding to  $30^\circ$  and  $60^\circ$ ). This options between two possible values for each of the angles would lead to an exponential blow-up in  $N$ . It is not clear how such a blow-up can be avoided, unless a richer formalism than the double-cross method would be used. If instead of working with crosses, we would work with star-shaped divisions that would divide the space around each vertex into 8 regions instead of 4, we could also apply the methods of this paper to obtain a polynomial-time decision procedure for  $30^\circ$ -polylines.

## Acknowledgements

The authors would like to thank the anonymous reviewer, whose comments helped to improve the presentation of the paper considerably.

- Adler, I., Beling, P. A., 1994. Polynomial algorithms for linear programming over the algebraic numbers. *Algorithmic* 12 (6), 436–457.
- Basu, S., Pollack, R., Roy, M.-F., 2006. *Algorithms in Real Algebraic Geometry (Algorithms and Computation in Mathematics)*. Springer-Verlag New York, Inc.
- Bochnak, J., Coste, M., Roy, M.-F., 1998. *Real Algebraic Geometry*. Vol. 36 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. Folge 3*. Springer-Verlag.
- Bookstein, F. L., 1986. Size and shape spaces for landmark data in two dimensions. *Statistical Science* 1, 181–242.
- Canny, J., 1988. Some algebraic and geometric computations in PSPACE. In: *Proceedings of the Twentieth Annual ACM Symposium on Theory of Computing (STOC 1988)*. ACM, pp. 460–467.
- Collins, G., 1975. Quantifier elimination for real closed fields by cylindrical algebraic decomposition. In: Brakhage, H. (Ed.), *Automata Theory and Formal Languages*. Vol. 33 of *Lecture Notes in Computer Science*. Springer, pp. 134–183.
- Cox, D. A., Little, J., O’Shea, D., 1997. *Ideals, varieties, and algorithms - an introduction to computational algebraic geometry and commutative algebra (2. ed.)*. Undergraduate texts in mathematics. Springer.
- Dolzmann, A., Sturm, T., 1997. Redlog: Computer algebra meets computer logic. *ACM SIGSAM Bulletin* 32 (2), 2–9, (see also [redlog.dolzmann.de/](http://redlog.dolzmann.de/)).
- Dryden, I., Mardia, K. V., 1998. *Statistical Shape Analysis*. Wiley.
- Forbus, K. D., 1990. Qualitative physics: past present and future. In: *Readings in qualitative reasoning about physical systems*. Morgan Kaufmann, pp. 11–39.
- Freksa, C., 1992. Using orientation information for qualitative spatial reasoning. In: *Spatio-Temporal Reasoning (GIS’92)*. pp. 162–178.
- Gero, J. S., 1999. Representation and reasoning about shapes: cognitive and computational studies in visual reasoning in design. In: et al., K. (Ed.),

- Proceedings of the International Conference on Spatial Information Theory (COSIT'99). Vol. Lecture Notes in Computer Science, 2825. Springer, pp. 315–330.
- Giannotti, F., Pedreschi, D., 2008. *Mobility, Data Mining and Privacy*. Springer.
- Giusti, M., Heintz, J., 2001. Kronecker's smart, little black boxes. In: DeVore, R., A., I., Suli, E. (Eds.), *Foundations of Computational Mathematics*. Cambridge University Press, Cambridge, pp. 69–104.
- Güting, R. H., Schneider, M., 2005. *Moving Objects Databases*. Morgan Kaufmann.
- Heintz, J., Kuijpers, B., Paredes, A. R., 2013. Software engineering and complexity in effective algebraic geometry. *Journal of Complexity* 29 (1), 92–138.
- Hong, H., 2000. QEPCAD. [www.usna.edu/CS/qepcadweb/B/QEPCAD.html](http://www.usna.edu/CS/qepcadweb/B/QEPCAD.html).
- Kent, J. T., Mardia, K. V., 1986. Shape, procrustes tangent projections and bilateral symmetry. *Biometrika* 88, 469–485.
- Khovanskii, A., 1991. *Fewnomials*. Translations of mathematical monographs. American Mathematical Society.
- Kuijpers, B., Moelans, B., Van de Weghe, N., 2006. Qualitative polyline similarity testing with applications to query-by-sketch, indexing and classification. In: *Proceedings of the 14th annual ACM International Symposium on Advances in Geographic Information Systems*. ACM, pp. 11–18.
- Lang, S., 1986. *Algebraic Number Theory* (2nd ed.). No. 110 in Graduate Texts in Mathematics. Springer.
- Matousek, J., Gärtner, B., 2007. *Understanding and Using Linear Programming*. No. v. 36 in Universitext. Springer.
- Mokhtarian, F., Mackworth, A. K., 1992. A theory of multiscale, curvature-based shape representation for planar curves. *IEEE Transactions on Pattern Analysis and Machine Intelligence (TPAMI)* 14, 789–805.

- Renz, J., Nebel, B., 2007. Qualitative spatial reasoning using constraint calculi. In: Aiello, M., Pratt-Hartmann, I., van Benthem, J. (Eds.), *Handbook of Spatial Logics*. Springer.
- Scivos, A., Nebel, B., 2001. Double-crossing: Decidability and computational complexity of a qualitative calculus for navigation. In: *Spatial Information Theory: Foundations of Geographic Information Science*, International Conference, COSIT 2001. pp. 431–446.
- Tarski, A., 1951. *A Decision Method for Elementary Algebra and Geometry*. University of California Press.
- Wolfram Research, 2015. *Mathematica 9*. [www.wolfram.com](http://www.wolfram.com).
- Zimmermann, K., Freksa, C., 1996. Qualitative spatial reasoning using orientation, distance, and path knowledge. *Applied Intelligence* 6 (1), 49–58.