Made available by Hasselt University Library in https://documentserver.uhasselt.be

Blood pressure in young adulthood and residential greenness in the early-life environment of twins Peer-reviewed author version

BIJNENS, Esmee; NAWROT, Tim; Loos, Ruth J. F.; Gielen, Marij; Vlietinck, Robert; Derom, Catherine & Zeegers, Maurice P. (2017) Blood pressure in young adulthood and residential greenness in the early-life environment of twins. In: ENVIRONMENTAL HEALTH, 16, p. 1-8 (Art N° 53).

DOI: 10.1186/s12940-017-0266-9 Handle: http://hdl.handle.net/1942/24269

Blood pressure in young adulthood and residential exposure of greenness in the early-life environment of twins

Esmée M Bijnens^{a,b}, PhD; Tim S Nawrot^{a,c}, PhD; Ruth JF Loos^d, MD, PhD; Marij Gielen^b, MD, PhD; Robert Vlietinck^e, MD, PhD; Catherine Derom^{e,f}, PhD; Maurice P Zeegers^{b,g}, PhD

11 12	^a Centre for Environmental Sciences, Hasselt University, Agoralaan Building D, 3590 Diepenbeek, Belgium							
13 14 15	^b Department of Complex Genetics, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre, P.O. Box 616, 6200 MD Maastricht, The Netherlands							
16	^c CAPHR School for Public Health and Primary Care. Maastricht University. The Netherlands							
17	^d The Genetics of Obesity and Related Metabolic Traits Program. The Charles Bronfman Institute for							
18	Personalized Medicine, The Mindich Child Health and Development Institute, The Icahn School of Medicine et Meuret Singi 1468 Medicen Ave. New York, NY 10020, United States							
20	^e Centre of Human Constice. University Hospitals Leuven, Herestraat 49, 3000 Leuven, Belgium							
20	^f Department of Obstatrics and Gynaecology, Chent University Hespital, De Pintelaan 185, Chent							
21	Belgium							
22	g Department of Public Health, Leuven University (KULLeuven), Kapucijnenvoer 35, 3000 Leuven							
23	Belgium							
25	Beigium							
26	Email addresses of the authors are respectively:							
27	esmee bijnens@ubasselt be tim nawrot@ubasselt be ruth loos@mssm edu							
28	marii.gielen@maastrichtuniversity.nl robert.vlietinck@med.kuleuven.be c.derom@telenet.be							
29	m.zeegers@maastrichtuniversity.nl							
30								
31								
32								
33								
34								
35	Corresponding author: Dr. Tim Nawrot, Agoralaan Building D, 3590 Diepenbeek, Belgium. Email:							
36	tim.nawrot@uhasselt.be. Tel: +3211268382.							

37 Abstract

38 Background

Previous research shows that, besides risk factors in adult life, the early-life environment can influence blood pressure and hypertension in adults. However, the effects of residential traffic exposure and residential greenness in the early-life on blood pressure in young adulthood are currently unknown.

42 Methods

Ambulatory (24-hour) blood pressures of 278 twins (132 pairs) of the East Flanders Prospective Twins Study were obtained at the age of 18 to 25 years. Prenatal and adulthood residential addresses were geocoded and used to assign prenatal and postnatal traffic and greenness indicators. Mixed modelling was performed to investigate blood pressure in association with greenness while adjusting for potential confounding factors.

48 Results

49 Night-time systolic blood pressure was inversely associated with greenness at the residential address 50 in twins living at the same address their entire life (non-movers, n=97, 34.9%). An interquartile increase 51 in residential greenness exposure (1000 m radius) was associated with a 3.59 mm Hg (95% CI: -6.0 to 52 -1.23; p=0.005) lower adult night systolic blood pressure. Among twins who were living at a different 53 address than their birth address at time of the measurement (n=181, 65.1%), night-time blood pressure 54 was inversely associated with residential surrounding greenness at adult age as well as with residential 55 greenness in early-life. However after additional adjustment for residential greenness exposure in 56 adulthood, only residential greenness exposure in early-life was significantly associated with night 57 systolic blood pressure. While no significant effect of adult residential greenness with adult blood 58 pressure was observed, while accounting for the early-life greenness exposure.

59 Conclusions

Lower residential greenness in the early-life environment was independently associated with a higher
 adult blood pressure. This indicates that residential greenness has persistent effects on blood pressure.

63 *Keywords:* Blood pressure; greenness; early-life environment.

65 Introduction

High blood pressure is the leading global risk factor for cardiovascular disease and mortality in the world [1, 2]. Long-term exposure to air pollution has been associated with increased blood pressure [3] and measures of atherosclerosis, including carotid-intima-media thickness [4]. In a meta-analysis of 113,926 participants of the ESCAPE study, a weak positive association was found between high traffic exposure in 100 m of the residence and increased systolic and diastolic blood pressure, and elevated risk of prevalent hypertension [5]. Blood pressure tracks over time and childhood blood pressure is closely related with adult blood pressure, increasing later cardiovascular risk [6].

73 Given that an unfavorable intrauterine environment may contribute to increased blood pressure [7], we 74 hypothesized that exposure to traffic related air pollution and residential greenness during early-life 75 might be associated with higher blood pressure in early adulthood. Prenatal exposure to air pollutants 76 may influence epigenetic changes in placental tissue, which might result in fetal growth disturbance, and 77 subsequently make children more susceptible to the development of cardiovascular pathologies and 78 disease later in life. Until now, studies on residential greenness and land cover on blood pressure are 79 limited. In the few studies conducted in healthy populations of children or young adults, childhood or 80 recent exposures to greenness has been associated with blood pressure [8-10]. A study in 2,076 81 German children showed that lower residential greenness was associated with higher blood pressure in 82 10 year-old children living in an urbanised area [8]. A Dutch study investigating the effect of 83 neighborhood-level environmental stressors on blood pressure among different ethnic groups observed 84 that living in a neighborhood with a high quality of green space was associated with a lower systolic 85 blood pressure and lower odds of hypertension in Moroccans [9]. Similar but non-significant associations 86 were observed among Dutch and Turkish ethnic groups [9]. In addition, among 3,416 pregnant women 87 of the city Kaunas, Lithuania an association was observed between proximity of the place of residence 88 to green spaces and lower blood pressure [10]. To understand the role of early-life traffic related 89 exposure and residential greenness in blood pressure tracking into adulthood, we investigated the 90 association between early-life exposure of these environmental exposures on blood pressure in young 91 adulthood.

93 Methods

94 Subjects

95 The East Flanders Prospective Twin Study (EFPTS), a population based register of multiple births in 96 the province of East-Flanders (Belgium), started in 1964 to enrol twins at birth [11]. The twin cohort in 97 our analysis was based on a previously designed twin study containing 424 twin pairs [12]. From this 98 population we could obtain 333 twins born between 1975 and 1982. We excluded twins born before 99 1975 since no major changes in the road network have occurred in East-Flanders since 1974. Their 100 addresses at birth and at adult age were geocoded. Twins with missing data were excluded; gestational 101 age (n=4), parental education level (n=17), BMI (n=1), 24h urinary sodium and potassium (n=6), 102 Gamma-Glutamyl Transferase (n=9), age mother (n=9), neigbourhood SES 1994 (n=3), blood pressure 103 (n=6), resulting in a final study population of 278 persons, 170 monozygotic and 108 dizygotic twins. 104 Informed consent was obtained and ethical approval was given by the Ethics Committee of the Faculty 105 of Medicine of the Katholieke Universiteit Leuven.

106 Collection maternal and neonatal data

107 Data recorded by the obstetrician at birth included gestational age, birth weight, sex of the twins and 108 parental ages. Gestational age was based on the mother's last menstruation and was calculated as the 109 number of completed weeks of pregnancy. Zygosity was determined by sequential analysis based on 110 sex, choriontype, blood groups based on umbilical cord blood, placental alkaline phosphatase, and, 111 since 1982, DNA fingerprints [13]. After DNA-fingerprinting, zygosity was determined with a 99.9% 112 probability. At a later stage, the parents of the twins filled out questionnaires. In this way, maternal 113 smoking during pregnancy and parental education were collected retrospectively. Educational level as 114 a proxy of socio economic status (SES) was categorized into three groups according to the Belgian 115 education system; no education or primary school, lower secondary education, and higher secondary education and tertiary education. In addition to individual SES data, we gathered information on 116 117 neighborhood SES. Based on their home address, all mothers were assigned to statistical sectors (average area = 1.55 km^2), the smallest administrative entity for which statistical data are produced by 118 119 the Belgian National Institute of Statistics (NIS). Belgian census data (FOD Economie/DG Statistiek) derived from the NIS were used to define neighborhood SES based on annual household income in theyear 1994.

122 Collection data adulthood

Biometric and laboratory measurements were obtained at the research centre during a 2 h morning session. Standing height and weight were measured as described in detail before [14]. BMI was calculated as body mass (in kg) divided by squared height (in m). The twins completed questionnaires to obtain information on smoking status and physical activity. Regarding habitual physical activity, the twins rated themselves on a 10-point scale after brief instructions, 1 represented very little and 10 very intensive physical activity.

Fasting blood samples were drawn; Gama-Glutamyl Transferase was measured on an Olympus AU600 Auto-Analyzer (Kyoto, Japan). Potassium and sodium were determined in 24 h urine with ion-selective electrodes (Olympus AU600 Auto-Analyzer). Before urine collection, participants were asked to select a day of normal routine, to drink normally and refrain from caffeine use, and to refrain from heavy exercise.

134 Ambulatory blood pressure

135 Ambulatory blood pressure was monitored using the SpaceLabs 90207 device (SpaceLabs, Inc), as 136 previously described [15]. The ambulatory blood pressure measurements were performed between 137 February 1997 and April 2000. The monitor was applied at home on the nondominant arm. They were 138 instructed to perform normal activity, but not to engage in vigorous physical exercise or contact sports. 139 Recording began between 6.00 and 9.00 AM and was finished 24 hours later. The recorders were taken 140 every 15 minutes during daytime (8.00 AM to 10.00 PM) and every 30 minutes during night-time (10.00 141 PM to 8.00 AM). Whenever a reading could not be successfully completed, the measurement was 142 repeated 2 minutes later. Readings were automatically rejected when systolic blood pressure was >220 143 mm Hg or <70 mm Hg and diastolic BP was >140 mm Hg or <40 mm Hg. In addition to the automatic 144 exclusion of readings by the monitor, individuals were excluded from further analysis when there were 145 no valid measurements in any 2-hour period. Night and day were defined with short fixed-clock time 146 periods that ranged from midnight to 6.00 and from 10.00 AM to 8.00 PM. We investigated systolic and 147 diastolic blood pressure during both night and day. Measurements during the morning (6.00 AM to 10.00 AM) and during the evening (8.00 PM to midnight) were excluded. Both periods may be subjected to additional variation due to different activity patterns between individuals.

150 Traffic exposure and land use data

Residential addresses of the mothers at time of birth of the twin and the residential addresses of the twin at time of the measurement were geocoded. Distance to the nearest major road, a proxy for trafficrelated exposure, was determined using Geographic Information System (GIS) functions. All GIS analyses were carried out using ArcGIS 10 software.

Semi-natural-, forested -, and agricultural areas (greenness), residential and industrial areas in a 5000 m radius from the residential address were estimated based on CORINE Land Cover 2000 (European Environment Agency). This was repeated for 4000, 3000, 2000, 1000, 500, 300 and 100 m buffers. CORINE, the acronym for 'coordination of information of information on the environment', was initiated by the European Union and has been taken over by the European Environment Agency. The land cover data is based on satellite data and is dived in 44 classes. It is presented as a cartographic product, at a scale of 1:1 000 000. This database is available for most areas of Europe.

162 Assessment of traffic noise

163 A GIS-based noise model including the Flemish street and railway networks was used to estimate traffic 164 noise levels at the residential address of the mother at time of birth in 5 dB(A)-intervals according to the 165 European Noise Directive (2002/49/EC) (The European Parliament and The Council of The European 166 Union, 2016). The modelling of road noise level included road traffic intensity, vehicle-type-specific traffic 167 density, type of street surface, small-scale topography of the area, and the presence or dimensions of 168 buildings and reflecting objects. Railway noise modelling included the amount of passing trains, type of 169 trains, speed, small-scale topography of the area, and the presence or dimensions of buildings and 170 reflecting objects. Weighted equivalent noise levels in dB(A) for traffic over day-time (based on the 171 weighted yearly average noise level between 7 a.m. to 7 p.m., and 7 p.m. to 11 p.m.,) and at night 172 (yearly average noise level between 11 p.m. and 7 a.m.) were modelled. Exposure to traffic noise was 173 categorized as \leq 55 dB, >55 to \leq 60 dB, and >60 dB.

175 Statistical analysis

For data management and statistical analyses, we used SAS software, version 9.3 (SAS Institute, Cary,
NC). All reported p-values are two-sided and were considered statistically significant when p < 0.05.

178 Mixed modeling was performed to investigate blood pressure in association with distance to major road 179 and land-use indicators. The twins were analyzed as individuals in a multilevel regression analysis to 180 account for relatedness between twin members by adding a random intercept to the model. The 181 variance-covariance structure was allowed to differ between the three zygosity-chorionicity groups 182 including dizygotic dichorionic, monozygotic dichorionic, and monozygotic monochorionic. Covariates 183 were selected a priori including sex, gestational age, birth weight, birth year, zygosity-chorionicity group, 184 maternal age, age, smoking, physical activity, BMI, 24h sodium and potassium, gamma-glutamyl 185 transferase, indicators of socioeconomic status (maternal education and neighborhood household 186 income), smoking during pregnancy and noise exposure during the night. To distinguish between 187 exposure early and late in life, we divided the twins in two groups; twins who were living at the same 188 address their whole life (non-movers) and twins who were living at a different address than their birth 189 address at time of the measurement (movers). In the twins who moved during life, we additionally 190 adjusted for land use/distance to major road in adulthood/early-life. Correlation coefficients between 191 greenness in adulthood and early-life are shown in supplement (Table S1). Variance Inflation Factors 192 (VIF) were less than 5, this suggest that the potential issue of multi-collinearity is rather limited (Table 193 S2).

In addition, we tested potential effect-modifications of the association between land-use indicators and
blood pressure by zygosity–chorionicity group.

196

197 Results

The average age of mothers was 27.4 years at the time twins were born and 14.4% of the mothers smoked during pregnancy (Table 1). At time the blood pressure measurements were taken, the twins had a mean (SD) age of 21 (1.9) years, 31% of the twins were current smokers and 5 subjects (1.8%) used antihypertensive medication. The average 24 h urine sodium and potassium excretion was respectively 128.1 \pm 59.4 mmol/day and 63.2 \pm 28.4 mmol/day and the fasting blood concentration of 203 gamma-glutamyl transferase was 16.6 ± 8.54 U/L. In our final analysis, 95% (n=264) of the participants 204 included both twins from each twin pair, whereas the remaining 5% (n=14) only had one participating 205 twin from each twin pair. A total of 181 (65.1 %) of the twins moved between birth address and young 206 adulthood (Table 2). No significant interaction by zygosity-chorionicity (p≥0.81) was observed in the 207 model of land-use indicators and blood pressure.

In 97 twins living at the same address their entire life (non-movers), significant associations were observed between systolic blood pressure during the night and all land use indicators in a 1000 m buffer after adjustment for previously mentioned covariates (Figure 1 and figure S2 and S3). An interquartile increase in residential greenness exposure within 1000 m residential radius was associated with a decrease of 3.59 mm Hg (95% CI: -6.0 to -1.23; p=0.005) in night-time systolic blood pressure. Diastolic blood pressure during night and day was also inversely associated with greenness in a 300 m radius (-4.0 mm Hg for IQR increase; 95% CI: -6.6 to -1.3; p=0.006, -3.8; 95% CI: -6.8 to -0.9; p=0.01).

215 In the twins who had moved since birth, an increased systolic blood pressure during night was 216 significantly associated with lower greenness and higher residential area in early-life after adjustment 217 for previously mentioned covariates (Table S1). Exposure to land use indicators in early-life remained 218 significant associated with systolic blood pressure during the night, even after additional adjustment for 219 land use indicators in adulthood (Figure 1, figure S2 and figure S3). The association with residential 220 greenness was significant for all buffer sizes (residential greenness radius between 300m and 5000m, 221 Table S3). An interquartile increase in residential greenness exposure in early-life (5000 m radius) was 222 associated with a -2.47 mm Hg (95 CI: -4.7 to -0.2%; p=0.04) decrease in adult night systolic blood 223 pressure, after additional adjustment for residential greenness exposure in adulthood. Diastolic blood 224 pressure during the night was significantly associated with residential greenness in a 300m buffer (-2.4 mm Hg for IQR increase; 95% CI: -4.2 to -0.6; p=0.01; figure 1). No associations were observed between 225 226 blood pressure and distance to major roads in early-life (Figure 2). Residential greenness indicators 227 were robust for mutually adjustment for distance to major roads, which is a proxy for traffic related 228 exposures.

In the movers, night-time systolic blood pressure was inversely associated with residential surrounding greenness at adult age when no adjustments for early-life exposure were made (Figure S1). We found no significant effect of adult land use indicators with adult blood pressure, while accounting for the earlylife land use indicators (Figure 1). Only a significant and positive association between diastolic blood
pressure during the night and industrial area in a 5000 m buffer remains (-2.4; 95% CI: -4.5 to -0.2;
p=0.03; figure S3).

235

236 Discussion

237 The key finding of our paper is that in this twin population, increased early-life exposure to residential 238 greenness was associated with lower blood pressure in young adulthood. These results lend further 239 evidence in support of the developmental origins of disease hypothesis for blood pressure [16] and the 240 role of the residential landscape on the blood pressure life trajectory. Changes in blood pressure in 241 association with land-use exposure were noted during the night and not during the day. Due to physical 242 and mental activity, blood pressure is more variable during the day than during the night. Blood pressure 243 during the night is more related to basal blood pressure and is a better predictor of mortality and coronary 244 heart disease than day-time blood pressure [17].

Our findings are in line with previous studies, showing an association between high residential greenness and low blood pressure in children and adults [8, 9]. However, no studies are available on greenness exposure during pregnancy in association with blood pressure in the newborn. Although, it is reasonable to expect that maternal exposure to greenness also affects newborn blood pressure since a US cohort study of 1059 mothers and their newborn infants shows that maternal systolic blood pressure during the third trimester of pregnancy is associated with the systolic blood pressure among newborns [18].

252 Residential green space might act via three routes. First, we postulate that the underlying mechanism 253 between surrounding greenness in early-life and blood pressure in young adulthood could be stress. 254 Higher levels of green space in residential neighborhoods, are linked with lower perceived stress and a 255 healthier cortisol levels in women in a deprived urban population [19]. Moreover, the presence of multiple 256 psychosocial stressors in women during pregnancy are associated with 1.5 mm Hg higher systolic and 257 diastolic blood pressure in the offspring at the age of 5-7 years [20]. In addition, the prenatal maternal 258 psychological status also does influence blood pressure response to stress in the child when aged 7-9 259 years [21]. Meta-analysis including 50 cohorts and consisting of 617 data points for systolic blood pressure and 547 data points for diastolic blood pressure shows that blood pressure in childhood persists in later life [6]. The blood pressure tracking was greater for systolic than for diastolic blood pressure [6]. Second, our association between residential green space indicators early in life and adult blood pressure might be mediated or partly mediated by air pollution

264 It might be possible that low air pollution concentrations in areas with more surrounding greenness 265 result in a lower blood pressure. Studies in adults show that a long-term exposure to air pollution is 266 associated with higher blood pressure [3, 22-26]. The underlying pathway between particulate air 267 pollution and blood pressure may be oxidative stress and inflammation which can affect placental 268 function and fetal growth [27]. Third, besides air pollution, green space reduce exposure to noise. Noise 269 may play a role in the association between residential greenness and blood pressure since noise 270 exposure can result in a higher blood pressure [28, 29]. However, the models were robust for mutually 271 adjustment for noise exposure during the night.

272 Studies on early-life stressors, such as ambient air pollution, noise and stress, that may influence blood 273 pressure in adulthood are limited. One study shows that prenatal exposure to ambient air pollutants are 274 associated with an increased risk for adverse cardiovascular health outcomes [30]. Another study in 275 adults noted a negative association between prenatal life stress and blood pressure in young adulthood 276 [31]. To our knowledge no studies have investigated prenatal noise exposure in association with adult 277 blood pressure. Besides these early-life stressors, other factors including smoking behaviour are 278 associated with adult blood pressure. With this regard, maternal smoking during pregnancy has been 279 associated with an increase in offspring blood pressure in adulthood [32, 33].

A major advantage of this study is that we individually estimated greenness exposure of each participant. Our study is the first to include information on greenness exposure in the early-life when investigating adult blood pressure. Since we cannot differentiate between prenatal and postnatal exposure to residential greenness, our study is limited to define the exact time period of the exposure. Nevertheless, residential greenness in early-life remained significant even after additional adjustment for residential greenness exposure in adulthood. We found no significant effect of adult residential greenness with adult blood pressure, while accounting for the early-life greenness exposure.

287 Strengths include the use of ambulatory blood pressure and the possibility to control for potential 288 confounding factors and covariates of blood pressure such as age, gender, smoking status, physical

289 activity, BMI, sodium and potassium intake and alcohol consumption. Ambulatory blood pressure was 290 measured at home during usual daily activities and is, compared to office measurements, a better 291 predictor of all cause and cardiovascular mortality and has a moderate-to-relatively high tracking stability 292 and predictability over time [34, 35]. There are a number of limitations to our study. The twins were born 293 from 1975 till 1982, although during the last 30 years traffic volume did increase no major geographical 294 changes did appear since no major changes in the road network have occurred in East-Flanders since 295 1974. In addition, we used land-cover data from 2000 as no earlier satellite data were available. 296 However strong correlations have been shown over time between 2000 and 2006 [36]. A major limitation 297 of our study is that information on residential location is only available for the first and last residence. 298 This results in more accurate exposure estimations for non-movers than for the movers because no data 299 is available on moment of moving and frequency of moving.

Since high blood pressure is the leading global risk factor for cardiovascular disease and mortality in the world [1], these findings have public health implications. Previous studies have shown that a small reduction in blood pressure at the population level could have a major impact in reducing morbidity and mortality [37]. Even, a 2 mm Hg lower systolic blood pressure at the population level would result in an 10% overall reduction in mortality due to stroke mortality and about 7% lower mortality from ischemic heart disease or other vascular causes in middle age [37].

306 Conclusions

Our findings indicate that more surrounding greenness at the residential address in early-life shows
 lower blood pressure in adult life. This may be especially relevant for policy makers and urban planners
 for designing healthier urban environments.

310 Declarations

- 311 Ethics approval and consent to participate
- 312 Informed consent was obtained and ethical approval was given by the Ethics Committee of the Faculty
- 313 of Medicine of the Katholieke Universiteit Leuven.
- 314 Consent for publication
- 315 Not applicable
- 316 Availability of data and material
- 317 The datasets used and/or analysed during the current study are available from the corresponding author
- 318 on reasonable request.
- 319 Competing interests
- 320 The authors declare that they have no conflict of interests.
- 321 Funding
- 322 This investigation is supported by the EU research council "project ENVIRONAGE" (ERC-2012-StG
- 323 310890) and Flemish Scientific Fund (G073315N). Since its start, the East Flanders Prospective Twin
- 324 Study has been partly supported by grants from the Fund of Scientific Research Flanders and Twins, a
- 325 non-profit Association for Scientific Research in Multiple Births (Belgium).
- 326 Authors' contributions
- 327 R.V., C.D., T.S.N., M.P.Z., M.G. and E.B. designed the study; C.D. and R.J.F.L. did data collection;
- 328 T.S.N., M.P.Z., M.G. and E.B. analyzed the data; E.B. and T.S.N. wrote the first draft of paper. All
- 329 authors critically revised and approved the final version of the manuscript.
- 330 Acknowledgements
- 331 Not applicable
- 332

Reference list

- 1. Colin Mathers GS, Maya Mascarenhas, Global Health Risks: Mortality and burden of disease attributable to selected major risks. 2009, World Health Organization 70.
- 2. WHO, Global atlas on cardiovascular disease prevention and control Policies, strategies and interventions. 2011 1+4.
- 3. Fuks K, Moebus S, Hertel S, Viehmann A, Nonnemacher M, Dragano N, Mohlenkamp S, Jakobs H, Kessler C, Erbel R, et al. Long-term urban particulate air pollution, traffic noise, and arterial blood pressure. Environ Health Perspect. 2011; 119(12):1706-11.
- 4. Provost EB, Madhloum N, Int Panis L, De Boever P, and Nawrot TS. Carotid intima-media thickness, a marker of subclinical atherosclerosis, and particulate air pollution exposure: the meta-analytical evidence. PLoS One. 2015; 10(5):e0127014.
- 5. Fuks KB, Weinmayr G, Foraster M, Dratva J, Hampel R, Houthuijs D, Oftedal B, Oudin A, Panasevich S, Penell J, et al. Arterial blood pressure and long-term exposure to traffic-related air pollution: an analysis in the European Study of Cohorts for Air Pollution Effects (ESCAPE). Environ Health Perspect. 2014; 122(9):896-905.
- 6. Chen X and Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic review and meta-regression analysis. Circulation. 2008; 117(25):3171-80.
- 7. Hogberg L, Cnattingius S, Lundholm C, D'Onofrio BM, Langstrom N, and Iliadou AN. Effects of maternal smoking during pregnancy on offspring blood pressure in late adolescence. J Hypertens. 2012; 30(4):693-9.
- 8. Markevych I, Thiering E, Fuertes E, Sugiri D, Berdel D, Koletzko S, von Berg A, Bauer CP, and Heinrich J. A cross-sectional analysis of the effects of residential greenness on blood pressure in 10-year old children: results from the GINIplus and LISAplus studies. BMC Public Health. 2014; 14:477.
- 9. Agyemang C, van Hooijdonk C, Wendel-Vos W, Ujcic-Voortman JK, Lindeman E, Stronks K, and Droomers M. Ethnic differences in the effect of environmental stressors on blood pressure and hypertension in the Netherlands. BMC Public Health. 2007; 7:118.
- 10. Grazuleviciene R, Dedele A, Danileviciute A, Vencloviene J, Grazulevicius T, Andrusaityte S, Uzdanaviciute I, and Nieuwenhuijsen MJ. The influence of proximity to city parks on blood pressure in early pregnancy. Int J Environ Res Public Health. 2014; 11(3):2958-72.
- 11. Derom C, Thiery E, Peeters H, Vlietinck R, Defoort P, and Frijns JP. The East Flanders Prospective Twin Survey (EFPTS): an actual perception. Twin Research and Human Genetics. 2013; 16(1):58-63.
- 12. Loos RJ, Beunen G, Fagard R, Derom C, and Vlietinck R. The influence of zygosity and chorion type on fat distribution in young adult twins consequences for twin studies. Twin Research. 2001; 4(5):356-364.
- 13. Vlietinck R, Determination of the zygosity of twins. 1986, Leuven 1-123.
- 14. Loos RJ, Beunen G, Fagard R, Derom C, and Vlietinck R. Birth weight and body composition in young women: a prospective twin study. Am J Clin Nutr. 2002; 75(4):676-82.
- 15. Loos RJ, Fagard R, Beunen G, Derom C, and Vlietinck R. Birth weight and blood pressure in young adults: a prospective twin study. Circulation. 2001; 104(14):1633-8.
- 16. de Boo HA and Harding JE. The developmental origins of adult disease (Barker) hypothesis. Aust N Z J Obstet Gynaecol. 2006; 46(1):4-14.
- 17. Fagard RH, Celis H, Thijs L, Staessen JA, Clement DL, De Buyzere ML, and De Bacquer DA. Daytime and nighttime blood pressure as predictors of death and cause-specific cardiovascular events in hypertension. Hypertension. 2008; 51(1):55-61.
- 18. Gillman MW, Rich-Edwards JW, Rifas-Shiman SL, Lieberman ES, Kleinman KP, and Lipshultz SE. Maternal age and other predictors of newborn blood pressure. J Pediatr. 2004; 144(2):240-5.

- 19. Roe JJ, Thompson CW, Aspinall PA, Brewer MJ, Duff EI, Miller D, Mitchell R, and Clow A. Green space and stress: evidence from cortisol measures in deprived urban communities. Int J Environ Res Public Health. 2013; 10(9):4086-103.
- 20. van Dijk AE, van Eijsden M, Stronks K, Gemke RJ, and Vrijkotte TG. The association between prenatal psychosocial stress and blood pressure in the child at age 5-7 years. PLoS One. 2012; 7(8):e43548.
- 21. Fan F, Zou Y, Tian H, Zhang Y, Zhang J, Ma X, Meng Y, Yue Y, Liu K, and Dart AM. Effects of maternal anxiety and depression during pregnancy in Chinese women on children's heart rate and blood pressure response to stress. J Hum Hypertens. 2015:1-6.
- 22. Bilenko N, van Rossem L, Brunekreef B, Beelen R, Eeftens M, Hoek G, Houthuijs D, de Jongste JC, van Kempen E, Koppelman GH, et al. Traffic-related air pollution and noise and children's blood pressure: results from the PIAMA birth cohort study. Eur J Prev Cardiol. 2015; 22(1):4-12.
- 23. Sorensen M, Hoffmann B, Hvidberg M, Ketzel M, Jensen SS, Andersen ZJ, Tjonneland A, Overvad K, and Raaschou-Nielsen O. Long-term exposure to traffic-related air pollution associated with blood pressure and self-reported hypertension in a Danish cohort. Environ Health Perspect. 2012; 120(3):418-24.
- 24. Foraster M, Basagana X, Aguilera I, Rivera M, Agis D, Bouso L, Deltell A, Marrugat J, Ramos R, Sunyer J, et al. Association of long-term exposure to traffic-related air pollution with blood pressure and hypertension in an adult population-based cohort in Spain (the REGICOR study). Environ Health Perspect. 2014; 122(4):404-11.
- 25. Dong GH, Qian ZM, Xaverius PK, Trevathan E, Maalouf S, Parker J, Yang L, Liu MM, Wang D, Ren WH, et al. Association between long-term air pollution and increased blood pressure and hypertension in China. Hypertension. 2013; 61(3):578-84.
- 26. Chan SH, Van Hee VC, Bergen S, Szpiro AA, DeRoo LA, London SJ, Marshall JD, Kaufman JD, and Sandler DP. Long-Term Air Pollution Exposure and Blood Pressure in the Sister Study. Environ Health Perspect. 2015; 123(10):951-8.
- 27. Kannan S, Misra DP, Dvonch JT, and Krishnakumar A. Exposures to airborne particulate matter and adverse perinatal outcomes: a biologically plausible mechanistic framework for exploring potential effect modification by nutrition. Environ Health Perspect. 2006; 114(11):1636-42.
- 28. Foraster M, Kunzli N, Aguilera I, Rivera M, Agis D, Vila J, Bouso L, Deltell A, Marrugat J, Ramos R, et al. High blood pressure and long-term exposure to indoor noise and air pollution from road traffic. Environ Health Perspect. 2014; 122(11):1193-200.
- 29. Liu C, Fuertes E, Tiesler CM, Birk M, Babisch W, Bauer CP, Koletzko S, von Berg A, Hoffmann B, and Heinrich J. The associations between traffic-related air pollution and noise with blood pressure in children: results from the GINIplus and LISAplus studies. Int J Hyg Environ Health. 2014; 217(4-5):499-505.
- 30. Breton CV, Mack WJ, Yao J, Berhane K, Amadeus M, Lurmann F, Gilliland F, McConnell R, Hodis HN, Kunzli N, et al. Prenatal Air Pollution Exposure and Early Cardiovascular Phenotypes in Young Adults. PLoS One. 2016; 11(3):e0150825.
- 31. Bhat SK, Beilin LJ, Robinson M, Burrows S, and Mori TA. Contrasting effects of prenatal life stress on blood pressure and body mass index in young adults. J Hypertens. 2015; 33(4):711-9; discussion 719.
- 32. Cupul-Uicab LA, Skjaerven R, Haug K, Melve KK, Engel SM, and Longnecker MP. In utero exposure to maternal tobacco smoke and subsequent obesity, hypertension, and gestational diabetes among women in the MoBa cohort. Environ Health Perspect. 2012; 120(3):355-60.
- 33. Bucher BS, Tschumi S, and Simonetti GD. [Childhood's determinants for high blood pressure in adulthood]. Ther Umsch. 2012; 69(5):295-8.
- 34. Hansen TW, Jeppesen J, Rasmussen S, Ibsen H, and Torp-Pedersen C. Ambulatory blood pressure and mortality: a population-based study. Hypertension. 2005; 45(4):499-504.

- 35. Li Z, Snieder H, Harshfield GA, Treiber FA, and Wang X. A 15-year longitudinal study on ambulatory blood pressure tracking from childhood to early adulthood. Hypertens Res. 2009; 32(5):404-10.
- 36. Bijnens E, Zeegers MP, Gielen M, Kicinski M, Hageman GJ, Pachen D, Derom C, Vlietinck R, and Nawrot TS. Lower placental telomere length may be attributed to maternal residential traffic exposure; a twin study. Environment International. 2015; 79:1-7.
- 37. Lewington S, Clarke R, Qizilbash N, Peto R, and Collins R. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. Lancet. 2002; 360(9349):1903-13.

Table 1 Study population characteristics

Characteristic

Maternal	(n=146)	Adulthood	(n=278)		
Maternal age, years	27.4 ± 4.5	Age, years	20.9 ± 1.9		
Socioeconomic status: maternal educ	cation	Body mass index, kg/m ²	21.1 ± 2.8		
Low	59 (40.4)	Smokers, n	87 (31.3)		
Middle	32 (21.9)	Physical activity score	4.73 ± 2.2		
High	55 (37.7)	Potassium excretion, mmol/day	63.2 ± 28.4		
Neighborhood income, euro	19,447 ± 4,202	Sodium excretion, mmol/day	128.1 ± 59.4		
Smoking during pregnancy	21 (14.4)	Gamma-Glutamyl Transferase, U/L	16.6 ± 8.54		
		Zygosity - Chorionicity			
Birth	(n=278)	Dizygotic-Dichorial	108 (38.8)		
Gestational age, weeks	37.0 ± 2.4	Monozygotic-Dichorial	83 (29.9)		
Neonate birth weight, g	2,524 ± 497	Monozygotic-Monochorial	87 (31.3)		
Sex		Complete-pair in final study			
Male	149 (53.6)	One twin	14 (5.0)		
Female	129 (46.4)	Both twins	264 (95.0)		
Twin birth year	1978 ± 2.0	Moved since birth	181 (65.1)		

Data presented are means ± standard deviation or number (percentage)

Table 2 Distribution of the traffic and land use indicators

	Residential address at birth				Residential address at adult age							
	Geo		Percentile		Geo			Percentile				
	metric mean	5 th	25 th	50 th	75 th	95 th	metric mean	5 th	25 th	50 th	75 th	95 th
Distance to major road, m	234	23	98	266	591	1881	300	31	134	334	786	1940
Greenness, %:												
5000 m buffer	58	22	54	70	74	83	62	25	59	71	75	85
1000 m buffer	36	0	21	44	67	85	62	7	29	55	70	88
300 m buffer	17	0	0	19	43	83	41	0	2	25	54	94
Residential area, %:												
5000 m buffer	28	13	22	27	37	58	27	13	20	26	35	57
1000 m buffer	44	14	33	51	69	89	38	12	28	43	63	85
300 m buffer	63	18	51	78	98	100	60	3	46	68	87	100
Industrial area, %:												
5000 m buffer	2.80	0.00	1.08	2.90	6.33	9.01	2.48	0.00	0.79	2.67	5.41	9.08
1000 m buffer	7.72	0.00	0.00	0.00	5.52	19.6	7.14	0.00	0.00	0.00	2.43	18.47
300 m buffer, %	12.88	0.00	0.00	0.00	70.00	18.8	11.18	0.00	0.00	0.00	0.00	11.55

Figure 1: Ambulatory blood pressure (mm Hg) in association with residential greenness exposure. A) Night systolic, B) Night diastolic, C) Day systolic and D) Day diastolic blood pressure. Adjusted for sex, gestational age, birth weight, birth year, zygosity–chorionicity group, maternal age, age, smoking, physical activity, BMI, 24h sodium and potassium, gamma-glutamyl transferase, indicators of socioeconomic status (maternal education and neighborhood household income), smoking during pregnancy and noise exposure during the night. In the twins who moved during life, we additionally adjust for greenness exposure in adulthood/early-life. Vertical lines denote 95% confidence intervals. *indicates significant (p < 0.05) change in blood pressure for an IQR increase in land use indicators.

Figure 2: Ambulatory blood pressure (mm Hg) in association with distance to major road. A) Night systolic, B) Night diastolic, C) Day systolic and D) Day diastolic blood pressure. Adjusted for sex, gestational age, birth weight, birth year, zygosity–chorionicity group, maternal age, age, smoking, physical activity, BMI, 24h sodium and potassium, gamma-glutamyl transferase, indicators of socioeconomic status (maternal education and neighborhood household income), smoking during pregnancy and noise exposure during the night. In the twins who moved during life, we additionally adjust for distance to major road in adulthood/early-life. Vertical lines denote 95% confidence intervals. *indicates significant (p < 0.05) change in blood pressure for a doubling in distance.