

The role of salt for immune cell function and disease

Peer-reviewed author version

WILLEBRAND, Ralf & KLEINWIETFIELD, Markus (2018) The role of salt for immune cell function and disease. In: IMMUNOLOGY, 154 (3). p. 346-353.

DOI: 10.1111/imm.12915

Handle: <http://hdl.handle.net/1942/25961>

Article type : Review

## The role of salt for immune cell function and disease

Ralf Willebrand<sup>1</sup> & Markus Kleinewietfeld<sup>1</sup>

<sup>1</sup>VIB Laboratory of Translational Immunomodulation, Center for Inflammation Research, Biomedical Research Institute, Hasselt University, Diepenbeek, Belgium

Address for correspondence: Markus Kleinewietfeld, VIB Laboratory of Translational Immunomodulation, Center for Inflammation Research, Hasselt University, Agoralaan building C, 3590 Diepenbeek, Belgium. markus.kleinewietfeld@vib-uhasselt.be

key words: diet, Th17, Treg, autoimmunity, sodium chloride

### Abbreviations

BMDM	-	Bone marrow-derived macrophage
BMI	-	Body mass index
CNS	-	Central nervous system
DC	-	Dendritic cell
EAE	-	Experimental autoimmune encephalitis
FoxP3	-	Forkhead box protein
GF	-	Germ-free
GM-CSF	-	Granulocyte macrophage-colony stimulating factor
GvHD	-	Graft versus host disease
HSD	-	High-salt diet
ILC2	-	Group 2 innate lymphoid cell
LCFA	-	Long-chain fatty acids
LP	-	Lamina propria
LPS	-	Lipopolysaccharide

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/imm.12915

This article is protected by copyright. All rights reserved.

Accepted Article

MS	-	Multiple Sclerosis
mTOR	-	Mammalian target of rapamycin
NaCl	-	Sodium chloride
NFAT	-	Nuclear factor of activated T cells
SCFA	-	Short-chain fatty acids
SGK	-	Serum glucocorticoid-regulated kinase
Tfh	-	Follicular follicular helper T cells
T <sub>H</sub> 1	-	T helper cell 1
T <sub>H</sub> 2	-	T helper cell 2
T <sub>H</sub> 17	-	T helper cell 17
TNF- $\alpha$	-	Tumor necrosis factor- $\alpha$
Treg	-	Regulatory T cell

### **Abstract**

The immune system evolved to protect organisms from invading pathogens. A network of pro- and anti-inflammatory cell types equipped with special effector molecules guarantees efficient elimination of intruders like viruses and bacteria. However, imbalances can lead to an exceeding response of effector cells incurring autoimmune or allergic diseases. An interplay of genetic and environmental factors contributes to autoimmune diseases and recent studies provided evidence for an impact of dietary habits on the immune status and related disorders. Western societies underwent a change in lifestyle associated with changes in food consumption. Salt (sodium chloride) is one component prevalent in processed food frequently consumed in western countries. Here we summarize recent advances in understanding the mechanisms behind the effects of sodium chloride on immune cells like regulatory T cells (Tregs) and T helper (T<sub>H</sub>) 17 cells and its implication as a risk factor for several diseases.

## Introduction

Innate and adaptive immune cells are armed with a diverse arsenal of effector molecules that provide effective defence mechanism against pathogens. T cells can provide host protection via soluble mediators including cytokines. Interferon (IFN)- $\gamma$  producing T helper ( $T_H$ ) 1 cells contribute to the elimination of intracellular bacteria and viruses (1). In contrast  $T_H$ 2 cells are characterized by interleukin (IL)-4, IL-5 and IL-13 production and play important roles to combat extracellular parasites (2). However,  $T_H$ 2 cells are also associated with allergic disorders (2). IL-17 producing T helper cells have been identified as a distinct lineage and subsequently referred as  $T_H$ 17 cells (3, 4).  $T_H$ 17 cells provide protection from fungal infection and extracellular bacteria (5). However, exacerbating  $T_H$ 17 responses are also associated with autoimmune diseases (5). Altogether, T helper cells provide essential effector molecules to fight pathogens but can also contribute to immune pathology.  $CD25^+$   $FoxP3^+$  regulatory T cells (Tregs) are important modulators of immune reactions due to their capacity to suppress effector cells (6). The absence of regulatory T cells, either through deletion of the transcription factor Forkhead box protein 3 (*FoxP3*) or antibody mediated depletion in mice, results in severe autoimmunity. Further, mice with a mutation in the *FoxP3* gene (Scurfy mice) develop autoimmune syndroms (7). In humans, mutations in the *FoxP3* gene manifest as IPEX syndrome (immunodysregulation polyendocrinopathy enteropathy X-linked syndrome) (8, 9). Tregs are constitutively produced in the thymus and referred as natural Tregs (nTregs) (10). However, it has been demonstrated that Tregs can also be induced in peripheral tissues e.g. by TGF- $\beta$  and Retinoic acid mediated *FoxP3* upregulation, thus these Tregs were subsequently named induced Tregs (iTregs) (10). Furthermore, other  $FoxP3^+$   $CD4^+$  T cells with suppressive potential have been identified. For instance, the so called Type 1 regulatory T (Tr1) cells can produce large amounts of the suppressive cytokine IL-10 (11, 12). The equilibrium between effector cells and Tregs is important to maintain immunity without harmful side effects. Especially at protective barriers, like the skin, the mucosal tissue of the lung and the gastrointestinal tract, a balance between tolerance to harmless environmental antigens and immunity to pathogens needs to be maintained.

A large fraction of innate and adaptive immune cells resides in the lamina propria (LP) of the small intestine. Hence, it is of interest how these cells are influenced by dietary components and the gut microbiota. The gut microbiome comprises a total of  $>10^{14}$  microorganisms composed of  $>1000$  species of bacteria and fungi (13-15). These microorganisms are supportive in dietary food digestion. The gut microbiota and its metabolites have a profound impact on the local and systemic immune system (16, 17). Insights in how the gut microbiota shapes the immune system were derived from studies in germ free (GF) mice. GF mice have reduced numbers of  $T_H17$  cells in the lamina propria of the small intestine (18). Furthermore, GF mice are protected from experimental autoimmune encephalomyelitis (EAE) a mouse model for multiple sclerosis (MS) (19). In addition, it has been observed that the suppressive capacity of Tregs isolated from GF mice is impaired (20). A critical role for the gut microbiota on Treg modulation is further underlined by the fact that the transfer of a single polysaccharide derived from *Bacteroides fragilis* was sufficient to ameliorate disease score in mice suffering from EAE due to induction of IL-10 producing Tregs (21). In line with this notion, GF mice reconstituted with a specific mixture of bacteria showed ameliorated colitis scores attributed to Treg induction by specific members of the group Clostridium (22). In contrast, reconstitution of GF mice with segmented filamentous bacteria (SFB) promote  $T_H17$  cell development that protects against infection by pathogenic *Citrobacter rodentium* (18). A critical contribution of the microbiota to autoimmune diseases was recently underpinned by the observation that transfer of MS patients microbiota into mice could modulate EAE. GF mice reconstituted with MS patient microbiota developed more severe EAE attributed to a diminished generation of IL-10 producing regulatory T cells compared to GF mice reconstituted with microbiota derived from healthy controls (23). A finding shared by another study using GM free RR mice as recipients of MS patient microbiota. In this model, relapsing-remitting (RR) mice develop spontaneous EAE due to the expression of a transgenic MOG-specific T cell receptor. Interestingly, transfer of MS patient microbiota accelerated spontaneous EAE development in these mice as well (24).

The composition of the gut microbiota is highly influenced by the diet. Human studies have shown that the microbiota rapidly adapts to short-term dietary changes towards plant-based or animal-based nutrition (25). Specific fiber-rich diets are usually associated with enhanced generation of short chain fatty acids (SCFA), metabolized by gut commensals, that have been shown to promote immune tolerance (26, 27). For instance, feeding mice with SCFA ameliorates EAE symptoms accompanied by an expansion of Tregs and a decline of T<sub>H</sub>17 cell numbers in the small intestine, the spleen and the CNS. In contrast, long chain fatty acids (LCFA) promote intestinal and systemic T<sub>H</sub>17 cell induction leading to more severe disease score (26). Of note, a recent study has shown that dietary salt could affect the gut microbiota as well. Enhanced sodium chloride consumption depletes particularly *Lactobacillus murinus* in line with enhanced EAE development and induction of pathogenic T<sub>H</sub>17 cells in mice. Interestingly, oral gavage of *L. murinus* protected high-salt diet (HSD) fed mice from exacerbated EAE, attributed to diminished T<sub>H</sub>17 responses. This effect was suspected to be correlated to the ability of *L. murinus* to produce certain indole-metabolites that had the capability to block murine T<sub>H</sub>17 differentiation *in vitro* (28).

Intestinal T<sub>H</sub>17 cells are enabled to enter distant organs as recently demonstrated with the kaede reporter mouse (29, 30). Photoconversion of the kaede protein from green to a red fluorescence upon violet light exposure of the colon, allows tracking of intestinal cells by flow cytometry. Of note, kaede-red T<sub>H</sub>17 cells were found in the spleens of arthritis mice indicating recruitment of gut-derived T<sub>H</sub>17 cells during extraintestinal autoimmune reactions (29). A similar approach using kaede mice revealed that intestinal imprinted T<sub>H</sub>17 cells egress the gut in a S1P-receptor-1 dependent manner and migrate to kidneys via Chemokine (C-C motif) ligand Ccl20 / chemokine receptor Ccr6 interactions. In the kidneys these gut derived T<sub>H</sub>17 cells promote glomerulonephritis an autoimmune inflammation of the kidneys (30). In summary, the gut-immune cell axis seems to regulate systemic immune balance (31), hence it is important to know the impact of dietary habits on it. In western countries a trend toward consumption of processed food enriched in salt has been observed.

This article is protected by copyright. All rights reserved.

In this review we discuss how salt as one component of typical western diets shapes the gut-immune cell axis and may thereby increase the risk for many diseases.

### **Effects of salt on innate and adaptive immune cells**

The effects of salt on immune cells were studied *in vitro* by enriching the cell culture medium with additional 40mM NaCl. This mirrors the excess concentration detected in the interstitial fluids of rats fed a HSD compared to rats on control diet (32). *In vitro* differentiation of naive T cells towards T<sub>H</sub>17 cells was strongly enhanced under high-salt conditions. In line with that, high-salt induces a particularly pathogenic phenotype of T<sub>H</sub>17 cells manifested by enhanced TNF- $\alpha$  and GM-CSF production, as shown for both murine and human T cells (33, 34), enhanced expression of CCR6 and the receptor for IL-23 (IL-23R). Application of small molecule inhibitors as well as short-hairpin RNA (shRNA) mediated gene silencing, revealed the involvement of p38 mitogen-activated protein kinase (MAPK), nuclear factor of activated T cells (NFAT)-5 and serum glucocorticoid-regulated kinase-1 (SGK-1) (33, 34). SGK-1 is upregulated in salt exposed naïve T cells and mediates Forkhead box protein O (FOXO)-1 phosphorylation that leads to retinoic acid related orphan receptor (ROR)- $\gamma$ T release from the complex with Foxo1 and subsequent ROR- $\gamma$ T-dependent IL-23R expression. In line herewith, sodium effects on *in vitro* T<sub>H</sub>17 cultures were more pronounced when IL-23 was supplied (34). It has previously been shown that IL-23 augments the pathogenicity of T<sub>H</sub>17 cells (35, 36). Tregs are also sensitive to sodium chloride. It was demonstrated that *in vitro* exposure of human peripheral blood Tregs to increased sodium concentrations reduces their suppressive capacity, along with higher expression of pro-inflammatory cytokines. More specifically, salt-cultured Tregs particularly produce higher amounts of IFN- $\gamma$  and antibody-mediated neutralization of IFN- $\gamma$  restores the suppressive activity of Tregs in co-cultures with effector T cells. This was further confirmed by Treg specific IFN- $\gamma$  gene silencing using lentiviral transduced shRNA (37). The effect is dependent on SGK-1; similarly to T<sub>H</sub>17 cells,

Accepted Article

salt-exposed human Tregs upregulate SGK-1 and show enhanced phosphorylation of FOXO proteins. Silencing of *SGK1* expression in Tregs by lentiviruses containing *SGK1* targeting shRNA restores the suppressive capacity under high-salt conditions in line with reduced IFN- $\gamma$  production (37). Besides its direct effects on T cells, high-salt can also alter the function of macrophages. *In vitro* experiments with murine bone marrow-derived macrophages (BMDM) and human monocyte-derived macrophages have shown that salt increased the expression of pro-inflammatory genes while decreasing anti-inflammatory markers introducing the concept of M(Na) macrophages (38). Furthermore, salt influences the polarization of murine and human macrophages *in vitro*, promoting pro-inflammatory LPS/IFN- $\gamma$  induced M1-type macrophages while suppressing the function of IL-4-polarized regulatory M2-type macrophages (38-41). M1-type macrophages produce higher amounts of the pro-inflammatory cytokines IL-6, IL-8, TNF- $\alpha$  when differentiated under high-salt and are more efficient in bacterial clearance (38-40). In line with this, the anti-leishmania activity of M1-type macrophages is increased under excessive sodium conditions, attributed to elevated production of nitric oxid (40). In contrast, M2-type macrophage effector molecules like Fizz-1, Ym-1, Arg-1 and PD-L2 are suppressed under high-salt (38, 41). M2-type macrophages can also suppress T cell proliferation, a function that is abrogated under high-salt conditions (41). Sodium chloride may also acts as chemoattractant for macrophages. Murine macrophages are attracted by a 40mM excess NaCl concentration in *in vitro* transwell assays. Interestingly, this effect was less pronounced with 80mM excess NaCl concentration (42). Sgk-1 is required for NaCl mediated effects on T cell polarization as shown by T cell specific deletion of the *Sgk1* gene in mice (34). However, it seems to be dispensable for mediating high-salt effects on LPS stimulated macrophages, since application of a Sgk-1 specific inhibitor does not block the salt effects in BMDM (39). On the other hand, silencing or overexpression experiments for *Nfat5* revealed a critical role for this transcription factor as mediator of the salt response of murine macrophages (40). Recently, it was shown that bone marrow-derived dendritic cells react to enhanced sodium concentration in a Sgk-1- and p38-MAPK-dependent manner resulting in elevated IL-23 secretion (43). In summary, high-salt

Accepted Article

conditions could particularly affect the functionality of macrophages, dendritic cells, T<sub>H</sub>17 cells and Tregs (Table1).

## **Salt mediated immune cell modulation and its role in disease**

### ***Hypertension***

To decipher the effects of enhanced dietary salt *in vivo*, animals are fed with a diet supplemented with 4% or higher NaCl compared to control mice that receive 0,5% NaCl containing food, a classical protocol used for salt-induced hypertension since decades. The first evidence for the involvement of immune cells in Na<sup>+</sup> sensing *in vivo* was derived from feeding rats on high-salt diet. It was shown that excessive Na<sup>+</sup> accumulates in the skin where it activates macrophages to release nitric oxid and vascular endothelial growth factor (VEGF)-C (44). Depletion of macrophages impairs clearing mechanism of excessive Na<sup>+</sup> leading to hyperplasia of lymph capillaries and increased blood pressure (hypertension) (44-46). Interestingly, there is accumulating evidence that the adaptive immune system plays a critical role in hypertension as well (47). Mice with a deficient B cell and T cell compartment (*Rag*<sup>-/-</sup>) are resistant to angiotensin II-induced hypertension. Only T cell transfer into *Rag*<sup>-/-</sup> mice induces hypertension symptoms as observed in wild type controls (48). Recently, it was demonstrated that mice with CD4<sup>+</sup> T cell restricted deficiency in *Sgk1* are resistant to angiotensin II-induced hypertension and vascular inflammation (49). Similar experiments in *Il17a* deficient mice suggest involvement of T<sub>H</sub>17 cells in promoting hypertension (50). IL-17 and IFN-γ were found to be elevated in patients with symptoms of coronary artery atherosclerosis, supporting a role for T cells as mediators of vessel damage in humans (51).

### ***Autoimmune diseases***

Experimental animal studies have provided strong evidence that excessive dietary sodium could also influence induction and severity of several autoimmune disease models for e.g. colitis, systemic lupus erythematosus (SLE) and multiple sclerosis (MS). Indeed, lupus prone MRL/lpr mice, a mouse model for SLE, develop severe disease on HSD accompanied by

Accepted Article

elevated T<sub>H</sub>17 responses and increased serum titers of autoantibodies (52). Furthermore, PBMCs from SLE patients cultured *ex vivo* under high-salt conditions tend to develop more T<sub>H</sub>17 cells compared to healthy controls. The observation that HSD deteriorates lupus symptoms was confirmed by others in the same mouse model. However, this study attributes the disease outcome to an increase in the frequency of follicular helper T cells (T<sub>fh</sub>) promoting autoantibody production by B cells. Additionally, increased numbers of CD4<sup>+</sup> T cells with a signature of T<sub>fh</sub> cells, expressing the programmed death receptor (PD)-1 and the chemokine receptor CXCR5, were found in human PBMC cultures when supplied with excessive sodium chloride. The hydroxyltransferase TET2 was shown to mediate NaCl induced T<sub>fh</sub> differentiation via hypomethylation of the *spn* and *STAT5b* gene (53).

In line with this, it has been demonstrated that HSD exacerbates colitis in mice, an autoimmune inflammation of the gastrointestinal tract that can also occur in humans. This is associated with an elevated T<sub>H</sub>17 response in the LP (43, 54, 55). Additionally, HSD induces LP innate lymphoid cells (ILCs) to produce IL-17 while small intestinal Tregs produce less IL-10 and are impaired in their suppressive activity (55). Accordingly, co-transfer of *in vitro* NaCl pre-treated Tregs and effector T cells leads to accelerated colitis in *Rag*<sup>-/-</sup> recipients (37). The inflammatory response in the colitis model is dependent on GM-CSF sensing eosinophils (56). Thus, it is possible that salt incensed T<sub>H</sub>17 cells are the source of increased GM-CSF, however, this still needs to be experimentally proven. Disturbances of the gut-immune axis could be responsible for neuroinflammatory diseases. Hence, the hypothesis that dietary salt might have an impact on EAE progress has been tested experimentally in mice. Indeed, several studies have shown that HSD fed mice succumb to more severe EAE (33, 34, 39, 57-59). This is caused by a more pronounced CNS infiltration of pathogenic T<sub>H</sub>17 cells (33, 34). Moreover, mice with CD4<sup>+</sup> T cell specific deletion of the *Sgk-1* gene is protected from HSD induced EAE exacerbation. These mice failed to mount a similar T<sub>H</sub>17 response as wild type controls under HSD. The important role for T<sub>H</sub>17 cells was further corroborated with mice genetically manipulated to deplete *Sgk-1* exclusively in IL-17 producing cells (*IL-17*<sup>Cre</sup> *Sgk-1*<sup>fl/fl</sup>). These mice were protected from severe EAE

compared to control mice (34). The observation that T<sub>H</sub>17 cells produce GM-CSF, when differentiated *in vitro* under high-salt, argues for a role of this cytokine in exacerbating EAE. Supporting evidence for this hypothesis derives from the observation that GM-CSF activated microglia cells promote EAE development (60). This is further corroborated by the finding that mice with a deletion of the GM-CSF receptor in monocytes developed a less severe EAE phenotype (61). Moreover, high-salt diet affects monocyte recruitment and activation toward an pro-inflammatory phenotype in the CNS of EAE mice, supporting a role for the monocyte/macrophage lineage in committing augmented EAE scores in HSD mice. In agreement with that, transfer of BMDM pre-treated with 40mM NaCl in EAE mice exacerbated disease severity (39). Studies in humans also suggest a role for myeloid cells as possible mediators of MS disease severity induced by excessive salt intake (62, 63). Volunteers were subjected to constant dietary salt intake ranging from either 6,9 up to 12 gramm per day (g/d) for several months. Monocyte counts in the blood peaked after the 12g/d interval and decreased to normal levels with dietary salt adjustment to 6 g/d. Similarly, highest serum concentrations of the pro-inflammatory cytokines IL-6 and IL-23 were measured at the end of the 12g/d intervall. In line with this, serum IL-17 concentration was elevated in the 12g/d group (63). In summary, these studies in mice and humans suggest dietary salt as an environmental risk factor for autoimmune diseases (Figure1).

### ***Transplantation***

Deleterious effects of excessive salt intake were also observed in a murine allogeneic transplantation model. Cardiac transplants were faster rejected in MHC class II-mismatched recipients kept under HSD compared to control feeding. Immune tolerance to transplants is normally committed by functional Tregs. However, Treg numbers were reduced in allograft draining lymph nodes of HSD fed recipients. Interestingly, transplant tolerance was improved in HSD fed mice with a *Sgk-1* gene-deficient CD4<sup>+</sup> T cell compartment including Tregs (64). The *in vivo* relevance for impaired function of Tregs under high-salt was further demonstrated in a xenogeneic graft-versus-host disease (xGvHD) model. In this model

human PBMCs depleted of Tregs are transferred in immune-deficient *NOD-scid IL-2Rg<sup>null</sup>* mice where they cause severe host tissue destruction as evident by body weight loss. Hosts that receive human PBMCs and Tregs showed ameliorated disease score under normal diet. The suppressive effect of co-transferred Tregs was abrogated when recipient mice were kept on a high-salt diet. These mice showed severe disease score similar to control mice that received Treg depleted PBMCs, indicating that sodium diminishes the suppressive capacity of human Tregs *in vivo* (37).

### **Infection**

Sodium-induced amplification of effector cell activation has been proved to be a critical factor in pathogen clearance in infectious diseases. It has been demonstrated that excessive sodium triggers inflammasome activation in macrophages *in vitro*, initiating a caspase-1 dependent release of pro-inflammatory cytokines. Mice with a targeted deletion of the *caspase-1* gene were compared to wild-type mice after LPS/OVA immunization under HSD. *Caspase-1* deficient mice failed to mount a similar OVA-specific T<sub>H</sub>17 response as wild-type mice. Therefore sodium might act as an additional stimuli to amplify Toll-like receptor signalling induced by pathogen associated molecular patterns (PAMPs) (60). Accordingly, independent research data proved that HSD mice controlled *Leishmania major* infection more efficiently than control diet fed mice. This study further demonstrates that *L. major* infection leads to an accumulation of Na<sup>+</sup> in infected skin lesions. The local Na<sup>+</sup> accumulation enhances M1 macrophage activation and promoted parasite clearance (40). In contrast, skin wound repair responses are delayed under HSD that is associated with decreased M2-type macrophage activation at wound sites. NaCl excess leads to epigenetic modifications at the transcription start sites of M2-type macrophage prototype genes. It has been observed that H3K4me3 marks are reduced under high-salt conditions, suppressing IL-4/IL-13 induced M2 gene expression. Moreover, high-salt promotes metabolic reprogramming of M2-type macrophages manifested by reduced glycolysis and mitochondrial metabolic output (41).

## **Obesity & Cancer**

Obesity rates have increased in the last decades in developed countries worldwide (65). Obesity is a consequence of high caloric diet intake, enriched in sugar and saturated fatty acids (SFA), that causes diseases like metabolic syndrome characterised by insulin resistance and hypertension (66). Salt intake has been determined as a risk factor for obesity in humans (67, 68). It has been shown that elevated dietary sodium intake correlates with a high body mass index (BMI) and waist circumference (67). This was supported by the observation that patients with metabolic syndrome tend to have a higher salt intake compared to healthy controls (66). Furthermore, studies in rats have shown that HSD feeding promotes an increase of adipose tissue mass (69). Metabolic homeostasis is regulated by an immune cell circuitry involving Tregs and cells associated with a type 2 immune response e.g. M2-type macrophages, eosinophils and ILC2 cells (70-75). Interestingly, adiposity is accompanied by a decline and phenotypical changes of these cells in fat tissue (70, 72, 76-79). Obesity is associated with a shift toward a pro-inflammatory environment indicated by the dominance of M1-type macrophages (76, 78). Mice fed a HSD were shown to have higher serum levels of the pro-inflammatory cytokines TNF- $\alpha$  and IL-6 (80). Additionally, HFD fed mice also showed a bias to develop T<sub>H</sub>17 cells that manifests in more severe EAE (81). Thus, it is tempting to speculate that high-salt intake could further enhance the inflammatory environment and subsequently worsens obesity and associated diseases. However, this has to be analysed in detail in future studies.

Obesity also correlates with the development of cancer (82). People with a high BMI have a 1,5 to 1,6-fold higher risk to die from cancer (83). This was further confirmed in animal studies. Sodium chloride was shown to be a risk factor for colonic cancer development (84, 85). Gerbils that were fed a HSD succumbed to *Helicobacter pylori* induced Gastric Carcinogenesis. The authors found higher levels of *Helicobacter pylori* pro-cancerous factor cagA in HSD mice (86). Moreover, feeding mice on HSD results in pronounced gastric colonization with *Helicobacter pylori* along with alterations in mucus production favouring a pro-carcinogenic environment (87). Additive effects of dietary salt on tumor growth were also

described for chemically induced tumors in gerbils (88). The findings that HSD favors tumor growth are surprising, since sodium chloride increases the pathogenicity of helper T cells as well as Tregs, and given that T cells contribute to tumor rejection by production of cytokines e.g. TNF- $\alpha$  and IFN- $\gamma$  (89, 90).

### **Allergic Diseases**

Allergic asthma is another example of an imbalanced immune system with high incidence in the western world that incurs high costs for health insurances (91). It has been shown that western diet rather accelerates allergic diseases, whereas a Mediterranean diet has a protective effect (92, 93). Allergies are driven by an amplified type 2 immune response including T<sub>H</sub>2 cells, ILC2s, Immunoglobulin (Ig) E producing B cells, eosinophils and basophils and M2-type macrophages. It was shown that T cells are the critical source of the disease promoting IL-4/13 in the Alum/OVA mouse model of asthma (94). T cell-derived IL-4/13 is further non-redundant in protecting mice from fatal schistosomiasis (95). However, a type 2 response can also have a protective effect ameliorating autoimmunity symptoms (96). Research data related to the effect of salt on type 2 responses are limited. A recently published study shows that T<sub>H</sub>2 cell differentiation is impaired *in vitro* under high-salt conditions (57). However, *in vivo* data are not available to our knowledge yet.

### **Conclusions**

Recent research gave insights on how diet may influence the immune system. A deeper understanding of how single dietary components affect subtypes of immune cells is the milestone for developing dietary based therapies to efficiently treat diseases. Salt, a component enriched in the western diet turned out to be harmful in several aspects. We now know that salt has an impact on the function of immune cells with negative effects in onset and progression of diseases like hypertension, obesity, cancer, metabolic syndrome and autoimmune diseases (Figure1). Animal studies support a role for dietary salt intake as risk factor for EAE. Human studies linking dietary salt intake to MS are contradictory. It was

shown that MS patients with the highest salt intake have 3,95 fold higher disease scores and a 3,4 fold increased risk to develop new MS lesions (97). However, other studies failed to link dietary salt intake to increased risk of MS (98-100). Conclusions derived from questionnaires regarding the dietary habits of MS patients compared to healthy controls. Furthermore, Na<sup>+</sup> excretion was measured from 12-16 spot urine samples over 5 years to monitor dietary sodium intake. Recent data from long-term studies with volunteers subjected to constant sodium intake have shown that urine Na<sup>+</sup> concentrations oscillates between day to day measurements (101, 102). Studies using multi resonance imaging (MRI) revealed the skin interstitium and muscles as a Na<sup>+</sup> depot (40, 103-105). Hence, the methods to be applied for precisely measuring sodium intake are under debate. Moreover, the dogma that high sodium intake is balanced by enhanced fluidic intake as counterbalance mechanism has been challenged (106, 107). It has been demonstrated that high-salt feeding incurs metabolic reprogramming in liver, muscle and blood vessels tissue representing a complex mechanism to produce osmolytes for body water conservation (107). Immune cells are equipped with molecular mechanism to sense increasing salt concentrations *in vitro*, leading to cell type specific alterations of effector functions (Table1). Most of the *in vivo* studies linking high-salt intake to immune cell mediated diseases focused on T<sub>H</sub>17 dependent models. We further need to acquire more knowledge of the effects of salt on immune reactions during infections. HSD has an advantageous effects in containing *L. major* spread attributed to an amplified M1-type macrophage activation (Figure1 & Table1). T<sub>H</sub>1 responses against viral infection have not been studied under high-salt conditions to our knowledge yet.

### **Acknowledgments**

MK was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (640116), by a SALK-grant from the government of Flanders, Belgium and by an Odysseus-grant of the Research Foundation Flanders, Belgium (FWO).

Conflict-of-interest disclosure: The authors declare no competing financial interest.

## References

1. Mosmann TR, Sad S. The expanding universe of T-cell subsets: Th1, Th2 and more. *Immunol Today*. 1996;17(3):138-46.
2. Wynn TA. Type 2 cytokines: mechanisms and therapeutic strategies. *Nat Rev Immunol*. 2015;15(5):271-82.
3. Harrington LE, Hatton RD, Mangan PR, Turner H, Murphy TL, Murphy KM, et al. Interleukin 17-producing CD4<sup>+</sup> effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nat Immunol*. 2005;6(11):1123-32.
4. Park H, Li Z, Yang XO, Chang SH, Nurieva R, Wang YH, et al. A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17. *Nat Immunol*. 2005;6(11):1133-41.
5. Kleinewietfeld M, Hafler DA. The plasticity of human Treg and Th17 cells and its role in autoimmunity. *Semin Immunol*. 2013;25(4):305-12.
6. Takahashi T, Kuniyasu Y, Toda M, Sakaguchi N, Itoh M, Iwata M, et al. Immunologic self-tolerance maintained by CD25<sup>+</sup>CD4<sup>+</sup> naturally anergic and suppressive T cells: induction of autoimmune disease by breaking their anergic/suppressive state. *Int Immunol*. 1998;10(12):1969-80.
7. Brunkow ME, Jeffery EW, Hjerrild KA, Paepers B, Clark LB, Yasayko SA, et al. Disruption of a new forkhead/winged-helix protein, scurfy, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet*. 2001;27(1):68-73.
8. Bennett CL, Ochs HD. IPEX is a unique X-linked syndrome characterized by immune dysfunction, polyendocrinopathy, enteropathy, and a variety of autoimmune phenomena. *Curr Opin Pediatr*. 2001;13(6):533-8.
9. Bennett CL, Christie J, Ramsdell F, Brunkow ME, Ferguson PJ, Whitesell L, et al. The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. *Nat Genet*. 2001;27(1):20-1.
10. Ohkura N, Kitagawa Y, Sakaguchi S. Development and maintenance of regulatory T cells. *Immunity*. 2013;38(3):414-23.
11. Vieira PL, Christensen JR, Minaee S, O'Neill EJ, Barrat FJ, Boonstra A, et al. IL-10-secreting regulatory T cells do not express Foxp3 but have comparable regulatory function to naturally occurring CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cells. *J Immunol*. 2004;172(10):5986-93.
12. Mayne CG, Williams CB. Induced and natural regulatory T cells in the development of inflammatory bowel disease. *Inflamm Bowel Dis*. 2013;19(8):1772-88.
13. Booiijink CC, Zoetendal EG, Kleerebezem M, de Vos WM. Microbial communities in the human small intestine: coupling diversity to metagenomics. *Future Microbiol*. 2007;2(3):285-95.
14. Zoetendal EG, Vaughan EE, de Vos WM. A microbial world within us. *Mol Microbiol*. 2006;59(6):1639-50.
15. Zoetendal EG, Rajilic-Stojanovic M, de Vos WM. High-throughput diversity and functionality analysis of the gastrointestinal tract microbiota. *Gut*. 2008;57(11):1605-15.
16. Grigg JB, Sonnenberg GF. Host-Microbiota Interactions Shape Local and Systemic Inflammatory Diseases. *J Immunol*. 2017;198(2):564-71.
17. Belkaid Y, Hand TW. Role of the microbiota in immunity and inflammation. *Cell*. 2014;157(1):121-41.
18. Ivanov, II, Atarashi K, Manel N, Brodie EL, Shima T, Karaoz U, et al. Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell*. 2009;139(3):485-98.
19. Berer K, Mues M, Koutrolos M, Rasbi ZA, Boziki M, Johner C, et al. Commensal microbiota and myelin autoantigen cooperate to trigger autoimmune demyelination. *Nature*.

2011;479(7374):538-41.

20. Ostman S, Rask C, Wold AE, Hultkrantz S, Telemo E. Impaired regulatory T cell function in germ-free mice. *Eur J Immunol.* 2006;36(9):2336-46.
21. Ochoa-Reparaz J, Mielcarz DW, Wang Y, Begum-Haque S, Dasgupta S, Kasper DL, et al. A polysaccharide from the human commensal *Bacteroides fragilis* protects against CNS demyelinating disease. *Mucosal Immunol.* 2010;3(5):487-95.
22. Atarashi K, Tanoue T, Shima T, Imaoka A, Kuwahara T, Momose Y, et al. Induction of colonic regulatory T cells by indigenous *Clostridium* species. *Science.* 2011;331(6015):337-41.
23. Cekanaviciute E, Yoo BB, Runia TF, Debelius JW, Singh S, Nelson CA, et al. Gut bacteria from multiple sclerosis patients modulate human T cells and exacerbate symptoms in mouse models. *Proc Natl Acad Sci U S A.* 2017;114(40):10713-8.
24. Berer K, Gerdes LA, Cekanaviciute E, Jia X, Xiao L, Xia Z, et al. Gut microbiota from multiple sclerosis patients enables spontaneous autoimmune encephalomyelitis in mice. *Proc Natl Acad Sci U S A.* 2017;114(40):10719-24.
25. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature.* 2014;505(7484):559-63.
26. Haghikia A, Jorg S, Duscha A, Berg J, Manzel A, Waschbisch A, et al. Dietary Fatty Acids Directly Impact Central Nervous System Autoimmunity via the Small Intestine. *Immunity.* 2016;44(4):951-3.
27. Marino E, Richards JL, McLeod KH, Stanley D, Yap YA, Knight J, et al. Gut microbial metabolites limit the frequency of autoimmune T cells and protect against type 1 diabetes. *Nat Immunol.* 2017;18(5):552-62.
28. Wilck N, Matus MG, Kearney SM, Olesen SW, Forslund K, Bartolomaeus H, et al. Salt-responsive gut commensal modulates TH17 axis and disease. *Nature.* 2017;551(7682):585-9.
29. Morton AM, Sefik E, Upadhyay R, Weissleder R, Benoist C, Mathis D. Endoscopic photoconversion reveals unexpectedly broad leukocyte trafficking to and from the gut. *Proc Natl Acad Sci U S A.* 2014;111(18):6696-701.
30. Krebs CF, Paust HJ, Krohn S, Koyro T, Brix SR, Riedel JH, et al. Autoimmune Renal Disease Is Exacerbated by S1P-Receptor-1-Dependent Intestinal Th17 Cell Migration to the Kidney. *Immunity.* 2016;45(5):1078-92.
31. Fung TC, Olson CA, Hsiao EY. Interactions between the microbiota, immune and nervous systems in health and disease. *Nat Neurosci.* 2017;20(2):145-55.
32. Titze J, Shakibaei M, Schafflhuber M, Schulze-Tanzil G, Porst M, Schwind KH, et al. Glycosaminoglycan polymerization may enable osmotically inactive Na<sup>+</sup> storage in the skin. *Am J Physiol Heart Circ Physiol.* 2004;287(1):H203-8.
33. Kleiweietfeld M, Manzel A, Titze J, Kvakana H, Yosef N, Linker RA, et al. Sodium chloride drives autoimmune disease by the induction of pathogenic TH17 cells. *Nature.* 2013;496(7446):518-22.
34. Wu C, Yosef N, Thalhammer T, Zhu C, Xiao S, Kishi Y, et al. Induction of pathogenic TH17 cells by inducible salt-sensing kinase SGK1. *Nature.* 2013;496(7446):513-7.
35. El-Behi M, Ciric B, Dai H, Yan Y, Cullimore M, Safavi F, et al. The encephalitogenicity of T(H)17 cells is dependent on IL-1- and IL-23-induced production of the cytokine GM-CSF. *Nat Immunol.* 2011;12(6):568-75.
36. Gaffen SL, Jain R, Garg AV, Cua DJ. The IL-23-IL-17 immune axis: from mechanisms to therapeutic testing. *Nat Rev Immunol.* 2014;14(9):585-600.
37. Hernandez AL, Kitz A, Wu C, Lowther DE, Rodriguez DM, Vudattu N, et al. Sodium chloride inhibits the suppressive function of FOXP3<sup>+</sup> regulatory T cells. *J Clin Invest.* 2015;125(11):4212-22.
38. Zhang WC, Zheng XJ, Du LJ, Sun JY, Shen ZX, Shi C, et al. High salt primes a specific activation state of macrophages, M(Na). *Cell Res.* 2015;25(8):893-910.
39. Hucke S, Eschborn M, Liebmann M, Herold M, Freise N, Engbers A, et al. Sodium chloride promotes pro-inflammatory macrophage polarization thereby aggravating CNS autoimmunity. *J*

Autoimmun. 2016;67:90-101.

40. Jantsch J, Schatz V, Friedrich D, Schroder A, Kopp C, Siegert I, et al. Cutaneous Na<sup>+</sup> storage strengthens the antimicrobial barrier function of the skin and boosts macrophage-driven host defense. *Cell Metab.* 2015;21(3):493-501.
41. Binger KJ, Gebhardt M, Heinig M, Rintisch C, Schroeder A, Neuhofer W, et al. High salt reduces the activation of IL-4- and IL-13-stimulated macrophages. *J Clin Invest.* 2015;125(11):4223-38.
42. Muller S, Quast T, Schroder A, Hucke S, Klotz L, Jantsch J, et al. Salt-dependent chemotaxis of macrophages. *PLoS One.* 2013;8(9):e73439.
43. Tubbs AL, Liu B, Rogers TD, Sartor RB, Miao EA. Dietary Salt Exacerbates Experimental Colitis. *J Immunol.* 2017;199(3):1051-9.
44. Machnik A, Neuhofer W, Jantsch J, Dahlmann A, Tammela T, Machura K, et al. Macrophages regulate salt-dependent volume and blood pressure by a vascular endothelial growth factor-C-dependent buffering mechanism. *Nat Med.* 2009;15(5):545-52.
45. Wiig H, Schroder A, Neuhofer W, Jantsch J, Kopp C, Karlsen TV, et al. Immune cells control skin lymphatic electrolyte homeostasis and blood pressure. *J Clin Invest.* 2013;123(7):2803-15.
46. Machnik A, Dahlmann A, Kopp C, Goss J, Wagner H, van Rooijen N, et al. Mononuclear phagocyte system depletion blocks interstitial tonicity-responsive enhancer binding protein/vascular endothelial growth factor C expression and induces salt-sensitive hypertension in rats. *Hypertension.* 2010;55(3):755-61.
47. Trott DW, Harrison DG. The immune system in hypertension. *Adv Physiol Educ.* 2014;38(1):20-4.
48. Guzik TJ, Hoch NE, Brown KA, McCann LA, Rahman A, Dikalov S, et al. Role of the T cell in the genesis of angiotensin II induced hypertension and vascular dysfunction. *J Exp Med.* 2007;204(10):2449-60.
49. Norlander AE, Saleh MA, Pandey AK, Itani HA, Wu J, Xiao L, et al. A salt-sensing kinase in T lymphocytes, SGK1, drives hypertension and hypertensive end-organ damage. *JCI Insight.* 2017;2(13).
50. Madhur MS, Lob HE, McCann LA, Iwakura Y, Blinder Y, Guzik TJ, et al. Interleukin 17 promotes angiotensin II-induced hypertension and vascular dysfunction. *Hypertension.* 2010;55(2):500-7.
51. Eid RE, Rao DA, Zhou J, Lo SF, Ranjbaran H, Gallo A, et al. Interleukin-17 and interferon-gamma are produced concomitantly by human coronary artery-infiltrating T cells and act synergistically on vascular smooth muscle cells. *Circulation.* 2009;119(10):1424-32.
52. Yang X, Yao G, Chen W, Tang X, Feng X, Sun L. Exacerbation of lupus nephritis by high sodium chloride related to activation of SGK1 pathway. *Int Immunopharmacol.* 2015;29(2):568-73.
53. Wu H, Huang X, Qiu H, Zhao M, Liao W, Yuan S, et al. High salt promotes autoimmunity by TET2-induced DNA demethylation and driving the differentiation of Tfh cells. *Sci Rep.* 2016;6:28065.
54. Monteleone I, Marafini I, Dinallo V, Di Fusco D, Troncone E, Zorzi F, et al. Sodium chloride-enriched Diet Enhanced Inflammatory Cytokine Production and Exacerbated Experimental Colitis in Mice. *J Crohns Colitis.* 2017;11(2):237-45.
55. Wei Y, Lu C, Chen J, Cui G, Wang L, Yu T, et al. High salt diet stimulates gut Th17 response and exacerbates TNBS-induced colitis in mice. *Oncotarget.* 2017;8(1):70-82.
56. Griseri T, Arnold IC, Pearson C, Krausgruber T, Schiering C, Franchini F, et al. Granulocyte Macrophage Colony-Stimulating Factor-Activated Eosinophils Promote Interleukin-23 Driven Chronic Colitis. *Immunity.* 2015;43(1):187-99.
57. Hammer A, Schliep A, Jorg S, Haghikia A, Gold R, Kleinewietfeld M, et al. Impact of combined sodium chloride and saturated long-chain fatty acid challenge on the differentiation of T helper cells in neuroinflammation. *J Neuroinflammation.* 2017;14(1):184.
58. Jorg S, Kissel J, Manzel A, Kleinewietfeld M, Haghikia A, Gold R, et al. High salt drives Th17 responses in experimental autoimmune encephalomyelitis without impacting myeloid dendritic

cells. *Exp Neurol.* 2016;279:212-22.

59. Kremontsov DN, Case LK, Hickey WF, Teuscher C. Exacerbation of autoimmune neuroinflammation by dietary sodium is genetically controlled and sex specific. *FASEB J.* 2015;29(8):3446-57.
60. Ponomarev ED, Shriver LP, Maresz K, Pedras-Vasconcelos J, Verthelyi D, Dittel BN. GM-CSF production by autoreactive T cells is required for the activation of microglial cells and the onset of experimental autoimmune encephalomyelitis. *J Immunol.* 2007;178(1):39-48.
61. Croxford AL, Lanzinger M, Hartmann FJ, Schreiner B, Mair F, Pelczar P, et al. The Cytokine GM-CSF Drives the Inflammatory Signature of CCR2+ Monocytes and Licenses Autoimmunity. *Immunity.* 2015;43(3):502-14.
62. Zhou X, Zhang L, Ji WJ, Yuan F, Guo ZZ, Pang B, et al. Variation in dietary salt intake induces coordinated dynamics of monocyte subsets and monocyte-platelet aggregates in humans: implications in end organ inflammation. *PLoS One.* 2013;8(4):e60332.
63. Yi B, Titze J, Rykova M, Feuerecker M, Vassilieva G, Nichiporuk I, et al. Effects of dietary salt levels on monocytic cells and immune responses in healthy human subjects: a longitudinal study. *Transl Res.* 2015;166(1):103-10.
64. Safa K, Otori S, Borges TJ, Uehara M, Batal I, Shimizu T, et al. Salt Accelerates Allograft Rejection through Serum- and Glucocorticoid-Regulated Kinase-1-Dependent Inhibition of Regulatory T Cells. *J Am Soc Nephrol.* 2015;26(10):2341-7.
65. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2014;384(9945):766-81.
66. Hoffmann IS, Cubeddu LX. Salt and the metabolic syndrome. *Nutr Metab Cardiovasc Dis.* 2009;19(2):123-8.
67. Ma Y, He FJ, MacGregor GA. High salt intake: independent risk factor for obesity? *Hypertension.* 2015;66(4):843-9.
68. Larsen SC, Angquist L, Sorensen TI, Heitmann BL. 24h urinary sodium excretion and subsequent change in weight, waist circumference and body composition. *PLoS One.* 2013;8(7):e69689.
69. Fonseca-Alaniz MH, Brito LC, Borges-Silva CN, Takada J, Andreotti S, Lima FB. High dietary sodium intake increases white adipose tissue mass and plasma leptin in rats. *Obesity (Silver Spring).* 2007;15(9):2200-8.
70. Feuerer M, Herrero L, Cipolletta D, Naaz A, Wong J, Nayer A, et al. Lean, but not obese, fat is enriched for a unique population of regulatory T cells that affect metabolic parameters. *Nat Med.* 2009;15(8):930-9.
71. Cipolletta D, Feuerer M, Li A, Kamei N, Lee J, Shoelson SE, et al. PPAR-gamma is a major driver of the accumulation and phenotype of adipose tissue Treg cells. *Nature.* 2012;486(7404):549-53.
72. Wu D, Molofsky AB, Liang HE, Ricardo-Gonzalez RR, Jouihan HA, Bando JK, et al. Eosinophils sustain adipose alternatively activated macrophages associated with glucose homeostasis. *Science.* 2011;332(6026):243-7.
73. Brestoff JR, Kim BS, Saenz SA, Stine RR, Monticelli LA, Sonnenberg GF, et al. Group 2 innate lymphoid cells promote beiging of white adipose tissue and limit obesity. *Nature.* 2015;519(7542):242-6.
74. Lee MW, Odegaard JI, Mukundan L, Qiu Y, Molofsky AB, Nussbaum JC, et al. Activated type 2 innate lymphoid cells regulate beige fat biogenesis. *Cell.* 2015;160(1-2):74-87.
75. Nguyen KD, Qiu Y, Cui X, Goh YP, Mwangi J, David T, et al. Alternatively activated macrophages produce catecholamines to sustain adaptive thermogenesis. *Nature.* 2011;480(7375):104-8.
76. Lumeng CN, Bodzin JL, Saltiel AR. Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *J Clin Invest.* 2007;117(1):175-84.

77. Lumeng CN, DelProposto JB, Westcott DJ, Saltiel AR. Phenotypic switching of adipose tissue macrophages with obesity is generated by spatiotemporal differences in macrophage subtypes. *Diabetes*. 2008;57(12):3239-46.
78. Lumeng CN, Deyoung SM, Bodzin JL, Saltiel AR. Increased inflammatory properties of adipose tissue macrophages recruited during diet-induced obesity. *Diabetes*. 2007;56(1):16-23.
79. Lumeng CN, Deyoung SM, Saltiel AR. Macrophages block insulin action in adipocytes by altering expression of signaling and glucose transport proteins. *Am J Physiol Endocrinol Metab*. 2007;292(1):E166-74.
80. Bastard JP, Jardel C, Bruckert E, Blondy P, Capeau J, Laville M, et al. Elevated levels of interleukin 6 are reduced in serum and subcutaneous adipose tissue of obese women after weight loss. *J Clin Endocrinol Metab*. 2000;85(9):3338-42.
81. Winer S, Paltser G, Chan Y, Tsui H, Engleman E, Winer D, et al. Obesity predisposes to Th17 bias. *Eur J Immunol*. 2009;39(9):2629-35.
82. Font-Burgada J, Sun B, Karin M. Obesity and Cancer: The Oil that Feeds the Flame. *Cell Metab*. 2016;23(1):48-62.
83. Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N Engl J Med*. 2003;348(17):1625-38.
84. Dungal N, Sigurjonsson J. Gastric cancer and diet. A pilot study on dietary habits in two districts differing markedly in respect of mortality from gastric cancer. *Br J Cancer*. 1967;21(2):270-6.
85. Kelley JR, Duggan JM. Gastric cancer epidemiology and risk factors. *J Clin Epidemiol*. 2003;56(1):1-9.
86. Gaddy JA, Radin JN, Loh JT, Zhang F, Washington MK, Peek RM, Jr., et al. High dietary salt intake exacerbates *Helicobacter pylori*-induced gastric carcinogenesis. *Infect Immun*. 2013;81(6):2258-67.
87. Fox JG, Dangler CA, Taylor NS, King A, Koh TJ, Wang TC. High-salt diet induces gastric epithelial hyperplasia and parietal cell loss, and enhances *Helicobacter pylori* colonization in C57BL/6 mice. *Cancer Res*. 1999;59(19):4823-8.
88. Kato S, Tsukamoto T, Mizoshita T, Tanaka H, Kumagai T, Ota H, et al. High salt diets dose-dependently promote gastric chemical carcinogenesis in *Helicobacter pylori*-infected Mongolian gerbils associated with a shift in mucin production from glandular to surface mucous cells. *Int J Cancer*. 2006;119(7):1558-66.
89. Kammertoens T, Friese C, Arina A, Idel C, Briesemeister D, Rothe M, et al. Tumour ischaemia by interferon-gamma resembles physiological blood vessel regression. *Nature*. 2017;545(7652):98-102.
90. Schuler T, Blankenstein T. Cutting edge: CD8+ effector T cells reject tumors by direct antigen recognition but indirect action on host cells. *J Immunol*. 2003;170(9):4427-31.
91. Accordini S, Corsico AG, Braggion M, Gerbase MW, Gislason D, Gulsvik A, et al. The cost of persistent asthma in Europe: an international population-based study in adults. *Int Arch Allergy Immunol*. 2013;160(1):93-101.
92. Julia V, Macia L, Dombrowicz D. The impact of diet on asthma and allergic diseases. *Nat Rev Immunol*. 2015;15(5):308-22.
93. Nurmatov U, Devereux G, Sheikh A. Nutrients and foods for the primary prevention of asthma and allergy: systematic review and meta-analysis. *J Allergy Clin Immunol*. 2011;127(3):724-33 e1-30.
94. Oeser K, Maxeiner J, Symowski C, Stassen M, Voehringer D. T cells are the critical source of IL-4/IL-13 in a mouse model of allergic asthma. *Allergy*. 2015;70(11):1440-9.
95. Schwartz C, Oeser K, Prazeres da Costa C, Layland LE, Voehringer D. T cell-derived IL-4/IL-13 protects mice against fatal *Schistosoma mansoni* infection independently of basophils. *J Immunol*. 2014;193(7):3590-9.
96. Eberl G. Immunity by equilibrium. *Nat Rev Immunol*. 2016;16(8):524-32.
97. Farez MF, Fiol MP, Gaitan MI, Quintana FJ, Correale J. Sodium intake is associated with

- increased disease activity in multiple sclerosis. *J Neurol Neurosurg Psychiatry*. 2015;86(1):26-31.
98. Nourbakhsh B, Graves J, Casper TC, Lulu S, Waldman A, Belman A, et al. Dietary salt intake and time to relapse in paediatric multiple sclerosis. *J Neurol Neurosurg Psychiatry*. 2016;87(12):1350-3.
99. McDonald J, Graves J, Waldman A, Lotze T, Schreiner T, Belman A, et al. A case-control study of dietary salt intake in pediatric-onset multiple sclerosis. *Mult Scler Relat Disord*. 2016;6:87-92.
100. Cortese M, Yuan C, Chitnis T, Ascherio A, Munger KL. No association between dietary sodium intake and the risk of multiple sclerosis. *Neurology*. 2017.
101. Lerchl K, Rakova N, Dahlmann A, Rauh M, Goller U, Basner M, et al. Agreement between 24-hour salt ingestion and sodium excretion in a controlled environment. *Hypertension*. 2015;66(4):850-7.
102. Rakova N, Juttner K, Dahlmann A, Schroder A, Linz P, Kopp C, et al. Long-term space flight simulation reveals infradian rhythmicity in human Na(+) balance. *Cell Metab*. 2013;17(1):125-31.
103. Kopp C, Beyer C, Linz P, Dahlmann A, Hammon M, Jantsch J, et al. Na<sup>+</sup> deposition in the fibrotic skin of systemic sclerosis patients detected by <sup>23</sup>Na-magnetic resonance imaging. *Rheumatology (Oxford)*. 2017;56(4):674.
104. Wang P, Deger MS, Kang H, Ikizler TA, Titze J, Gore JC. Sex differences in sodium deposition in human muscle and skin. *Magn Reson Imaging*. 2017;36:93-7.
105. Titze J, Lang R, Ilies C, Schwind KH, Kirsch KA, Dietsch P, et al. Osmotically inactive skin Na<sup>+</sup> storage in rats. *Am J Physiol Renal Physiol*. 2003;285(6):F1108-17.
106. Rakova N, Kitada K, Lerchl K, Dahlmann A, Birukov A, Daub S, et al. Increased salt consumption induces body water conservation and decreases fluid intake. *J Clin Invest*. 2017;127(5):1932-43.
107. Kitada K, Daub S, Zhang Y, Klein JD, Nakano D, Pedchenko T, et al. High salt intake reprioritizes osmolyte and energy metabolism for body fluid conservation. *J Clin Invest*. 2017;127(5):1944-59.

Table 1: Effects of sodium chloride on different immune cells

<i>celltype</i>	<i>effector function under high-salt</i>	<i>molecular mechanism</i>	<i>reference</i>
M1-type macrophage	↑	p38MAPK, NFAT-5	(38, 40)
M2-type macrophage	↓	mTOR, Akt	(41)
Treg	↓	Sgk-1	(37)
T <sub>H</sub> 17	↑	p38MAPK, NFAT-5, Sgk-1	(33, 34)
T <sub>H</sub> 2	↓	N.D.	(57)
T <sub>H</sub> 1	↑	N.D.	(33)
Tfh	↑	TET2	(53)
DC	↑	p38MAPK, Sgk-1	(43)

N.D. not determined

### Figure legend

#### Figure1: Effects of increased sodium chloride consumption on health.

High-salt diet (HSD) is considered a risk factor for increased blood pressure, atherosclerosis (blood vessel damage), obesity, cancer, and autoimmunity, while it could negatively impact wound healing processes. In contrast, HSD could enhance protection against parasite infections.

