

A non-commutative Bertini theorem

Non Peer-reviewed author version

Rennemo, Jorgen Vold; Segal, Ed & VAN DEN BERGH, Michel (2019) A non-commutative Bertini theorem. In: JOURNAL OF NONCOMMUTATIVE GEOMETRY, 13(2), p. 609-616.

DOI: 10.4171/JNCG/334

Handle: <http://hdl.handle.net/1942/29172>

A NON-COMMUTATIVE BERTINI THEOREM

JØRGEN VOLD RENNEMO, ED SEGAL, AND MICHEL VAN DEN BERGH

ABSTRACT. We prove a version of the classical ‘generic smoothness’ theorem with smooth varieties replaced by non-commutative resolutions of singular varieties. This in particular implies a non-commutative version of the Bertini theorem.

1. INTRODUCTION

Let k be a field of characteristic 0 and let X be a smooth variety over k , equipped with a morphism $f : X \rightarrow \mathbb{P}^n$ for some n . The classical Bertini Theorem, in one of its formulations, states that for a general hyperplane $H \subset \mathbb{P}^n$ the fibre-product $X_H = X \times_{\mathbb{P}^n} H$ will still be smooth. The aim of this short note is to establish a non-commutative analogue of this theorem, where instead of assuming X is smooth, we assume it comes provided with a *non-commutative resolution*.

For an affine variety $X = \operatorname{Spec} S$, we use the following definition of non-commutative resolution.

Definition 1.1 ([VdB04, ŠVdB15]). *Let S be a normal Noetherian domain. A non-commutative resolution of S is an S -algebra of finite global dimension of the form $A = \operatorname{End}_S(M)$ where M is a non-zero finitely generated reflexive S -module. The resolution is said to be crepant if in addition S is Gorenstein and A is a maximal Cohen–Macaulay S -module.*

The idea is that the category $A\text{-mod}$ should be a reasonable substitute for the category of coherent sheaves on a geometric resolution of $\operatorname{Spec} S$. The most important property is the requirement that A has finite global dimension, this is ‘smoothness’ of the resolution. The fact that A is ‘birational’ to S is also encoded in this definition: M must be locally-free (and non-zero) over some Zariski open subset $U \subset \operatorname{Spec} S$, then $A_U = \operatorname{End}(M_U)$ is a trivial Azumaya algebra and is Morita equivalent to S_U . In fact if the resolution is crepant one can prove that M must be locally-free over the smooth locus in $\operatorname{Spec}(S)$. This is no longer true for non-crepant resolutions, see [DII⁺16].

We abbreviate *non-commutative resolution* and *non-commutative crepant resolution* to NCR and NCCR respectively. When X is not affine, we simply replace the single algebra A with a sheaf of algebras.

Definition 1.2. *Let X be a normal variety. An NCR (resp. NCCR) of X is a coherent sheaf of algebras A on X , such that for any affine open subset $U = \operatorname{Spec} S \subset X$, the algebra $A|_U$ is an NCR (resp. NCCR) of S .*

Our path to the Bertini theorem follows a standard approach, similar to the commutative case, and passes through some other results which are interesting in their own right. The starting point is a non-commutative version of “generic smoothness”: the geometrically intuitive result that if $f : X \rightarrow Y$ is a map of varieties and X is nonsingular, then for a generic point $y \in Y$ the fibre $f^{-1}(y)$ is nonsingular.

Theorem 1.3. *Let k be a field of characteristic zero, let R be a finitely-generated Noetherian domain over k , and let S be a commutative finitely-generated normal R -algebra. Let A be a non-commutative resolution of S . Then for a general point $p \in \operatorname{Spec}(R)$, the algebra $A_{k(p)}$ is an NCR of $S_{k(p)}$, which is moreover crepant if A is.*

Here $k(p)$ denotes the residue field at p . When we say ‘for a general point’ here we mean that there is a Zariski open subset $U \subset \operatorname{Spec}(R)$ such that $A_{k(p)}$ is an NCR for every $p \in U$.

This theorem is the technical contribution of this paper, our other results are easy corollaries. Its proof is not difficult, but neither is it purely formal, since it does require that the ground field k has characteristic 0. Indeed, without further assumptions, none of the results are true in characteristic p even in the commutative setting. There is an elementary and well-known counterexample: set $A = S = k[x]$ and $R = k[t]$, and consider the homomorphism $t \mapsto x^p$. If $\operatorname{char}(k) = p$ then for every $y \in \operatorname{Spec} R$ the fibre A_y has infinite global dimension.

From Theorem 1.3, we easily deduce an ‘affine non-commutative Bertini’s theorem’. Stating an affine version of Bertini’s theorem takes a little care, even for complex varieties. There are examples of resolutions $\pi : X \rightarrow \operatorname{Spec} S$ where S is local, such that for any $f \in S$ lying in the maximal ideal the slice $\{\pi^*f = 0\}$ is singular.

Theorem 1.4. *Let S be a normal Noetherian domain over a field k of characteristic zero, and let A be an NCR of S . Fix a finite-dimensional k -vector space $V \subset S$ with $1 \in V$. Then for a general element $f \in V$, the algebra A/fA is an NCR of S/fS , which is crepant if A is.*

Finally, we turn to our main motivation, which is the non-commutative version of the projective Bertini theorem.

Let (X, A) be a normal k -variety equipped with a non-commutative resolution, and let $f : X \rightarrow \mathbb{P}^n$ be a morphism. Given a hyperplane $H \subset \mathbb{P}^n$, the fibre-product X_H carries a sheaf of algebras A_H by restriction.

Theorem 1.5. *For a general hyperplane $H \subset \mathbb{P}^n$, the pair (X_H, A_H) is a non-commutative resolution of X_H , which is crepant if (X, A) is.*

The first- and second-named authors came across this question in the context of a specific NCCR of a Pfaffian variety [RS16]; see that paper for an application of these results.

No doubt these results still hold if we replace A with an appropriate sheaf of smooth categories (or dg-categories) \mathcal{A} on X , rather than assuming that $\mathcal{A} = A\text{-mod}$ as we do here. This would allow for example modules over a sheaf of DGAs, or modules twisted by a gerbe. However, we didn’t feel the need to work in that level of generality for this paper.

1.1. Acknowledgements. We thank Michael Wemyss for helpful conversations. JVR would like to acknowledge the contribution of Icelandair, who stranded him in Reykjavik for a very productive 24 hours.

2. PROOF OF THEOREMS

The bulk of this (rather skinny) paper is the proof of the following proposition:

Proposition 2.1. *Let R be a Noetherian domain whose fraction field $K(R)$ is of characteristic 0, let S be a finitely generated commutative R -algebra, and suppose A is an S -algebra which is finite rank as an S -module.*

If A has finite global dimension, then for a generic $p \in \operatorname{Spec} R$, the algebra $A_{k(p)}$ has finite global dimension.

One can compare this to Schofield's result that for algebras which are finite-dimensional over a field, having finite global dimension is a Zariski open condition [Sch85].

Our proof works by flipping between 'finite global dimension' and the following property of 'non-commutative smoothness', whose definition appears to be due to Kontsevich. Recall that a module is perfect if it has finite projective dimension.

Definition 2.2. Let R be a commutative ring, and let A be an R -algebra. We declare that A is smooth over R if A is flat over R and A is perfect as a module over $A^{op} \otimes_R A$.

In other words, if we work relative to R then the diagonal bimodule is perfect. The flatness assumption is just to avoid needing to work with the derived tensor product $A^{op} \otimes_R^L A$.

Proof of Proposition 2.1. Here we just give the outline of the argument, the necessary supporting lemmas are in Section 3. Let $K = K(R)$ and let \overline{K} be its algebraic closure. We prove our claim via a chain of implications $(i) \Rightarrow (i+1)$. Observe that all the algebras involved, A, A^{op}, A_K, \dots are finite rank over a commutative Noetherian ring so they are Noetherian (Lemma 3.1).

- (1) A has finite global dimension.
- (2) A_K has finite global dimension.

This is Lemma 3.2.

- (3) $A_{\overline{K}}$ has finite global dimension.

This follows from Lemma 3.3 – the extension \overline{K}/K is separable because $\operatorname{char}(K) = 0$.

- (4) $A_{\overline{K}}$ is smooth over \overline{K} .

By Lemma 3.4 the algebra $A_{\overline{K}}^{op}$ has finite global dimension. Then $A_{\overline{K}}^{op} \otimes_{\overline{K}} A_{\overline{K}}$ also has finite global dimension by Lemma 3.5, in particular the diagonal bimodule is perfect.

- (5) A_K is smooth over K .

This is Lemma 3.6.

- (6) There is an $f \in R$ such that A_f is smooth over R_f .

By Corollary 3.9 (with $T = S \otimes_R S$, $\Lambda = A^{op} \otimes_R A$ and $M = A$) there is an $f \in R$ such that A_f is perfect over $A_f^{op} \otimes_{R_f} A_f$. By generic freeness (Lemma 3.7) we may choose f so that A_f is flat over R_f , and so A_f is smooth over R_f .

- (7) $A_{k(p)}$ is smooth over $k(p)$ for a general $p \in \operatorname{Spec} S$.

Take f as above and let $U = \operatorname{Spec}(R_f)$. Take a bounded resolution of A_f by finitely generated projective $A_f^{op} \otimes_{R_f} A_f$ modules. By shrinking U further we may assume that all these modules are flat over U , then restricting to $p \in U$ we have a bounded projective resolution of $A_{k(p)}$.

- (8) $A_{k(p)}$ has finite global dimension for a general $p \in \operatorname{Spec} S$.

We can get a projective resolution of any $A_{k(p)}$ -module by tensoring it with our projective resolution of the diagonal $A_{k(p)}$ -bimodule.

□

Assume now that we are in the setting of Theorem 1.3:

- k is a field of characteristic zero,
- R is a finitely-generated Noetherian domain over k ,
- S is a commutative finitely-generated normal algebra over R , and

- M is a reflexive S -module such that $A = \text{End}_S(M)$ is an NCR of S .

We want to prove that for a general point $p \in \text{Spec } R$, the algebra $A_{k(p)}$ is an NCR of $S_{k(p)}$, which is crepant if A is. We've already discussed the most important aspect – the finiteness of the global dimension – but there are a few remaining details.

Lemma 2.3. *For a general point $p \in \text{Spec } R$, $S_{k(p)}$ is normal, $M_{k(p)}$ is a reflexive $S_{k(p)}$ -module, and $A_{k(p)} = \text{End}_{S_{k(p)}}(M_{k(p)})$.*

If S is Gorenstein, then so is $S_{k(p)}$, and if A is a maximal Cohen–Macaulay S -module, then $A_{k(p)}$ is a maximal Cohen–Macaulay $S_{k(p)}$ -module.

Proof. Take a presentation for M , i.e. let $M = \text{cok } \phi$ for $\phi: S^m \rightarrow S^n$. The fact that $A = \text{End}_S(M)$ means that $A = \ker(\phi^T \otimes \text{id}_M)$, so there is an exact sequence

$$0 \rightarrow A \rightarrow M^n \rightarrow M^m \rightarrow N := \text{cok}(\phi^T \otimes \text{id}_M).$$

By generic freeness (Lemma 3.7), we may assume (after passing to an open set in $\text{Spec } R$) that A , M and N are flat R -modules. It follows that $A_{k(p)}$ is the kernel of $M_{k(p)}^n \rightarrow M_{k(p)}^m$, which means precisely that $A_{k(p)} = \text{End}_{S_{k(p)}}(M_{k(p)})$.

The other properties are all consequences of [FOV99, Thm. 3.3.10, Thm. 3.3.15]. \square

Proof of Theorem 1.3. Combine Lemma 2.3 and Prop. 2.1. \square

From here our ‘non-commutative Bertini’ theorems follow easily.

Proof of Theorem 1.4. Let S be an normal, finitely-generated k -algebra and let A be an NCR of S . Fix $V = \langle 1, f_1, \dots, f_n \rangle \subset S$. Now set $R = k[x_0, \dots, x_n]$, and let

$$S' = \frac{S \otimes_k k[x_0, \dots, x_n]}{(x_0 + x_1 f_1 + \dots + x_n f_n)} \cong S \otimes_k k[x_1, \dots, x_n].$$

Then S' is an integral normal algebra over R , and $A' = A \otimes_S S'$ is an NCR of S' . Given a point $f \in V = \text{Spec } R$, the fibres of S and A over f are S/fS and A/fA . Hence by Thm. 1.3, for a general element $f \in V$ the algebra A/fA is an NCR of S/fS , and it's an NCCR if A is an NCCR. \square

Proof of Theorem 1.5. A hyperplane in \mathbb{P}^n is a divisor defined by an equation $\sum_{i=0}^n a_i x_i$, with $a_i \in k$, so the result follows from Theorem 1.4 by seeing \mathbb{P}^n as a union of $n+1$ copies of \mathbb{A}^n . \square

3. TECHNICAL LEMMAS

In this section we provide the lemmas necessary for the proof of Proposition 2.1. The argument is spelled out in detail for the benefit of those readers who are not very familiar with non-commutative algebras (the first two authors of this paper themselves belonging to this group), and in the hope of making it clear where each hypothesis is used.

Lemma 3.1. *Let S be a Noetherian commutative ring, and let A be an S -algebra which has finite rank as an S -module. Then A is Noetherian.*

Proof. Any ideal of A must be a finitely-generated S -module, hence a finitely-generated A -module. \square

Lemma 3.2. *Let A be an algebra over an integral domain R such that A has finite global dimension. If $A_{K(R)}$ is Noetherian then it has finite global dimension.*

Proof. Let $K = K(R)$. Every finitely generated A_K -module M is the cokernel of some $\phi: A_K^m \rightarrow A_K^n$. Multiplying by denominators, we can ensure that ϕ is the localisation of a map $\phi': A^m \rightarrow A^n$, and so setting $M' = \text{cok}(\phi')$ we find $M'_K = M$. Since M' is perfect, so is M . \square

We say that an infinite field extension L/K is separable if it is algebraic and every finite subextension is separable.

Lemma 3.3. *Let A be an algebra over a field K . Let L be a separable field extension of K , and let $A_L = A \otimes_K L$. If A has finite global dimension, and A_L is Noetherian, then A_L has finite global dimension.*

This is mostly contained in [JL82, Thm. 2.4], but we give a complete proof here for convenience.

Proof. Say A has global dimension n , and let M be a finite-rank A_L -module. Since A_L is Noetherian, M is finitely presented, and hence defined over some finite extension $L' \subset L$ of K , i.e. $M = M' \otimes_{L'} L$ for an $A_{L'}$ -module M' . Separability implies that the map $L' \otimes_K L' \rightarrow L'$ splits as a map of L' -bimodules (see e.g. [Row88, Thm. 5.3.7, Prop. 5.3.16]), so M' is a direct $A_{L'}$ -module summand of

$$L' \otimes_K L' \otimes_{L'} M' = L' \otimes_K M' = A_{L'} \otimes_A M'.$$

We thus get the inequality of projective dimensions

$$\text{pd}_{A_{L'}}(M') \leq \text{pd}_{A_{L'}}(A_{L'} \otimes_A M') \leq \text{pd}_A(M') \leq n. \quad \square$$

Lemma 3.4. [Aus55, Cor. 5] *If both A and A^{op} are Noetherian, and A has finite global dimension, then A^{op} has finite global dimension.*

Recall that a *polynomial identity ring* is a ring A satisfying a polynomial identity, which means that there is a non-commutative polynomial $f \in k\langle x_1, \dots, x_n \rangle$ such that for any n elements $a_1, \dots, a_n \in A$ we have $f(a_1, \dots, a_n) = 0$. In particular this class includes commutative rings (by taking $f = x_1 x_2 - x_2 x_1$), and more generally any ring which is a finite module over its centre.

Lemma 3.5. [SVdB08, Lem. 4.2.] *Let K be an algebraically closed field, and let A_1, A_2 be two finitely-generated polynomial identity algebras over K . If A_1 and A_2 both have finite global dimension, then so does $A_1 \otimes_K A_2$.*

Lemma 3.6. *An algebra A over a field K is smooth over K if and only if $A_{\overline{K}}$ is smooth over \overline{K} .*

Proof. If A is a perfect $A^{op} \otimes_K A$ -module, then obviously $A_{\overline{K}}$ is perfect. For the reverse implication, suppose for a contradiction that A is not smooth. Then there exists a sequence of $A^{op} \otimes_K A$ -modules M_i and an increasing sequence of integers n_i such that $\text{Ext}_{A^{op} \otimes_K A}^{n_i}(A, M_i) \neq 0$. But then $\text{Ext}_{A^{op} \otimes_{\overline{K}} A}^{n_i}(A_{\overline{K}}, (M_i)_{\overline{K}}) \neq 0$, contradicting the assumption that $A_{\overline{K}}$ is smooth over \overline{K} . \square

Recall Grothendieck's generic freeness lemma [Eis95, Thm. 14.4]:

Lemma 3.7. *Let R be a Noetherian domain, and T a finitely generated commutative R -algebra. Let M be a finitely generated T -module. Then there is an element $f \in R$ such that M_f is a free R_f -module. In particular if $M_{K(R)} = 0$ then there exists an f such that $M_f = 0$.*

Lemma 3.8. *Let R be a Noetherian domain, let T be a finitely generated commutative R -algebra, and let Λ be a T -algebra which is finitely generated as a T -module. Let P be a finitely generated Λ -module such that $P_{K(R)}$ is a projective $\Lambda_{K(R)}$ -module. Then there exists an $f \in R$ such that P_f is a projective Λ_f -module.*

Proof. Let $K = K(R)$. Since P_K is projective, there exists a free module $F = \Lambda^{\oplus n}$ and maps $i: P_K \rightarrow F_K$ and $p: F_K \rightarrow P_K$, such that $pi = 1_{P_K}$. There exists an $f \in R$ such that the maps are defined over R_f , i.e. there are maps $i_f: P_f \rightarrow F_f$ and $p_f: F_f \rightarrow P_f$ which localise to i and p .

Now $p_f i_f - 1_{P_f}$ has a kernel L and a cokernel C , which are finitely generated T -modules. We have $L_K = C_K = 0$, and so by Lemma 3.7, there exists a $g \in R_f$ such that $L_g = C_g = 0$. This implies that $p_{fg} i_{fg} = 1_{P_{fg}}$, and so P_{fg} is a projective Λ_{fg} -module. \square

Corollary 3.9. *Let R, T and Λ be as in the previous lemma. If M is a finitely generated Λ -module such that $M_{K(R)}$ is a perfect $\Lambda_{K(R)}$ -module, then there exists an $f \in R$ such that M_f is a perfect Λ_f -module.*

Proof. Take a projective resolution $\cdots \rightarrow M^i \rightarrow M^{i-1} \rightarrow \cdots$ of M , and let C^n be the kernel of $M^n \rightarrow M^{n-1}$. Since Λ is Noetherian (Lemma 3.1), we may assume that all M^i and C^i are finitely presented. Since M_K has finite projective dimension, there is an n such that C_K^n is a projective Λ_K -module. Applying Lemma 3.8, we find an f such that C_f^n is projective, and so $C_f^n \rightarrow M_f^{n-1} \rightarrow \cdots \rightarrow M_f^0$ is a projective resolution of M_f . \square

REFERENCES

- [Aus55] Maurice Auslander. On the dimension of modules and algebras. III. Global dimension. *Nagoya Math. J.*, 9:67–77, 1955.
- [DII⁺16] Hailong Dao, Osamu Iyama, Srikanth B. Iyengar, Ryo Takahashi, Michael Wemyss, and Yuji Yoshino. Noncommutative resolutions using syzygies. *arXiv:1609.04842 [math]*, 2016.
- [Eis95] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [FOV99] H. Flenner, L. O’Carroll, and W. Vogel. *Joins and intersections*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1999.
- [JL82] Christian U. Jensen and Helmut Lenzing. Homological dimension and representation type of algebras under base field extension. *Manuscripta Math.*, 39(1):1–13, 1982.
- [Row88] Louis H. Rowen. *Ring theory. Vol. II*, volume 128 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1988.
- [RS16] Jørgen Vold Rennemo and Ed Segal. Hori-mological projective duality. *arXiv:1609.04045 [hep-th]*, 2016.
- [Sch85] A. Schofield. Bounding the global dimension in terms of the dimension. *Bull. London Math. Soc.*, 17(4):393–394, 1985.
- [SVdB08] J. T. Stafford and M. Van den Bergh. Noncommutative resolutions and rational singularities. *Michigan Math. J.*, 57:659–674, 2008. Special volume in honor of Melvin Hochster.
- [ŠVdB15] Špela Špenko and Michel Van den Bergh. Non-commutative resolutions of quotient singularities. *arXiv:1502.05240 [math]*, 2015. To appear in *Invent. Math.*
- [VdB04] Michel Van den Bergh. Non-commutative crepant resolutions. In *The legacy of Niels Henrik Abel*, pages 749–770. Springer, Berlin, 2004. Revised version at *arXiv:math/0211064 [math.RA]*.

ALL SOULS COLLEGE, OXFORD, OX1 4AL, UK
E-mail address: jvrennemo@gmail.com

IMPERIAL COLLEGE LONDON, 180 QUEEN’S GATE, LONDON, SW7 2AZ, UK
E-mail address: edward.segal04@imperial.ac.uk

UNIVERSITEIT HASSELT, UNIVERSITAIRE CAMPUS, B-3590 DIEPENBEEK, BELGIUM
E-mail address: michel.vandenbergh@uhasselt.be