Made available by Hasselt University Library in https://documentserver.uhasselt.be

An evaluation of species distribution models to estimate tree diversity at genus level in a heterogeneous urban-rural landscape Peer-reviewed author version

Stas, Michiel; AERTS, Raf; Hendrickx, Marijke; Dendoncker, Nicolas; Dujardin, Sebastien; Linard, Catherine; NAWROT, Tim; Van Nieuwenhuyse, An; Aerts, Jean-Marie; Van Orshoven, Jos & Somers, Ben (2020) An evaluation of species distribution models to estimate tree diversity at genus level in a heterogeneous urban-rural landscape. In: LANDSCAPE AND URBAN PLANNING, 198 (Art N° 103770).

DOI: 10.1016/j.landurbplan.2020.103770 Handle: http://hdl.handle.net/1942/31345



institutional repository and make this publicly available after an embargo period has expired.



20

21 22

Creative Commons Attribution-Noncommercial-Share Alike 2.0 Belgium LicensePermissions
 beyond the scope of this license may be available at
 <u>http://www.elsevier.com/wps/find/authorsview.authors/copyright#whatrights</u>

An evaluation of species distribution models to estimate tree diversity at genus level in a heterogeneous urban-rural landscape

28

29 Abstract

Trees provide ecosystem services that improve the environment and human health. The magnitude of these improvements may be related to tree diversity within green spaces, yet spatially explicit diversity data necessary to investigate such associations are often missing. Here, we evaluate two methods to model tree diversity at genus level based on environmental covariates and presence point data. We want to identify the drivers and suitable methods for urban and rural tree diversity models in the heterogeneous region of Flanders, Belgium.

We stratified our research area into dominantly rural and dominantly urban areas and developed distribution models for 13 tree genera for both strata as well as for the area as a whole. Occurrence data were obtained from an open-access presence-only database of validated observations of vascular plants. These occurrence data are combined with environmental covariates in MaxEnt models. Tree diversity is modelled by adding up the individual species distribution models.

42 Models in the dominantly rural areas are driven by soil characteristics (soil texture and drainage class). Models in the dominantly urban areas are driven by environmental covariates explaining 43 urban heterogeneity. Nevertheless, the stratification into urban and rural did not contribute to 44 45 a higher model quality. Generic tree diversity estimates were better when presences derived from distribution models were simply added up (binary stacking, True Positive Rate of 0.903). 46 The application of macro-ecological constraints resulted in an underestimation of generic tree 47 diversity (probability stacking, True Positive Rate of 0.533). We conclude that summing 48 presences derived from species distribution models (binary stacking) is a suitable approach to 49 increase knowledge on regional diversity. 50

51

52 HIGHLIGHTS:

53	Rural species distribution models (SDMs) are driven by soil characteristics.
54	Urban SDMs are driven by environmental covariates explaining urban heterogeneity.
55	Summing presences derived from SDMs is suitable to assess diversity.
56	Summing habitat suitability derived from SDMs underestimates diversity.
57	

58 **1. Introduction**

Trees in urban environments deliver important ecosystem services. Street trees help to cool 59 cities (Konarska et al., 2016; Scholz et al., 2018), partially mitigating the urban heat island 60 effect and human heat stress (Lee et al., 2016). Trees can also remove particulate matter from 61 the atmosphere (Scholz et al., 2018; Selmi et al., 2016). Through climate regulation, air 62 pollution mitigation and aesthetic and cultural values, trees contribute to better human well-63 being (Salmond et al., 2016). Individual trees in the city also contribute to indirect nature 64 experiences, which benefits human health (Cox et al., 2019). The ecosystem services delivered 65 66 by trees are tree species specific (Donovan et al., 2005). Increasing biodiversity is expected to result in better ecosystem functioning and may yield more stable ecosystem services through 67 time (Cardinale et al., 2012). 68

69 Tree diversity has been studied at genus level before (Hoover et al., 2017; Hope et al., 2003) and might become more popular with increasing availability of observations from citizen 70 science (Dobbs et al., 2018). Observations of trees through citizen science initiatives have been 71 72 found to be more accurate at genus level than at species level (Roman et al., 2017). Plant genuslevel diversity is strongly linked to plant species-level diversity (O'Brien et al., 1998). 73 Additionally, interactions with host plants often occur at genus level and therefore genus level 74 diversity is also relevant for insect diversity (Kemp and Ellis, 2017), or ectomycorrhizal fungal 75 diversity (Gao et al., 2013). Thomsen et al. (2016) emphasize that a healthy urban tree 76 77 population requires a high generic diversity. Modelling tree diversity at genus level is thus of high value because it encompasses more reliable observation data and allows for diverse 78 applications. 79

Genus-level presence-only data can be used in species distribution models (SDMs). SDMs
correlate species observations and environmental covariates to predict habitat suitability (Elith *et al.*, 2006). Applications of SDMs are manifold: identifying species distributions, studying

impact of climate and land use change scenarios (Dyderski *et al.*, 2018) and identifying areas
of interest for conservation (McCune, 2016). Applications of SDMs in urbanized areas are not
common (Della Rocca *et al.*, 2017), yet they have been successfully applied in urban green
spaces (Milanovich *et al.*, 2012) and human-dominated landscapes (McCune, 2016).

Species richness can be modelled by stacking individual SDMs on top of one another to yield 87 a total richness. Stacking of SDMs is most commonly done after thresholding the continuous 88 probability output of the individual SDMs, a method known as binary stacking (Calabrese et 89 al., 2014). Nevertheless, discretizing continuous probabilities using fixed thresholds (for 90 91 example considering all cases with a modelled probability of p > 0.55 as being present) is generally discouraged (Merow et al., 2013). Instead, species-specific threshold rules can be 92 applied (Cao et al., 2013). Still, the literature suggests that binary stacking tends to 93 94 overestimate species richness because biotic limitations are not accounted for (Calabrese et al., 2014; Gavish et al., 2017; Guisan and Rahbek, 2011). Nonetheless, combining binary SDMs 95 is the most straightforward method to create species richness maps (Trotta-Moreu & Lobo, 96 2010). Combining continuous probability data, which is called probability stacking, is an 97 alternative stacking approach (Calabrese et al., 2014), although interpretations are less 98 straightforward. Guisan & Rahbek (2011) have proposed a framework for spatially explicit 99 species assemblage modelling (SESAM). In the SESAM framework, a macro-ecological model 100 101 limits the number of species that can co-occur in one cell. One way of defining the macro-102 ecological constraint is by stacking the probabilities of SDMs (probability stacking). D'Amen et al. (2015) reduced overestimation by applying probability stacking successfully in the Alps 103 of western Switzerland at a fine spatial resolution. 104

In this study, we test a stratified approach, in which we run separate SDMs for rural and urban areas in a mosaic landscape, to model tree diversity at genus level. We evaluate which environmental covariates drive the urban and rural models. First, we hypothesize that different

environmental covariates drive the urban and rural models. We expect that soil nutrients and 108 soil moisture determine vegetation in rural areas, because this vegetation resembles the 109 potential natural vegetation more closely (Walthert and Meier, 2017). For urban areas we 110 expect an anthropogenic influence on the vegetation composition (Bourne and Conway, 2014). 111 Second, we hypothesize that the application of a macro-ecological constraint to take biotic 112 interactions into account would improve models for rural areas, but not for urban areas. We 113 expect that biotic interactions are more relevant in rural areas (D'Amen et al., 2015). Third, we 114 hypothesize that binary stacking performs sufficiently well in urban areas. We expect that 115 116 biotic interactions are less relevant in urban areas because of human intervention.

117

118 **2. Materials and methods**

119 *2.1. Study area and stratification*

Flanders is the northernmost of the three administrative regions of Belgium with an area of 120 13,522 km² and a population density of 482 inhabitants per km². The area has a north-south 121 soil gradient of decreasing fraction of sand and increasing fraction of silt. The climate 122 according to Köppen is a maritime temperate climate (Cfb) (Peel et al., 2007). The 123 Organization for Economic Co-operation and Development (OECD) considers Flanders 124 entirely as urbanized (Vervoort, 2016). Nevertheless, based on the Urban Audit of 2018 125 published by Eurostat, core cities and functional urban areas (FUAs) are delineated for 126 127 Flanders. The delineation is used to distinguish dominantly urban areas from more rural areas in Flanders (Fig. 1). FUAs define a metropolitan area outside the geographical city boundaries, 128 taking into account demographic, economic and environmental factors (Khalili et al., 2018). In 129 130 Flanders, the FUAs are located around the capitals of each province, except for Hasselt (Fig. 1). Hasselt is a provincial capital located in the east of the region where forest cover is higher. 131 As only 10.6% of the study area consists of forest (De Keersmaeker *et al.*, 2015), other (urban) 132

133green spaces are of high importance for biodiversity (Aronson et al., 2017; Lepczyk et al.,1342017). Green space is most commonly defined as a vegetated area (Taylor and Hochuli, 2017).135We will focus on vegetated areas containing woody vegetation. The Belgian and Luxemburg136Institute for Floristics (IFBL) developed a regular grid of 4×4 km squares further divided into137 1×1 km squares covering Belgium and Luxemburg. This grid is used as a reference for mapping138species distributions (Van Rompaey, 1943). In this study the 1×1 km IFBL grid is used to mask139observation data and environmental covariates.



140

Fig. 1: Stratification of the Region of Flanders into functional urban areas and rural areas according to the Urban Audit of Eurostat (2018). The provincial capital cities are labeled and the forest cover in the region is visualized.

144 2.2. Species distribution modelling

Species distribution models (SDMs) estimate the relationship between recorded occurrences at sites (samples) and the environmental and/or spatial characteristics of those sites (environmental covariates) (Elith *et al.*, 2011). This relationship is then used to predict occurrences elsewhere. In the stratified approach, separate SDMs are run for the urban and rural strata. Subsequently, the urban and rural data are combined in an integrated approach to form a model that covers the whole region of Flanders. The models are validated with independent field data to evaluate their utility in the urban and rural strata. The workflow for the integrated approach, which is parallel for the stratified approach, is visualized in Fig. 2 and explained in the following sections.

2.2.1. Occurrence data

The occurrence data of thirteen tree genera were included in the study: Aesculus (horse 154 chestnut), Alnus (alder), Betula (birch), Carpinus (hornbeam), Corylus (hazel), Fagus (beech), 155 Fraxinus (ash), Juglans (walnut), Platanus (plane), Populus (poplar), Quercus (oak), Salix 156 (willow) and Tilia (linden). Presence-records of these genera were extracted from Florabank1 157 (Van Landuyt & Brosens, 2017) available on GBIF.org. The Florabank is an open-access 158 presence-only database of validated observations of vascular plants, from checklists, literature 159 and herbarium specimen information. The observations are georeferenced and attributed to the 160 centers of 14317 1km×1km IFBL grid cells (Van Landuyt et al., 2012). 161

162

163



PROBABILITY STACKED SDM

BINARY STACKED SDM

Fig. 2: Mapping tree diversity at the genus level from presence-only and environmental data. Presence-only data from Florabank (a) and environmental data from various sources (b) are combined in a Species Distribution Model (SDM) using MaxEnt. The continuous output of MaxEnt, probability models (c), can be converted to presence-absence models (binary SDMs) by applying genus specific thresholds (d). Stacking the probability models and applying the probability ranking rule results in a probability stacked SDM (e). Aggregating the thresholded SDMs results in a binary stacked SDM (f).

164 *2.2.2. Environmental covariates*

Soil texture class, soil drainage class, mean lowest and highest groundwater table depth, land 165 cover type and habitat type were the environmental covariates used. In Belgium, natural plant 166 communities are primarily determined by variation in soil nutrient content and soil moisture 167 (Cornelis et al., 2009). Thus, soil texture and drainage class were extracted as categorical soil 168 variables from the Belgian soil map (Dondeyne et al., 2014). This vector geodataset was first 169 resampled to a raster, using the IFBL grid as the mask layer and the cell assignment type 170 'maximum combined area'. The 'maximum combined area' rule selects the attribute value from 171 172 the polygon with the largest total area overlapping with the grid cell (ESRI, 2017). Mean highest and lowest groundwater tables data were obtained from a soil hydrology raster 173 (ECOPLAN, 2014) and resampled to the IFBL grid. 174

Land cover data were obtained from one of the base layers in the ECOPLAN ecosystem services information system (ECOPLAN, 2014). The geodataset contains a basic land cover classification (the list of classes is available in Appendix 1). The grid with a spatial resolution of 5m was resampled to the IFBL grid, retaining the land cover with the largest area in the grid cell.

Habitat data were obtained from the Biological Valuation Map (BVM), a geodataset of habitat types with attribute information on the ecological context and value of the delineated areas (Vriens *et al.*, 2011). The BVM contains information about heterogeneity of urban areas, such as the density and context of built-up areas, industrial areas and recreational areas. The classes of the BVM are listed in Appendix 1. The BVM is a vector geodataset and was resampled to the IFBL grid using 'maximum combined area' as the cell assignment type. Resampling and masking of the environmental geodatasets were performed in ArcGIS 10.5.1-software (ESRI,

187 Redlands, CA, 2017).

188 *2.2.3. Probability models*

Probability models of the spatial distribution of each of the 13 genera were developed using 189 MaxEnt version 3.3.3k. MaxEnt is a machine-learning algorithm highly suitable to develop 190 models from presence-only data (Elith et al., 2006; Phillips et al., 2006). The algorithm is based 191 on the principles of maximum entropy and finds an optimal probability distribution using a 192 combination of occurrence data and environmental data (Elith et al., 2011). MaxEnt is known 193 to perform well even when environmental covariates are linearly correlated (De Marco and 194 195 Nóbrega, 2018). The logistic output of MaxEnt is an attempt at expressing the raw output as a probability of presence (Elith et al., 2011). A 10-fold cross validation was applied. Model 196 197 performance was assessed with the area under the receiver operating characteristic curve (AUC) statistic, ranging between 0 and 1. When AUC values are higher than 0.5, the model 198 performs better than a random distribution. For every genus, three models were developed: one 199 using the entire dataset (integrated approach), then one for the rural and one for the urban areas 200 (stratified approach). To evaluate the driving factors in these models, we determined the 201 202 environmental covariate with the highest percentage of contribution to the model.

203

204 *2.2.4. Binary stacking*

We applied the '10 percentile training presence' rule on the MaxEnt-output (Ficetola *et al.*, 2009; Pearson *et al.*, 2006; Skowronek *et al.*, 2017), for every genus and model approach separately, resulting in a threshold value above which 90% of the training samples are correctly classified. Thus, a unique threshold value is used for every genus to create a binary output (0 = absence, 1 = presence). Binary stacking is the process of adding up the individual binary models, resulting in a generic tree diversity varying from 0 to 13 genera.

211

212 2.2.5. Probability stacking

As a cell-specific macro-ecological constraint we summed the MaxEnt-probabilities per grid 213 cell (D'Amen et al., 2015), resulting in a possible generic tree diversity range between 1.96 214 and 8.76. To determine which genera occur in the constrained cells we used the 'probability 215 ranking' rule. The genera are assigned to the cell according to decreasing order of probability 216 of presence determined from the SDMs (2.2.3), until the cell-specific macro-ecological 217 constraint is reached. Probability ranking as described in the SESAM framework is 218 incorporated in the package 'ecospat' available for R (Di Cola et al., 2017) and was executed 219 with R software 3.4.3 (R Core Team, 2017). 220

221

222 2.3. Validation

The probability models (2.2.3) were cross-validated before they were stacked. In addition, the 223 224 stacked models were validated with independent field data. The independent field data consisted of recordings of the genus's occurrence around 208 randomly selected point 225 locations, with a search effort per point of ten minutes with two observers. The sampling 226 protocol is derived from the timed-meander sampling protocol (Goff et al., 1982), which is 227 applied in various fields of ecology (Threlfall et al., 2017) and is favored because of its cost-228 effectiveness (Hamm, 2013). The 208 point locations are distributed over 130 IFBL cells. 229 There are 87 rural cells and 43 urban cells. A genus is present in a cell if it is observed in at 230 least one of the random point locations within the cell. The field data are assumed to provide 231 232 the true condition that is compared to the predicted condition provided by the SDMs at genus level. True condition data and predicted condition data were compared in a confusion matrix, 233 describing true positive (TP), true negative (TN), false positive (FP) and false negative (FN) 234 outcomes. Based on the values in the confusion matrix, we evaluated the model performance 235 by calculating the true positive rate (TPR). TPR is the number of true positives divided by the 236 total of positive cases, the sum of true and false positives. The TPR informs simultaneously 237

about the presences that are correctly predicted and about those that were incorrectly identified 238 as positives. A TPR of 80% would indicate that 80% of the presences are true positives while 239 20% are false positives. However, in the present study, the false positives are not necessarily 240 false as the species could have been missed during the validation field work. Therefore we 241 focus on the true positives when interpreting the TPR. Additionally, the percentage of false 242 negatives is included in the evaluation, because this percentage provides information on the 243 underestimation of the stacking method. The higher the percentage of false negatives, the more 244 the tree diversity at genus level is underestimated. 245

246

247 2.4 Compare model outcomes

To compare model outcomes we calculated average modelled tree diversity at genus level and 248 249 95% confidence intervals for binary stacked vs. probability stacked models and for integrated vs. stratified approaches and for urban vs. rural areas. We used the paired sample t-test (with a 250 statistical cutoff value of 0.05) to test whether overall average modelled tree diversity at genus 251 level differed between binary stacked and probability stacked models. We then used the paired 252 sample t-test to test whether modelled generic tree diversity differed between integrated and 253 stratified approaches within the binary stacked models, both for the entire dataset and for a 254 dataset stratified in urban vs. rural areas. 255

256

257 **3. Results**

258 *3.1 Probability models*

The species distribution models outperformed the random spatial distribution (all AUC > 0.5; Table 1), for the stratified approach as well as the integrated approach. On average the AUC is 0.60 with a standard deviation of 0.007. The strength of the strongest predictor ranges from 32.7-85.7 percent of contribution (Table 1). For 11 out of 13 urban models, the strongest predictor is the Biological Valuation Map (BVM), containing information on urban heterogeneity. For the rural model as well as the integrated model, we found that for some genera the strongest predictors were the soil variables texture class and drainage class.

Urban stratum **Integrated model Rural stratum** AUC SD SP PC AUC SD SP PC AUC SD SP PC Genus presences Aesculus 0.03 1162 0.63 **BVM** 54.3 0.62 0.04 **BVM** 61.2 0.64 0.03 BVM 55.1 Alnus 6918 0.55 0.01 0.54 0.02 0.54 0.01 39.9 BVM 38.0 **BVM** 40.7 **BVM** Betula 0.01 8888 0.60 **BVM** 40.9 0.58 0.02 BVM 0.58 0.01 BVM 47.7 45.6 Carpinus 0.02 42 0.02 3029 0.64 texture class 0.63 0.02 **BVM** 29.2 0.63 texture class 41.3 Corylus 0.02 5490 0.60 texture class 45.8 0.59 0.02 **BVM** 34.5 0.59 0.01 texture class 47.1 Fagus 0.02 drainage class 38.1 0.02 0.02 drainage class 37.2 2710 0.64 0.64 BVM 32.7 0.64 Fraxinus 0.01 6635 0.58 texture class 67.6 0.55 0.02 texture class 42.6 0.57 0.01 texture class 71.2 Juglans 0.03 1833 0.61 texture class 44.9 0.59 0.03 BVM 0.61 0.02 texture class 46.5 40.3 Platanus 50 0.63 0.34 BVM 0.90 0.06 **BVM** 81.7 0.81 0.10 BVM 85.9 47.5 Populus 0.02 drainage class 55.9 0.01 drainage class 47.1 0.02 5380 0.57 0.56 BVM 41.8 0.56 Quercus 0.01 **BVM** 0.01 8182 0.57 36.2 0.55 0.02 texture class 38.4 0.55 BVM 38.7 Salix 8349 0.01 drainage class 65.0 0.02 drainage class 46.4 0.53 0.52 BVM 39.9 0.52 0.01 Tilia 0.04 BVM 0.63 0.04 1171 0.61 42.6 BVM 54.3 0.64 0.03 BVM 43.8

Table 1: Summary of the species distribution models for each genus. Reporting the number of grid cells occupied by an observation (presences), the average area under the curve (AUC) as a measure to evaluate the models, the standard deviation of the AUC (SD), the strongest predictor (SP) and the percent of contribution (PC) of this strongest predictor to the MaxEnt model. (BVM = Biological Valuation Map)

3.2 Stacked species distribution models

For the integrated as well as the stratified approach binary stacking resulted in a generic tree diversity varying between 0 and 13. Probability stacking resulted in a lower generic tree diversity between 2 and 9 (Fig. 3). Spatial differences in generic tree diversity between the integrated and the stratified approach are not strongly pronounced.



Fig. 3: Tree diversity at genus level determined by binary (upper) and probability (lower) stacking of the MaxEnt models developed in an integrated (left) and stratified (right) modelling approach. The cell size is 1km×1km.

3.3 Validation

The binary stacking method, with an overall TPR of 0.90, performed better than the probability stacking method, with a considerably lower overall TPR of 0.52-0.53 (Table 2). Overall, the integrated and the stratified approach performed equally well. The binary stacking method had a lower percentage of false negatives (6.0 - 6.1 %) than the probability stacking method (29.3 -29.9 %) (Table 3).

 Table 2: Validation results: true positive rate (TPR) derived from the confusion matrix.

			TPR		
Model	Approach	Overall	Urban	Rural	
Binary stacked	Integrated	0.90	0.94	0.89	
	Stratified	0.90	0.95	0.88	
Probability stacked	Integrated	0.53	0.54	0.53	
	Stratified	0.52	0.53	0.53	

Table 3: Validation results: percentage of false negatives (%) derived from the confusion matrix.

			FN (%)	
Model	Approach	Overall	Urban	Rural
Binary stacked	Integrated	6.1	4.3	7.0
	Stratified	6.0	3.3	7.3
Probability stacked	Integrated	29.3	30.8	28.6
	Stratified	29.9	31.5	29.2

3.4.1 Binary stacking vs. probability stacking

The overall average tree diversity at genus level was higher for binary stacked models (9.8-9.9) than for probability stacked models (5.6) (Table 4). The diversity based on the validation data was 8.1, in line with results from the binary stacking approach. There was a significant mean difference of 4.3 (95% CI 4.2-4.3) between binary stacked and probability stacked models for the integrated approach (paired t-test t = 200.1, df = 13458, p < 0.001). There was a significant mean difference of 4.2 (95% CI 4.1-4.3) between binary and probability stacked models for the stratified approach (paired t-test t = 153.7, df = 13458, p < 0.001).

Table 4: Average modelled tree diversity at genus level based on binary and probability stacked models, for integrated and stratified approaches and for urban and rural areas.

		T	ree diversity (95% C	CI)
Model	Approach	Overall	Urban	Rural
Binary stacked	Integrated	9.9 (9.8-9.9)	10.1 (10.0-10.1)	9.8 (9.7-9.8)
	Stratified	9.8 (9.7-9.8)	9.5 (9.4-9.6)	10.0 (9.9-10.0)
Probability stacked	Integrated	5.6 (5.6-5.6)	5.7 (5.7-5.8)	5.6 (5.5-5.6)
	Stratified	5.6 (5.6-5.6)	5.5 (5.5-5.6)	5.6 (5.6-5.6)
Validation data		8.1 (7.6-8.7)	8.6 (7.8-9.5)	7.9 (7.3-8.6)

3.4.2 Binary stacking: rural vs. urban areas

For binary stacked models, the integrated approach yielded a statistically significant higher overall estimated tree diversity at genus level [mean difference integrated vs. stratified 0.09 (95% CI 0.06-0.11), t = 6.72, df = 13458, p<0.001]. However, subtracting the stratified result from the integrated result revealed a spatial differentiation (Fig. 4). The stratified approach resulted in a significantly higher diversity in the rural areas, thus a negative mean difference of -0.18 (95% CI -0.21 - -0.15) (t = -12.5, df = 8359, p<0.001). The integrated approach, however, resulted in a significantly higher diversity in the urban areas, thus a positive mean difference of 0.52 (95% CI 0.48-0.57) (t = 22.3, df = 5098, p<0.001).

Green pixels (Fig. 4) represent a higher diversity obtained with the integrated approach. These green pixels are often clustered within the FUAs (black outline, Fig. 4). Conversely, red pixels represent a higher diversity obtained with the stratified approach. These red pixels are observed in clusters outside the FUAs, especially in extremely rural areas such as: 'De Westhoek' in the west and 'Haspengouw' in the south-east of Flanders (Fig. 4).



Fig. 4: The difference in the binary stacking results from the integrated and stratified approach. Cell size is 1km×1km.

4. Discussion and conclusions

4.1 Environmental covariates

Rural and urban models were driven by different environmental covariates, which confirms the first hypothesis. As expected, soil variables, such as texture class and drainage class, were of high importance to explain the distribution of native trees in rural areas. Texture class was an important covariate for *Corylus*, for example, because this genus requires richer loamy soils (Ozenç, 2001). Salix and Populus can tolerate wet soils (Zalesny and Bauer, 2007) and as a result drainage class was an important environmental covariate in their SDMs. Earlier studies demonstrated that including soil factors in plant SDMs results in improved predictions (Buri et al., 2017). For urban areas we expected that the vegetation would be determined by anthropogenic influences. The urban heterogeneity, which is better described in the Biological Valuation Map (BVM), was the most important covariate in the urban SDMs (Table 1). In the BVM seven urban/built up types are included (ua, ud, un, ur, uv, uc and ui), while the the other land cover map (ECOPLAN) contains only three urban land cover types (9101, 9201, 9202) (see Appendix 1). It has been emphasized in the past that including environmental covariates that account for the diverse functions of urban areas is important to understand urban plant species patterns (Godefroid and Koedam, 2007). Future developments of species distribution models in urban areas need to include covariates that address the variety of anthropogenic influence.

4.2 Stacking methods

At a fine spatial resolution, as in the work of D'Amen *et al.* (2015), binary stacking overestimates species diversity in a natural environment, because dispersal limitations and biotic interactions are not taken into account. In this study, however, we worked at a spatial resolution of 1km×1km and biotic interactions are less important at this relatively coarse resolution (Thuiller *et al.*, 2015). The probability stacking method should prevent from overestimating tree diversity at genus level. Nevertheless, the low true positive rate (Table 2)

and the high percentage of false negatives (Table 3) showed that the probability stacking method underestimated tree diversity in Flanders. Additionally, in urbanized areas, human decision-making and management most likely override natural species selection and biotic interactions are therefore less likely to drive species composition. Therefore, binary stacking is the preferred method for biodiversity modelling at 1km×1km resolution in both urban areas and rural areas. Nevertheless, the scale-dependent applicability of a macro-ecological constraint needs further research as there is, to our knowledge, no literature on this topic.

4.3 Comparison of model outcomes

Binary stacking resulted in a significantly higher diversity at genus level compared to probability stacking (Table 4). This difference is not an overestimation of binary stacking, but due to an underestimation of probability stacking. Using the binary stacking method and the stratified approach, higher diversity was clustered in the rural areas (Table 4 and Fig. 4). Rural areas are thus more prone to overestimation and would probably benefit more from applying macro-ecological constraints, compared to urban areas. Nevertheless, applications of macro-ecological constraints seem to be of higher relevance in more natural areas (D'Amen *et al.*, 2015), far less urbanized than the rural areas in Flanders.

To conclude, we find that binary stacking is most suitable for both urban and rural areas in Flanders. Stratification of the study area did not improve model quality considerably, but confirmed that different environmental covariates contributed to the models of urban and rural areas. Probability stacking is to be considered in natural areas, but does not perform well in urbanized areas, especially at the moderate spatial resolution of 1km×1km.

4.4 Limitations

All Species Distribution Models (SDMs) had relatively low Area Under the Curve values (average AUC: 0.60 ± 0.01), but all performed better than random distributions (Table 1). By

stacking SDMs, errors in individual species models accumulate and degrade predictions of species diversity (D'Amen *et al.*, 2015; Pottier *et al.*, 2013). The importance of the BVM as an environmental covariate emphasizes the relevance of including urban heterogeneity in SDMs. Unfortunately, at a moderate resolution of 1×1 km relevant intra-urban variation of the tree canopy (Weinberger *et al.*, 2016) cannot be observed.

4.5 Applications

The model resulting from this study can be expanded by stacking more binary SDMs, by producing species-level models or by producing models of other plant groups. Spatially-explicit biodiversity data are vital for emerging environmental health studies (McInnes *et al.*, 2017), for example to study relationships between residential and dynamic exposure and human health outcomes (Cox *et al.*, 2017; Shanahan *et al.*, 2016). Hjort et al. (2016) present a concept to calculate individual long-term or life time exposure to pollen with geographic information systems. Landscape and urban planners could also use tree diversity maps to identify areas with low diversity and optimize the delivery of ecosystem services or decrease potential social inequalities in access to biodiverse green space by increasing biodiversity in focus areas (Wolch *et al.*, 2014). Finally, when subsets of models for allergenic species are used, diversity maps could be interpreted as allergy risk maps and inform pollen allergy patients about pollen allergy risks (McInnes *et al.*, 2017).

References

- Alvey, A.A., 2006. Promoting and preserving biodiversity in the urban forest. Urban For. Urban Green. 5, 195–201. https://doi.org/10.1016/J.UFUG.2006.09.003
- Aronson, M.F., Lepczyk, C.A., Evans, K.L., Goddard, M.A., Lerman, S.B., MacIvor, J.S., Nilon, C.H., Vargo, T., 2017. Biodiversity in the city: key challenges for urban green space management. Front. Ecol. Environ. 15, 189–196. https://doi.org/10.1002/fee.1480
- Buri, A., Cianfrani, C., Pinto-Figueroa, E., Yashiro, E., Spangenberg, J.E., Adatte, T., Verrecchia, E., Guisan, A., Pradervand, J.-N., 2017. Soil factors improve predictions of plant species distribution in a mountain environment. Prog. Phys. Geogr. 41, 703–722. https://doi.org/10.1177/0309133317738162
- Calabrese, J.M., Certain, G., Kraan, C., Dormann, C.F., 2014. Stacking species distribution models and adjusting bias by linking them to macroecological models. Glob. Ecol. Biogeogr. 23, 99–112. https://doi.org/10.1111/geb.12102
- Cao, Y., DeWalt, R.E., Robinson, J.L., Tweddale, T., Hinz, L., Pessino, M., 2013. Using Maxent to model the historic distributions of stonefly species in Illinois streams: The effects of regularization and threshold selections. Ecol. Modell. 259, 30–39. https://doi.org/10.1016/J.ECOLMODEL.2013.03.012
- Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., Narwani, A., Mace, G.M., Tilman, D., Wardle, D.A., Kinzig, A.P., Daily, G.C., Loreau, M., Grace, J.B., Larigauderie, A., Srivastava, D.S., Naeem, S., 2012. Biodiversity loss and its impact on humanity. https://doi.org/10.1038/nature11148
- Cornelis, J., Hermy, M., Roelandt, B., De Keersmaeker, L., Vandekerkhove Kris, 2009. Bosplantengemeenschappen in Vlaanderen, eEen typologie van bossen gebasseerd op de kruidlaag. INBO.M.2009.5. Agentschap voor Natuur en Bos en Intituut voor Natuur- en Bosonderzoek, Brussel.
- Cox, D.T.C., Bennie, J., Casalegno, S., Hudson, H.L., Anderson, K., Gaston, K.J., 2019. Skewed contributions of individual trees to indirect nature experiences. Landsc. Urban

Plan. 185, 28–34. https://doi.org/10.1016/J.LANDURBPLAN.2019.01.008

- D'Amen, M., Dubuis, A., Fernandes, R.F., Pottier, J., Pellissier, L., Guisan, A., 2015. Using species richness and functional traits predictions to constrain assemblage predictions from stacked species distribution models. J. Biogeogr. 42, 1255–1266. https://doi.org/10.1111/jbi.12485
- De Keersmaeker, L., Onkelinx, T., De Vos, B., Rogiers, N., Vandekerkhove, K., Thomaes, A., De Schrijver, A., Hermy, M., Verheyen, K., 2015. The analysis of spatio-temporal forest changes (1775–2000) in Flanders (northern Belgium) indicates habitat-specific levels of fragmentation and area loss. Landsc. Ecol. 30, 247–259. https://doi.org/10.1007/s10980-014-0119-7
- De Marco, P., Nóbrega, C.C., 2018. Evaluating collinearity effects on species distribution models: An approach based on virtual species simulation. PLoS One 13, e0202403. https://doi.org/10.1371/journal.pone.0202403
- Della Rocca, F., Bogliani, G., Milanesi, P., 2017. Patterns of distribution and landscape connectivity of the stag beetle in a human-dominated landscape. Nat. Conserv. 19, 19–37. https://doi.org/10.3897/natureconservation.19.12457
- Di Cola, V., Broennimann, O., Petitpierre, B., Breiner, F.T., D'Amen, M., Randin, C., Engler, R., Pottier, J., Pio, D., Dubuis, A., Pellissier, L., Mateo, R.G., Hordijk, W., Salamin, N., Guisan, A., 2017. ecospat: an R package to support spatial analyses and modeling of species niches and distributions. Ecography (Cop.). 40, 774–787. https://doi.org/10.1111/ecog.02671
- Dobbs, C., Hernández, A., de la Barrera, F., Miranda, M.D., Paecke, S.R., 2018. Integrating Urban Biodiversity Mapping, Citizen Science and Technology, in: Urban Biodiversity.
 Routledge, pp. 236–247. https://doi.org/10.9774/GLEAF.9781315402581_16
- Dondeyne, S., Vanierschot, L., Langohr, R., Ranst, E. Van, Deckers, J., 2014. The soil map of the Flemish region converted to the 3rd edition of the World Reference Base for soil resources 139.

- Donovan, R.G., Stewart, H.E., Owen, S.M., MacKenzie, A.R., Hewitt, C.N., 2005.
 Development and Application of an Urban Tree Air Quality Score for Photochemical Pollution Episodes Using the Birmingham, United Kingdom, Area as a Case Study. Environ. Sci. Technol. 39, 6730–6738. https://doi.org/10.1021/ES050581Y
- Dyderski, M.K., Paź, S., Frelich, L.E., Jagodziński, A.M., 2018. How much does climate change threaten European forest tree species distributions? Glob. Chang. Biol. 24, 1150–1163. https://doi.org/10.1111/gcb.13925
- ECOPLAN, 2014. ECOPLAN Monitor [WWW Document]. URL http://www.ecosysteemdiensten.be/cms/
- Elith, J., H. Graham, C., P. Anderson, R., Dudík, M., Ferrier, S., Guisan, A., J. Hijmans, R., Huettmann, F., R. Leathwick, J., Lehmann, A., Li, J., G. Lohmann, L., A. Loiselle, B., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., McC. M. Overton, J., Townsend Peterson, A., J. Phillips, S., Richardson, K., Scachetti-Pereira, R., E. Schapire, R., Soberón, J., Williams, S., S. Wisz, M., E. Zimmermann, N., 2006. Novel methods improve prediction of species' distributions from occurrence data. Ecography (Cop.). 29, 129–151. https://doi.org/10.1111/j.2006.0906-7590.04596.x
- Elith, J., Phillips, S.J., Hastie, T., Dudík, M., Chee, Y.E., Yates, C.J., 2011. A statistical explanation of MaxEnt for ecologists. Divers. Distrib. 17, 43–57. https://doi.org/10.1111/j.1472-4642.2010.00725.x
- ESRI, 2017. ESRI, Redlands, US. http://www.esri.com/software/arcgis. last accessed 23/09/2017.
- Ficetola, G.F., Thuiller, W., Padoa-Schioppa, E., 2009. From introduction to the establishment of alien species: bioclimatic differences between presence and reproduction localities in the slider turtle. Divers. Distrib. 15, 108–116. https://doi.org/10.1111/j.1472-4642.2008.00516.x
- Gao, C., Shi, N.-N., Liu, Y.-X., Peay, K.G., Zheng, Y., Ding, Q., Mi, X.-C., Ma, K.-P., Wubet, T., Buscot, F., Guo, L.-D., 2013. Host plant genus-level diversity is the best predictor of

ectomycorrhizal fungal diversity in a Chinese subtropical forest. Mol. Ecol. 22, 3403–3414. https://doi.org/10.1111/mec.12297

- Gavish, Y., Marsh, C.J., Kuemmerlen, M., Stoll, S., Haase, P., Kunin, W.E., 2017. Accounting for biotic interactions through alpha-diversity constraints in stacked species distribution models. Methods Ecol. Evol. 8, 1092–1102. https://doi.org/10.1111/2041-210X.12731
- Godefroid, S., Koedam, N., 2007. Urban plant species patterns are highly driven by density and function of built-up areas. Landsc. Ecol. 22, 1227–1239. https://doi.org/10.1007/s10980-007-9102-x
- Goff, F.G., Dawson, G.A., Rochow, J.J., 1982. Site examination for threatened and endangered plant species. Environ. Manage. 6, 307–316. https://doi.org/10.1007/BF01875062
- Guisan, A., Rahbek, C., 2011. SESAM a new framework integrating macroecological and species distribution models for predicting spatio-temporal patterns of species assemblages.
 J. Biogeogr. 38, 1433–1444. https://doi.org/10.1111/j.1365-2699.2011.02550.x
- Hamm, C.A., 2013. Estimating abundance of the federally endangered Mitchell's satyr butterfly using hierarchical distance sampling. Insect Conserv. Divers. 6, 619–626. https://doi.org/10.1111/icad.12017
- Hjort, J., Hugg, T.T., Antikainen, H., Rusanen, J., Sofiev, M., Kukkonen, J., Jaakkola, M.S.,
 Jaakkola, J.J.K., 2016. Fine-Scale Exposure to Allergenic Pollen in the Urban
 Environment: Evaluation of Land Use Regression Approach. Environ. Health Perspect.
 124, 619–26. https://doi.org/10.1289/ehp.1509761
- Hoover, J.D., Kumar, S., James, S.A., Liesz, S.J., Laituri, M., 2017. Modeling hotspots of plant diversity in New Guinea. Trop. Ecol. 58, 623–640.
- Hope, D., Gries, C., Zhu, W., Fagan, W.F., Redman, C.L., Grimm, N.B., Nelson, A.L., Martin,C., Kinzig, A., 2003. Socioeconomics drive urban plant diversity, PNAS.
- Hüse, B., Szabó, S., Deák, B., Tóthmérész, B., 2016. Mapping an ecological network of green habitat patches and their role in maintaining urban biodiversity in and around Debrecen city (Eastern Hungary). Land use policy 57, 574–581.

https://doi.org/10.1016/J.LANDUSEPOL.2016.06.026

- Kemp, J.E., Ellis, A.G., 2017. Significant local-scale plant-insect species richness relationship independent of abiotic effects in the temperate cape floristic region biodiversity hotspot. PLoS One 12. https://doi.org/10.1371/journal.pone.0168033
- Khalili, A., van den Besselaar, P., de Graaf, K.A., 2018. Using Linked Open Geo Boundaries for Adaptive Delineation of Functional Urban Areas. Springer, Cham, pp. 327–341. https://doi.org/10.1007/978-3-319-98192-5_51
- Konarska, J., Uddling, J., Holmer, B., Lutz, M., Lindberg, F., Pleijel, H., Thorsson, S., 2016.
 Transpiration of urban trees and its cooling effect in a high latitude city. Int. J.
 Biometeorol. 60, 159–172. https://doi.org/10.1007/s00484-015-1014-x
- Lee, H., Mayer, H., Chen, L., 2016. Contribution of trees and grasslands to the mitigation of human heat stress in a residential district of Freiburg, Southwest Germany. Landsc. Urban Plan. 148, 37–50. https://doi.org/10.1016/j.landurbplan.2015.12.004
- Lepczyk, C.A., Aronson, M.F.J., Evans, K.L., Goddard, M.A., Lerman, S.B., MacIvor, J.S., 2017. Biodiversity in the City: Fundamental Questions for Understanding the Ecology of Urban Green Spaces for Biodiversity Conservation. Bioscience 67, 799–807. https://doi.org/10.1093/biosci/bix079
- McCune, J.L., 2016. Species distribution models predict rare species occurrences despite significant effects of landscape context. J. Appl. Ecol. 53, 1871–1879. https://doi.org/10.1111/1365-2664.12702
- McInnes, R.N., Hemming, D., Burgess, P., Lyndsay, D., Osborne, N.J., Skjøth, C.A., Thomas, S., Vardoulakis, S., 2017. Mapping allergenic pollen vegetation in UK to study environmental exposure and human health. Sci. Total Environ. 599–600, 483–499. https://doi.org/10.1016/J.SCITOTENV.2017.04.136
- Merow, C., Smith, M.J., Silander, J.A., 2013. A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. Ecography (Cop.). 36, 1058–1069. https://doi.org/10.1111/j.1600-0587.2013.07872.x

- Milanovich, J.R., Peterman, W.E., Barrett, K., Hopton, M.E., 2012. Do species distribution models predict species richness in urban and natural green spaces? A case study using amphibians. Landsc. Urban Plan. 107, 409–418. https://doi.org/10.1016/J.LANDURBPLAN.2012.07.010
- O'Brien, E.M., Whittaker, R.J., Field, R., 1998. Climate and woody plant diversity in southern Africa: relationships at species, genus and family levels. Ecography (Cop.). 21, 495–509. https://doi.org/10.1111/j.1600-0587.1998.tb00441.x
- Özenç, D.B., 2001. Methods of determining lime requirements of soils in the eastern black sea hazelnut growing region. Acta Hortic. 335–342. https://doi.org/10.17660/ActaHortic.2001.556.50
- Pearson, R.G., Raxworthy, C.J., Nakamura, M., Townsend Peterson, A., 2006. Predicting species distributions from small numbers of occurrence records: a test case using cryptic geckos in Madagascar. J. Biogeogr. 34, 102–117. https://doi.org/10.1111/j.1365-2699.2006.01594.x
- Peel, M.C., Finlayson, B.L., Mcmahon, T.A., 2007. Updated world map of the Köppen-Geiger climate classification, Hydrology and Earth System Sciences Discussions, European Geosciences Union. https://doi.org/<hal-00305098>
- Phillips, S.J., Anderson, R.P., Schapire, R.E., 2006. Maximum entropy modeling of species geographic distributions. Ecol. Modell. 190, 231–259. https://doi.org/10.1016/j.ecolmodel.2005.03.026
- Pottier, J., Dubuis, A., Pellissier, L., Maiorano, L., Rossier, L., Randin, C.F., Vittoz, P., Guisan,
 A., 2013. The accuracy of plant assemblage prediction from species distribution models
 varies along environmental gradients. Glob. Ecol. Biogeogr. 22, 52–63.
 https://doi.org/10.1111/j.1466-8238.2012.00790.x
- R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Roman, L.A., Scharenbroch, B.C., Östberg, J.P.A., Mueller, L.S., Henning, J.G., Koeser, A.K.,

Sanders, J.R., Betz, D.R., Jordan, R.C., 2017. Data quality in citizen science urban tree inventories. Urban For. Urban Green. 22, 124–135. https://doi.org/10.1016/J.UFUG.2017.02.001

- Salmond, J.A., Tadaki, M., Vardoulakis, S., Arbuthnott, K., Coutts, A., Demuzere, M., Dirks, K.N., Heaviside, C., Lim, S., Macintyre, H., McInnes, R.N., Wheeler, B.W., 2016. Health and climate related ecosystem services provided by street trees in the urban environment. Environ. Heal. 15, S36. https://doi.org/10.1186/s12940-016-0103-6
- Scholz, T., Hof, A., Schmitt, T., 2018. Cooling Effects and Regulating Ecosystem Services Provided by Urban Trees—Novel Analysis Approaches Using Urban Tree Cadastre Data. Sustainability 10, 712. https://doi.org/10.3390/su10030712
- Selmi, W., Weber, C., Rivière, E., Blond, N., Mehdi, L., Nowak, D., 2016. Air pollution removal by trees in public green spaces in Strasbourg city, France. Urban For. Urban Green. 17, 192–201. https://doi.org/10.1016/J.UFUG.2016.04.010
- Skowronek, S., Ewald, M., Isermann, M., Van De Kerchove, R., Lenoir, J., Aerts, R., Warrie, J., Hattab, T., Honnay, O., Schmidtlein, S., Rocchini, D., Somers, B., Feilhauer, H., 2017.
 Mapping an invasive bryophyte species using hyperspectral remote sensing data. Biol. Invasions 19, 239–254. https://doi.org/10.1007/s10530-016-1276-1
- Taylor, L., Hochuli, D.F., 2017. Defining greenspace: Multiple uses across multiple disciplines.
 Landsc. Urban Plan. 158, 25–38.
 https://doi.org/10.1016/J.LANDURBPLAN.2016.09.024
- Thomsen, P., Bühler, O., Kristoffersen, P., 2016. Diversity of street tree populations in larger
 Danish municipalities. Urban For. Urban Green. 15, 200–210. https://doi.org/10.1016/J.UFUG.2015.12.006
- Threlfall, C.G., Mata, L., Mackie, J.A., Hahs, A.K., Stork, N.E., Williams, N.S.G., Livesley, S.J., 2017. Increasing biodiversity in urban green spaces through simple vegetation interventions. J. Appl. Ecol. 54, 1874–1883. https://doi.org/10.1111/1365-2664.12876

Thuiller, W., Pollock, L.J., Gueguen, M., Münkemüller, T., 2015. From species distributions to

meta-communities. Ecol. Lett. 18, 1321–1328. https://doi.org/10.1111/ele.12526

- Trotta-Moreu, N., Lobo, J.M., 2010. Deriving the Species Richness Distribution of Geotrupinae (Coleoptera: Scarabaeoidea) in Mexico From the Overlap of Individual Model Predictions. Environ. Entomol. 39, 42–49. https://doi.org/10.1603/EN08179
- Van Landuyt, W., Vanhecke, L., Brosens, D., 2012. Florabank1: a grid-based database on vascular plant distribution in the northern part of Belgium (Flanders and the Brussels Capital region). PhytoKeys 12, 59–67. https://doi.org/10.3897/phytokeys.12.2849
- Van Rompaey, E., 1943. Cartes Floristiques. Bull. la Société R. Bot. Belgique / Bull. van K. Belgische Bot. Ver. 75, 48–56. https://doi.org/10.2307/20791920
- Vervoort, P., 2016. A healthy urban future for Flanders? Reducing the gap in knowledge for spatial policy and outlining consequences for governance, in: 10th AESOP YA Conference. AESOP, Ghent.
- Vriens, L., De Knijf, G., De Saeger, S., Guelinckx, R., Oosterlynck, P., Van Hove, M., Paelinckx, D., 2011. De Biologische Waarderingskaart. Biotopen en hun verspreiding in Vlaanderen en het Brussels Hoofdstedelijk Gewest., in: Mededelingen van Het Instituut Voor Natuur- En Bosonderzoek. INBO.M.2011.1, Brussels, p. 416.
- Weinberger, K.R., Kinney, P.L., Robinson, G.S., Sheehan, D., Kheirbek, I., Matte, T.D., Lovasi, G.S., 2016. Levels and determinants of tree pollen in New York City. J. Expo. Sci. Environ. Epidemiol. 28, 119. https://doi.org/10.1038/jes.2016.72
- Wolch, J.R., Byrne, J., Newell, J.P., 2014. Urban green space, public health, and environmental justice: The challenge of making cities 'just green enough.' Landsc. Urban Plan. 125, 234–244. https://doi.org/10.1016/J.LANDURBPLAN.2014.01.017
- Zalesny, R.S., Bauer, E.O., 2007. Selecting and utilizing Populus and Salix for landfill covers: Implications for leachate irrigation. Int. J. Phytoremediation 9, 497–511. https://doi.org/10.1080/15226510701709689

Appendix 1:

Table 1: Ecoplan land cover classification (in Dutch). Retrieved from: <u>https://www.uantwerpen.be/nl/onderzoeksgroep/ecoplan/ecoplan-tools/ecoplan-geoloket/</u>

Complex	Code	Dutch name
water	10101	Stilstaand water
	10201	Getijde mesohaline
	10202	Getijde Oligohalien
	10203	Getijde zoet
	10204	Zoet
	10301	Zee
Bush land	1101	Ruigten en pioniersvegetatie
	1201	Struweel
	1301	Struiken boomgaard
	1302	Laagstam
	1303	Hoogstam
	1401	Ander hooggroen
	1402	Ander laaggroen
Forest complexes	2101	Berk
r orest comprenes	2102	Benk
	2102	Beuk – naaldhout
	2103	Eik
	2101	Eik – naaldhout
	2106	Populier
	2100	Populier – naaldhout
	2107	Ander loofhout
	2109	Ander loofhout – naaldhout
	2201	lork
	2202	Lork – loofhout
	2203	Fiinspar
	2204	Fiinspar – loofhout
	2205	Zwarte den
	2206	Zwarte den – loofhout
	2207	Grove den
	2208	Grove den – loofhout
	2209	Ander naaldhout
	2210	Ander naaldhout – loofhout
Grasslands	3101	Voedselarm grassland
	3102	Voedselrijk grassland
	3201	Voedselrijk grasland
Heath lands	4101	Droge heide
	4201	Vochtige heide
Bare soils	5101	Kale bodem
	5201	Duinen
	5301	Strand
	5401	Niet verharde wegen
Agricultural land	6101	Aardappel
	6102	Mais
	6103	Graan
	6104	Zaden
	6105	Peulvruchten
	6106	Suikerbiet

	6107	Groenten
	6108	Kruiden
	6109	Fruit
	6110	Vlas en hennep
	6111	Voedergewassen
	6112	Andere akker
	6201	Eénjarige sierplanten
	6202	Meerjarige sierplanten
	6203	Kerstbomen
Wet vegetation types	7101	Moeras
	7201	Rietland
	7301	Alluvial bos
	8101	Slikken
	8201	Schorren
Built up area	9101	Gebouwen
	9201	Verhard
	9202	Verhard andere

Table 2:	Biological	valuation	map	classification	of habitat types	in (in	Dutch).	Retreived	from:
https://wv	ww.geopunt.b	e/catalogu	s/data	setfolder/bf31d	5c7-e97d-4f71-a	453-	5584371e	7559	

Complex	Code(s)	Dutch name	
Stagnant water	ad	Bezinkingsbekken	
	ae, aer, aev	Eutroof	
	ap, apo, app	Diep of zeer diep water	
	ao, aoo, aom	Oligotroof tot mesotroof water	
	ah	Brak of zilt water	
Swamps	ms	Zuur laagveen	
	mm	Galigaanvegetatie	
	mk	Alkalisch laagveen	
	mc	Grote zeggenvegetatie	
	mz	Brak tot zilt moeras	
	mr	Rietland en andere Phragmition vegetaties	
	md	Drijfzoom en/of drijftil	
Grasslands	ha	Struisgrasvegetatie	
	hc	Dotterbloemgrasland	
	hk	Kalkgrasland	
	hm, hmm, hme	Vochtig schraalgrasland	
	hmo	Vochtig heischraalgrasland	
	hn	Droog heischraalgrasland	
	hu	Mesofiel hooiland	
	hj	Vochtig grasland gedomineerd door russen	
	hp×, hpr×	Soortenrijk premanent cultuurgrasland	
	hpr(x)+da, hp(x)+da,	Soortenrijk premanent cultuurgraslandmet zilte	
	h+da	elementen	
	hp	Soortenarm permanent cultuurgrasland	
	hx	Zeer soortenarm, vaak tijdelijk grasland	
	hf, hfc, hft	Moerasspirearuigte	
	hr	verruigd grasland	
	hz	grasland op zware metalen vergiftiged bodems	
	hpr	weidelandcomplex met veel sloten of microreliëf	

High fenn	t	hoogveen
Heath lands	cg	Droge struikheivegetatie
	cv	Droge heide met bosbes
	ce, ces	Vochtige tot natte dopheivegetatie
	cm	Gedegradeerde heide met dominantie van
		pijpenstrootje
	ср	Gedegradeerde heide met dominantie van
		adelaarsvaren
	cd	Gedegradeerde heide met dominantie van
Dunes and tidal flats	ds	Slikken
	da	Schorre
	dd	Stuifduinen aan de kust
	dl	Strand
	dz	Zandbank
Bush land	sd(b)	Duinstruweel
Dubhi hund	su(c)	Doornstruweel
	sk	Struweel op kalkriike bodem
	sg. sgu, sgb	Brem- en gaspeldoornstruweel
	sg, sga, sge	Opslag van allerlei aard
	sf	Vochtig wilgenstruweel op voedselrijke bodem
	50	Vochtig wilgenstruweel op venige of zure grond
	sm	Gagelstruweel
	se	Kapylakte
Beech forests	fe	Beukenbos met wilde hyacint
	fa	Beukenbos met vooriaarsflora, zonder wilde
		hyacint
	fm	Beukenbos met parelgras en lievevrouwebedstro
	fk	Beukenbos op mergel
	fl	Beukenbos met witte veldbies
	fs	Zuur beukenbos
Oak forests	qe	Eiken-haagbeukenbos met wilde hyacint
	qa	Eiken-haagbeukenbos
	qk	Eiken-haagbeukenbos op mergel
	ql	Eikenbos met witte veldbies
	qs	Zuur eikenbos
	qb	Eikenberkenbos
Wet forests	vc	Bronbos
	va	Alluviaal elzen-essenbos
	vf	Elzen-eikenbos
	vn	Nitrofiel alluviaal elzenbos
	vm	Elzenbroek
	VO	Oligotroof elzenbroek met veenmossen
	vt	Berkenbroek
Ruderal forests	ru, rud	ruderaal olmenbos
Coniferous forests	pi, ppi, pa, ppa	Naaldhoudsbestand zonder ondergroei
	pmh, pms, pmb, ppmh,	Naaldhoutbestand met ondergroei
	ppms, ppmb	
Poplar forests	lhi, lhb, lsi, lsb, lsh	Populiersbestand

Other deciduous forests	n	Loofhout aanplant (exclusief populier)
Agricultural fields	bk, bl, bs, bu	Akker
Urban and built up	ua, ud, un, ur	Bebouwing
areas	uv, uc	Recreatiegebied
	ui	Industrie
Small landscape	kj	Hoogstamboomgaard
elements	kb	Bomenrij
	kh	Houtkant
	khw	Houtwal
	k	lijnvormige begroeiing van perceelsranden, sloten en bermen
	kk	Doline, ingang ondergrondse mergelgroeve
	km	Muurvegetatie
	kn	Veedrinkpoel
	kt	Talud
	kw	Holle weg
Other mapped	ko	Stort
elements	kr	Groeve
	kf	Voormalig militair fort
	kg	Terril
	kz	Opgehoogd terrein
	ki	Vliegveld
	kg	Kwekerij of Serre
	ka	Eendenkooi
	kr	Rots
	kd	Dijk
	ks	Verlaten spoorweg
	kl	Laagstamboomgaard
	kp	Park
	kpa	Arboretum
	kpk	Kasteelpark