Made available by Hasselt University Library in https://documentserver.uhasselt.be

Non-commutative crepant resolutions for some toric singularities. Il Non Peer-reviewed author version

SPENKO, Spela & VAN DEN BERGH, Michel (2020) Non-commutative crepant resolutions for some toric singularities. II. In: JOURNAL OF NONCOMMUTATIVE GEOMETRY, 14 (1), p. 73 -103.

DOI: 10.4171/JNCG/359 Handle: http://hdl.handle.net/1942/31940

NON-COMMUTATIVE CREPANT RESOLUTIONS FOR SOME TORIC SINGULARITIES II

ŠPELA ŠPENKO AND MICHEL VAN DEN BERGH

ABSTRACT. Using the theory of dimer models Broomhead proved that every 3-dimensional Gorenstein affine toric variety Spec R admits a toric noncommutative crepant resolution (NCCR). We give an alternative proof of this result by constructing a tilting bundle on a (stacky) crepant resolution of Spec R using standard toric methods. Our proof does not use dimer models.

1. INTRODUCTION

Throughout k is an algebraically closed base field of characteristic zero. Let R be a normal Gorenstein domain. A non-commutative crepant resolution (NCCR) [DITV15, Leu12, ŠVdB17a, VdB04, Wem16] of R is an R-algebra of finite global dimension of the form $\Lambda = \text{End}_R(M)$ which in addition is Cohen-Macaulay as R-module and where M is a non-zero finitely generated reflexive R-module. In [ŠVdB17a] we studied NCCRs of rings of invariants which are given by modules of covariants. More precisely, we assumed $R = \text{Sym}(W)^G$ where G is a reductive group and W is a representation of G and $M = (U \otimes_k \text{Sym}(W))^G$ for a representation U of G. When G is a product of a torus and a finite abelian group such invariant rings are coordinate rings of affine toric varieties and in that case NCCRs given by modules of covariants are called "toric" NCCRs. See e.g. [Boc12].

In loc. cit. we were able to construct toric NCCRs in many cases (e.g. when W is self-dual). However the following beautiful result of Broomhead remained outside the scope of our methods.

Theorem 1.1. [Bro12, Theorem 8.6] The coordinate ring of a 3-dimensional Gorenstein affine toric variety admits a toric NCCR.

Broomhead proves this result by exploiting the close relationship between 3dimensional Gorenstein affine toric varieties and so-called "dimer models" (certain bipartite graphs embedded in a real toric surface). More precisely, it is well-known that the fan corresponding to a 3-dimensional Gorenstein affine toric variety is a

²⁰¹⁰ Mathematics Subject Classification. 13A50, 14M25, 32S45.

Key words and phrases. Toric varieties, tilting bundle, noncommutative resolution.

The first author is a FWO [PEGASUS]² Marie Skłodowska-Curie fellow at the Free University of Brussels (funded by the European Union Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 665501 with the Research Foundation Flanders (FWO)). During part of this work she was also a postdoc with Sue Sierra at the University of Edinburgh.

The second author is a senior researcher at the Research Foundation Flanders (FWO). While working on this project he was supported by the FWO grant G0D8616N: "Hochschild cohomology and deformation theory of triangulated categories".

cone over a lattice polygon and an algorithm by Gulotta [Gul08] (or an alternative algorithm by Ishii and Ueda [IU15]) associates to every lattice polygon P a "consistent" dimer model whose lattice polygon of "perfect matchings" coincides with P. Any dimer model also possesses an associated superpotential algebra and this yields the required toric NCCR.

Instead of Broomhead's algebraic approach one may try to construct the NCCR as the endomorphism algebra of a tilting bundle on a (stacky) crepant resolution of singularities. In fact in [ŠVdB17b] we achieved this in a relatively straightforward way by modifying the main method of [VdB04]. The resulting NCCRs are however generally non-toric and so we do not obtain a proof of Theorem 1.1 in this way.

If we want to obtain toric NCCRs then the tilting bundle has to be *split* (= sum of line bundles). In [IU15] Ishii and Ueda give indeed a proof of Theorem 1.1 by constructing a (tautological) split tilting bundle on a toric crepant resolution which is the solution of a suitable moduli problem related to dimer models (see [IU15, Theorem 1.4], [IU08, Theorem 6.4]).

The purpose of this note is to give a proof of Theorem 1.1 which *avoids* dimer models altogether and which uses traditional toric methods instead. In the interest of full disclosure we mention that our proof still depends on the initial combinatorial input provided by Gulotta and Ishii-Ueda. A distinguishing feature of our approach is that we focus on "small" toric resolutions, i.e. those without exceptional divisors. Except for the most simple cases such resolutions cannot be schemes and must be Deligne-Mumford stacks. In particular they cannot be realized as moduli spaces of vector bundles (in contrast to the projective crepant resolutions used by Ishii and Ueda) as they generally have non-connected stabilizers.

The version of our proof starting from Gulotta's approach is the least combinatorially demanding, so we discuss it first. Let Spec R be a 3-dimensional Gorenstein affine toric variety¹ and let P be the lattice polygon corresponding to R (see §3). Gulotta's "algorithm" [Gul08] first embeds P into a rectangle P_0 and then reconstructs P from P_0 by iteratively removing corner triangles (see Figure 7.1). After appropriately subdividing the removed triangles we obtain a triangulation of $P_0 - P$ and we complete it to a triangulation of P_0 by adding diagonals to P. The affine toric variety Spec R_0 corresponding to P_0 is a so-called "generalized conifold" and it has a standard toric NCCR (see Remark 7.2). We show that the latter is in fact obtained from a split tilting bundle on the stacky crepant resolution \mathcal{Y}_0 of Spec R_0 corresponding to the triangulation of P_0 (see Appendix A). The restriction to the open substack \mathcal{Y} of \mathcal{Y}_0 corresponding to the triangulation of P is then shown to be a tilting bundle on \mathcal{Y} . Note that this is not a formality as being tilting is not a local property.

Another version of our proof starts with the Ishii-Ueda approach (see §8). In this case we embed P in a triangle P_0 and we reconstruct P from P_0 by iteratively removing corner vertices and taking convex hulls of the remaining lattice points. See Figure 8.1. This yields again a natural triangulation of $P_0 - P$ and we continue in a similar manner as above.

¹As in [CLS11] we assume throughout that a normal toric affine variety is of the form $\operatorname{Spec}(k[\sigma^{\vee} \cap M])$ for a rational strongly convex polyhedral cone in N (see e.g. [CLS11, Remark 1.2.19(1), Theorem 1.3.5]).

Besides the above results we also discuss some side results in Appendices B, C (Appendix A is devoted to a technical result necessary for the paper).

In Proposition B.3 we give a useful combinatorial criterion for recognizing NC-CRs of three-dimensional toric Gorenstein singularities.

In Appendix C we elaborate on the relationship between tilting bundles and NCCRs. In [IU16] it was established by Ishii and Ueda that if X is an arbitrary *projective* crepant resolution of Spec R then every NCCR of Spec R is the endomorphism ring of some tilting bundle on X. As the proof depends on realizing X as a GIT moduli space it appears to use the projectivity hypothesis in an essential way and we do not know if the result is true otherwise. We note however that there certainly exist particular instances of non-projective crepant resolutions for which the result remains true. In appendix C we discuss such an example using our combinatorial techniques.

2. Acknowledgement

The authors thank Seung-Jo Jung, Martin Kalck, Sasha Kuznetsov and Michael Wemyss for interesting discussions concerning the material in this paper.

3. Preliminaries and notation

Let $N = \mathbb{Z}^n$, $M = \text{Hom}(N,\mathbb{Z})$ and let \langle , \rangle the natural pairing between Mand N. Let $\{n_i \in N \mid 1 \leq i \leq k\}$ be a set of vectors which generates $N_{\mathbb{R}}$ such that

$$\sigma = [n_1, \dots, n_k] = \{ r_1 n_1 + \dots + r_k n_k \in N_{\mathbb{R}} \mid r_i \ge 0 \}$$

is a rational strongly convex polyhedral cone in $N_{\mathbb{R}}$. We do not assume that $\{n_1, \ldots, n_k\}$ is a minimal set of generators of σ . Let

$$\sigma^{\vee} = \{ m \in M_{\mathbb{R}} \mid \langle m, n_i \rangle \ge 0 \text{ for all } 1 \le i \le k \}$$

be the dual cone of σ . We denote by $R_{\sigma} = k[\sigma^{\vee} \cap M]$ the associated semigroup algebra, and set $X_{\sigma} = \operatorname{Spec} R_{\sigma}$.

Let $\rho : M \hookrightarrow \mathbb{Z}^k$ be defined by $m \mapsto (\langle m, n_1 \rangle, \dots, \langle m, n_k \rangle)$. We set $G = \operatorname{Hom}(\mathbb{Z}^k/\rho(M), k^*) \subseteq (k^*)^k$ so that $X(G) = \mathbb{Z}^k/\rho(M)$.

Let e_i be the *i*-th generator of \mathbb{Z}^k , and let β_i be its image in X(G). Then $M \cong \rho(M) = \{(a_i) \in \mathbb{Z}^k \mid \sum_i a_i \beta_i = 0\}$, and $\sigma^{\vee} \cap M \cong \rho(\sigma^{\vee} \cap M) = \{(a_i) \in \mathbb{N}^k \mid \sum_i a_i \beta_i = 0\}$. Thus $R_{\sigma} \cong k[x_1, \ldots, x_k]^G$ where G acts on x_i by β_i , and $X_{\sigma} = Y/\!\!/G$ for $Y = k^k$ with the G-weights $(-\beta_i)_{i=1}^k$.

For $b \in \mathbb{Z}^k$ let

(3.1)
$$M_b = \{ m \in M \mid \langle m, n_i \rangle \ge -b_i \text{ for all } 1 \le i \le k \}, \quad \mathsf{M}_b = kM_b.$$

Note that $R_{\sigma} = \mathsf{M}_0 = kM_0$ and $b - b' \in \rho(M)$ implies $M_b \cong M_{b'}$ as modules over M_0 . Denote the image of b in X(G) by χ_b . To $m \in M_b$ we associate $(a_i)_i =$ $(\langle m, n_i \rangle + b_i)_i \in \mathbb{N}^k$. With this identification we get $M_b \cong \rho(M_b) + b = \{(a_i) \in \mathbb{N}^k \mid \sum_i a_i \beta_i = -\chi_b\}$. Thus M_b is isomorphic to the module of covariants $M(\chi_{-b}) = (k[x_1, \ldots, x_k] \otimes \chi_{-b})^G$.

If $[n_1, \ldots, n_k]$ is a minimal presentation of σ then the action of G on Y is generic in the sense of [ŠVdB17a, Def.§1.3.4] (see [ŠVdB17a, §11.6.1]). In that case $\chi_{-b} \mapsto M(\chi_{-b}) = M_b$ provides isomorphism $X(G) = \operatorname{Cl}(R_{\sigma})$ (see e.g. [CLS11, Section 5.1] or [ŠVdB17a, Lemma 4.1.3]). In particular M_b is reflexive and

(3.2)
$$\operatorname{Hom}_{R_{\sigma}}(\mathsf{M}_{b},\mathsf{M}_{b'})=\mathsf{M}_{b'-b}.$$

We will often consider the special case where the $(n_i)_i$ are the generators of the 1-dimensional cones in a simplicial fan Σ with $|\Sigma| = \sigma$. In that case we will denote the corresponding stacky fan² $(\Sigma, (n_i)_{i=1}^k)$ by Σ (see [BH06]). We denote by X_{Σ} (resp. X_{Σ}) the corresponding smooth toric DM stack (resp. (possibly) singular toric variety). As Σ is simplicial, X_{Σ} is an orbifold and X_{Σ} is its coarse moduli space.

We have that X_{Σ} (resp. X_{Σ}) equals Y_{Σ}/G (resp. $Y_{\Sigma}//G$) for an open subvariety $Y_{\Sigma} \subset Y$ which consists of all points $z = (z_1, \ldots, z_k)$ such that the set of n_i for which $z_i = 0$ is contained in a cone of Σ (see [BH06, Section 2]). In the following diagram we assemble the varieties/stacks we have defined and we give names to some of the canonical maps between them.



Performing the construction $b \mapsto \mathsf{M}_b$ for each of the cones in Σ we obtain a reflexive rank 1 $\mathcal{O}_{X_{\Sigma}}$ -module which we denote by $\mathcal{M}_{\Sigma,b}$. Recall the following result.

Lemma 3.1. Put

$$\mathcal{M}_{\Sigma,b} = \mu_{s,*}(\chi_{-b} \otimes \mathcal{O}_{Y_{\Sigma}/G})$$

Then $\mathcal{M}_{\Sigma,b}$ is a line bundle on X_{Σ} and

$$\mathbb{Z}^k \mapsto \operatorname{Pic}(X_{\Sigma}) : b \mapsto \mathcal{M}_{\Sigma,b}$$

is a morphism of abelian groups. We have

(3.4)
$$\mathcal{O}(D_b) = \mathcal{M}_{\Sigma,b} = \pi_{s,*} \mathcal{M}_{\Sigma,b}$$

where $D_b = \sum_{i=1}^{k} b_i D_{n_i}$ is the equivariant Weil divisor associated to b as in [CLS11, §4.1]. Moreover

(3.5)
$$\Gamma(X_{\Sigma}, \mathcal{M}_{\Sigma, b}) = \mathsf{M}_{b}.$$

Proof. Since μ_s is an isomorphism the first two claims are obvious. The first equality in (3.4) is [CLS11, Proposition 5.3.7]. The last equality in (3.4) is local on X_{Σ} and can be verified on the open covering given by the maximal cones using Proposition A.1. Finally (3.5) follows from [CLS11, Proposition 4.3.2].

Furthermore, for $m \in M$ we have by [CLS11, Theorem 9.1.3]

(3.6)
$$H^p(X_{\Sigma}, \mathcal{O}(D_b))_m = \tilde{H}^{p-1}(V_{D_b,m}, k),$$

where

(3.7)
$$V_{D_b,m} = \bigcup_{\sigma \in \Sigma} \operatorname{conv} \{ v \in \sigma(1) \mid \langle m, v \rangle < -b_v \}$$
$$= \bigcup_{\sigma \in \Sigma} \operatorname{conv} \{ v \in \sigma(1) \mid s_v^b(m) = -\},$$

 $^{^{2}}$ A stacky fan is a simplicial fan together with generators for the one dimensional cones.

and where we denote

(3.8)
$$s_v^b(m) = \text{sign of } (\langle m, v \rangle + b_v)$$

Convention 3.2. Except for §5,6 we assume that there is $m \in M$ such that $\langle m, n_i \rangle = 1$ for all $1 \leq i \leq k$. In particular X_{σ} is Gorenstein (see e.g. [CLS11, Proposition 11.4.12]). Changing bases we may assume $m = (0, \ldots, 0, 1)$ and hence σ is a cone over a convex lattice polyhedron $P \subseteq \mathbb{R}^{n-1} \times \{1\} \subset \mathbb{R}^n = M_{\mathbb{R}}$, and we write $\sigma = \sigma_P$. We abbreviate $R_{\sigma_P}, X_{\sigma_P}$ as R_P, X_P , respectively.

3.1. Tilting bundles and NCCRs. Recall that a tilting bundle on an algebraic stack \mathcal{Y} is a vector bundle \mathcal{T} such that the following conditions hold:

- (T1) \mathcal{T} generates $D_{\text{Qch}}(\mathcal{Y})$;
- (T2) $\operatorname{Ext}^{i}_{\mathcal{V}}(\mathcal{T},\mathcal{T}) = 0 \text{ for } i > 0.$

We then have $D_{\text{Qch}}(\mathcal{Y}) \cong D(\Lambda)$ where $\Lambda = \text{End}_{\mathcal{Y}}(\mathcal{T})$. If \mathcal{Y} is a noetherian Deligne Mumford stack then we write $\mathcal{D}(\mathcal{Y})$ for $D^b_{\text{coh}}(\mathcal{Y})$. Similarly if Λ is a noetherian ring then we write $\mathcal{D}(\Lambda) = D^b_f(\Lambda)$. If \mathcal{Y} is a smooth separated DM-stack then $D_{\text{Qch}}(\mathcal{Y})$ is compactly generated by $\mathcal{D}(\mathcal{Y})$ (see [HR14, Theorem B]) and hence if \mathcal{T} is a tilting bundle on \mathcal{Y} then \mathcal{T} "classically generates" $\mathcal{D}(\mathcal{Y})$ (see [BVdB03]) and moreover if $\Lambda = \text{End}_{\mathcal{Y}}(\mathcal{T})$ is noetherian then $\mathcal{D}(\mathcal{Y}) \cong \mathcal{D}(\Lambda)$. Recall the following

Proposition 3.3. Let P be as in Convention 3.2. Choose a triangulation of P without extra vertices³ and let Σ be the corresponding fan. Let \mathcal{T} be a tilting bundle on X_{Σ} . Then $\Lambda = \operatorname{End}_{X_{\Sigma}}(\mathcal{T})$ is an NCCR for R_P corresponding to $M = \Gamma(X_{\Sigma}, \mathcal{T})$. If \mathcal{T} is toric of the form $\mathcal{T} = \bigoplus_{b \in S} \mathcal{M}_{\Sigma,b}$ then $\Lambda = \operatorname{End}_{R_P}(\bigoplus_{b \in S} M_b)$.

Proof. We proceed similarly as in the proof of [ŠVdB17b, Corollary 2.4]. Since X_{Σ} is smooth and $\mathcal{D}(X_{\Sigma}) \cong \mathcal{D}(\Lambda)$, gl dim $\Lambda < \infty$. It follows from Proposition A.1 (c.f. proof of Lemma A.2) that π_s is identity in codimension 1. Hence $(\pi_s)_*$ defines a (monoidal) equivalence between categories of reflexive sheaves on X_{Σ} and X_{Σ} . As τ does not contract a divisor by the hypothesis on the triangulation, τ_* defines a (monoidal) equivalence between categories of reflexive sheaves on X_{Σ} and X_{Γ} . Thus, $\operatorname{End}_{X_{\Sigma}}(\mathcal{T}) \cong \operatorname{End}_{R_P}(\Gamma(X_{\Sigma}, \mathcal{T}))$ as $\Gamma(X_P, (\tau \pi_s)_* \mathcal{T}) = \Gamma(X_{\Sigma}, \mathcal{T})$.

To show that Λ is Cohen-Macaulay one can proceed as in [SVdB17b, Corollary 2.8] thanks to Lemma A.2. The last claim follows from (3.5).

4. Strategy

As alluded to in the introduction our main steps towards the construction of an NCCR of R_P (for 2-dimensional P) are the following:

(1) Embed P in a rectangle (resp. lattice triangle) P_0 , and choose a triangulation of P_0 which contains a triangulation of P without extra vertices. To obtain these triangulations we use Gulotta's (see §7) (resp. the Ishii-Ueda (see §8)) inductive procedures.

Let Σ_0 , Σ be the fans corresponding to the triangulations of P_0 and P.

- (2) Construct a split tilting bundle \mathcal{T}_0 on the stacky crepant resolution X_{Σ_0} of R_{P_0} .
- (3) Restrict \mathcal{T}_0 to X_{Σ} to obtain a tilting bundle \mathcal{T} on X_{Σ} (this is not a formal step as the property of being tilting is not local).

³The existence of such triangulations follows from the theory of secondary fans [GKZ89, GKZ08]. See [CLS11, Proposition 15.2.9].

(4) Now $\Lambda = \operatorname{End}_{X_{\Sigma}}(\mathcal{T})$ yields an NCCR of R_P by Proposition 3.3.

The vanishing properties (T2) for \mathcal{T}_0 , \mathcal{T} in (2)(3) are proved using standard toric geometry (see Lemmas 7.1, 8.3), while the generation property (T1) for \mathcal{T} in (3) follows formally from the corresponding property of \mathcal{T}_0 in (2) (generation is compatible with restriction). The verification of the latter constitutes the heart of our proof. We proceed as follows:

- (a) We start with a standard NCCR Λ_0 of R_{P_0} given by $\bigoplus_{b \in S_0} \mathsf{M}_b$, $S_0 = \{(b_v)_{v \in V'_0}\}$, where V'_0 is the set of the vertices of P_0 (V'_0 yields the *minimal* presentation of σ_{P_0}). Clearly, depending on whether P_0 is a rectangle or a triangle, we have $|V'_0| = 4$ or 3.
- (b) Let V_0 be the set of vertices in the triangulation in (1) (V_0 gives a non-minimal presentation of σ_{P_0}). We develop an inductive procedure called *compatible convex induction* (see §5) which extends the collection $S_0 = \{(b_v)_{v \in V'_0}\}$ from (a) to a collection $S = \{(\tilde{b}_v)_{v \in V_0}\}$ (thus S_0 and S are in bijection via (-)) such that for $b, b' \in S_0$:

(4.1)
$$M_b = M_{\tilde{b}} \text{ and } M_{b-b'} = M_{\tilde{b}-\tilde{b}'}.$$

The construction of b from b is is done inductively and matching the inductive construction of the triangulation of P_0 (see (1)).

- (c) We define T₀ := ⊕_{b∈S} M_{Σ0,b} and we verify using the standard results on the cohomology of rank one reflexive sheaves on 3-dimensional toric varieties (see Lemmas 7.1, 8.3) that T₀ satisfies (T2). This verification is done inductively.
- (d) We use (4.1) to deduce that $\Lambda_0 \cong \operatorname{End}_{X_{\Sigma_0}}(\mathcal{T}_0)$.
- (e) We show that $\mathcal{D}(X_{\Sigma_0})$ has no non-trivial semi-orthogonal decompositions using the fact that it has trivial relative Serre functor over X_P (see Lemma A.3, Corollary A.5).
- (f) Since $\operatorname{Perf}(\Lambda_0) \cong \operatorname{Perf}(\operatorname{End}_{X_{\Sigma_0}}(\mathcal{T}_0))$ is fully faithfully embedded in $\mathcal{D}(X_{\Sigma_0})$ by (T2) (see (c)), and since gl dim $\Lambda_0 < \infty$ by (a) we have that $\operatorname{Perf}(\Lambda_0) = \mathcal{D}(\Lambda_0)$ and moreover the embedding is admissible by [PVdB15, Lemma 1.1.1]. So it is a factor in semi-orthogonal decomposition of $\mathcal{D}(X_{\Sigma_0})$. We now use (e) to conclude that $\mathcal{D}(\Lambda_0) \cong \mathcal{D}(X_{\Sigma_0})$ and hence \mathcal{T}_0 satisfies (T1).

Our constructions were inspired by Bocklandt's result [Boc12, Corollary 4.7] showing that Cohen-Macaulay modules of covariants are "preserved under projection". In fact using Bocklandt's result we may prove that Λ is Cohen-Macaulay, without (re)invoking toric geometry (see Appendix B for more details). However we have been unable to prove directly that Λ is an NCCR.

5. Convex induction

For $t = (t_1, ..., t_k), 0 \le t_i \le 1, \sum_i t_i = 1$, we set

(5.1)
$$b_{t,-} = (b_1, \dots, b_k, \left\lfloor \sum_i t_i b_i \right\rfloor), \quad b_{t,+} = (b_1, \dots, b_k, \left\lfloor \sum_i t_i b_i \right\rfloor),$$

where $\lfloor x \rfloor$, $\lceil x \rceil$ denote the largest (resp. the smallest) integer not greater (resp. smaller) than x.

Lemma 5.1. If σ is presented as $[n_1, \ldots, n_k, n_{k+1}]$, where $n_{k+1} = \sum_i t_i n_i \in N$ for some $t = (t_1, \ldots, t_k)$, $0 \le t_i \le 1$, $\sum_i t_i = 1$, then $M_b = M_{b_{t,-}} = M_{b_{t,+}}$.

Proof. The inclusions $M_{b_{t,-}}, M_{b_{t,+}} \subseteq M_b$ are trivial, while for $m \in M_b$ we have $\langle m, n_i \rangle \geq -b_i$ for all i, and thus $\langle m, n_{k+1} \rangle \geq -\sum_i t_i b_i \geq -\lceil \sum_i t_i b_i \rceil$, so $m \in M_{b_{t,+}}$, and as $\langle m, n_{k+1} \rangle \in \mathbb{Z}$ we also have $\langle m, n_{k+1} \rangle \geq -\lfloor \sum_i t_i b_i \rfloor$, implying $m \in M_{b_{t,-}}$.

If $a \in \mathbb{R}$ then by convention its "sign" will be + if $a \ge 0$ and - otherwise. For $b = (b_v)_{v \in S} \in \mathbb{Z}^l$, with $S \subset N$, and $m \in M$ recall (from (3.8))

(5.2)
$$s_v^b(m) = \text{sign of } (\langle m, v \rangle + b_v).$$

If we use indexed vectors $(v_i)_i$ then we also write $s_i^b(m)$ for $s_{v_i}^b(m)$.

Remark 5.2. The signs $s_v^b(m)$ appear often in toric geometry. See e.g. (3.6) and Proposition B.1 below.

Let notation be as in Lemma 5.1. Applying (5.2) with $(v_1, \ldots, v_k, v_{k+1}) = (n_1, \ldots, n_k, n_{k+1})$ we have $s_i^b(m) = s_i^{b_{t,+}}(m) = s_i^{b_{t,-}}(m)$ for $1 \le i \le k$ since $(b_{t,\pm})_i = b_i$ for $1 \le i \le k$. We will need the following simple lemma.

Lemma 5.3. We have $s_{k+1}^{b_{t,-}}(m), s_{k+1}^{b_{t,+}}(m) \in \{s_i^b(m) \mid 1 \le i \le k, t_i \ne 0\}.$

Proof. We may assume that all signs $s_i^b(m)$ for $1 \le i \le k$, $t_i \ne 0$ are the same, otherwise there is nothing to prove. Assume that they are all -; i.e. $\langle m, n_i \rangle + b_i \le -1$ for $1 \le i \le k$, $t_i \ne 0$. Then $\langle m, n_{k+1} \rangle + \sum_i t_i b_i \le -1$. Applying [?], [?] yields the desired conclusion. The case where all signs are + is similar.

Definition 5.4. Let $k, l \in \mathbb{N}$. An *induction datum* $(t_j^{(i)})_{ij} \in \mathbb{R}$ is a collection of vectors $t^{(i)} = (t_j^{(i)})_j \in \mathbb{R}^{i-1}$ for $k < i \le k+l$, such that $0 \le t_j^{(i)} \le 1$, $\sum_{j=1}^{i-1} t_j^{(i)} = 1$.

Definition 5.5. Let $[n_1, \ldots, n_k] \in N^k$. We say that $[n_1, \ldots, n_{k+l}] \in N^{k+l}$ is convexly induced from $[n_1, \ldots, n_k] \in N^k$ with induction datum $(t_j^{(i)})_{i,j}$ if for i > k

$$n_i = \sum_{j=1}^{i-1} t_j^{(i)} n_j.$$

Definition 5.6. Let $\mathfrak{s} \in \{\pm\}^l$ be a sign sequence. We say that an integer vector $\tilde{b} \in \mathbb{Z}^{k+l}$ is convexly induced from $b \in \mathbb{Z}^k$ with sign sequence \mathfrak{s} and induction datum $(t_i^{(i)})_{i,j}$ if \tilde{b} is obtained from b via $\tilde{b}_i = (\tilde{b}_{\leq i-1})_{t^{(i)},\mathfrak{s}_i}$ (see (5.1)) for i > k; i.e.

$$\tilde{b}_i = \begin{cases} b_i & \text{if } 1 \le i \le k, \\ \lfloor \sum_{j=1}^{i-1} t_j^{(i)} \tilde{b}_j \rfloor & \text{if } i > k \text{ and } \mathfrak{s}_i = -, \\ \lceil \sum_{j=1}^{i-1} t_j^{(i)} \tilde{b}_j \rceil & \text{if } i > k \text{ and } \mathfrak{s}_i = +. \end{cases}$$

We say that \hat{b} is convexly induced from b if it is convexly induced for *some* sign sequence (and a given induction datum).

Let $K \in \mathbb{N}$. We say that $\{\tilde{b}^j \mid 1 \leq j \leq K\} \subset \mathbb{Z}^{k+l}$ is compatibly convexly induced from $\{b^j \mid 1 \leq j \leq K\} \subset \mathbb{Z}^k$ if for each j, \tilde{b}^j is convexly induced from b^j for a fixed sign sequence and a fixed induction datum (i.e. they are both independent of j).

Below we will assume the induction datum is specified once and for all and we will usually not mention it afterwards.

Corollary 5.7. If \tilde{b} is convexly induced from b and I is a subsequence of $1, \ldots, k+l$ containing $1, \ldots, k$ then $M_b = M_{b'}$ for $b' = (\tilde{b}_i)_{i \in I}$.

Proof. Repeated application of Lemma 5.1 yields $M_{\tilde{b}} = M_b$. This implies $M_b = M_{\tilde{b}} \subset M_{b'} \subset M_b$ which gives the desired conclusion.

Lemma 5.8. If $\{\tilde{b}_1, \tilde{b}_2\}$ is compatibly convexly induced from $\{b, b'\}$ then $\tilde{b}_1 - \tilde{b}_2$ is convexly induced from b - b' (for the same induction datum and a possibly different sign sequence). In consequence, $M_{b'_1-b'_2} = M_{b_1-b_2}$ with b'_1, b'_2 like b' in Corollary 5.7.

Proof. By definition \tilde{b}_1 , \tilde{b}_2 are convexly induced from b_1, b_2 , respectively, with the same sign sequence. Let * be a sign. We do not necessarily have $(\tilde{b}_{1,\leq i-1})_{t^{(i)},*} - (\tilde{b}_{2,\leq i-1})_{t^{(i)},*} = ((\tilde{b}_1 - \tilde{b}_2)_{\leq i-1})_{t^{(i)},*}$. However, as $\lfloor x \rfloor - \lfloor y \rfloor, \lceil x \rceil - \lceil y \rceil \in \{\lfloor x - y \rfloor, \lceil x - y \rceil\}$ for any $x, y \in \mathbb{R}$, we still have $(\tilde{b}_{1,\leq i-1})_{t^{(i)},*} - (\tilde{b}_{2,\leq i-1})_{t^{(i)},*} = (\tilde{b}_{1,\leq i-1} - \tilde{b}_{2,\leq i-1})_{t^{(i)},*}$ for a possibly different sign *'. This is sufficient by Corollary 5.7. \Box

5.1. **Interval convex induction.** A simple version of convex induction consists of lattice points on an interval with the induction datum only referring to the nearest neighbours.

Definition 5.9. Let $n_0, n_{r+1} \in N$ be end points of an interval *I*. Let $\{n_1, \ldots, n_r\} \in N \cap I$ be such that $n_0, n_1, \ldots, n_r, n_{r+1}$ are consecutive distinct points in *I*. Assume furthermore we are given another ordering of these points

(5.3)
$$n'_1, \dots, n'_{r+2} = n_0, n_{r+1}, n_{i_1}, \dots, n_{i_r}.$$

Then for any $3 \leq j \leq r+2$ there exist unique j', j'' < j such that $[n'_{j'}, n'_{j''}[\cap \{n'_1, \ldots, n'_j\} = \{n'_j\}$. In particular have

(5.4)
$$n'_{j} = (1-t)n'_{j'} + tn'_{j''}$$

for some $t \in]0,1[$. The *interval induction datum* t associated to the ordering (5.3) is the induction datum obtained from (5.4).

We say that $(b_{n_0}, b_{n_{r+1}}, b_{n_1}, \dots, b_{n_r}) \in \mathbb{Z}^{r+2}$ is obtained by *interval convex induc*tion from $(b_{n_0}, b_{n_{r+1}}) \in \mathbb{Z}^2$ if it is obtained by convex induction from $(b_{n_0}, b_{n_{r+1}})$ for an interval induction datum (associated to some ordering of $(n_i)_{i=0,\dots,r+1}$ as in (5.3)) and an arbitrary sign sequence.

Lemma 5.10. Let $(n_i)_i$ be as above. Assume $(b_{n_0}, b_{n_{r+1}}, b_{n_1}, \dots, b_{n_r}) \in \mathbb{Z}^{r+2}$ is obtained by interval convex induction from $(b_{n_0}, b_{n_{r+1}}) \in \mathbb{Z}^2$. Then for all $m \in M$ the signs $s_{n_j}^b(m)$, $j = 0, \dots, r+1$, follow the pattern $+ \dots + - \dots -$ (possibly reflected and perhaps with no + or - present).

Proof. This follows by repeated application of Lemma 5.3.

6. TRIANGLE CONVEX INDUCTION

In this section we analyze a type of the convex induction that we will use for constructing a tilting bundle on a stacky resolution via the Ishii-Ueda approach. It is similar in spirit, but considerably more involved, than interval convex induction and it will be used only in §8.

⁴This innocent looking formula is in fact the key ingredient of our proofs!

Definition 6.1. Let $v_{-1}, v_0, v_{r+1} \in N$ be vertices of a solid triangle T such that $|v_{-1}, v_0[, v_{r+1}|$ contain no lattice points and let C be the convex hull of $N \cap (T \setminus \{v_{-1}\})$. Put $\{v_1, \ldots, v_r\} = N \cap \partial C \cap \operatorname{int} T$ such that $v_0, v_1, \ldots, v_r, v_{r+1}$ are consecutive boundary points of C.

For $j = 1, ..., r, v_j$ is in the convex hull of v_{-1}, v_{j-1}, v_{r+1} . In other words there exist unique $p_j, q_j, r_j \in [0, 1]$ such that $p_j + q_j + r_j = 1$ and

(6.1)
$$v_j = p_j v_{-1} + q_j v_{j-1} + r_j v_{r+1}.$$

The triangle induction datum \mathfrak{t} is the induction datum obtained from (6.1).

We say that $(b_{-1}, b_0, b_{r+1}, b_1, \ldots, b_r) \in \mathbb{Z}^{r+3}$ is obtained by triangle convex induction from (b_{-1}, b_0, b_{r+1}) if it is obtained by convex induction from (b_{-1}, b_0, b_{r+1}) for t and a sign sequence corresponding entirely of -'s; i.e., for $1 \leq j \leq r$

(6.2)
$$b_j = b_{v_j} = \lfloor p_j b_{-1} + q_j b_{j-1} + r_j b_{r+1} \rfloor$$

The following is our main result concerning triangle convex induction. It will play a similar role as Lemma 5.10.

Proposition 6.2 (see §6.3). Let $m \in M$ and let b, b' be obtained by triangle convex induction from (b_{-1}, b_0, b_{r+1}) , $(b'_{-1}, b'_0, b'_{r+1})$. Put c = b' - b. Then the possible sign patterns for $s^c_{v_{-1}}(m), s^c_{v_0}(m), \ldots, s^c_{v_{r+1}}(m)$ are

$$-\underbrace{+\cdots+}_{p}\underbrace{-\cdots-}_{q}\underbrace{+\cdots+}_{r} \quad or \ +\underbrace{-\cdots-}_{p}\underbrace{+\cdots+}_{q}\underbrace{-\cdots-}_{r}$$

where p, q and r may be equal to zero.

6.1. An explicit expression for the triangle induction datum.

6.1.1. Generalities. Assume $(a_l)_{l \in \mathbb{Z}} \in \mathbb{R}$. Below we will consider solutions $(x_l)_l \in \mathbb{R}$ to the second order recursion relation

$$(6.3) x_{l+1} = a_l x_l - x_{l-1}.$$

The following result follows easily from (6.3).

Lemma 6.3. Assume $(x_l)_l, (y_l)_l \in \mathbb{R}$ are solutions to (6.3). Then

$$\begin{vmatrix} x_l & x_{l+1} \\ y_l & y_{l+1} \end{vmatrix}$$

is independent of l.

We define $(q_{s,t})_{s,t\in\mathbb{Z}}$ in such a way that for all s, $(q_{s,t})_t$ is a solution to (6.3) with initial conditions $q_{s,s} = 0$, $q_{s,s+1} = 1$. Note that $q_{s,s-1} = -1$, $q_{s,s+2} = a_{s+1}$. The following equation shows that $(q_{s,t})_s$ is also a solution to (6.3) and hence the indices s, t of $q_{s,t}$ play symmetric roles:

(6.4)
$$q_{s+1,t} = a_s q_{s,t} - q_{s-1,t}.$$

Indeed: both sides of (6.4) are solutions to (6.3) and they are equal for t = s, s + 1 (using $q_{s-1,s+1} = a_s$). For $(x_t)_t$ a solution to (6.3) we get

(6.5)
$$x_t = q_{s,t} x_{s+1} - q_{s+1,t} x_s$$

by a similar argument. Both sides of this equation are solutions to (6.3) and moreover they agree for t = s, s + 1. Applying (6.5) with $(x_t)_t = (q_{s,t})_t$ (and the index s replaced by k) we get

(6.6)
$$q_{s,t} = q_{s,k+1}q_{k,t} - q_{s,k}q_{k+1,t}.$$

6.1.2. Continued fractions. Put $u_i := v_i - v_{-1}$. After choosing a suitable basis for the plane spanned by u_0 , u_{r+1} and performing a translation we may assume $N = \mathbb{Z}^2$ and

(6.7)
$$v_{-1} = (0,0), \quad v_0 = (0,1), \quad v_{r+1} = (n,-q)$$

where n > 0, $0 \le q < n$ and gcd(n,q) = 1 (see e.g. [CLS11, Proposition 10.1.1]). We will assume n > q > 0, since otherwise r = 0, and Proposition 6.2 holds trivially. Let

$$\frac{a_{q}}{q} = [a_{1}, \dots, a_{r}] := a_{1} - \frac{1}{a_{2} - \frac{1}{\dots - \frac{1}{a_{r}}}}$$

be the Hirzebruch-Jung continued fraction expansion of n/q where $a_i \ge 2$. The a_j , $j \ge 1$, may be computed by an Euclidean style algorithm starting from $i_0 = n$, $i_1 = q$, and inductively determining a_j, i_{j+1} via division with remainder:

(6.8)
$$i_{j-1} = a_j i_j - i_{j+1}, \quad 0 < i_{j+1} < i_j,$$

until we obtain $i_r = 1$. Then we set $a_r = i_{r-1}$ and $i_{r+1} = 0$. The vectors $u_0, u_1, \ldots, u_{r+1}$ are related by the equations

(6.9)
$$a_j u_j = u_{j-1} + u_{j+1},$$

 $1 \le j \le r$, (see [CLS11, Theorem 10.2.8(b), Theorem 10.2.5]).

Below we extend $[a_1, \ldots, a_r]$ to a doubly infinite sequence $(a_i)_{i \in \mathbb{Z}}$ by putting $a_i = 0$ for $i \notin \{1, \ldots, r\}$. We let $q_{s,t}$ be as in §6.1.1.

Remark 6.4. By (6.8), (6.9) $x_l = i_l$, $x_l = u_l$ are solutions to (6.3). We silently extend them to doubly infinite solutions to avoid having to keep track of cumbersome restrictions on indices in some formulas.

Remark 6.5. The $q_{s,t}$, for suitable s, t, appear frequently in combinatorics of Hirzebruch-Jung continued fractions. In [CLS11, Proposition 10.2.2], $q_{0,t+1}$, $q_{1,t+1}$ are denoted by P_t , Q_t , respectively, for $0 \le t \le r$. For example, they provide expressions for the truncated Hirzebruch-Jung continued fractions: $[b_s, \ldots, b_t] =$ $q_{s-1,s+t}/q_{s,s+t}$ for $1 \le s \le t \le r$. The formula is proved in loc. cit. for right truncations, i.e. for s = 1, and it is not difficult to see that it holds also for general truncations.

6.1.3. Application.

Lemma 6.6. Assume $0 \le s \le s' \le t' \le t \le r+1$ and $(s', t') \ne (s, t)$. Then

$$(6.10) 0 \le q_{s',t'} < q_{s,t}.$$

Proof. This follows easily by induction on s, t using the fact that $a_i \ge 2$ for $i = 1, \ldots, r$.

Lemma 6.7. We have

(6.11)
$$u_j = \frac{q_{j,l}}{q_{j-1,l}} u_{j-1} + \frac{1}{q_{j-1,l}} u_l$$

for $1 \leq j \leq l \leq r+1$.

Proof. By Remark 6.4 and Lemma 6.3, $C_l := q_{j-1,l}u_j - q_{j,l}u_{j-1}$ does not depend on j. Specializing to j = l we see that $C_l = u_l$.

NON-COMMUTATIVE CREPANT RESOLUTIONS FOR SOME TORIC SINGULARITIES II 11

Substituting $u_j = v_j - v_{-1}$ in (6.11) we obtain

(6.12)
$$v_{j} = \frac{q_{j-1,l} - q_{j,l} - 1}{q_{j-1,l}} v_{-1} + \frac{q_{j,l}}{q_{j-1,l}} v_{j-1} + \frac{1}{q_{j-1,l}} v_{l}$$

which gives a concrete expression for the coefficients in (6.1) (and also for the induction datum in Proposition 6.10 below). Note that we have

as both sides satisfy (6.3) by Remark 6.4 and are equal for t = r, r+1. Substituting this in (6.14) we obtain

(6.14)
$$v_j = \frac{i_{j-1} - i_j - 1}{i_{j-1}} v_{-1} + \frac{i_j}{i_{j-1}} v_{j-1} + \frac{1}{i_{j-1}} v_{r+1},$$

and hence by (6.2)

(6.15)
$$b_j = \left\lfloor \frac{i_{j-1} - i_j - 1}{i_{j-1}} b_{-1} + \frac{i_j}{i_{j-1}} b_{j-1} + \frac{1}{i_{j-1}} b_{r+1} \right\rfloor.$$

6.2. An expression for b. We give an explicit expression for the solutions to (6.15) in case $(b_{-1}, b_0, b_{r+1}) = (0, 0, d), 0 \le d < n$ (see Lemma 6.9 below with l = r + 1). Since $i_1 > \cdots > i_r = 1$ successive division with remainder yields a unique representation

$$d = \sum_{t=1}^{r} i_t d_t$$

with $d_t \in \mathbb{N}$ and

_

(6.16)
$$\sum_{t=k+1}^{r} i_t d_t < i_k \text{ for any } 0 \le k \le r$$

(for k = 0 this formula becomes d < n which is true by hypothesis). For $0 \leq j \leq r+1$ put

(6.17)
$$c_j = \sum_{t=1}^{j-1} q_{t,j} d_t$$

Using (6.13) and interpreting an empty sum as zero we have in particular

$$(6.18) c_0 = 0, c_{r+1} = d.$$

There is bound for the tail sums in (6.17) similar to (6.16).

Lemma 6.8. We have for
$$0 \le k \le j - 1 \le r$$

(6.19) $\sum_{t=k+1}^{j-1} q_{t,j} d_t < q_{k,j}.$

Proof. For k = j - 1 there is nothing to prove so we assume k < j - 1. According to [IU15, Lemma 3.5] d_t has the following properties

- (i) $0 \le d_t \le a_t 1$.
- (ii) If $d_s = a_s 1$, $d_t = a_t 1$ for s < t then there is s < l < t such that $d_l = a_l - 3.$

By (6.10) the sequence $(q_{t,j})_t$ is descending. Using this it is easy to see that among the sequences $(d_t)_t$ which satisfy (i), (ii) the one that gives the maximum value for $\sum_{t=k+1}^{j-1} q_{t,j}d_t$ is given by $d_{k+1} = a_{k+1} - 1$, $d_{k+2} = a_{k+2} - 2, \ldots, d_{j-1} = a_{j-1} - 2$. Hence the left-hand side of (6.19) is bounded by

$$q_{k+1,j} + \sum_{t=k+1}^{j-1} q_{t,j}(a_t - 2).$$

We compute

$$q_{k+1,j} + \sum_{t=k+1}^{j-1} q_{t,j}(a_t - 2) = q_{k+1,j} + \sum_{t=k+1}^{j-1} (q_{t-1,j} - q_{t,j} + q_{t+1,j} - q_{t,j}) \quad (6.4)$$
$$= q_{k+1,j} + q_{k,j} - q_{j-1,j} - q_{k+1,j} + q_{j,j}$$
$$= q_{k,j} - 1$$

which implies (6.19).

Lemma 6.9. We have for $1 \le j \le l \le r+1$

(6.20)
$$c_j = \left\lfloor \frac{q_{j,l}}{q_{j-1,l}} c_{j-1} + \frac{1}{q_{j-1,l}} c_l \right\rfloor.$$

Proof. We compute the right-hand side of (6.20). We have

$$\frac{q_{j,l}}{q_{j-1,l}}c_{j-1} + \frac{1}{q_{j-1,l}}c_l = \frac{q_{j,l}}{q_{j-1,l}}\sum_{t=1}^{j-2} q_{t,j-1}d_t + \frac{1}{q_{j-1,l}}\sum_{t=1}^{l-1} q_{t,l}d_t$$

$$= \sum_{t=1}^{j-2} \frac{q_{j,l}q_{t,j-1} + q_{t,l}}{q_{j-1,l}}d_t + d_{j-1} + \frac{1}{q_{j-1,l}}\sum_{t=j}^{l-1} q_{t,l}d_t$$

$$= \sum_{t=1}^{j-1} q_{t,j}d_t + \frac{1}{q_{j-1,l}}\sum_{t=j}^{l-1} q_{t,l}d_t$$
(6.17)
(6.17)

Using (6.17), (6.19) we see that the right-hand side of (6.20) equals c_j .

6.3. **Proof of Proposition 6.2.** Assume $b = (b_{-1}, b_0, b_{r+1}, b_1, \ldots, b_r)$ is obtained by triangle convex induction from (b_{-1}, b_0, b_{r+1}) . Then it is trivially true that for $0 \le l \le r, (b_{-1}, b_l, b_{r+1}, b_{l+1}, \ldots, b_r)$ is obtained by triangle convex induction from (b_{-1}, b_l, b_{r+1}) .

The following result shows that this is also true when we "truncate" b on the right. This is a combinatorial fact for which we have no conceptual explanation.

Proposition 6.10. For $1 \leq l \leq r+1$, $(b_{-1}, b_0, b_l, b_1, \ldots, b_{l-1})$ is obtained by triangle convex induction from (b_{-1}, b_0, b_l) .

Proof. In view of (6.12) we have to prove for $1 \le j < l \le r+1$

(6.21)
$$b_{j} = \left\lfloor \frac{q_{j-1,l} - q_{j,l} - 1}{q_{j-1,l}} b_{-1} + \frac{q_{j,l}}{q_{j-1,l}} b_{j-1} + \frac{1}{q_{j-1,l}} b_{l} \right\rfloor$$

To prove (6.21) we may add to b an affine function of the form $\langle m, -\rangle + k$ for $m \in M$ and $k \in \mathbb{Z}$ as such an altered b still satisfies the recursion (6.15). Using (6.7) it is therefore easy to see that we may assume $(b_{-1}, b_0, b_{r+1}) = (0, 0, d)$ for $0 \leq d < n$. Comparing (6.15) (and (6.13)) with (6.20) for l = r + 1 together with (6.18) we find that the normalized b satisfy $b_j = c_j$. Then (6.21) follows by invoking (6.20) for arbitrary l.

Remark 6.11. It is clear that in the proof of Proposition 6.10 we made essential use of the fact that in (6.15) we used $\lfloor - \rfloor$, rather than $\lfloor - \rceil$.

Proof of Proposition 6.2. It is enough to prove that $s_{v_{-1}}^c(m) = s_{v_{j-1}}^c(m) = s_{v_l}^c(m) \neq s_{v_j}^c(m), 1 \leq j < l$, cannot occur. This follows from Lemmas 5.3, 5.8 (using Proposition 6.10).

7. A TILTING BUNDLE ON A STACKY RESOLUTION USING GULOTTA'S APPROACH

From now on we assume that n = 3. We fix a triangulation of P without extra vertices and we let Σ be the corresponding fan. In this section we will construct a tilting bundle on X_{Σ} . This yields an NCCR for R_P by Proposition 3.3.

To start we embed P in a (minimal) rectangle $P_0 = [0, c] \times [0, d] \times \{1\}$. Then $P_0 - \operatorname{int} P$ is the union of at most four (non-convex) polygons $\{Q_i\}$, one polygon for every corner vertex in P_0 not occurring in P. Each of these polygons can be divided into triangles by successively cutting off "corner triangles" of maximal size whose internal edge has slope determined by the Farey tree, traversed row by row (see [Gul08, Figure 15]). E.g. if $Q \in \{Q_i\}$ is the polygon in the upper left or lower right corner then we use the slope sequence⁵



If $Q \in \{Q_i\}$ is the polygon in the upper right or lower left corner then we use the corresponding negative slopes. See Figure 7.1 for an example.



FIGURE 7.1. A corner triangles, / extra diagonal

⁵The slopes 0/1 and 1/0 which are sometimes considered part of the Farey tree are missing as they correspond to the vertical and horizontal boundary of Q.

Let P_i be the polygon that we obtain after removing *i* triangles from P_0 and let $P_l = P$. As explained above P_i is obtained by "sawing off" a corner from P_{i-1} . The thus created edge in P_i will be called the *cutting edge*. Remarkably, because of the clever slope choices during the cutting, all P_i are lattice polygons (cf. proof of [Gul08, Theorem 6.1]).

We add extra diagonals to the triangle $P_{i-1} - P_i$ connecting the unique vertex of P_{i-1} not in P_i to the points on the cutting edge in P_i which are vertices of P_j for $j \ge i$.⁶ Combining the resulting triangulation of $P_i - P$ with the one of P we obtain compatible triangulations of $P_0 \supset P_1 \supset \cdots \supset P_l = P$ where $P_{i-1} - P_i$ is a subdivided triangle. Let $V_0 \supset V_1 \supset \cdots \supset V_l$ be the corresponding sets of vertices. Note that $|V_{i-1} - V_i| = 1$ and moreover V_i is the union of the vertices of P_j for $j \ge i$.

Let $n_1 = (0, 0, 1)$, $n_2 = (c, 0, 1)$, $n_3 = (c, d, 1)$, $n_4 = (0, d, 1)$ be the vertices of P_0 and let V'_i be the union of the vertices of P_j for $0 \le j \le i$. For each i, $V'_i - V'_{i-1}$ consists of endpoints of the cutting edge in P_i . More precisely $V'_i - V'_{i-1}$ consists of 0, 1 or 2 elements of ∂P_{i-1} (depending on how many of the endpoints of the cutting edge are already vertices of P_{i-1}). If v is one of those elements then

(7.1)
$$v = (1-t)v' + tv'$$

for $t \in [0, 1[$ and v', v'' vertices of P_{i-1} . We use this to order $V_0 = V'_l = [n_1, \ldots, n_{4+l'}]$ in such a way that it is convexly induced from $[n_1, n_2, n_3, n_4]$ in the sense of Definition 5.5, with the induction datum $\mathbf{t} = (t_j^{(i)})_{i,j}$ being obtained from the expressions (7.1).

We now fix $b_0 \in \mathbb{Z}^4$. We also fix a sign sequence and we let $(b_v)_{v \in V_0} := b$ be convexly induced from b_0 with respect to this sign sequence (and the induction datum t). In down to earth terms this means that whenever v, v', v'' are as in (7.1) then $b_v = \lfloor (1-t)b_{v'} + tb_{v''} \rfloor$ or $b_v = \lceil (1-t)b_{v'} + tb_{v''} \rceil$ depending on the chosen sign sequence.

Let Σ_i be the fan on the cone over P_i corresponding to its triangulation. The elements of V_i are the minimal generators of the one-dimensional cones in Σ_i and we let $\Sigma_i = (\Sigma_i, V_i)$ be the corresponding stacky fan as in §3. Let $\tilde{b}_i = (b_v)_{v \in V_i}$.

Lemma 7.1. We have

(7.2)
$$\Gamma(X_{\Sigma_0}, \mathcal{M}_{\Sigma_0, \tilde{b}}) = \mathsf{M}_{b_0}.$$

If $b_0 \in \mathbb{Z}^4$ moreover satisfies

(7.3)
$$\forall m \in M : [s_1^{b_0}(m), s_2^{b_0}(m), s_3^{b_0}(m), s_4^{b_0}(m)] \notin \{[+-+-], [-++-]\}$$

(cfr. (5.2) for notation) then $H^j(X_{\Sigma_i}, \mathcal{M}_{\Sigma_i, \tilde{b}_i}) = 0$ for j > 0 and $i = 0, \ldots, l$.

Proof. We start with the first claim. By Lemma 3.1 we have

$$\Gamma(X_{\Sigma_0}, \mathcal{M}_{\Sigma_0, \tilde{b}}) = \mathsf{M}_{\tilde{b}}.$$

Since \tilde{b} is convexly induced from b_0 , the conclusion follows from Corollary 5.7.

Now we discuss the last claim. As $\pi_s: X_{\Sigma_i} \to X_{\Sigma_i}$ is finite we have by Lemma 3.1

$$H^{j}(X_{\Sigma_{i}}, \mathcal{M}_{\Sigma_{i}, \tilde{b}_{i}}) = H^{j}(X_{\Sigma_{i}}, \mathcal{M}_{\Sigma_{i}, \tilde{b}_{i}}) = H^{j}(X_{\Sigma_{i}}, \mathcal{O}(D_{\tilde{b}_{i}})).$$

⁶The extra diagonals are necessary to obtain a triangulation of P_0 in the sense of algebraic topology, i.e. a homeomorphism $|K| \cong P_0$ where K is a combinatorial simplicial complex.

We will prove that $V_{D_{\tilde{b}_i},m}$ (see (3.6), (3.7)) is either empty or contractible.⁷ This implies what we want.

Step 1. Let $u_0, u_1, \ldots, u_{r+1} \in V_i$ be the vertices on an edge in P_i (ordered in an arbitrary direction). We claim that the signs $s_{u_i}^b(m)$ follow the pattern $+ \cdots + - \cdots -$ (possibly reflected and perhaps with no + or - present). This follows from the easy observation that $(\tilde{b}_{u_0}, \tilde{b}_{u_{r+1}}, \tilde{b}_{u_1}, \ldots, \tilde{b}_{u_r})$ is obtained by interval convex induction from $(\tilde{b}_{u_0}, \tilde{b}_{u_{r+1}})$ together with Lemma 5.10.

Step 2. Let w_1, w_2, \ldots be the elements of $\partial P_i \cap V_i$ (ordered in an arbitrary direction). We claim that the signs $s_{w_i}^b(m)$ follow the pattern $+ \cdots + - \cdots -$ (up to cyclic permutation and possibly with no + or - present).

By Step 1 it is sufficient to prove this for w_1, w_2, \ldots being the vertices of P_i . We note that it is true for P_0 by (7.3). An easy case by case analysis using Lemma 5.3 for removing a single triangle then proves the claim by induction.

Step 3. $V_{D_{\tilde{b}_l},m}$ is either empty or contractible. This follows from Step 2 applied with i = l. We leave the verification as an exercise to the reader.

Step 4. If the sign pattern in Step 2 is $+\cdots +$ then $V_{D_{\tilde{b}_i},m} = \emptyset$. Indeed, it follows from the inductive construction and repeated application of Lemma 5.3 that $s_v^b(m) = +$ for all $v \in V_i$ in this case.

Step 5. Again by a case by case analysis of the same nature as in Step 2 one shows that the following possibilities can occur for the relation between $V_{D_{\tilde{b}_i},m}$ and $V_{D_{\tilde{b}_{i-1},m}}$.

(1) They are the same.

(2) $V_{D_{\tilde{b}_i},m} = \emptyset$ and $V_{D_{\tilde{b}_{i-1},m}}$ is a single point (this needs in particular Step 4).

(3) $V_{D_{\tilde{b}_{i-1},m}}$ is obtained from $V_{D_{\tilde{b}_i},m}$ by gluing an interval at an end point.

(4) $V_{D_{\tilde{b}_{i-1},m}}$ is obtained from $V_{D_{\tilde{b}_i},m}$ by gluing a triangle at one of its edges.

In each of the cases the property of being empty or contractible is preserved, finishing the proof. $\hfill \Box$

We set

$$\begin{split} S_0 &= \{(0,i,i+j,j) \mid 0 \leq i < \mathtt{c}, 0 \leq j < \mathtt{d}\} \cup \{(0,i,i+j+1,j) \mid 0 \leq i < \mathtt{c}, 0 \leq j < \mathtt{d}\}.\\ Remark 7.2. \text{ Note that } X_{P_0} &= Y/\!\!/ G \text{ where } Y = (k^{\oplus 4})^{\vee} \text{ and } G = k^* \times \mathbb{Z}_{\mathtt{c}} \times \mathbb{Z}_{\mathtt{d}} \text{ acts} \\ \text{on } k^{\oplus 4} \text{ with weights } \chi_1 \xi_{\mathtt{c}}^{-1} \xi_{\mathtt{d}}^{-1}, \chi_{-1} \xi_{\mathtt{c}}, \chi_1, \chi_{-1} \xi_{\mathtt{d}} \text{ (see §3). Furthermore this action} \\ \text{is generic in the sense of [ŠVdB17a, Def.§1.3.4]. Since } S_0 \text{ can be naturally identified} \\ \text{with } \{0,1\} \times \mathbb{Z}_{\mathtt{c}} \times \mathbb{Z}_{\mathtt{d}} = ((1/2)[-2,2]_{\epsilon} \cap X(k^*)) \times X(\mathbb{Z}_{\mathtt{c}} \times \mathbb{Z}_{\mathtt{d}}) \text{ for } \epsilon > 0, \text{ it follows} \\ \text{by [ŠVdB17a, Theorem 1.6.2] combined with [ŠVdB17a, Lemma 4.5.1] (see also [VdB04, §7]) that } \Lambda_0 := \operatorname{End}_{R_{P_0}}(\bigoplus_{b \in S_0} \mathsf{M}_b) \text{ is a (well-known) NCCR of } R_{P_0}. \end{split}$$

For each vertex in V_0 we choose a sign and we compatibly convexly induce the elements $S_0 = \{(b_v)_{v \in V'_0}\}$ using those signs to obtain a set $S = S_l = \{(b_v)_{v \in V_0}\}$ (S_l and S_0 are in bijection).

Let $\overline{S}_l^i \subset \prod_{v \in V_i} \mathbb{Z}$ be the image of S_l under the projection $\prod_{v \in V_0} \mathbb{Z} \to \prod_{v \in V_i} \mathbb{Z}$. Put $\mathcal{T}_i = \bigoplus_{b \in \overline{S}_l^i} \mathcal{M}_{\Sigma_i, b}$ (see (3.3) for notation). This is a vector bundle on the toric DM-stack X_{Σ_i} .

⁷Note that the distinction is relevant as contractible spaces have no reduced cohomology but $\tilde{H}^{-1}(\emptyset, k) = k$.

Corollary 7.3. \mathcal{T}_i has no higher self-extensions.

Proof. It is easy to verify that b - b' for $b, b' \in S_0$ satisfy (7.3). As S_l is compatibly convexly induced from S_0 , Lemma 5.8 enables us to apply Lemma 7.1.

Lemma 7.4. \mathcal{T}_0 is a classical generator (see [BVdB03]) of $\mathcal{D}(X_{\Sigma_0})$.

Proof. By (7.2), Lemma 5.8 and Remark 7.2 (combined with (3.2)) we obtain $\tilde{\Lambda}_0 := \operatorname{End}_{X_{\Sigma_0}}(\mathcal{T}_0) \cong \Lambda_0$.

Since Λ_0 is an NCCR for R_{P_0} by Remark 7.2, we have in particular gl dim $\Lambda_0 < \infty$, and hence the same is true for $\tilde{\Lambda}_0$. By Corollary 7.3 we have a full and faithful embedding of Perf $(\tilde{\Lambda}_0)$ in $\mathcal{D}(X_{\Sigma_0})$. As gl dim $\tilde{\Lambda}_0 < \infty$, we have Perf $(\tilde{\Lambda}_0) = \mathcal{D}(\tilde{\Lambda}_0)$ and moreover $\mathcal{D}(\tilde{\Lambda}_0)$ is admissible in $\mathcal{D}(X_{\Sigma_0})$. Since $\mathcal{D}(X_{\Sigma_0})$ does not have a nontrivial semi-orthogonal decomposition by Corollary A.5, we have $\mathcal{D}(\tilde{\Lambda}_0) \cong \mathcal{D}(X_{\Sigma_0})$.

Theorem 7.5. \mathcal{T}_i is a tilting bundle on X_{Σ_i} .

Proof. By Corollary 7.3 it suffices to show that \mathcal{T}_i is a classical generator of $\mathcal{D}(X_{\Sigma_i})$. Since X_{Σ_i} is isomorphic to an open substack of X_{Σ_0} and \mathcal{T}_i is a restriction of \mathcal{T}_0 to X_{Σ_i} this follows by Lemma 7.4.

8. TILTING BUNDLE ON A STACKY RESOLUTION USING THE ISHII-UEDA APPROACH

We start in exactly the same way as in §7. In this section we will construct a tilting bundle on X_{Σ} using a procedure inspired by [IU15]. This will yield again an NCCR of R_P using Proposition 3.3.

Now we embed P in a triangle P_0 with vertices $(0, 0, 1), (0, \mathbf{c}, 1), (\mathbf{d}, 0, 1)$. To obtain P_i , i > 0, from P_{i-1} one removes a (corner) vertex of P_{i-1} and takes the convex hull of the remaining lattice points in P_{i-1} . Let l be such that $P_l = P$. We triangulate P_0 as in §7 by first triangulating P (without extra vertices) and then triangulating $P_{i-1} - P_i$ by adding edges from the unique vertex in P_{i-1} which is not on P_i to the vertices of P_i , $j \geq i$ which are not on ∂P_{i-1} . See Figure 8.1.



FIGURE 8.1. • corner vertices

The objects Σ_i , Σ_i , V_i , V'_i are defined in the same way as in §7. We set

$$S_0 = \{ (i, j, 1) \mid 0 \le i < c, 0 \le j < d \}.$$

Note that $X_{P_0} \cong Y/\!\!/G$, where $Y = k^3$, $G = \mathbb{Z}_{\mathsf{c}} \times \mathbb{Z}_{\mathsf{d}}$ acts on k^3 with weights $\xi_{\mathsf{c}}^{-1}\xi_{\mathsf{d}}^{-1}, \xi_{\mathsf{c}}, \xi_{\mathsf{d}}$ (see §3). This action is generic. Since G is a finite abelian group and $S_0 \cong X(G)$, it is classical (see e.g. [Aus86], [Leu12, §J], [ŠVdB17a, §1.2], [Wem14, §1.4]) that $\Lambda_0 = \operatorname{End}_{R_{P_0}}(\bigoplus_{b \in S_0} \mathsf{M}_b)$ is an NCCR of R_{P_0} .

Below we will inductively extend $S_{i-1} = \{(b_i)_{i \in V'_{i-1}}\}$ to $S_i = \{(b_i)_{i \in V'_i}\}$ using compatible convex induction so that the required tilting bundle will be obtained from $S_l = \{(b_i)_{i \in V_0}\}$ (note $V'_l = V_0$). However here the induction process appears to be more delicate and in particular our proof depends on the choice of a specific sign sequence (see §5) in order to obtain an analogue for Lemma 7.1. See Remark 6.11.

8.1. Definition of S_i . Let $v_{-1} \in V'_{i-1}$ be the vertex of P_{i-1} that we remove to construct P_i , and let $v_0, v_{r+1} \in \partial P_{i-1} \cap \partial P_i \cap \partial (P_{i-1} \setminus P_i)$. Let v_1, \ldots, v_r be the other lattice points on $\partial (P_{i-1} \setminus P_i)$ such that $v_0, v_1, \ldots, v_r, v_{r+1}$ are consecutive.

Remark 8.1. Note that $V'_{i-1} \cup \{v_0, v_1, \ldots, v_{r+1}\}$ may be strictly larger than V'_i since some of the v_j may not be vertices of P_i (they will of course be elements of ∂P_i). For a simple example how this can happen look at [CLS11, Figure 10.3].

Let $b \in S_{i-1}$. We describe how to induce b to V'_i . This will be done for all $b \in S_{i-1}$ in a compatible way.

- (i) We first convexly induce b to $V'_{i-1} \cup \{v_0, v_{r+1}\}$ for a fixed sign sequence. We keep denoting the result by b.
- (ii) We induce up b to $V'_{i-1} \cup \{v_0, \ldots, v_{r+1}\}$ by triangle convex induction (see §6), and set $b^{(i)} = (b_v)_{v \in V'}$. We let S_i be the collection of $b^{(i)}$ so obtained.

We also adopt the notation \overline{S}_l^i from §7. We can now state the following analogue of Theorem 7.5.

Theorem 8.2. Put $\mathcal{T}_i^{\triangle} := \bigoplus_{b \in \overline{S}_i^i} \mathcal{M}_{\Sigma_i, b}$. Then $\mathcal{T}_i^{\triangle}$ is a tilting bundle on X_{Σ_i} .

This is proved using Proposition 8.3 below, which allows us to suitably adapt the proof of Corollary 7.3, in exactly the same way as Theorem 7.5.

Proposition 8.3. Let $\tilde{b}_i = b - b'$ for $b, b' \in \overline{S}_l^i$. Then $H^j(X_{\Sigma_i}, \mathcal{O}(D_{\tilde{b}_i})) = 0$ for j = 1, 2.

Proof. With the aid of Proposition 6.2 the proof follows similar lines as the proof of Lemma 7.1, therefore we only point out the necessary adjustment. Note that Step 1 is relevant because of (i). In Step 2 we need to additionally use Proposition 6.2, in Step 5 (4) a polygon is glued at a (contractible) string of edges. \Box

9. Example

As a sanity check we have verified our procedures for constructing toric NCCR's for 3-dimensional toric singularities on an explicit moderately complex example. Let P be as in Figure 7.1. We have $G = G_m^2$ and $R_P \cong (\text{Sym } W)^G$, where G acts on W with weights (-1,0), (3,-1), (-2,3), (-2,-3), (2,1) (see Figure 9.1). The Cohen-Macaulay modules of covariants can be deduced by [VdB93] or by [Boc12] (see Proposition B.1 below).

If we follow Gulotta's (resp. the Ishii-Ueda) procedure as shown in Figure 7.1 (resp. 8.1) for a sign sequence $\mathfrak{s} = (-)_i$ we get a module of covariants M corresponding to the grey weights in the left (resp. right) Figure 9.1. We observe the following comforting facts.

- Both sets of grey weights do indeed contain the same number of elements (16).
- The endomorphism rings of the corresponding modules of covariants are Cohen-Macaulay.
- The number 16 is also equal to 2 Vol(P), and so me may use Proposition B.3 below to double check that the endomorphism rings are indeed NCCRs.



FIGURE 9.1. • G-weights, • CM weights, \bigcirc Gulotta (left), Ishii-Ueda (right)

APPENDIX A. TORIC DM STACKS AND SEMI-ORTHOGONAL DECOMPOSITIONS

As usual we assume $M = N = \mathbb{Z}^n$. In this appendix we show that the derived category of a smooth toric DM stack X_{Σ} (see [BH06]) for $\Sigma = (\Sigma, (n_i)_{i=1}^k)$, where $|\Sigma|$ is a (rational strongly convex) polyhedral cone over a convex lattice polyhedron $P \times \{1\} \subset \mathbb{R}^n$ and $n_i \in P \times \{1\}$ are generators of 1-dimensional cones in Σ , does not admit any nontrivial semi-orthogonal decomposition.

We first recall a result from [BCS05]. Here Σ_{max} denotes the set of maximal cones in Σ .

Proposition A.1. [BCS05, Proposition 4.3 and its proof] Let σ be a maximal cone in Σ . If $\sigma = [n_{i_1}, \ldots, n_{i_n}]$, then X_{σ} , $\sigma = (\sigma, (n_{i_j})_{j=1}^n)$, corresponds to an open substack of X_{Σ} and $X_{\Sigma} = \bigcup_{\sigma \in \Sigma_{\max}} X_{\sigma}$. Moreover $X_{\sigma} \cong k^n/G_{\sigma}$ for a finite subgroup $G_{\sigma} \subset G$ (with G is as in §3) such that the action of G_{σ} on k^n is unimodular and faithful.

Recall that we denote $X_P = \operatorname{Spec}(R_{\sigma_P})$.

Lemma A.2. Let $\pi_s: X_{\Sigma} \to X_{\Sigma}$ be the canonical map. Then

$$\mathcal{O}_{X_{\Sigma}} \cong \omega_{X_{\Sigma}} \cong \pi_s^* \omega_{X_{\Sigma}} \cong \pi_s^* \mathcal{O}_{X_{\Sigma}}.$$

Proof. Let $\sigma \in \Sigma_{\text{max}}$. By Proposition A.1 the smooth points in X_{σ} correspond exactly to G_{σ} -orbits of points in k^n with trivial stabilizer. Indeed, since G_{σ} is unimodular the stabilizer groups cannot contain pseudo-reflections, and therefore the corresponding quotient (of the G_{σ} -orbit of a point in k^n with nontrivial stabilizer) cannot be regular by the Chevalley-Shephard-Todd theorem.

Thus, as both $\omega_{X_{\Sigma}}$ and $\omega_{X_{\Sigma}}$ are reflexive and $\pi_s : X_{\Sigma} \to X_{\Sigma}$ is the identity on the smooth locus of X_{Σ} (whose complement has codimension ≥ 2 as X_{Σ} is a normal variety) we have $\pi_s^* \omega_{X_{\Sigma}} = \omega_{X_{\Sigma}}$. Moreover, $\tau : X_{\Sigma} \to X_P$ is crepant; i.e., $\tau^* \omega_{X_P} = \omega_{X_{\Sigma}}$ (see e.g. [CLS11, Example 11.2.6]) and as X_P is Gorenstein it follows $\omega_{X_{\Sigma}} \cong \mathcal{O}_{X_{\Sigma}}$.

Lemma A.3. The relative Serre functor⁸ of $\mathcal{D}(X_{\Sigma})$ with respect to $\mathcal{D}(X_P)$ is the identity; i.e. for all $\mathcal{F}, \mathcal{G} \in \mathcal{D}(X_{\Sigma})$ we have

(A.1)
$$\operatorname{RHom}_{X_{\Sigma}}(\mathcal{F}, \mathcal{G}) \cong \operatorname{RHom}_{R_P}(\operatorname{RHom}_{X_{\Sigma}}(\mathcal{G}, \mathcal{F}), R_P).$$

Proof. Since $\pi_s^* \omega_{X_{\Sigma}} = \omega_{X_{\Sigma}}$ by Lemma A.2, it follows from [Abu16, Theorem 3.0.17] and its proof that the relative Serre functor of $\mathcal{D}(X_{\Sigma})$ with respect to $\mathcal{D}(X_{\Sigma})$ is the identity. It is easy to check that it is thus enough to show that the identity is a relative Serre functor of $\mathcal{D}(X_{\Sigma})$ with respect to $\mathcal{D}(X_P)$. Since $\theta : X_{\Sigma} \to X_P$ is crepant (see proof of Lemma A.2) and X_{Σ} , X_P are Gorenstein, this follows by the same proof as in [IW14b, Lemma 4.13].

Lemma A.4. $\mathcal{D}(X_{\Sigma})$ is indecomposable.

Proof. Let σ be a maximal cone in Σ . Then $X_{\sigma} \cong k^n/G_{\sigma}$ for a finite group G_{σ} , which acts faithfully on k^n by Proposition A.1. Thus, $\mathcal{D}(X_{\sigma})$ is indecomposable by [BKR01, Lemma 4.2]. Assume that we have an orthogonal decomposition $\mathcal{D} = \mathcal{D}(X_{\Sigma}) = \langle \mathcal{D}_1, \mathcal{D}_2 \rangle$, $\operatorname{Hom}(\mathcal{D}_1, \mathcal{D}_2) = \operatorname{Hom}(\mathcal{D}_2, \mathcal{D}_1) = 0$. Since $\mathcal{D}(X_{\sigma})$ is a Verdier quotient of $\mathcal{D}(X_{\Sigma})$ (as X_{Σ} is a noetherian stack by [LMB99, Prop. 15.4]) and if $C = C_1 \oplus C_2$, $C_i \in \mathcal{D}_i$, has support in $X_{\Sigma} \setminus U_{\Sigma}$, the same is true for C_1, C_2 , the orthogonal decomposition is preserved when passing to an open substack. For $\sigma \in \Sigma_{\max}$ we thus have $j_{\sigma}^* \mathcal{D}_1 = 0$ or $j_{\sigma}^* \mathcal{D}_2 = 0$ (where $j_{\sigma} : X_{\sigma} \hookrightarrow X_{\Sigma}$). Choose a point x in X_{Σ} which is in the intersection of the X_{σ} and also in the locus where π_s is an isomorphism. Then we can view \mathcal{O}_x as an object in \mathcal{D} . Since \mathcal{O}_x is indecomposable we can assume that $\mathcal{O}_x \in \mathcal{D}_1$. Since the restriction of \mathcal{O}_x to X_{σ} is nonzero for every $\sigma \in \Sigma_{\max}$, we get that $j_{\sigma}^* \mathcal{D}_2 = 0$ for every $\sigma \in \Sigma_{\max}$, which easily implies that $\mathcal{D}_2 = 0$.

Corollary A.5. $\mathcal{D}(X_{\Sigma})$ does not admit any nontrivial semi-orthogonal decompositions.

Proof. It follows by (A.1) that $\operatorname{RHom}_{X_{\Sigma}}(\mathcal{G}, \mathcal{F}) = 0$ implies $\operatorname{RHom}_{X_{\Sigma}}(\mathcal{F}, \mathcal{G}) = 0$. Thus, if $\mathcal{D}(X_{\Sigma})$ has a non-trivial semi-orthogonal decomposition then it is decomposable. This contradicts Lemma A.4.

APPENDIX B. COHEN-MACAULAY MODULES

In this section we show how one can extend a useful criterion of Bocklandt [Boc12] for recognising Cohen-Macaulay modules of covariants in dimension 3. It can be used to give a direct proof of the Cohen-Macaulayness of the endomorphism algebra $\Lambda = \operatorname{End}_{R_P}(\bigoplus_{b \in \overline{S}_i^t} \mathsf{M}_b)$ in §7. Moreover, we will explain that Theorem 1.1

could be deduced from this fact provided we knew that $|\overline{S}_l^i| = 2 \operatorname{Vol}(P)$.

Recall the following result.⁹

⁸See [Kuz08].

⁹The result may also be deduced from (3.6) by calculating the cohomology of M_b restricted to the punctured spectrum of Spec R_P . The latter is the toric variety associated to the fan spanned by ∂P .

Proposition B.1. [Boc12, Lemma 4.5] Let $\sigma = [n_1, \ldots, n_k]$, with n_1, \ldots, n_k oriented counter-clockwise, be a minimal presentation of σ , and let $b \in \mathbb{Z}^k$. The R_P -module M_b is Cohen-Macaulay if and only for every $m \in M$ the signs $s_i^b(m)$ follow the pattern $+ \cdots + - \cdots -$ (up to cyclic permutation and possibly with no + or - present).

Let $P \subset \mathbb{R}^2 \times 1$ be a lattice polygon with vertices n_1, \ldots, n_k and let P' denote a lattice polygon obtained from P by removing a triangle with vertex n_k . We denote the other vertices of the triangle by n_{k+1}, n_{k+2} . They lie on the boundary of P (and may, or may not, be vertices). Then $[n_1, \ldots, n_k, n_{k+1}, n_{k+2}]$ is convexly induced from $[n_1, \ldots, n_k]$ with an induction datum $\mathfrak{t} = (t^{(k+1)}, t^{(k+2)})$, where $t_j^{(k+1)} = 0$ for $j \neq k - 1, k, t_j^{(k+2)} = 0$ for $j \neq 1, k$. For $b \in \mathbb{Z}^k$ and $\mathfrak{s} \in \{+, -\}^2$ we let $b_{\mathfrak{s}}$ be convexly induced from b with the sign sequence \mathfrak{s} (and the induction datum \mathfrak{t}). By $\overline{b_{\mathfrak{s}}}$ we denote the projection of $b_{\mathfrak{s}}$ to \mathbb{Z}^{k+1} omitting the k-th component.

Using the above criterion one can use a simplified version of Step 2 in the proof of Lemma 7.1 (with $\{w_1, w_2, \ldots\} = \{n_1, \ldots, n_{k+2}\}$) to obtain Corollary B.2 below. This is a generalization of [Boc12, Corollary 4.7] relaxing the assumption that $\{n_{k+1}, n_{k+2}\} \subset \{n_1, \ldots, n_{k-1}\}$ (in the latter case $\overline{b_s}$ is just the projection of b and in particular the sign sequence \mathfrak{s} is irrelevant).

Corollary B.2. If M_b is a Cohen-Macaulay R_P -module then $M_{\overline{b_s}}$ is a Cohen-Macaulay $R_{P'}$ -module. Moreover, if $\operatorname{End}_{R_P}(\bigoplus_{b\in S} M_b)$ is a Cohen-Macaulay R_P -module then the $R_{P'}$ -module $\operatorname{End}_{R_{P'}}(\bigoplus_{b\in S} M_{\overline{b_s}})$ is also Cohen-Macaulay.

If we could prove that using the Gulotta "cutting" procedure combined with Corollary B.2 one gets a Cohen Macaulay R_P -module M such that $\operatorname{End}_{R_P}(M)$ is Cohen-Macaulay and such that M has precisely $2 \operatorname{Vol}(P)$ non-isomorphic indecomposable summands then one obtains by the following proposition (see also [Boc12, proof of Theorem 7.1]) a proof of Theorem 1.1 which does not require the construction of a tilting bundle on a resolution. Unfortunately we have not been able to solve this purely combinatorial problem.

Proposition B.3. Let M be a graded R_P -module (for some connected grading on R_P) such that M contains at least $2 \operatorname{Vol}(P)$ non-isomorphic graded indecomposable summands and let $\Lambda = \operatorname{End}_{R_P}(M)$. If Λ is a Cohen-Macaulay R_P -module, then Λ is an NCCR of R_P .

Proof. Since R_P has an NCCR A by [ŠVdB17b, Proposition 1.1, Corollary 3.2], it has a CCR which is derived equivalent to A by [VdB04, Theorem 6.3.1]. By [Bri02], all crepant resolutions of R_P are derived equivalent. There is a crepant resolution of R_P given by X_{Σ} corresponding to a triangulation of P with triangles of area 1/2. By [BH06] (see [ŠVdB17b, Theorem A.1]), rank $K(X_{\Sigma}) = 2 \operatorname{Vol}(P)$, thus rank $K_0(A) = 2 \operatorname{Vol}(P)$.

Since M is a modifying module, it is a direct summand of a graded¹⁰ "maximal modifying module" $M \oplus M'$ by [IW14a, Proposition 1.19] and we may assume that M and M' have no common indecomposable summands. Since R_P has an NCCR, the "maximal modification algebra" $\Lambda' := \operatorname{End}_{R_P}(M \oplus M')$ is also an NCCR by [IW14a, Proposition 1.13]. As all NCCRs of R_P are derived equivalent by [IW13, Theorem 1.5], rank $K_0(\Lambda') = \operatorname{rank} K_0(A) = 2\operatorname{Vol}(P)$. Let m, m' be the number

¹⁰The results of [IW14a] are valid in the graded context.

of non-isomorphic indecomposable summands of M, M'. By Lemma B.4 below applied to Λ' we now have $2 \operatorname{Vol}(P) = \operatorname{rank} K_0(\Lambda') \ge m + m' \ge m \ge 2 \operatorname{Vol}(P)$ (the last inequality by the hypotheses) and hence m' = 0. Therefore M' = 0 and $\Lambda' = \Lambda$. Hence Λ is an NCCR of R_P .

We have used the following lemma.

Lemma B.4. Let R be a commutative finitely generated connected graded ring and let M be be a finitely generated graded R-module with m non-isomorphic indecomposable summands. Put $\Lambda = \operatorname{End}_R(M)$. Then rank $K_0(\Lambda) \ge m$.

Proof. Let rad Λ be the graded radical of Λ . Then the morphism between Grothendieck groups $K_0(\Lambda) \to K_0(\Lambda/\operatorname{rad} \Lambda)$ is surjective (by lifting of idempotents) and moreover rank $K_0(\Lambda/\operatorname{rad} \Lambda) = m$, finishing the proof.

Appendix C. Tilting bundles vs NCCRs

In dimension 3 (split) tilting bundles on crepant resolutions and (toric) NC-CRs are intimately connected. The following theorem summarizes the relation. While the first part (see [VdB04]) of the theorem is standard, the converse (see [IU16]) is quite involved and requires a deep study of the delicate interplay between crepant resolutions, dimer models, and moduli spaces of representations of the corresponding quivers, and moreover NCCRs (see [Boc12]). Since such moduli spaces are projective over affine, the converse in addition requires that the resolution is projective.

Theorem C.1. [VdB04], [IU16, Corollary 1.2], [Boc12, Theorem 7.1.-7.3] Let Rbe the coordinate ring of a 3-dimensional Gorenstein affine toric variety and let $\pi : X \to \text{Spec } R$ be a crepant resolution. If \mathcal{T} is a tilting bundle on X then $\text{End}_X(\mathcal{T})$ is an NCCR of R. Moreover, if π is projective and Λ is a toric NCCR of R, then there exists a split tilting bundle \mathcal{T} on X such that $\Lambda \cong \text{End}_X(\mathcal{T})$.

Currently the full generality of this result seems out of reach of the methods used in this paper. This being said, the convex induction approach makes it often possible to find the tilting bundles whose existence is asserted in Theorem C.1 in a rather direct way. Interestingly, as we show below, convex induction is sometimes able to provide tilting bundles even if the resolution is not projective. This situation is not covered by Theorem C.1.

We list some examples where the convex induction method yields split tilting bundles on a crepant resolution of an affine toric variety for any toric NCCR. Note that any such crepant resolution is toric and is obtained from the fan associated to a lattice triangulation with triangles of area 1/2 of the polygon P corresponding to R (see [DHZ01, Proposition 2.4], such a triangulation is called "unimodular").

- (1) One way to triangulate P is by using the Ishii-Ueda method (successively removing corner vertices and taking the convex hull of the remaining lattice points). For this type of triangulation we can induce the b's defined on the vertices of P, corresponding to the toric NCCR, to the vertices of the triangulation using triangle convex induction as in §8, and define the corresponding split tilting bundle.
- (2) Convex induction may still work even if the triangulation is not as in (1). For example, the star triangulation of a (minimal) regular lattice hexagon (see Figure C.1) is not of the above form, but one may use convex induction

to find a split tilting bundle. Let $\Lambda = \operatorname{End}_R(\bigoplus_{b \in S} M_b)$ be a toric NCCR. Let n_1, \ldots, n_6 be vertices of the polygon and let n_7 be the middle point. We fix some induction datum $(t_j^{(7)})_j$ (e.g. $t_j^{(7)} = 1/6$ for $j = 1, \ldots, 6$) and let \tilde{S} be compatibly convexly induced from S. We claim that $\bigoplus_{\tilde{b} \in \tilde{S}} \mathcal{O}(D_{\tilde{b}})$ is a tilting bundle on X. We need to show (see (3.1), (3.7)) that $\tilde{H}^{\geq 0}(V_{D_{\tilde{b}-\tilde{b}'},m},k) = 0$ for $\tilde{b}, \tilde{b}' \in \tilde{S}$ and $m \in M$. It follows from the fact that Λ is Cohen-Macaulay (see Proposition B.1) that any sign sequence on the boundary of the polygon will be of the form $+, \ldots, +, -, \ldots, -$ (with possibly no + or - appearing), therefore $V_{D_{\tilde{b}-\tilde{b}'},m}$ would not be empty or contractible only if the sign pattern on the boundary is $-, \ldots, -$ and there is + in the middle. However, this is impossible by the convex induction (using Lemma 5.8).



FIGURE C.1. A crepant resolution of a hexagon.

The combinatorial argument used in the last example can be used in more general situations via the following lemma.

Lemma C.2. Let $(n_i)_{i \in I}$ be the vertices of a unimodular triangulation of P and assume that the $(n_i)_i$ are obtained by convex induction from the vertices of P for a chosen induction datum. Assume that every set of signs $(s_i)_{i \in I} \in \{\pm\}^I$ compatible with Lemma 5.3 whose the restriction to the vertices of P is of the form $+, \ldots, +, -, \ldots, -$ (possibly reflected and with at least one + and one -), has the property that every n_i is connected with a constant sign path to ∂P . Then R satisfies the converse of Theorem C.1 for the (not necessarily projective) crepant resolution of Spec R given by the triangulation.

Proof. Let $(n_j)_{j\in J}$ be the vertices of P and let let $S = \{(b_j)_{j\in J}\}$ be a collection of b's defining an NCCR of R and let $\tilde{S} = \{(\tilde{b}_i)_{i\in I}\}$ be obtained from S by compatible convex induction for an arbitrary sign sequence and the given induction datum. Let $\tilde{b} = \tilde{b}' - \tilde{b}''$ for some $\tilde{b}', \tilde{b}'' \in \tilde{S}$. Then \tilde{b} is obtained by convex induction from b := b' - b'' by Lemma 5.8.

For $m \in M$ put $s_i = s_i^b(m)$. Then $(s_i)_{i \in I}$ is compatible with Lemma 5.3. We will show that $V_{D_{\bar{i}},m}$ is either empty or contractible.

Since S defines an NCCR the restriction of $(s_i)_{i \in I}$ to the vertices of P will be of the form $+, \ldots, +, -, \ldots, -$ (with possibly no + or - appearing) (see Proposition B.1). By Lemma 5.3 the same will be true for the restriction to the vertices of the triangulation in ∂P .

If the sequence restricted to ∂P is $+, \ldots, +$ then by Lemma 5.3: $V_{\tilde{b},m} = \emptyset$. If the sequence is $-, \ldots, -$ then $V_{D_{\tilde{b}},m} = P$.

Assume now that the restricted sequence is $+, \ldots, +, -, \ldots, -$. Since every n_i has a constant sign path to the boundary the set of -'s is connected and moreover no cycle of -'s can surround a +. It now suffices to apply Lemma C.3 below. \Box

Lemma C.3. Assume we are given a triangulation of P with vertices $(n_i)_{i \in I}$. Let $b \in \mathbb{Z}^I$, $m \in M$ and assume in addition that $(s_i^b(m))_i$ is such that the set of -'s is connected and furthermore that no cycle of -'s surrounds a +. Then $V_{D_b,m}$ is either empty or contractible.

Proof. Put $V = V_{D_b,m}$. Assume $V \neq \emptyset$. Then the hypotheses imply $|\pi_i V| = 1$ for i = 0, 1. Furthermore as V is planar, Zastrow's theorem [Zas98, Theorem A] asserts that $|\pi_i V| = 1$ for $i \geq 2$. Since V is a CW-complex this implies that V is contractible by Whitehead's theorem.

(3) Projective crepant resolutions correspond to regular triangulation of P (see e.g. [CLS11, Definition 15.2.8]). However, there are also non-regular triangulations for which the convex induction method works. As an example (see Figure C.2) we take the triangulation [Log08, §5.2] refining the iconic (minimal) example of a non-regular triangulation [San06, Example 1.4].



FIGURE C.2. A non-regular triangulation

We induce $[n_1, n_2, n_3, n_4, \ldots, n_{10}]$ from $[n_1, n_2, n_3]$ with the induction datum $(t_j^{(i)})_{ij}$ satisfying $t_1^{(4)} = t_2^{(4)} = t_2^{(5)} = t_3^{(5)} = t_1^{(6)} = t_3^{(6)} = 1/2$, $t_1^{(7)} = t_2^{(7)} = t_3^{(7)} = 1/3$, $t_1^{(8)} = t_7^{(8)} = t_2^{(9)} = t_7^{(9)} = t_3^{(10)} = t_7^{(10)} = 1/2$, and $t_j^{(i)} = 0$ otherwise. We now use Lemma C.2. Let $(s_i)_i$ be as in the lemma; thus in particular $\{s_1, s_2, s_3\} = \{\pm\}$. We have to verify that every n_j in the interior of the triangle has a constant sign path to the boundary. Indeed, n_7 is connected to the n_j for j = 1, 2, 3 if $s_j = s_7$ and n_8 is also connected to a boundary point as otherwise $s_1 = s_6 = s_3 \neq s_8$, and therefore $s_8 = s_7 = s_2 = s_9$, a contradiction which establishes the claim by symmetry.

References

- [Abu16] R. Abuaf, Categorical crepant resolutions for quotient singularities, Math. Z. 282 (2016), no. 3-4, 679–689.
- [Aus86] M. Auslander, Isolated singularities and existence of almost split sequences, Proc. ICRA IV, Lecture Notes in Mathematics, vol. 1178, Springer Verlag, 1986, pp. 194– 241.

[BCS05] L. Borisov, L. Chen, and G. Smith, The orbifold Chow ring of toric Deligne-Mumford stacks, Journal of the American Mathematical Society 18 (2005), no. 1, 193–215. [BH06] L. A. Borisov and R. P. Horja, On the K-theory of smooth toric DM stacks, Snowbird lectures on string geometry, Contemp. Math., vol. 401, Amer. Math. Soc., Providence, RI, 2006, pp. 21-42. [BKR01] T. Bridgeland, A. King, and M. Reid, The McKay correspondence as an equivalence of derived categories, J. Amer. Math. Soc. 14 (2001), no. 3, 535–554 (electronic). [Boc12] R. Bocklandt, Generating toric noncommutative crepant resolutions, J. Algebra 364 (2012), 119-147.[Bri02] T. Bridgeland, Flops and derived categories, Invent. Math. 147 (2002), no. 3, 613-632. [Bro12] N. Broomhead, Dimer models and Calabi-Yau algebras, Mem. Amer. Math. Soc. 215 (2012), no. 1011, viii+86. [BVdB03] A. Bondal and M. Van den Bergh, Generators and representability of functors in commutative and noncommutative geometry, Mosc. Math. J 3 (2003), no. 1, 1-36. [CLS11] D. A. Cox, J. B. Little, and H. K. Schenck, Toric varieties, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. [DHZ01] D. I. Dais, C. Haase, and G. M. Ziegler, All toric local complete intersection singularities admit projective crepant resolutions, Tohoku Math. J. (2) 53 (2001), no. 1, 95 - 107.[DITV15] H. Dao, O. Iyama, R. Takahashi, and C. Vial, Non-commutative resolutions and Grothendieck groups, J. Noncommut. Geom. 9 (2015), no. 1, 21-34. [GKZ89] I. M. Gel'fand, M. M. Kapranov, and A. V. Zelevinsky, Newton polyhedra of principal A-determinants, Dokl. Akad. Nauk SSSR 308 (1989), no. 1, 20-23. [GKZ08] _, Discriminants, resultants and multidimensional determinants, Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2008, Reprint of the 1994 edition. [Gul08] D. R. Gulotta, Properly ordered dimers, R-charges, and an efficient inverse algorithm, J. High Energy Phys. (2008), no. 10, 14–31. [HR14] J. Hall and D. Rydh, Perfect complexes on algebraic stacks, arXiv:1405.1887 [math.AG], to appear Compositio Math., 2014. [IU08] A. Ishii and K. Ueda, On moduli spaces of quiver representations associated with dimer models, Higher dimensional algebraic varieties and vector bundles, RIMS Kôkyûroku Bessatsu, B9, Res. Inst. Math. Sci. (RIMS), Kyoto, 2008, pp. 127–141. [IU15] _, Dimer models and the special McKay correspondence, Geom. Topol. 19 (2015), no. 6, 3405-3466. [IU16] _, Dimer models and crepant resolutions, Hokkaido Math. J. 45 (2016), no. 1, 1 - 42O. Iyama and M. Wemyss, On the noncommutative Bondal-Orlov conjecture, J. Reine [IW13] Angew. Math. 683 (2013), 119-128. [IW14a] , Maximal modifications and Auslander-Reiten duality for non-isolated singularities, Invent. Math. 197 (2014), no. 3, 521-586. [IW14b] _, Singular derived categories of \mathbb{Q} -factorial terminalizations and maximal modification algebras, Adv. Math. 261 (2014), 85-121. [Kuz08] A. Kuznetsov, Lefschetz decompositions and categorical resolutions of singularities, Selecta Math. (N.S.) 13 (2008), no. 4, 661-696. [Leu12] G. J. Leuschke, Non-commutative crepant resolutions: scenes from categorical geometry, Progress in commutative algebra 1, de Gruyter, Berlin, 2012, pp. 293-361. [LMB99] G. Laumon and L. Moret-Bailly, Champs algébriques, vol. 39, Springer Science & Business Media, 1999. [Log08]T. Logvinenko, Derived McKay correspondence via pure-sheaf transforms, Math. Ann. **341** (2008), no. 1, 137–167. [PVdB15] A. Polishchuk and M. Van den Bergh, Semiorthogonal decompositions of the categories of equivariant coherent sheaves for some reflection groups, 2015, arXiv:1503.04160[math.AG]. F. Santos, Geometric bistellar flips: the setting, the context and a construction, Inter-[San06] national Congress of Mathematicians. Vol. III, Eur. Math. Soc., Zürich, 2006, pp. 931-962.

NON-COMMUTATIVE CREPANT RESOLUTIONS FOR SOME TORIC SINGULARITIES II 25

- [ŠVdB17a] Š. Špenko and M. Van den Bergh, Non-commutative resolutions of quotient singularities, Invent. Math. (2017), 1–65.
- [ŠVdB17b] _____, Noncommutative crepant resolutions for some toric singularities I, arXiv:1701.0525 [math.AG], 2017.
- [VdB93] M. Van den Bergh, Cohen-Macaulayness of semi-invariants for tori, Trans. Amer. Math. Soc. 336 (1993), no. 2, 557–580.
- [VdB04] _____, Non-commutative crepant resolutions, The legacy of Niels Henrik Abel, Springer, Berlin, 2004, pp. 749–770.
- [Wem14] M. Wemyss, Flops and clusters in the homological minimal model program, arXiv:1411.7189 [math.AG], to appear Invent. Math., 2014.
- [Wem16] _____, Noncommutative resolutions, Noncommutative algebraic geometry, Math. Sci. Res. Inst. Publ., vol. 64, Cambridge Univ. Press, New York, 2016, pp. 239–306.
- [Zas98] A. Zastrow, *Planar sets are aspherical*, 1997–1998, Habilitationsschrift, Bochum.

DEPARTEMENT WISKUNDE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, B-1050 ELSENE *E-mail address*: spela.spenko@vub.ac.be

DEPARTEMENT WNI, UNIVERSITEIT HASSELT, UNIVERSITAIRE CAMPUS, B-3590 DIEPENBEEK *E-mail address*: michel.vandenbergh@uhasselt.be