# **Made available by Hasselt University Library in https://documentserver.uhasselt.be**

A categorical framework for glider representations

Non Peer-reviewed author version

HENRARD, Ruben & VAN ROOSMALEN, Adam-Christiaan (2020) A categorical framework for glider representations.

Handle: http://hdl.handle.net/1942/32601

# A CATEGORICAL FRAMEWORK FOR GLIDER REPRESENTATIONS

RUBEN HENRARD AND ADAM-CHRISTIAAN VAN ROOSMALEN

Abstract. Glider representations are a generalization of filtered modules over filtered rings. Given a Γ-filtered ring FR and a subset  $\Lambda \subseteq \Gamma$ , we provide a category Glid<sub> $\Lambda$ </sub> FR of glider representations, and show that it is a complete and cocomplete deflation quasi-abelian category. We discuss its derived category, and the subcategories of natural gliders and Noetherian gliders.

If R is a bialgebra over a field k and FR is a filtration by bialgebras, we show that Glid<sub>Λ</sub> FR is a monoidal category which is derived equivalent to the category of representations of a semi-Hopf category (in the sense of Batista, Caenepeel, and Vercruysse). We show that the monoidal category of glider representations associated to the one-step filtration  $k \cdot 1 \subseteq R$  of a bialgebra R is sufficient to recover the bialgebra R by recovering the usual fiber functor from Glid<sub>Λ</sub> FR. When applied to group algebras, this shows that the monoidal category Glid<sub>Λ</sub>  $F(kG)$  alone is sufficient to distinguish even isocategorical groups.

**CONTENTS** 



<sup>2010</sup> Mathematics Subject Classification. 16W70; 18E10, 18E35.



#### 1. INTRODUCTION

Filtered rings and their representations occur naturally in many areas of mathematics and physics, for example in the study of D-modules and quantum groups (see, for example, [2, 26, 27, 32]). The following example illustrates that sometimes one is interested in truncated filtered representations. Consider the Z-filtered ring  $\mathbb{R}[t]$  with  $\deg(t) = 1$ . For each  $i \geq 0$ , write  $C^i(\mathbb{R})$  for the *i*-times continuously differentiable functions on R, and consider the chain

$$
\cdots \subseteq C^{i+1}(\mathbb{R}) \subseteq C^i(\mathbb{R}) \subseteq \cdots \subseteq C^1(\mathbb{R}) \subseteq C^0(\mathbb{R}).
$$

Whenever  $i \geq n$ ,  $t^n$  induces an action  $t^n: C^i(\mathbb{R}) \to C^{i-n}(\mathbb{R}): f \mapsto \frac{d^n f}{dx^n}$ . In this way, the differential operator  $\frac{d}{dx}$  (which is only densely defined on  $C^0(\mathbb{R})$ ) is encoded algebraically into this filtration.

This type of truncated filtered representations were studied by Nawal and Van Oystaeyen in 1995 (see [29]) under the name of *fragments* (as the definition of fragment has changed over the years, we will refer to this concept as a *prefragment*). Given a ring R and a positive ring filtration  $S = F_0R \subseteq F_1R \subseteq F_2R \subseteq F_3$  $\cdots \subseteq F_n R \subseteq \ldots$  one can consider an FR-fragment M, i.e. an  $\mathbb{Z}^{\leq 0}$ -filtered S-module M together with S-submodules

$$
\ldots \subseteq M_{-i} \subseteq M_{-i+1} \subseteq \ldots \subseteq M_0 = M
$$

and "fragmented actions"  $\phi: F_n R \times M_{-n} \to M$ . Going deeper into the chain, one gradually sees more of the R-action. The precise definition is recalled in definition 2.3.

Looking at the definition of an  $FR$ -prefragment, it is not clear to what extent one should require the fragmented actions to be compatible with each other. Over the course of several papers, the original definition of an  $FR$ -prefragment has been amended, requiring more associativity conditions on the partial actions (compare, for example, the definition in [29] to the one in [14]). One way to circumvent the associativity issue is to require that all partial actions of an  $FR$ -prefragment  $M$  are induced by some enveloping R-module  $\Omega_M \supseteq M$ . Prefragments satisfying this additional condition are called *glider representations*. Example 4.12 below shows that  $\{C^i(\mathbb{R})\}_{i\leq 0}$  is, in fact, a glider representation.

Recently, the theory of glider representations has regained some attention (see for example [9, 11, 12] and the book [14]). Despite these new developments, a categorical framework for glider representations is missing (see [14] or remark 2.6).

We construct a category of glider representations over a filtered ring  $FR$  as a localization of the category of pregliders. For the purpose of this introduction, we sketch the construction of the categories of glider and preglider representations. Let Γ be an ordered group and fix any subset  $\Lambda \subseteq \Gamma$ . Let FR be a Γfiltration of a ring R and let  $\Omega$  be any R-module. We choose subgroups  $\{M_{\lambda}\}_{{\lambda}\in\Lambda}$  of  $\Omega$  such that the module action  $R \times \Omega \to \Omega$  restricts to fragmented actions  $F_{\lambda\mu^{-1}}R \times M_{\mu} \to M_{\lambda}$  (for all  $\lambda, \mu \in \Lambda$ ). We refer to  $\Omega$  together with the subgroups  $\{M_{\lambda}\}_{\lambda\in\Lambda}$  as a preglider. A morphism  $(\{M_{\lambda}\}_{\lambda\in\Lambda},\Omega_M)\to (\{N_{\lambda}\}_{\lambda\in\Lambda},\Omega_N)$ between pregliders is an R-module morphism  $f: \Omega_M \to \Omega_N$  such that  $f(M_\lambda) \subseteq N_\lambda$ , for each  $\lambda \in \Lambda$ .

We then define the category of glider representations as the localization (Preglid<sub>Λ</sub>  $FR$ )[ $\Sigma^{-1}$ ] where Σ consists of those morphisms such that the all induced maps  $f_\lambda: M_\lambda \to N_\lambda$  are isomorphisms (but  $f: M \to N$  itself need not be an isomorphism). The set  $\Sigma$  is a right multiplicative system in Preglid<sub>Λ</sub> FR, so that the localization can be described using roofs (see proposition 3.12).

The category Prefrag<sub>Λ</sub> FR of prefragments does, in general, not recover sufficient structure of the filtered ring: this is illustrated in example 9.24. In contrast, the category Preglid<sub>Λ</sub> FR contains too much information: we see from example 3.15 that the category of pregliders does not reduce to the usual category of filtered modules. The category Glid<sub>Λ</sub> FR of glider representations lies in between the pregliders and the prefragments in the following way. We can consider the restriction map  $j^*$ : Preglid<sub>A</sub>  $FR \to \text{Prefrag}_{\Lambda} FR$ which forgets the ambient R-module  $\Omega$ . The category Glid<sub>Λ</sub> FR is obtained from the usual factorization of  $j^*$ : Preglid<sub>A</sub>  $FR \to \text{Glid}_{\Lambda} FR \to \text{Prefix}_{\Lambda} FR$  where  $\text{Glid}_{\Lambda} FR = \Sigma^{-1} \text{Prefix}_{\Lambda} FR$  and  $\Sigma$  consists of all morphisms  $\sigma$  between pregliders such that the restriction  $j^*(\sigma)$  is invertible.

The following theorem provides different interpretations of the category of glider representations (see proposition 4.8 and corollary 4.10).

**Theorem 1.1.** *Let*  $\Gamma$  *be an ordered group and let*  $\Lambda \subseteq \Gamma$ .

- *(1) The category* Glid<sub>Λ</sub> FR *is a reflective subcategory of* Prefrag<sub>Λ</sub> FR, and
- *(2) the category* Glid<sub>Λ</sub> FR *is a coreflective subcategory of* Preglid<sub>Λ</sub> FR.

The second statement implies that a glider representation admits a canonical ambient R-module. The first statement indicates that a glider representation is a prefragment satisfying additional properties, rather than possessing additional structure. In proposition 4.11, we provide some criterion to decide whether a preframent is a glider representation.

In order to describe the rich structure of the category of glider representations, we will describe the relations with several other categories. Figure 1 below showcases the categories involved as well as some adjunctions. Here, the preadditive category  $\mathcal{F}_{\Lambda}R$  is given by taking as objects the set  $\Lambda$ ; the Hom-spaces are given by the the filtered ring FR. Likewise,  $\overline{\mathcal{F}}_{\Lambda}R$  is given by adjoining an extra object with R as endomorphism ring (see definition 3.1 for details). We refer to  $\mathcal{F}_{\Lambda}R$  and  $\overline{\mathcal{F}}_{\Lambda}R$  as the *filtered companion category* and the *extended filtered companion category*.



Figure 1. Overview diagram

The categories Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR are torsion-free classes in Mod  $\overline{\mathcal{F}}_0R$  and Mod  $\mathcal{F}_0R$ , respectively (proposition 4.2). As such, they are quasi-abelian categories (that is, they admit kernels and cokernels, and the class of all kernel-cokernel pairs endows them with the structure of a Quillen exact category). In contrast, the category Glid<sub>Λ</sub> FR has kernels and cokernels, but the class of kernel-cokernel pairs of  $\text{Glid}_{\Lambda} FR$  is not closed under pushouts (but it is closed under pullbacks). Such a structure has been called a *left almost abelian category* in [36] (we will refer to it as a *deflation quasi-abelian category*).

The following theorem is shown in §5.

- **Theorem 1.2.** (1) The categories  $\text{Preglid}_{\Lambda} FR$  and  $\text{Prefrag}_{\Lambda} FR$  are complete and cocomplete quasi*abelian categories.*
	- *(2) The category* Glid<sup>Λ</sup> F R *is a complete and cocomplete deflation quasi-abelian category.*

In fact, one shows that these categories are Grothendieck (deflation) quasi-abelian (see §5.3).

The conflation structure on Glid<sub>Λ</sub> FR is compatible with the localization Q: Preglid<sub>Λ</sub> FR  $\rightarrow$  Glid<sub>Λ</sub> FR and the embedding  $\phi: \text{Glid}_{\Lambda} FR \to \text{Prefix}_{\Lambda} FR$  in the following sense: it is the minimal conflation structure such that Q preserves conflations, and the maximal conflation structure such that  $\phi$  preserves conflations (see corollary 5.6 and proposition 5.8).

In §5, we will also use the following interpretation of the category  $\text{Glid}_{\Lambda} FR$ : it is given as the quotient Preglid<sub>Λ</sub>  $FR/i_*(\text{Mod }R)$  in the sense of [24]. This is an example of a one-sided exact quotient of a two-sided exact category.

As Glid<sub>Λ</sub> FR is a deflation quasi-abelian category, it admits a meaningful (bounded) derived category  $\mathbf{D}^{\mathrm{b}}(\mathcal{E})$  (see [4, 23]). Using the main result of [23], we obtain a Verdier localization sequence

$$
\mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(R)) \to \mathbf{D}^{\mathrm{b}}(\mathrm{Preglid}_{\Lambda}(FR)) \to \mathbf{D}^{\mathrm{b}}(\mathrm{Glid}_{\Lambda}(FR)).
$$

Moreover, in section 8, we show that the categories in each column of figure 1 are derived equivalent. Hence, the above Verdier localization sequence is equivalent to the sequence

 $\mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(R)) \to \mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(\overline{\mathcal{F}}_{\Lambda}R)) \to \mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(\mathcal{F}_{\Lambda}R)).$ 

In particular, as in [37] we find that  $\mathbf{D}^{\mathrm{b}}(\text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)) \simeq \mathbf{D}^{\mathrm{b}}(\text{Glid}_{\Lambda}(FR))$  are triangle equivalent.

In section §9 we consider a filtration FB of a k-bialgebra B such that each  $F_nB$  is itself a k-coalgebra. For these type of filtrations the companion categories,  $\mathcal{F}_{\Lambda}R$  and  $\overline{\mathcal{F}}_{\Lambda}R$ , are *semi-Hopf categories* in the sense of [3, 16]. It follows that the category  $\mathrm{Glid}_{\Lambda} FB$  is a monoidal category. It is well-known that  $\text{Mod } B$  alone is not sufficient to reconstruct the bialgebra B. On the other hand, the following theorem indicates that the monoidal category Glid<sub>Λ</sub> FB (or, as we will assume that B is finite-dimensional, the full subcategory glid $\Lambda$  FB consisting of the noetherian objects) contains enough information to reconstruct the bialgebra B.

**Theorem 1.3.** Let B and B' be bialgebras and consider the trivial filtrations  $k \cdot 1_B \subseteq B$  and  $k \cdot 1_{B'} \subseteq B$ B'. The categories  $\text{glid}_{\Lambda}(FB)$  and  $\text{glid}_{\Lambda}(FB')$  are monoidally equivalent if and only if B and B' are *isomorphic as bialgebras.*

In particular, the above theorem allows to distinguish even isocategorical groups (see [19]) from the monoidal structure of the category glid<sub>Λ</sub>  $F(k)$  (without referring to the symmetric structure). This provides a conceptual explanation as to why the generalized character ring (which is related to the representation ring of the category glid<sub>Λ</sub>  $F(kG)$ ) discussed in [10] is capable of recovering more of the group structure than the ordinary character ring.

Structure of the paper. We now turn to an overview of the paper. Section §2 is preliminary in nature. We recall some definitions and results that will be used throughout the paper.

In section §3 we construct the category Glid<sub>Λ</sub> FR of glider representations. We consider general Γfiltered rings (where Γ is a filtered group) and consider  $\Lambda \subseteq \Gamma$ . We encode the information of  $\Lambda$  and of the Γ-filtered ring FR in the componanion categories  $\mathcal{F}_{\Lambda}R$  and  $\overline{\mathcal{F}}_{\Lambda}R$ , and use these to define the categories of fragments and (pre)glider representations.

In section §4 we provide a framework in which the category of glider representations fits naturally. We construct the diagram given in figure 1 and show that the top row is a recollement of abelian categories. The second row is then a restriction of the top row, and the category glider representations occurs naturally via a factorization of the restriction functor  $j^*$ : Preglid<sub>A</sub>  $FR \to \text{Prefix}_{\text{A}} FR$ .

In section §5 we study the homological properties of the categories in figure 1. In particular, we show that the categories  $\text{Preglid}_{\Lambda} FR$  and  $\text{Preglid}_{\Lambda} FR$  are Grothendieck quasi-abelian and that  $\text{Glid}_{\Lambda}(FR)$  is a Grothendieck deflation quasi-abelian category.

In sections §7 and §6 we look at some interesting subcategories of the category Glid<sub>Λ</sub> FR, namely the category NGlid<sub>Λ</sub> FR of natural glider representations and the category glid<sub>Λ</sub> FR of noetherian glider representations. We show that both of these still carry the structure of deflation-exact categories.

Section 8 deals with the (bounded) derived categories of all categories involved. In particular, we show that the localization sequence  $Mod(R) \rightarrow Preglid_A FR \rightarrow Glid_A FR$  induces a Verdier localization sequence  $\mathbf{D}^{\text{b}}(\text{Mod}(R)) \to \mathbf{D}^{\text{b}}(\text{Pred}_{\Lambda} FR) \to \mathbf{D}^{\text{b}}(\text{Glid}_{\Lambda} FR)$ . Moreover, this Verdier localization sequence is equivalent to the Verdier localization sequence  $\mathbf{D}^{\mathrm{b}}(\text{Mod}(R)) \to \mathbf{D}^{\mathrm{b}}(\text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)) \to \mathbf{D}^{\mathrm{b}}(\text{Mod}(\mathcal{F}_{\Lambda}R)).$ 

Finally, in section 9 we show that given a k-bialgebra B together with a filtration  $FB$  by bialgebras, the companion categories  $\mathcal{F}_{\Lambda}B$  and  $\overline{\mathcal{F}}_{\Lambda}B$  are semi-Hopf categories. In particular, the categories  $\text{Mod}_k(\mathcal{F}_{\Lambda}B)$ and  $\text{Mod}_k(\overline{\mathcal{F}}_{\Lambda}B)$  inherit a natural tensor structure. Moreover, the derived equivalences of the previous section are compatible with the tensor structure as well. We end by showing theorem 1.3 which shows that considering the monoidal category of glider representations of a filtered object retains enough information to reconstruct the original object.

Acknowledgements The authors would like to thank Francesco Genovese, Geoffrey Janssens, Tao Lu, and Wenqing Tao for helpful discussions and comments. We especially want to thank Frederik Caenepeel and Fred Van Oystaeyen for many useful discussions and ideas. We are grateful to Geoffrey Janssens and Frederik Caenepeel for sharing an early version of [10]; section 9.2 and the results therein were motivated by those discussions and results. The second author is currently a postdoctoral researcher at FWO (12.M33.16N).

#### 2. Preliminaries

This section is preliminary in nature. Throughout, we assume that all rings and ringhomomorphisms are unital.

2.1. Filtered rings. Let  $\Gamma$ ,  $\leq$  be an ordered group, i.e.  $\Gamma$  is a group,  $\leq$  is an ordering on  $\Gamma$ , satisfying the following property:

$$
\forall \alpha, \beta, \gamma, \delta \in \Gamma : (\alpha \le \beta \land \gamma \le \delta) \Rightarrow \alpha \gamma \le \beta \delta.
$$

**Remark 2.1.** If  $\Gamma$ ,  $\leq$  is an ordered group, then the poset category of  $\Gamma^+ := \{ \gamma \in \Gamma \mid e \leq \gamma \}$  is filtered.

Let R be a unital ring. A Γ-*filtration* on R is a Γ-indexed set of additive subgroups  $\{F_{\gamma}R\}_{\gamma\in\Gamma}$  of R satisfying:

(1) 
$$
1_R \in F_e R
$$
,  
\n(2)  $\forall \alpha, \beta \in \Gamma : \alpha \leq \beta \Rightarrow F_\alpha R \subseteq F_\beta R$ ,  
\n(3)  $\forall \alpha, \beta \in \Gamma : (F_\alpha R)(F_\beta R) \subseteq F_{\alpha\beta} R$ .

A Γ*-filtered ring* is a ring R together with a Γ-filtration on R. We write F R for the Γ-filtered ring  ${F_{\gamma}R}_{\gamma\in\Gamma}.$ 

Remark 2.2. Even though we will assume Γ is a group, it is straightforward to generalize our results to the case where  $\Gamma$  is a cancellative monoid.

2.2. Fragments and gliders. The definition of a fragment over a filtered ring has changed since its original definition in [29]. To avoid confusion with the terminology used in [9, 11, 12, 14, 15], we refer to the objects defined in [29] as prefragments. We start by recalling the definition.

**Definition 2.3.** Let FR be an N-filtered ring with subring  $S = F_0R$ . A *(left) prefragment over* FR is a left S-module together with a descending chain of subgroups

$$
M = M_0 \supseteq M_{-1} \supseteq \cdots \supseteq M_{-i} \supseteq \ldots
$$

satisfying the following properties.

 $f_1$  For every  $i \in \mathbb{N}$  there is a map  $\phi_i \colon F_i R \times M_{-i} \to M \colon (\lambda, m) \mapsto \lambda \cdot m$  such that

$$
\begin{array}{rcl}\n\lambda \cdot (m+n) & = & \lambda \cdot m + \lambda \cdot n, \\
(\lambda + \mu) \cdot m & = & \lambda \cdot m + \mu \cdot n, \\
1 \cdot m & = & m,\n\end{array}
$$

for all  $\lambda, \mu \in F_i R$  and  $m, n \in M_i$ .

 $f_2$  For every  $i, j \in \mathbb{N}$  with  $j + i \leq 0$ , there is a commutative diagram

$$
\begin{CD} M & \xrightarrow{\hspace{12mm}} M_{j-i} \xrightarrow{\hspace{12mm}} M \\ \uparrow{\hspace{12mm}} \phi_i & \uparrow{\hspace{12mm}} \phi_j \\ F_iR \times M_{-i} \xleftarrow{\hspace{12mm}} i_F \xrightarrow{\hspace{12mm}} F_jR \times M_{-i} \xleftarrow{\hspace{12mm}} i_M \xrightarrow{\hspace{12mm}} F_jR \times M_{-j} \xleftarrow{\hspace{12mm}} i_M \xleftarrow{\hspace{12mm}} i_H K \times M_{-j} \xleftarrow{\hspace{12mm}} i_M \xleftarrow{\hspace{12mm}} i_H K \times M_{-j} \xleftarrow{\hspace{12mm}} i_M K \times M_{-j} \xleftarrow{\hspace{12mm}} i_M K \times M_{-j} \xleftarrow{\hspace{12mm}} i_H K \times M_{-j} \xleftarrow{\hspace{12mm}} i_M K \times M_{-j} \xleftarrow{\hspace{12mm}} i_M
$$

 $f_3$  For every  $i, j \in \mathbb{N}$ , there is a commutative diagram

$$
F_i R \times F_j R \times M_{-(i+j)} \xrightarrow{m \times 1_{M_{-(i+j)}}} F_{i+j} R \times M_{-(i+j)}
$$
  
\n
$$
\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow
$$
  
\n
$$
F_i R \times M_{-i} \xrightarrow{\phi_i} M
$$

Here m denotes the multiplication in R and the left vertical arrow is defined using  $f_2$ .

Let M and N be FR-fragments. A morphism  $f: M \to N$  of FR-fragments is an S-linear map  $f: M \to N$  such that  $f(M_i) \subseteq N_i$  for all  $i \in \mathbb{N}$  and  $f(r \cdot m) = r \cdot f(m)$  for all  $r \in F_i R$  and  $m \in M_i$ .

**Remark 2.4.** The prefragments over a filtered ring  $FR$  form an additive category.

**Definition 2.5.** Let  $\{M_i\}_i$  be a prefragment over a filtered ring FR. If the fragmented scalar multiplications  $F_iR \times M_{-i} \to M$  are induced from an R-module  $\Omega \supseteq M$ , we say that M is a *glider representation*.

Let M and N be glider representations. A morphism  $f: M \to N$  of prefragments is called a *morphism of glider representations* if there exist R-modules  $\Omega_M$  and  $\Omega_N$  such that  $M \subseteq \Omega_M$  and  $N \subseteq \Omega_N$  exhibiting that M, N are glider representations and there exists an R-linear map  $F: \Omega_M \to \Omega_N$  such that the following diagram commutes:



(thus, the map between the prefragments  $M \to N$  needs to be induced by an R-module morphism  $\Omega_M \to \Omega_N$ ).

Remark 2.6. Despite remark 2.4, it is not clear that the glider representations form a category. Indeed, let  $f: A \to B$  and  $q: B \to C$  be morphisms of FR-glider representations. By definition, there are ambient R-modules  $\Omega_A, \Omega_B, \Omega'_B, \Omega_C$  and R-module maps  $F: \Omega_A \to \Omega_B, G: \Omega'_B \to \Omega_C$  such that the diagrams



commute. However, it is not clear whether the composition  $g \circ f$  defines a morphism of FR-glider representations.

Remark 2.7. The definitions of prefragments and glider representations can be adjusted to accommodate more general filtered rings. For prefragments, this will be done in definition 3.3. In definition 3.8, we will give a different definition for a morphism of gliders. It will follow from proposition 4.8 and 4.9 that this new definition is compatible with the one in definition 2.5.

2.3. Localizations of categories. We recall some basic results about localizations of categories. The material of this section is based on [21].

**Definition 2.8.** Let C be a small category and let  $\Sigma \subseteq \text{Mor } \mathcal{C}$  be a subset of morphisms of C. The *localization of* C with respect to  $\Sigma$  is a functor  $Q: \mathcal{C} \to \mathcal{C}[\Sigma^{-1}]$ , universal with respect to the property such that  $Q(s)$  is invertible, for all  $s \in \Sigma$ .

**Remark 2.9.** The universality in definition 2.8 means that, for every category  $D$ , the functor  $(Q \circ$  $-$ ): Fun( $\mathcal{C}[\Sigma^{-1}], \mathcal{D}$ ) → Fun( $\mathcal{C}, \mathcal{D}$ ) induces an isomorphism between Fun( $\mathcal{C}[\Sigma^{-1}], \mathcal{D}$ ) and the full subcategory of Fun( $\mathcal{C}, \mathcal{D}$ ) consisting of those functors  $F : \mathcal{C} \to \mathcal{D}$  which make every  $s \in \Sigma$  invertible.

The following proposition is standard (see [21, proposition I.1.3]).

**Proposition 2.10.** Let  $F: \mathcal{C} \to \mathcal{D}$  and  $G: \mathcal{D} \to \mathcal{C}$  be functors between small categories. We write  $\Sigma \subseteq \text{Mor } \mathcal{C}$  *for the set of all morphisms* f *for which*  $F(f)$  *is invertible. If* F *is left adjoint to* G, then the *following are equivalent:*

- *(1)* G *is fully faithful,*
- *(2) the counit*  $FG \Rightarrow 1_{\mathcal{D}}$  *is invertible.*
- *(3) the unique map*  $H: \mathcal{C}[\Sigma^{-1}] \rightarrow \mathcal{D}$  *making the diagram*



*commute, is an equivalence.*

We recall the following proposition from [21].

Proposition 2.11. *Consider functors* F, G, H *as in the diagram*



- *(1) Assume that*  $G: \mathcal{D} \to \mathcal{E}$  *is a localization.* 
	- *(a) If* HG *is left adjoint to* F*, then* H *is left adjoint to* GF*.*

*(b) If* F *is left adjoint to* HG*, then* GF *is left adjoint to* H*.*

- *(2)* Assume that  $G: \mathcal{D} \to \mathcal{E}$  is fully faithful.
	- *(a) If* H *is left adjoint to* GF*, then* HG *is left adjoint to* F
	- *(b) If* GF *is left adjoint to* H*, then* F *is left adjoint to* HG*.*

*Proof.* The statement (1a) is shown in [21, lemma I.1.3.1], and uses that  $\sim \text{G}$ : Fun $(\mathcal{E}, \mathcal{X}) \to \text{Fun}(\mathcal{D}, \mathcal{X})$ is fully faithful (as G is a localization). One can prove  $(1b)$  in a similar way. The other statements can be shown in a similar fashion, using that, for each category X, the functor  $G \circ -: \text{Fun}(\mathcal{X}, \mathcal{D}) \to \text{Fun}(\mathcal{X}, \mathcal{E})$ is fully faithful.  $\square$ 

In this paper, we often consider localizations with respect to right multiplicative systems.

**Definition 2.12.** Let C be a category and let  $\Sigma \subseteq \text{Mor } C$  be a subset of morphisms of C. We say that  $\Sigma$ is a *right multiplicative system* if the following properties are satisfied.

**RMS1** For every object A of C the identity  $1_A$  is contained in  $\Sigma$ , and  $\Sigma$  is closed under composition. RMS2 Every solid diagram



with  $s \in \Sigma$  can be completed to a commutative square as above, with  $t \in \Sigma$ . **RMS3** For every pair of morphisms  $f, g: X \to Y$  and every  $s \in \Sigma$  with source Y such that  $s \circ f = s \circ g$ there exists a  $t \in \Sigma$  with target X such that  $f \circ t = q \circ t$ .

For localizations with respect to a right multiplicative system  $\Sigma \subseteq \text{Mor }\mathcal{C}$ , we have the following description of the localization  $\mathcal{C}[\Sigma^{-1}].$ 

**Description 2.13.** Let C be a category and  $\Sigma$  a right multiplicative system in C. We define a category  $\Sigma^{-1}\mathcal{C}$  as follows:

- (1) We have  $Ob(\Sigma^{-1}\mathcal{C}) = Ob(\mathcal{C}).$
- (2) Let  $f_1: X_1 \to Y, s_1: X_1 \to X, f_2: X_2 \to Y, s_2: X_2 \to X$  be morphisms in C with  $s_1, s_2 \in \Sigma$ . We say that the pairs  $(f_1, s_1), (f_2, s_2) \in (\text{Mor } \mathcal{C}) \times \Sigma$  are equivalent (denoted by  $(f_1, s_1) \sim (f_2, s_2)$ ) if there exists a third pair  $(f_3: X_3 \to Y, s_3: X_3 \to X) \in (\text{Mor }\mathcal{C}) \times \Sigma$  and morphisms  $u: X_3 \to X_3$  $X_1, v \colon X_3 \to X_2$  such that



is a commutative diagram.

- (3) Hom<sub> $\Sigma^{-1}C(X,Y) = \{(f,s) | f \in \text{Hom}_{\mathcal{C}}(X',Y), s \colon X' \to X \text{ with } s \in \Sigma\} / \sim$ </sub>
- (4) The composition of  $(f: X' \to Y, s: X' \to X)$  and  $(g: Y' \to Z, t: Y' \to Y)$  is given by  $(g \circ h: X'' \to Y)$  $Z, s \circ u: X'' \to X$ ) where h and u are chosen to fit in a commutative diagram



which exists by RMS2.

The canonical functor  $Q: \mathcal{C} \to \Sigma^{-1} \mathcal{C}$  satisfies the conditions of definition 2.8.

2.4. Representations of small preadditive categories. Let  $k$  be a commutative ring. Let  $\mathfrak a$  be a small k-linear category. A *(left)*  $\mathfrak{a}$ *-module* is a covariant k-linear functor from  $\mathfrak{a}$  to Mod k, the category of all k-modules. The category of all left  $\alpha$ -modules is denoted by Mod  $\alpha$ . If we do not specify the ring k, we will take  $k = \mathbb{Z}$ .

It follows from the Yoneda lemma that, for every  $A \in \mathfrak{a}$ , the representable  $\mathfrak{a}\text{-module } \mathfrak{a}(A, -)$  is projective. We refer to such an a-module as a *standard projective* a*-module*. It is clear that every finitely generated projective is a direct summand of a finite direct sum of standard projectives. If a has finite direct sums and idempotents split in a, then every finitely generated projective is isomorphic to a standard projective.

If  $f: \mathfrak{a} \to \mathfrak{b}$  is a functor between small preadditive categories, then there is an obvious restriction functor

$$
(-)_\mathfrak{a}\colon \operatorname{Mod}(\mathfrak{b})\to \operatorname{Mod}(\mathfrak{a})
$$

which sends N to  $N \circ f$ . This restriction functor has a left adjoint

$$
\mathfrak{b}\otimes_{\mathfrak{a}}-:\,\mathrm{Mod}(\mathfrak{a})\rightarrow\mathrm{Mod}(\mathfrak{b})
$$

which is the right exact functor sending the projective generators  $a(A, -)$  in Mod(a) to  $b(f(A), -)$  in Mod(b). If  $f: \mathfrak{a} \to \mathfrak{b}$  is fully faithful, then the natural morphism  $M \to (\mathfrak{b} \otimes_{\mathfrak{a}} M)_{\mathfrak{a}}$  is an isomorphism, for all  $M \in \text{Mod } \mathfrak{a}$ . It follows from [21, Chapter I.3] that  $\text{Mod } \mathfrak{a} \simeq (\text{Mod } \mathfrak{b})/\text{ker}(-)_{\mathfrak{a}}$ .

2.5. Recollements. Recollements were introduced in a triangulated context in [5]. To establish notations, we recall the definition of a recollement in both an abelian and a triangulated setting.

**Definition 2.14.** Let A and B be categories and let L, R:  $\mathcal{B} \to \mathcal{A}$  and E:  $\mathcal{A} \to \mathcal{B}$  be functors. The triple  $(L, E, R)$  is called an *adjoint triple* if  $L \dashv E$  and  $E \dashv R$ .

Definition 2.15. A *recollement of abelian categories* is a triple of abelian categories A, B and C and a diagram

$$
\mathcal{A} \xleftarrow{i^* \atop i^!} \mathcal{B} \xleftarrow{j_!} \mathcal{C}
$$

with 6 additive functors satisfying the following conditions:

- (1) the triple  $(i^*, i_*, i^!)$  is an adjoint triple,
- (2) the triple  $(j_!, j^*, j_*)$  is an adjoint triple,
- (3) the functors  $i_*, j_*$  and  $j_!$  are fully faithful,
- (4)  $\text{im}(i_*) = \text{ker}(j^*).$
- **Remark 2.16.** (1) It follows from proposition 2.10 that  $C \simeq \Sigma^{-1} \mathcal{B}$  where  $\Sigma \subseteq \text{Mor } \mathcal{B}$  is the class of morphisms that become invertible under  $j^*$ . These are the morphisms with kernel and cokernel in i∗(A) ⊆ B.
	- (2) By proposition 2.10, the conditions in definition 2.15 are not minimal; it suffices to determine the adjoint triple  $(j_!, j^*, j_*)$  where either  $j_!$  or  $j_*$  is fully faithful (see, for example, [33, remark 2.3]).
	- (3) We will be interested in recollements of abelian categories where all three categories are module categories. It is shown in [34] that these recollements are classified by an idempotent (see also  $[25]$ ).

In the same fashion, one defines a recollement of triangulated categories.

**Definition 2.17.** A *recollement of triangulated categories* is a triple of triangulated categories  $\mathcal{T}'', \mathcal{T}$  and  $\mathcal{T}'$  and a diagram

$$
\mathcal{T}'' \xleftarrow[i]{i^* \atop i^!} \mathcal{T} \xleftarrow[j]{j_!} \mathcal{T}
$$

with 6 triangulated functors satisfying the following conditions:

- (1) the triple  $(i^*, i_*, i^!)$  is an adjoint triple,
- (2) the triple  $(j_!, j^*, j_*)$  is an adjoint triple,
- (3) the functors  $i_*, j_*$  and  $j_!$  are fully faithful,
- (4)  $j^*i_* = 0$ ,
- (5) for each  $X \in \mathcal{T}$ , there are two triangles in  $\mathcal{T}$ :

 $i_*i^!(X) \to X \to j_*j^*(X) \to \Sigma i_*i^!(X)$  and  $j_!j^*(X) \to X \to i_*i^*(X) \to \Sigma j_!j^*(X)$ .

Remark 2.18. The conditions in definition 2.17 are not minimal; it suffices to determine the adjoint triple  $(j_!, j^*, j_*)$  of exact functors where either  $j_!$  or  $j_*$  is fully faithful (see, for example, [22, proposition 1.14]).

2.6. One-sided exact categories and admissibly percolating subcategories. One-sided exact categories were introduced in [4, 36] as a framework for studying one-sided quasi-abelian categories [36]. We recall some definitions as well as some results concerning quotients of one-sided exact categories by percolating subcategories ([23, 24]).

**Definition 2.19.** Let  $C$  be an additive category. We say that a sequence  $A \xrightarrow{f} B \xrightarrow{g} C$  is a *kernel-cokernel pair* if  $f = \ker q$  and  $q = \text{coker } f$ . A *conflation category* C is an additive category with a chosen class of kernel-cokernel pairs, called *conflations*, closed under isomorphisms. Given a conflation  $A \xrightarrow{f} B \xrightarrow{g} C$ , the map f is called an *inflation* and the map g is called a *deflation*. Inflations are denoted by  $\rightarrow$  and deflations by  $\rightarrow$ . A map  $f: X \rightarrow Y$  is called *admissible* if it admits a deflation-inflation factorization, i.e. f factors as  $X \rightarrow Z \rightarrow Y$ .

Let C and D be conflation categories. An additive functor  $F: \mathcal{C} \to \mathcal{D}$  is called *exact* or *conflation-exact* if it maps conflations in  $\mathcal C$  to conflations in  $\mathcal D$ .

**Definition 2.20.** A conflation category  $\mathcal C$  is called *right exact* or *deflation-exact* if it satisfies the following axioms:

**R0** The identity morphism  $1_0: 0 \rightarrow 0$  is a deflation.

R1 The composition of two deflations is a deflation.

R2 Pullbacks along deflations exist and deflations are stable under pullbacks.

Dually, we call a conflation category  $\mathcal C$  *left exact* or *inflation-exact* if the opposite category  $\mathcal C^{op}$  is right exact. For completeness, an inflation-exact category is a conflation category such that the distinguished class of conflations satisfies the following axioms:

**L0** The identity morphism  $1_0: 0 \rightarrow 0$  is an inflation.

L1 The composition of two inflations is an inflation.

L2 Pushouts along inflations exist and inflations are stable under pushouts.

A conflation category which is both inflation-exact and deflation-exact is a (Quillen) *exact category*.

**Definition 2.21.** A category C is called *left quasi-abelian* or *deflation quasi-abelian* if it is a pre-abelian category, i.e. an additive category with kernels and cokernels, such that cokernels are stable under pullbacks. Dually, C is called *right quasi-abelian* or *inflation quasi-abelian* if it is a pre-abelian category such that kernels are stable under pushouts.

The category  $\mathcal C$  is called *quasi-abelian* if it is both left and right quasi-abelian.

- Remark 2.22. (1) A deflation quasi-abelian category can be given the structure of a deflation-exact category by choosing all kernel-cokernel pairs as the conflations. Dually, an inflation quasiabelian category can be endowed with the structure of an inflation-exact category by choosing all kernel-cokernel pairs as conflations. To avoid confusion, we prefer using "inflation-exact" and "deflation-exact" over "left exact" and "right exact".
	- (2) A quasi-abelian category has a natural exact structure (the conflations are given by all kernelcokernel pairs).

**Definition 2.23.** Let C be a deflation-exact category. A non-empty full subcategory  $\mathcal A$  of C satisfying the following three axioms is called an *admissibly deflation-percolating subcategory* of C.

A1 A is a *Serre subcategory*, that is:

If  $A' \rightarrowtail A \twoheadrightarrow A''$  is a conflation in C, then  $A \in Ob(\mathcal{A})$  if and only if  $A', A'' \in Ob(\mathcal{A})$ .

**A2** For all morphisms  $C \to A$  with  $C \in Ob(\mathcal{C})$  and  $A \in Ob(\mathcal{A})$ , there exists a commutative diagram



with  $A' \in Ob(\mathcal{A})$ .

**A3** If  $a: C \rightarrow D$  is an inflation and  $b: C \rightarrow A$  is a deflation with  $A \in Ob(A)$ , then the pushout of a along b exists and yields an inflation and a deflation, respectively, i.e.



**Remark 2.24.** (1) An admissibly deflation-percolating subcategory  $A$  of a deflation-exact category  $\mathcal C$  is automatically an abelian category (see [24, proposition 6.4]).

(2) When  $\mathcal{E}$  is an exact category, any full subcategory  $\mathcal{A} \subseteq \mathcal{E}$  satisfies axiom **A3** (this follows, for example, from [8, proposition 2.12]).

Definition 2.25. Let A be an admissibly deflation-percolating subcategory of a deflation-exact category C. A morphism f is called a *weak* A-isomorphism if it is admissible and ker(f), coker(f) ∈ A. A weak isomorphism will often be endowed with  $\sim$ , i.e. we write  $f: A \overset{\sim}{\to} B$  for a weak isomorphism.

**Remark 2.26.** A morphism  $f: X \to Y$  is a weak A-isomorphism if it has a kernel and a cokernel (both lying in A) and factors as  $X \rightarrow X/(\ker f) \rightarrow Y$ .

The following theorem summarizes various results from [24].

Theorem 2.27. *Let* A *be an admissibly deflation-percolating subcategory of a delation-exact category* C*.*

- *(1) The set*  $\Sigma_A$  *of weak* A-isomorphisms is a right multiplicative system.
- (2) The weakest conflation structure on  $\Sigma_{\mathcal{A}}^{-1}$ C for which  $Q: \mathcal{C} \to \Sigma_{\mathcal{A}}^{-1}$ C is conflation-exact, makes  $\Sigma_{\mathcal{A}}^{-1}\mathcal{C}$  *into a deflation-exact category.*
- *(3) The localization functor* Q *is a conflation-exact functor and is 2-universal among conflation-exact functors*  $F: \mathcal{C} \to \mathcal{D}$  *such that*  $F(\mathcal{A}) = 0$ *. Here,*  $\mathcal{D}$  *is any deflation-exact category.*
- *(4) The right multiplicative system*  $\Sigma_A$  *is saturated, i.e.*  $Q(f)$  *is an isomorphism if and only if*  $f \in \Sigma_A$ *.*
- *(5) Pullbacks alongs weak isomorphisms exist and weak isomorphisms are stable under pullbacks.*

**Remark 2.28.** Because of the above universal property, we often write  $C/\mathcal{A}$  for the category  $\Sigma_{\mathcal{A}}^{-1}\mathcal{C}$ .

2.7. Torsion and torsion-free classes. Let A be an abelian category. A *torsion theory* on A is a pair  $(\mathcal{T}, \mathcal{F})$  of full subcategories of A so that Hom<sub>A</sub> $(\mathcal{T}, \mathcal{F}) = 0$  and for every object  $A \in \mathcal{A}$  there is a short exact sequence

$$
0 \to T \to A \to F \to 0
$$

with  $T \in \mathcal{T}$  and  $F \in \mathcal{F}$ . This short exact sequence is necessarily unique up to isomorphism. The subcategory T is called the *torsion subcategory* and the category F is called a *torsion-free subcategory*. Any full subcategory of  $A$  satisfying the following properties is a torsion subcategory:

- (1)  $\mathcal F$  is closed under extensions and subobjects,
- (2) the embedding  $\mathcal{F} \to \mathcal{A}$  has a left adjoint.

The associated torsion subcategory is then given by

$$
\mathcal{T} = {}^{\perp_0} \mathcal{F} = \{ A \in \mathcal{A} \mid \text{Hom}(A, \mathcal{F}) = 0 \}.
$$

Let  $(\mathcal{T}, \mathcal{F})$  be a torsion theory. We say that  $(\mathcal{T}, \mathcal{F})$  is *tilting* if for every  $A \in \mathcal{A}$  there is a monomorphism  $A \to T$  for some  $T \in \mathcal{T}$ . Likewise, we say that  $(\mathcal{T}, \mathcal{F})$  is *cotilting* if for every  $A \in \mathcal{A}$  there is an epimorphism  $F \to A$  for some  $F \in \mathcal{A}$ .

**Remark 2.29.** If A has enough projectives, then a torsion theory  $(\mathcal{T}, \mathcal{F})$  is cotilting if and only if all projective objects lie in F.

We recall the following result from [6, proposition B.3].

Theorem 2.30. *Let* C *be an additive category. The following are equivalent.*

- *(1)* C *is a quasi-abelian category,*
- *(2) There is a cotilting torsion theory*  $(\mathcal{T}, \mathcal{F})$  *in an abelian category A with*  $\mathcal{C} \simeq \mathcal{F}$ .
- *(3) There is a tilting torsion theory*  $(\mathcal{T}, \mathcal{F})$  *in an abelian category A with*  $\mathcal{C} \simeq \mathcal{T}$ .

2.8. The derived category of a deflation-exact category. We recall the definition of the derived category of a deflation-exact category from [4, 23]. The definition is similar to the derived category of an exact category [8, 31]. For an additive category  $\mathcal{E}$ , we write  $\mathbf{C}(\mathcal{E})$  for the category of complexes and  $\mathbf{K}(\mathcal{E})$  for the homotopy category. We write  $\mathbf{C}^{b}(\mathcal{E})$  and  $\mathbf{K}^{b}(\mathcal{E})$  for the bounded variants.

**Definition 2.31.** Let  $\mathcal{E}$  be a deflation-exact category. A complex  $X^{\bullet} \in C^{b}(\mathcal{E})$  is called *acylic in degree* n if  $d_X^{n-1}$  factors as



where the deflation  $p^{n-1}$  is the cokernel of  $d_X^{n-2}$  and the inflation  $i^{n-1}$  is the kernel of  $d_X^n$ .

A complex  $X^{\bullet}$  is called *acyclic* if it is acylic in each degree. The full subcategory of  $\mathbf{C}^{\mathbf{b}}(\mathcal{E})$  of acyclic complexes is denoted by  $\mathbf{Ac}^b_{\mathbf{C}}(\mathcal{E})$ . We write  $\mathbf{Ac}^b_{\mathbf{K}}(\mathcal{E})$  for the full subcategory of acyclic complexes when viewed as a subcategory of  $\mathbf{K}^{b}(\mathcal{E})$ . We simply write  $\mathbf{Ac}^{b}(\mathcal{E})$  if there is no confusion.

In [4], it is shown that  $\mathbf{Ac}_{\mathbf{K}}(\mathcal{E})$  is a triangulated subcategory (not necessarily closed under isomorphisms) of  $\mathbf{K}(\mathcal{E})$ .

**Definition 2.32.** Let  $\mathcal{E}$  be a deflation-exact category. The bounded derived category  $\mathbf{D}^{\mathrm{b}}(\mathcal{E})$  is defined as the Verdier localization  $\mathbf{K}^{\mathrm{b}}(\mathcal{E})/\mathbf{A}\mathbf{c}^{\mathrm{b}}\mathbf{K}(\mathcal{E})$ .

The derived category  $\mathbf{D}^{b}(\mathcal{E})$  enjoys many standard properties as in the exact case. We refer the reader to [23] for details and precise statements. The following theorem is [23, theorem 1.4].

**Theorem 2.33.** Let  $\mathcal{E}$  be a deflation-exact category and let  $\mathcal{A} \subseteq \mathcal{E}$  be an admissibly deflation-percolating *subcategory. The following sequence is a Verdier localization sequence*

$$
\mathbf{D}^{\mathrm{b}}_\mathcal{A}(\mathcal{E}) \to \mathbf{D}^{\mathrm{b}}(\mathcal{E}) \to \mathbf{D}^{\mathrm{b}}(\mathcal{E}/\mathcal{A}).
$$

*Here,*  $D_{\mathcal{A}}^{b}(\mathcal{E})$  *is the thick triangulated subcategory of*  $D^{b}(\mathcal{E})$  *generated by*  $\mathcal{A} \subseteq D^{b}(\mathcal{E})$ *.* 

2.9. Semi-Hopf categories. Let  $V$  be a strict braided monoidal category and let  $\underline{\mathcal{C}}(\mathcal{V})$  be the category of coalgebra (or comonoid) objects in V with coalgebra morphisms. Note that  $\mathcal{C}(\mathcal{V})$  is itself a monoidal category and the unit object  $k$  of  $V$  is a coalgebra.

**Definition 2.34.** Let  $V$  be a strict braided monoidal category. A category C enriched over  $C(V)$  is called a *semi-Hopf category*. If  $\mathcal{V} = \text{Vec}_k$ , the category of vector spaces over a field k, a category C enriched over  $\mathcal{C}(\mathrm{Vec}_k)$  is called a k-linear semi-Hopf category.

When C is a k-linear semi-Hopf category, the category  $\text{Mod}_k \mathcal{C} = \text{Fun}_k(\mathcal{C}, \text{Vec}_k)$  of k-linear C-modules has an induced pointwise monoidal structure (see [3, proposition 3.2] for details).

Remark 2.35. The terminology of a semi-Hopf category has been introduced in [7, 16]. A semi-Hopf category with an antipode is called a *Hopf category*, and was earlier introduced in [3].

#### 3. The category of glider representations

Let Γ be an ordered group and let  $\Lambda \subseteq \Gamma$  be any subset. Let FR be a Γ-filtered ring. In this section, we collect all definitions necessary to define the category  $\text{Glid}_{\Lambda} FR$  of  $\Lambda$ -glider representations over a filtered ring FR. We refer to the diagram in figure 1 for an overview. We start by defining the categories  $\mathcal{F}_{\Lambda}R$ and  $\overline{\mathcal{F}}_0R$ , and proceed by defining Prefrag<sub>Λ</sub> FR and Preglid<sub>Λ</sub> FR as full subcategories of Mod  $\mathcal{F}_0R$  and Mod  $\overline{\mathcal{F}}_{\Lambda}R$ , respectively. The category Glid<sub> $\Lambda$ </sub> FR of glider representation is then defined as a localization of the category Preglid<sub>Λ</sub>  $FR$  of pregliders (see definition 3.8).

The categories Mod  $\mathcal{F}_{\Lambda}R$  and Prefrag<sub> $\Lambda$ </sub> FR do not occur in the definition of glider representations, but will occur in subsequent sections.

3.1. Companion categories over a filtered ring. Let  $(\Gamma, \leq)$  be an ordered group and let FR be a Γ-filtered ring.

**Definition 3.1.** Let  $FR$  be a Γ-filtered ring and let  $\Lambda \subseteq \Gamma$  be any subset.

(1) We define the  $\Lambda$ -*filtered companion category*  $\mathcal{F}_{\Lambda}R$  of FR as follows. The objects are given by  $Ob(\mathcal{F}_{\Lambda}R) = \Lambda$ ; the morphisms are given by

$$
\operatorname{Hom}_{\mathcal{F}_{\Lambda}R}(\alpha,\beta) = \begin{cases} F_{\beta\alpha^{-1}}R & \alpha \le \beta, \\ 0 & \text{otherwise.} \end{cases}
$$

The composition is given by the multiplication in R.

(2) We define the *extended*  $\Lambda$ -*filtered companion category*  $\overline{\mathcal{F}}_{\Lambda}R$  of FR as follows. The objects are given by  $Ob(\mathcal{F}_{\Lambda}R) = \Lambda \coprod \{\infty\}$ ; the morphisms are given by

$$
\text{Hom}_{\overline{\mathcal{F}}_{\Lambda}R}(\alpha,\beta) = \begin{cases} \text{Hom}_{\mathcal{F}_{\Lambda}R}(\alpha,\beta) & \alpha,\beta \in \Lambda, \\ R & \beta = \infty, \\ 0 & \text{otherwise.} \end{cases}
$$

The composition is given by the multiplication in R. We write  $j: \mathcal{F}_{\Lambda}R \to \overline{\mathcal{F}}_{\Lambda}R$  for the inclusion functor.

**Notation 3.2.** For each  $\alpha, \beta \in Ob(\overline{\mathcal{F}}_{\Lambda})$  such that  $\alpha \leq \beta$  (or  $\beta = \infty$ ), there is an element  $1_R \in Hom(\alpha, \beta)$ . We refer to this element as  $1_{\alpha,\beta}$ .

3.2. Pregliders and prefragments. Having introduced the categories  $\mathcal{F}_{\Lambda}R$  and  $\overline{\mathcal{F}}_{\Lambda}R$ , we can now define the categories  $\text{Preglid}_{\Lambda} FR$  and  $\text{Prefrag}_{\Lambda} FR$ .

**Definition 3.3.** Let FR be a Γ-filtered ring and let  $\Lambda \subseteq \Gamma$  be any subset. With definitions as above, we have:

(1) The category Prefrag<sub>Λ</sub> FR of FR-prefragments is the full additive subcategory of Mod( $\mathcal{F}_{\Lambda}R$ ) given by those  $M \in Mod(\mathcal{F}_{\Lambda}R)$  which satisfy:

for all  $\alpha \leq \beta$  in  $Ob(\mathcal{F}_{\Lambda}R)$ , the map  $M(1_{\alpha,\beta})$ :  $M(\alpha) \to M(\beta)$  is a monomorphism.

We write  $\eta$ : Prefrag<sub>Λ</sub>  $FR \hookrightarrow \text{Mod}(\mathcal{F}_{\Lambda}R)$  for the inclusion Prefrag<sub>Λ</sub>  $FR \subseteq \text{Mod}(\mathcal{F}_{\Lambda}R)$ .

(2) The category Preglid<sub>Λ</sub> FR of FR-pregliders is the full additive subcategory of Mod( $\overline{\mathcal{F}}_n$ R) given by those  $M \in Mod(\overline{\mathcal{F}}_{\Lambda}R)$  which satisfy:

for all  $\alpha < \beta$  in  $Ob(\overline{\mathcal{F}}_{\Lambda}R)$ , the map  $M(1_{\alpha,\beta})$ :  $M(\alpha) \to M(\beta)$  is a monomorphism.

We write  $\iota$ : Preglid<sub>Λ</sub>  $FR \hookrightarrow \text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)$  for the inclusion Preglid<sub>Λ</sub>  $FR \subseteq \text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)$ .

**Remark 3.4.** For an object  $M \in Ob(\text{Mod}\overline{\mathcal{F}}_h R)$  to be a preglider, it suffices to check that, for each  $\alpha \in \Lambda$ , the map  $M(1_{\alpha,\infty})$ :  $M(\alpha) \to M(\infty)$  is a monomorphism.

**Example 3.5.** When  $\Lambda = \Gamma$  and FR is a  $\Gamma^+$ -filtered ring, then Prefrag<sub> $\Lambda$ </sub> FR is equivalent to the category of filtered FR-modules in the sense of [37]. In particular, when  $\Gamma = \mathbb{Z}$ , we recover the usual notion of a Z-filtered module (see, for example, [32]).

**Remark 3.6.** Let  $\Gamma = \mathbb{Z}$  and let  $\Lambda = \mathbb{Z}^{\leq 0}$ . Let FR be a Z-filtered ring. The category Prefrag<sub>A</sub> FR is equivalent to the category of prefragments from definition 2.3. Indeed, given a prefragment

$$
M = M_0 \supseteq M_{-1} \supseteq \cdots \supseteq M_{-i} \supseteq \ldots
$$

in the sense of definition 2.3, we find a functor  $\mathcal{F}_{\Lambda}R \to A\mathbf{b}$  by  $-i \mapsto M_{-i}$ . The fragmented action  $F_iR \times M_{-i} \to M_0$  is given by the action of Hom $_{FR}(-i, 0) = F_iR$ . In particular, the action of  $1_{-i,0}$  on  $M_{-i}$  is given by the inclusion  $M_{-i} \subseteq M_0$ .

**Example 3.7.** Let  $\Gamma$  be a partially ordered group and  $\Lambda \subseteq \Gamma$ . Let FR be a  $\Gamma$ -filtered ring. For each  $\lambda \in \Lambda$ , the representable functor  $\mathcal{F}_{\Lambda}(\lambda, -)$  is a prefragment and the representable functor  $\mathcal{F}_{\Lambda}(\lambda, -)$  is a preglider. Indeed, the conditions in definition 3.3 are easy to verify in this case.

3.3. The category of glider representations. Let  $(\Gamma, \leq)$  be an ordered group and let FR be a Γfiltered ring. Let  $\Lambda \subseteq \Gamma$  be any subset. Consider the set  $\Sigma \subseteq \text{Mor}(\text{Preglid}_{\Lambda} FR)$  given by:

 $(f: N \to M) \in \Sigma \Leftrightarrow$  for all  $\lambda \in \Lambda$ , the map  $f_{\lambda}: N(\lambda) \to M(\Lambda)$  is an isomorphism.

Note that we do not require  $f_{\infty} : M(\infty) \to N(\infty)$  to be an isomorphism.

**Definition 3.8.** The category Glid<sub>Λ</sub> FR is defined as the localization of the category Preglid<sub>Λ</sub> FR with respect to  $\Sigma$ , i.e.  $\text{Glid}_{\Lambda} FR \coloneqq (\text{Preglid}_{\Lambda} FR)[\Sigma^{-1}].$ 

Notation 3.9. We denote the localization functor by  $Q: \text{Preglid}_{\Lambda} FR \to \text{Glid}_{\Lambda} FR$ .

We will now work to understand this localization better: we will show that  $\Sigma$  is a saturated right multiplicative system, so that morphisms in Glid<sub>Λ</sub> FR are described by "right roofs" as in description 2.13. We start with the following lemma.

**Lemma 3.10.** *Let*  $f: M \to N$  *be a morphism in* Mod  $\overline{\mathcal{F}}_{\Lambda}R$ *. Assume that* ker  $f_{\lambda} = 0$ *, for all*  $\lambda \in \Lambda$ *. If*  $N \in \text{Preglid}_{\Lambda} FR$ , then  $M \in \text{Preglid}_{\Lambda} FR$ .

*Proof.* As in remark 3.4, it suffices to show that, for each  $\lambda \in \Lambda$ , the map  $M(1_{\lambda,\infty})$ :  $M(\lambda) \to M(\infty)$  is a monomorphism. Consider the commutative diagram

$$
M(\lambda) \xrightarrow{M(1_{\lambda,\infty})} M(\infty)
$$
  

$$
\downarrow^{f_{\lambda}} \qquad \qquad \downarrow^{f_{\infty}}
$$
  

$$
N(\lambda) \xrightarrow{N(1_{\lambda,\infty})} N(\infty)
$$

given by the morphism  $f: M \to N$ . As  $f_{\lambda}$  is a monomorphism by assumption and N is a preglider, we know that the left-lower branch composes to a monomorphism. We now see that  $M(1_{\lambda,\infty})$ :  $M(\lambda) \to$  $M(\infty)$  is a monomorphism as well.

**Remark 3.11.** Note that every  $s \in \Sigma$  satisfies the conditions in lemma 3.10.

**Proposition 3.12.** *The set*  $\Sigma \subset \text{Mor}(\text{Preglid}_{\Lambda} FR)$  *is a saturated right multiplicative system.* 

*Proof.* We let  $\Theta = \{f \in \text{Mor}(\text{Mod }\overline{\mathcal{F}}_{\Lambda}R) \mid \forall \lambda \in \Lambda : \text{ker } f_{\lambda} = 0 = \text{coker } f_{\lambda}\},\$  thus a morphism  $f: M \to N$ is in Θ if and only if the restriction  $(f)_{\mathcal{F}_{\Lambda}R} = f \circ j: M \circ j \to N \circ j$  is invertible. We know that  $(\text{Mod}\,\overline{\mathcal{F}}_{\Lambda}R)[\Theta^{-1}] \cong \text{Mod}\,\mathcal{F}_{\Lambda}R$  and that  $\Theta$  is a saturated right (as well as a left) multiplicative system in  $Mod\overline{\mathcal{F}}_A R$ .

Note that  $\Sigma = \Theta \cap \text{Mor}(\text{Preglid}_{\Lambda} FR)$ . Using lemma 3.10, it is straightforward to show that  $\Sigma$  is a saturated right multiplicative system.

Corollary 3.13. *The category* Glid<sub>Λ</sub> FR *is an additive category.* 

*Proof.* This follows directly from proposition 3.12 (see [21, corollary I.3.3]).

**Remark 3.14.** In §5.2, we obtain a different proof of proposition 3.12 by interpreting the category of glider representations as the quotient  $\text{Preglid}_{\Lambda} FR/i_*(\text{Mod } R)$ .

**Example 3.15.** When  $\Gamma = \Lambda = \mathbb{Z}$ , then Prefrag<sub> $\Lambda$ </sub> FR is the usual category of filtered modules over the filtered ring FR. The objects in Prefrag<sub>Λ</sub> FR are given by Z-filtered modules  $\{M_i\}_{i\in\mathbb{Z}}$ ; the objects in Preglid<sub>Λ</sub> FR are given by Z-filtered modules  $\{M_i\}_{i\in\mathbb{Z}}$ , together with an R-module M and a monomorphism  $\lim_{i\in\mathbb{Z}}M_i\to M$ . The category Glid<sub>Λ</sub> FR of glider representations is equivalent to the category Prefrag<sub>A</sub>  $\widetilde{FR}$  of prefragments, which is itself equivalent to the usual category of Z-filtered  $FR$ -modules.

**Example 3.16.** Let  $\Gamma = \mathbb{Z}$  and  $\Lambda = \{0\}$ . Let  $R = k[t]$  for a commutative ring k, filtered in the usual way (with deg  $t = 1$ ). We consider the following pregliders  $M, N \in \text{Preglid}_{\Lambda} FR$ :

$$
M(i) = \begin{cases} k & i = 0 \\ k[t]/(t) & i = \infty \end{cases}
$$
 
$$
N(i) = \begin{cases} k & i = 0 \\ k[t]/(t-1) & i = \infty \end{cases}
$$

Note that  $\text{Hom}_{\text{Preglid}_{\Lambda}FR}(M, N) = 0$  while  $\text{Hom}_{\text{Prefrag}_{\Lambda}}(M \circ j, N \circ j) \cong k$  (here,  $j: \mathcal{F}_{\Lambda}R \to \overline{\mathcal{F}}_{\Lambda}R$  is the inclusion). Indeed, as  $\Lambda = \{0\}$ , we find that Prefrag<sub> $\Lambda$ </sub> FR  $\simeq$  Mod k and M ∘ j  $\simeq$  N ∘ j are simple in Prefrag<sub>Λ</sub> FR. In Glid<sub>Λ</sub> FR, an isomorphism  $Q(M) \to Q(N)$  is given by a roof:



4. Glider representations as prefragments and pregliders

Let Γ be an ordered group and let  $\Lambda \subseteq \Gamma$  be any subset. Let FR be a Γ-filtered ring. Having defined the categories in figure 1, we now turn our attention to the remaining functors in the same diagram. Our main goals are to introduce the (fully faithful) functors L and  $\phi$ , allowing us to interpret the category of glider representations as full subcategories of Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR, respectively (see propositions 4.8 and 4.9 below).

We start with completing the localization sequence in the top row of figure 1 to a recollement of abelian categories (proposition 4.1 below).

4.1. A recollement. We first consider the top row of figure 1:

$$
\operatorname{Mod} R \xrightarrow[i^*]{i^*} \operatorname{Mod} \overline{\mathcal{F}}_{\Lambda} R \xleftarrow[j^*]{j_!} \operatorname{Mod} \mathcal{F}_{\Lambda} R
$$

- The functor  $j^*$ :  $Mod(\overline{\mathcal{F}}_{\Lambda}R) \to Mod(\mathcal{F}_{\Lambda}R)$  is the restriction functor induced by the embedding  $\mathcal{F}_{\Lambda}R \subseteq \overline{\mathcal{F}}_{\Lambda}R.$
- The functor  $j_! \colon \text{Mod}(\mathcal{F}_{\Lambda}R) \to \text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)$  is the left adjoint of the restriction functor  $j^*$ . Explicitly,  $j_!$  is the right exact functor  $\mathcal{F}_{\Lambda}R\otimes_{\mathcal{F}_{\Lambda}R}$  — which maps the representable functor  $\text{Hom}_{\mathcal{F}_{\Lambda}R}(\lambda,-)$ to the representable functor  $\text{Hom}_{\overline{\mathcal{F}}_{\Lambda}R}(\lambda, -)$ , for each  $\lambda \in \Lambda$  (see §2.4).
- The functor  $j_*\colon \text{Mod}(\mathcal{F}_{\Lambda}R) \to \text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)$  is defined by

$$
(j_*M)(\lambda) = \begin{cases} M(\lambda) & \text{if } \lambda \in \Lambda, \\ 0 & \text{if } \lambda = \infty. \end{cases}
$$

It is straightforward to see that  $(j_!, j^*, j_*)$  is an adjoint triple.

• The functor  $i_*\colon \text{Mod}(R) \to \text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)$  is the kernel of  $j^*$ . Explicitly,  $i_*$  is given by

$$
(i_*M)(\lambda) = \begin{cases} 0 & \text{if } \lambda \in \Lambda, \\ M & \text{if } \lambda = \infty. \end{cases}
$$

- The functor  $i^! \colon Mod(\overline{\mathcal{F}}_{\Lambda}R) \to Mod(R)$  is defined by  $i^!(M) = M(\infty)$ . Equivalently,  $i^!(M) =$  $\ker(\mu_M)$  where  $\mu: Id_{\text{Mod}(\overline{\mathcal{F}}_A R)} \Rightarrow j_*j^*$  is the unit of the adjunction  $j^* \dashv j_*$ .
- The functor  $i^* \colon \text{Mod}(\overline{\mathcal{F}}_{\Lambda}R) \to \text{Mod}(R)$  is defined by  $i^*(M) = M(\infty)/j_!j^*(M)$ , thus  $i^*(M) =$  $\operatorname{coker}(\varepsilon_M)$  where  $\varepsilon: j_!j^* \Rightarrow Id_{\text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)}$  is the counit of the adjunction  $j_! \dashv j^*$ .

Again, the triple  $(i^*, i_*, i^!)$  is an adjoint triple.

Proposition 4.1. *The diagram*

$$
\operatorname{Mod} R \xrightarrow[i^*]{i^*} \operatorname{Mod} \overline{\mathcal{F}}_{\Lambda} R \xrightarrow[j^*]{j_!} \operatorname{Mod} \mathcal{F}_{\Lambda} R
$$

*is a recollement of abelian categories.*

*Proof.* Note that  $j_*$  is fully faithful. The result now follows from [33, remark 2.3 and example 2.13].  $\Box$ 

4.2. Pregliders and prefragments as torsion-free classes. Recall from definition 3.3 that we write η: Prefrag<sub>Λ</sub>  $FR \hookrightarrow \text{Mod}(\mathcal{F}_{\Lambda}R)$  and  $\iota$ : Preglid<sub>Λ</sub>  $FR \hookrightarrow \text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)$  for the inclusion functors.

**Proposition 4.2.** *(1) The category*  $\text{Preglid}_{\Lambda} FR$  *is a cotilting torsion-free subcategory of*  $\text{Mod}(\mathcal{F}_{\Lambda} R)$ . *(2) The category* Prefrag<sub>Λ</sub> FR *is a cotilting torsion-free subcategory of* Mod( $\overline{\mathcal{F}}_$ <sup>*NR*</sup>).

*Proof.* We only show the first statement, the second statement is analogous. Note that Preglid<sub>Λ</sub> FR is closed under subobjects, direct products and extensions in  $Mod(\overline{\mathcal{F}}_{\Lambda}R)$ . By [39, example V.2.2], the category Mod( $\overline{\mathcal{F}}_{\Lambda}R$ ) is a Grothendieck category. It now follows from [18, theorem 2.3] that Preglid<sub>Λ</sub> FR is a torsion-free class. Note that the standard projectives in  $Mod(\overline{\mathcal{F}}_{\Lambda}R)$  are in Preglid<sub>Λ</sub> FR (see example 3.7). As every projective  $\overline{\mathcal{F}}_{\Lambda}R$ -module is a direct summand of a (possibly infinite) direct sum of standard projectives, we find that all projective  $\overline{\mathcal{F}}_{{\Lambda}}R$ -modules are in Preglid<sub>Λ</sub> FR. Hence, Preglid<sub>Λ</sub> FR is a cotilting torsion-free class.  $\hfill \square$ 

By proposition 4.2, the functor  $\eta$  has a left adjoint  $\theta$  and the functor  $\iota$  has a left adjoint  $\kappa$ . We give an explicit description of the functor  $\kappa$ : Mod $(\overline{\mathcal{F}}_{\Lambda}R) \to \text{Preglid}_{\Lambda}FR$ .

**Proposition 4.3.** *The left adjoint functor*  $\kappa$ :  $\text{Mod}(\overline{\mathcal{F}}_{\Lambda}R) \to \text{Preglid}_{\Lambda}FR$  *is given by* 

 $\kappa(M): \lambda \mapsto \text{im} (M(1_{\lambda,\infty}))$ .

*Proof.* Note that  $\kappa(M) \in \text{Preglid}_{\Lambda} FR$ , for all  $M \in \text{Mod }\overline{\mathcal{F}}_{\Lambda} R$ . There is a natural map  $M \to \iota(\kappa(M))$ . It is now easy to verify that a morphism  $M \to \iota(N)$  (for any  $N \in \text{Preglid}_{\Lambda} FR$ ) factors uniquely as  $M \to \iota(\kappa(M)) \to \iota(N)$ . This shows that  $\kappa$  is left adjoint to  $\iota$ .

**Proposition 4.4.** *The functor*  $j^*$ : Mod  $\overline{\mathcal{F}}_{\Lambda}R \to \text{Mod }\mathcal{F}_{\Lambda}R$  *restricts to*  $j^*$ : Preglid<sub> $\Lambda$ </sub>  $FR \to \text{Prefix}_{\Lambda}FR$ . *Proof.* Straightforward. □

**Corollary 4.5.** The composition  $\kappa \circ j_!$ : Mod  $\mathcal{F}_{\Lambda}R \to \text{Preglid}_{\Lambda}FR$  is left adjoint to the composition  $\eta \circ j^* \colon \operatorname{Preglid}_{\Lambda} FR \to \operatorname{Mod} \mathcal{F}_{\Lambda} R.$ 

*Proof.* This follows directly from  $\eta \circ j^* \cong j$  $\overline{\phantom{a}}$   $\circ$   $\iota$ .

**Proposition 4.6.** *The functor*  $j^*$ :  $\text{Preglid}_{\Lambda} FR \to \text{Prefrag}_{\Lambda} FR$  *has a left adjoint*  $j_L = \kappa \circ j_l \circ \eta$ .

*Proof.* This follows from proposition 2.11(2a) with  $H = \kappa \circ j_1$ ,  $G = \eta$ , and  $F = j^*$ : Preglid<sub>A</sub>  $FR \rightarrow$  $\Gamma$ refrag $\Lambda$  FR.

4.3. Glid<sub>Λ</sub> FR as subcategory of Prefrag<sub>Λ</sub> FR. The universal property of the localization functor shows  $\operatorname{that} j^*\colon \operatorname{Preglid}_{\Lambda} FR \to \operatorname{Prefrag}_{\Lambda} FR$  factors as  $\operatorname{Preglid}_{\Lambda} FR \stackrel{Q}{\to} \operatorname{Glid}_{\Lambda} FR \stackrel{\phi}{\to} \operatorname{Prefrag}_{\Lambda} FR.$  We now show that  $\phi$  is fully faithful, so that Glid<sub>Λ</sub> FR can be interpreted as a full subcategory of Prefrag<sub>Λ</sub> FR.

**Remark 4.7.** The functor  $\phi$ : Glid<sub>Λ</sub>  $FR \rightarrow$  Prefrag<sub>Λ</sub>  $FR$  is given on objects by mapping a glider  $M \in$  $Ob(Glid_{\Lambda} FR) = Ob(Preglid_{\Lambda} FR)$  to the restriction  $j^*(M) = M \circ j$ .

Let  $f: M \to N$  be a morphism in Glid<sub>Λ</sub> FR, say that f is represented by the roof  $M \stackrel{s}{\leftarrow} M' \stackrel{g}{\to} N$  in Preglid<sub>Λ</sub> FR (with  $s \in \Sigma$ ). The corresponding morphism in  $\phi(f)$ :  $\phi(M) \to \phi(N)$  is given by  $\phi(f)(\lambda) =$  $g(\lambda) \circ s^{-1}(\lambda) \colon M(\lambda) \to N(\lambda)$ , for all  $\lambda \in \Lambda$ .

**Proposition 4.8.** *The functor*  $\phi$ : Glid<sub>Λ</sub>  $FR \rightarrow$  Prefrag<sub>Λ</sub>  $FR$  *is fully faithful and admits a left adjoint*  $\psi$ : Prefrag<sub>A</sub>  $FR \rightarrow$  Glid<sub>A</sub>  $FR$  given by  $\psi = Q \circ j_L$ .

*Proof.* To see that  $\phi$  is fully faithful, it suffices to prove that  $\eta \circ \phi$  is fully faithful. Recall that Mod  $\mathcal{F}_\Lambda R \simeq$  $\Theta^{-1} \text{Mod}\overline{\mathcal{F}}_{\Lambda}R$  where  $\Theta = \{f \in \text{Mor}(\text{Mod}\overline{\mathcal{F}}_{\Lambda}R) \mid j^*(f) \text{ is invertible}\}\$  (as in proposition 3.12). Using this equivalence to identify these categories, we can describe the map  $\eta \circ \phi$ : Glid<sub>A</sub>  $FR \to \Theta^{-1}$  Mod  $\overline{\mathcal{F}}_{\Lambda}R$  as mapping a roof  $M \xleftarrow{s} M' \to N$  (with  $s \in \Sigma$ ) in  $\Sigma^{-1}$  Preglid<sub>A</sub> FR to the same roof in  $\Theta^{-1}$  Mod  $\overline{\mathcal{F}}_{\Lambda}R$  (note that  $\Sigma = \Theta \cap \text{Mor}(\text{Preglid}_{\Lambda} FR)$ . Using lemma 3.10, it is straightforward to show that  $\eta \circ \phi$  is fully faithful.

It is shown in proposition 4.6 that  $j_L$  is left adjoint to  $j^* = \phi \circ Q$ . It now follows from proposition 2.11(1b) that  $\psi = Q \circ j_L$  is left adjoint to  $\phi$ .

4.4. Glid<sub>Λ</sub> FR as subcategory of Preglid<sub>Λ</sub> FR. In this subsection, we show that the localization Q: Preglid<sub>Λ</sub>  $FR \rightarrow \text{Glid}_{\Lambda} FR$  has a fully faithful left adjoint L. In this way,  $\text{Glid}_{\Lambda} FR$  can be interpreted as a coreflective subcategory of  $\text{Preglid}_{\Lambda} FR$ .

**Proposition 4.9.** *The localization functor* Q: Preglid<sub>Λ</sub>  $FR \rightarrow$  Glid<sub>Λ</sub>  $FR$  *admits a fully faithful left adjoint*  $L: \text{Glid}_{\Lambda} FR \to \text{Preglid}_{\Lambda} FR.$ 

*Proof.* Recall from proposition 4.6 that  $j_L$  is left adjoint to  $j^* = \phi \circ Q$ . It follows from proposition 2.11(2a) (with  $F = Q$ ,  $G = \phi$ , and  $H = j_L$ ) that  $L = j_L \circ \phi$  is left adjoint to Q. This uses that  $\phi$  is fully faithful, see proposition 4.8. The dual of proposition 2.10 implies that L is fully faithful.

**Corollary 4.10.** *The category*  $\text{Glid}_{\Lambda} FR$  *is a coreflective subcategory of*  $\text{Preglid}_{\Lambda} FR$ .

*Proof.* The embedding L: Glid<sub>A</sub>  $FR \rightarrow$  Preglid<sub>A</sub>  $FR$  has a right adjoint.

4.5. A criterion for gliders. We will now give a criterion to determine whether a given prefragment  $M \in \text{Prefix}_{\Delta} FR$  is a glider representation, i.e. whether M lies in the essential image of  $\phi$ : Glid<sub>A</sub>  $FR \rightarrow$  $Prefrag<sub>Λ</sub> FR.$ 

**Proposition 4.11.** *The following are equivalent for an*  $M \in \text{Prefix}_{\Delta} FR$ :

- *(1)* M *is a glider representation,*
- *(2) the counit of the adjunction*  $\phi \circ \psi(M) \to M$  *is an isomorphism,*
- (3) the counit of the adjunction  $j^* \circ j_L(M) \to M$  is an isomorphism,
- *(4)* j(o η(M) is a preglider (i.e. j(o η(M) lies in the essential image of  $\iota$ : Preglid<sub>A</sub>  $FR \to \text{Mod }\overline{\mathcal{F}}_{\Lambda}R$ ).

*Proof.* Note that M is a glider if and only if it lies in the essential image of  $\phi$ , or equivalently, in the essential image of  $j^*$ : Preglid<sub>A</sub>  $FR \to$  Prefrag<sub>A</sub>  $FR$ . The equivalence of the first three statements is now easy.

Assume now that (3) holds, so  $M \cong j^*(\bar{M})$  where  $\bar{M} = j_L(M) \in \text{Preglid}_{\Lambda} FR$ . We find  $j_{!}\eta(M) \cong$  $j_!\eta j^*(\bar{M}) \cong j_!\eta j^*(\bar{M})$ . As  $\iota(M)$  is a preglider, the counit  $j_!\eta j^*(\bar{M}) \to \iota(\bar{M})$  satisfies the conditions of lemma 3.10. Hence,  $j_!\eta(M)$  is a preglider.

Assume now that (4) holds, so  $j_! \eta(M) \cong \iota(M)$ , for some  $\overline{M} \in \text{Preglid}_{\Lambda} FR$ . Using that  $j^* \circ j_! \cong 1$ , we find:  $\eta(M) \cong j^* \circ j_! \circ \eta(M) \cong j^* \circ \iota(M) \cong \eta \circ j^*(\bar{M})$ . As  $\eta$  is fully faithful, we find  $M \cong j^*(\bar{M})$ , hence M is a glider representation.

We now revisit the example in the introduction.

**Example 4.12.** Let  $R = \mathbb{R}[t]$  and let FR be the usual Z-filtered ring with t is degree 1. Let  $\Lambda = \mathbb{Z}^{\leq 0}$ and let  $M \in Mod \mathcal{F}_{\Lambda}R$  be the functor given by  $-i \mapsto C^i(\mathbb{R})$  and  $t^n : C^i(\mathbb{R}) \to C^{i-n}(\mathbb{R}) : f \mapsto \frac{d^n}{dx^n}f$ . This is clearly a prefragment. To see that this is a glider representation, we consider the  $\mathbb{R}[t]$ -module

$$
\Omega = \oplus_{i \geq 0} C^0(\mathbb{R}) \cdot t^i.
$$

Let N be the submodule of  $\Omega$  generated by the elements  $f \cdot t^i - (\frac{d}{dx} f)t^{i-1}$  where  $f \in C^1(\mathbb{R}), i \geq 1$ . We can extend M to a preglider  $\bar{M}$  by setting  $\bar{M}(\infty) = \Omega/N$ . It is now clear that the restriction  $M = j^*(\bar{M})$ is a glider representation.

#### 5. Properties of the category of glider representations

Let Γ be an ordered group and let  $\Lambda \subseteq \Gamma$  be any subset. Let FR be a Γ-filtered ring. In this section, we establish some (homological) properties of the category of glider representations. We start by showing that Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR are complete and cocomplete. The category Glid<sub>Λ</sub> FR inherits these properties as it is a reflective subcategory of Prefrag<sub>Λ</sub> FR (via the functor  $\phi$ ) and a coreflective subcategory of Preglid<sub>Λ</sub>  $FR$  (via the functor L).

In §5.2, we endow the category Glid<sub>Λ</sub> FR with the structure of a deflation-exact category, induced by the natural exact structures on Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR. We subsequently show that the conflation

structure on Glid<sub>Λ</sub> FR is an inherent feature (i.e. it can be defined without referring to Preglid<sub>Λ</sub> FR or Prefrag<sub>Λ</sub> FR) by showing that Glid<sub>Λ</sub> FR is a (Grothendieck) deflation quasi-abelian category.

5.1. Limits and colimits in Glid<sub>Λ</sub> FR. We will describe limits and colimits in Glid<sub>Λ</sub> FR using the embeddings L and φ into the categories  $Preglid_A FR$  and  $Prefrag_A FR$ , respectively (see §4).

**Proposition 5.1.** *The categories* Preglid<sub>Λ</sub> FR *and* Prefrag<sub>Λ</sub> FR *are complete and cocomplete quasiabelian categories.*

*Proof.* It follows from theorem 2.30 and proposition 4.2 that Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR are quasiabelian. It follows directly from the definitions that  $Preglid<sub>Λ</sub> FR$  and  $Prefrag<sub>Λ</sub> FR$  are closed under arbitrary direct sums and products. We conclude that  $Preglid_\Lambda FR$  and  $Prefrag_\Lambda FR$  are complete and cocomplete.

**Remark 5.2.** Limits in Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR coincide with limits in Mod  $\overline{\mathcal{F}}_0R$  and Mod  $\mathcal{F}_0R$ , respectively; colimits, however, need not coincide.

**Proposition 5.3.** *The category*  $\text{Glid}_{\Lambda} FR$  *is complete and cocomplete. Specifically, let*  $D: J \to \text{Glid}_{\Lambda} FR$ *be a diagram.*

- *(1)* We have  $\lim_{D \to \infty} D \cong \psi(\lim_{D \to \infty} \phi D)$  *and* colim  $D \cong \psi(\text{colim } \phi D)$ .
- *(2)* We have  $\lim_{D \to \infty} D \cong Q(\lim(LD))$  *and* colim  $D \cong Q(\text{colim}(LD)).$

*Proof.* We have shown in proposition 5.1 that  $\text{Preglid}_{\Lambda} FR$  and  $\text{Prefrag}_{\Lambda} FR$  are complete and cocomplete. As Glid<sub>Λ</sub> FR is a reflective subcategory of Prefrag<sub>Λ</sub> FR (proposition 4.8) and a coreflective subcategory of Preglid<sub>Λ</sub> FR (corollary 4.10), the other statements follow.

Remark 5.4. As Glid<sub>Λ</sub> FR is a reflective subcategory of Prefrag<sub>Λ</sub> FR via  $\phi$ , limits in Glid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR coincide. Similarly, Glid<sub>Λ</sub> FR is a coreflective subcategory of Preglid<sub>Λ</sub> FR via L, and, as such, colimits in Glid<sub>Λ</sub> FR and Preglid<sub>Λ</sub> FR coincide.

5.2. Glid<sub>Λ</sub> FR is a deflation-exact category. The category Glid<sub>Λ</sub> FR naturally factors the map  $j^*$ : Preglid<sub>Λ</sub>  $FR \stackrel{Q}{\rightarrow}$  Glid<sub>Λ</sub>  $FR \stackrel{\phi}{\rightarrow}$  Prefrag<sub>Λ</sub>  $FR$ . As such, it can be endowed with a conflation structure, either as a localization of Preglid<sub>Λ</sub> FR or as a subcategory of Prefrag<sub>Λ</sub> FR. In this subsection, we start with the former (corollary 5.6) and show that it coincides with the latter (proposition 5.8).

In definition 3.8, we introduced the category of gliders via a localization functor  $Q$ : Preglid<sub>A</sub>  $FR \rightarrow$  $\Sigma^{-1}$  Preglid<sub>Λ</sub>  $FR = \text{Glid}_{\Lambda} FR$ . We will now interpret  $\text{Glid}_{\Lambda} FR$  as the quotient  $\text{Preglid}_{\Lambda} FR/i_*(\text{Mod } R)$ , as defined in §2.6.

**Proposition 5.5.** *Let*  $(\Gamma, \leq)$  *be an ordered group and let* FR *be a*  $\Gamma$ -filtered ring. Let  $\Lambda \subseteq \Gamma$  *be any subset.*

- *(1) With the embedding* i∗*, the category* Mod(R) *is an admissibly deflation-percolating subcategory of* Preglid<sup>Λ</sup> F R*.*
- *(2)* A morphism  $f: M \to N$  in Preglid<sub>A</sub> FR is a weak Mod(R)-isomorphism if and only if  $f \in \Sigma$ *(i.e. the induced maps*  $f_{\lambda}$ :  $M(\lambda) \rightarrow N(\lambda)$  *are isomorphisms for each*  $\lambda \neq \infty$ *).*
- *(3)* Glid<sub>Λ</sub>  $FR = \Sigma^{-1}$  Preglid<sub>Λ</sub>  $FR =$  Preglid<sub>Λ</sub>  $FR/i_*(Mod R)$ .

*Proof.* As Preglid<sub>Λ</sub> FR is an extension-closed subcategory of the abelian category Mod  $\overline{\mathcal{F}}_$ ΛR (see proposition 4.2), we find that Preglid<sub>A</sub> FR is an exact category (see [8, lemma 10.20]): the conflations of Preglid<sub>Λ</sub> FR are given by the short exact sequences in Mod  $\overline{\mathcal{F}}_$ <sup>R</sup>.

We now verify axioms A1, A2 and A3 of definition 2.23. As the category Preglid<sub>Λ</sub> FR is an exact category, axiom A3 is automatic (see remark 2.24). Axiom A1 is straightforward to see. For axiom **A2**, consider a map  $f: M \to A$  where  $M \in \text{Preglid}_{\Lambda} FR$  and  $A \in i^*(\text{Mod } R) \subseteq \text{Preglid}_{\Lambda} FR$ . Clearly  $f_{\lambda}: M(\lambda) \to 0$  for each  $\lambda \in \Lambda$ . The map  $f_{\infty}: M(\infty) \to A$  has an epi-mono factorization as  $Mod(R)$ is abelian. It readily follows that f is admissible in Preglid<sub>Λ</sub> FR. Hence, Mod(R) is an admissibly deflation-percolating subcategory of Preglid<sub>Λ</sub>  $FR$ .

For the next statement, let  $f: M \to N$  be a morphism in Preglid<sub>Λ</sub> FR. Assume that f is a weak  $Mod(R)$ -isomorphism. As f is admissible, we find that  $ker(f), coker(f) \in Mod(R)$ . It follows that  $\ker(f_\lambda) = 0 = \operatorname{coker}(f_\lambda)$  for each  $\lambda \neq \infty$ . Hence, each  $f_\lambda$  is an isomorphism. Conversely, assume that  $f_\lambda$ is an isomorphism for each  $\lambda \neq \infty$ . Clearly, ker(f) and coker(f) exist, and belong to Mod(R). The fact that f is admissible follows immediately from the fact that  $f_{\infty}$  admits an epi-mono factorization.

For the last statement, the first equality is the definition of  $\mathrm{Glid}_{\Lambda} FR$ , the second equality is the definition of  $\text{Preglid}_{\Lambda} FR/i_*(\text{Mod } R)$ .

It follows from proposition 5.5 that the localization  $\Sigma^{-1}$  Preglid<sub>Λ</sub> FR can be described using theorem 2.27.

Corollary 5.6. *The weakest conflation structure on* Glid<sub>Λ</sub>  $FR := \Sigma^{-1}$  Preglid<sub>Λ</sub>  $FR$  *for which the localization functor* Q: Preglid<sub>Λ</sub>  $FR \rightarrow$  Glid<sub>Λ</sub>  $FR$  *is conflation-exact, gives* Glid<sub>Λ</sub>  $FR$  *the structure of a deflation-exact category.*

**Remark 5.7.** Theorem 2.27 also implies that  $\Sigma$  is a saturated right multiplicative system, i.e. we recover proposition 3.12.

The following proposition connects the conflation structures of Glid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR using the embedding  $\phi$ .

**Proposition 5.8.** *The embedding*  $\phi$ : Glid<sub>Λ</sub>  $FR \rightarrow$  Prefrag<sub>Λ</sub>  $FR$  *reflects conflations, i.e. a sequence*  $K \xrightarrow{f} M \xrightarrow{g} N$  in Glid<sub>A</sub> FR is a conflation if and only if  $\phi(K) \xrightarrow{\phi(f)} \phi(M) \xrightarrow{\phi(g)} \phi(N)$  is a conflation in Prefrag<sub>Λ</sub> FR.

*Proof.* By construction,  $\phi$  is conflation-exact and hence maps conflations to conflations. Thus, let  $K \stackrel{f}{\to}$  $M \stackrel{g}{\to} N$  be a sequence in Glid<sub>A</sub> FR which maps to a conflation in Prefrag<sub>A</sub> FR. As  $\phi$  is fully faithful, we know that  $K \to M \to N$  is a kernel-cokernel pair in Glid<sub>Λ</sub> FR.

Using that L is a left adjoint, we find that  $L(g): L(M) \to L(N)$  is the cokernel of  $L(f)$ . Since Preglid<sub>Λ</sub> FR is a quasi-abelian category,  $L(g)$  is a deflation. Hence, so is  $g = Q \circ L(g)$ . This establishes that  $K \rightarrow M \rightarrow N$  is a conflation in Glid<sub>Λ</sub> FR.

**Proposition 5.9.** *For all*  $M \in \text{Prefix}_{\Delta} FR$ *, the reflection*  $M \to \phi \circ \psi(M)$  *is a deflation.* 

*Proof.* We write  $\Omega_M$  for the R-module  $j_! \circ \eta(M)(\infty)$ . As  $\psi = j_L \circ Q = \eta \circ j_! \circ \kappa \circ Q$ , we find  $\phi \circ \psi(M)(\lambda) \cong$  $im(M(1_{\lambda,\infty}): M(\lambda) \to \Omega_M)$ , and the reflection is given by maps  $r_M: M \to \phi \circ \psi(M)$  for which the following diagram commutes



It follows that each  $r_M$  is an epimorphism. As such the map  $r_M$  is an epimorphism in Mod  $\overline{\mathcal{F}}_h R$  and hence ker  $r_M \rightarrow M \rightarrow \phi \circ \psi(M)$  is a conflation.

**Remark 5.10.** Via the functor  $\phi$ , the category Glid<sub>Λ</sub> FR is an *epi-reflective* subcategory of Prefrag<sub>Λ</sub> FR, i.e. the fully faithful embedding has a left adjoint, and the unit  $1 \to \phi \circ \psi$  of the adjunction is a deflation. Hence, every prefragment  $M \in \text{Prefix}_{\Delta} FR$  has a largest glider quotient object  $\phi \circ \psi(M)$ .

**Corollary 5.11.** *The essential image of*  $\phi$  *is closed under subobjects.* 

*Proof.* Let  $M \in \text{Glid}_{\Lambda} FR$  and let  $f: N \hookrightarrow \phi M$  be a monomorphism in Prefrag<sub> $\Lambda$ </sub>FR. We find the following commutative diagram



where  $r: 1 \to \phi \circ \psi$  is the unit of the adjunction. As the top-right branch composes to a monomorphism, so does the left-lower branch. In particular,  $r_N: N \to \phi \psi N$  is a monomorphism. As  $r_N$  is a deflation (see proposition 5.9), we find that  $r_N$  is an isomorphism. Hence, N lies in the essential image of  $\phi$ .

5.3. Gliders as a Grothendieck deflation quasi-abelian category. We now proceed to showing that Glid<sub>Λ</sub> FR is a Grothendieck deflation quasi-abelian category (see definition 5.14 below). We start by showing that  $G \text{lid}_{\Lambda} FR$  is deflation quasi-abelian (see definition 2.21).

**Theorem 5.12.** *The category*  $\text{Glid}_{\Lambda} FR$  *is a complete and cocomplete deflation quasi-abelian category.* 

*Proof.* We have already established that Glid<sub>Λ</sub> FR is deflation-exact (see corollary 5.6), and is complete and cocomplete (see proposition 5.3). In particular, Glid<sub>Λ</sub> FR is pre-abelian. To see that Glid<sub>Λ</sub> FR is deflation quasi-abelian, we only need to verify that the conflation structure of Glid<sub>Λ</sub> FR consists of all

kernel-cokernel pairs. For this, it suffices to show that all cokernels are deflations. Let  $f: X \to Y$  be any map; we claim that  $q: Y \to \text{coker } f$  is a deflation. Applying the functor L and using that a left adjoint commutes with cokernels, we find that  $L(q)$  is the cokernel of  $L(f)$ . As Preglid<sub>Λ</sub> FR is a quasi-abelian category, this means that  $L(g)$  is a deflation. Finally, as Q maps deflations to deflations, we find that  $QL(q) \cong q$  is a deflation in Glid<sub>Λ</sub> FR.

**Remark 5.13.** It follows from theorem 5.12 that the conflation structure of Glid<sub>Λ</sub> FR is not additional structure but is inherent to the category: the class of conflations consists of all kernel-cokernel pairs.

We recall the definition of a Grothendieck quasi-abelian category from [40].

Let C be a category admitting all small filtered direct limits. We say that an object C is *finitely presentable* if, for all filtered diagrams  $D: J \to \mathcal{C}$ , the natural morphism  $\lim_{\epsilon \to 0} \lim_{\epsilon \to 0} C(D) \to \lim_{\epsilon \to 0} C(\lim_{\epsilon \to 0} D)$  $\mu$ -*sentable* it, for an intered diagrams  $D: J \to C$ , the natural morphism  $\lim_{\Delta t \to 0} \mu$  is  $\lim_{\Delta t \to 0} (\sigma, D)$  is an isomorphism. We say the category C is *locally presentable* if it has all small filtered direct limits and every object in  $\mathcal C$  is a filtered direct limit of finitely presentable objects.

Definition 5.14. A (deflation) quasi-abelian category is called a *Grothendieck (deflation) quasi-abelian category* if it is locally presentable and has exact filtered direct limits.

**Proposition 5.15.** *The categories* Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR are Grothendieck quasi-abelian cate*gories.*

*Proof.* It is shown in proposition 5.1 that Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR are cocomplete. As direct limits in Mod  $\overline{\mathcal{F}}_{\Lambda}R$  and Mod  $\overline{\mathcal{F}}_{\Lambda}R$  are exact and taken pointwise, we find that filtered direct limits are exact Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR as well. As the standard projectives in Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR form a strong generating set, it follows from [1, theorem 1.11] that Preglid<sub>Λ</sub> FR and Prefrag<sub>Λ</sub> FR are finitely presentable.

**Theorem 5.16.** Let  $\Gamma$  *be an ordered group.* Let  $\Lambda \subseteq \Gamma$  *be any subset and let* FR *be a*  $\Gamma$ -filtered ring. *The category* Glid<sub>Λ</sub> FR *of glider representations is a Grothendieck deflation quasi-abelian category.* 

*Proof.* Similar to the proof of proposition 5.15.

5.4. Some examples and counterexamples. We provide some examples and counterexamples to illustrate some properties in this section.

By proposition 5.1, the category  $Preglid_A FR$  admits kernels and cokernels. For sake of reference we describe these explicitly in the next proposition (see also [37, corollary 3.6] for a similar result).

**Proposition 5.17.** *Let*  $f: M \to N$  *be a morphism in* Preglid<sub>Λ</sub> FR. For each  $\lambda \in Ob(\overline{\mathcal{F}}_{\Lambda}R)$  *we have the following canonical isomorphisms:*

 $(a)$  (ker  $f$ )( $\lambda$ ) ≅ ker  $f(\lambda)$ ,

*(b)* (coker  $f(\lambda) \cong \text{im}(N(\lambda) \to \text{coker } f_{\infty}),$ 

- $(c)$  (im  $f$ )( $\lambda$ )  $\cong$  ker( $N(\lambda) \to$  coker  $f_{\infty}$ ),
- *(d)* (coim  $f(\lambda) \cong \text{im } f(\lambda)$ .

*Proof.* This follows from the fact that  $\iota$  commutes with kernels and that  $\kappa$  commutes with cokernels.  $\Box$ 

5.4.1. *The functor*  $\phi$  *need not commute with colimits.* In proposition 5.8, we showed that the functor  $\phi$ reflects conflations. However,  $\phi$  need not commute with colimits, as the following example shows. In particular, one needs to be careful in computing colimits using the functor  $\phi$ .

**Example 5.18.** Let  $R = \mathbb{C}[S_3]$  and consider the  $\mathbb{Z}^{\geq 0}$ -filtration

$$
\mathbb{C} \subseteq \mathbb{C}[S_3] \subseteq \mathbb{C}[S_3] \subseteq \dots
$$

and let  $\Lambda = \{-1, 0\}$ . Let V be a two-dimensional complex vector space with basis  $e_a, e_b$  and set  $e_c = -(e_a + e_b)$ . View  $S_3$  as the permutations on the set  $\{a, b, c\}$  and let  $S_3$  act on the vectors  $e_a, e_b, e_c$ accordingly. Thus, V is the standard representation of  $S_3$ . Consider the following conflation of prefragments:



The reader may verify that M and N belong to  $\phi(\text{Glid}_{\Lambda} FR)$ . However, S cannot be extended to a glider representation. Indeed, consider the transposition  $(ab) \in \text{Hom}_{\mathcal{F}_A} R(-1,0) = \mathbb{C}[S_3]$ . As  $q: M \to N$  is a natural transformation, the following diagram commutes:



Hence,  $S(ab)$  acts on  $e<sub>b</sub>$  as zero. It follows that S is not a glider representation as the partial actions are not induced by a  $\mathbb{C}[S_3]$ -module. Indeed, the transposition (ab) squares to the identity and thus cannot act as zero.

Following proposition 5.3, the cokernel of  $f$  as a morphism of glider representations is zero. In other words, the inflation f of prefragments is not an inflation of glider representations (as it is not the kernel of its cokernel).

5.4.2. *The functor* L need not be conflation-exact. The embedding L: Glid<sub>Λ</sub>  $FR \rightarrow$  Preglid<sub>Λ</sub>  $FR$  is right exact (as it is left adjoint to  $Q$ ), but need not commute with limits. The following example shows that L need not even be conflation-exact.

**Example 5.19.** Let  $\Gamma = \mathbb{Z}$ ,  $\Lambda = \{0\}$  and let FR be the  $\mathbb{Z}^{\geq 0}$ -filtration of  $\mathbb{C}[t, x]/(xt)$  given by

$$
\mathbb{C}[t] \subseteq \mathbb{C}[t,x]/(xt) \subseteq \mathbb{C}[t,x]/(xt) \subseteq \dots
$$

So,  $F_0R = \mathbb{C}[t]$ , and  $F_iR = \mathbb{C}[t, x]/(xt)$ , for all  $i \ge 1$ .

Consider the natural morphism  $M \stackrel{g}{\rightarrow} N$  in Preglid<sub>A</sub> FR determined by the solid part of the commutative diagram



Using proposition 5.17, one readily verifies that ker(g)  $\stackrel{f}{\to} M \stackrel{g}{\to} N$  is a conflation in Preglid<sub>A</sub> FR and thus descends to a conflation in  $\operatorname{Glid}_{\Lambda} FR$  as well.

Applying  $L \circ Q$  to f we obtain the following commutative diagram in Preglid<sub>Λ</sub> FR:



Note that multiplication by t is not an injection from  $\mathbb{C}[t, x]/(xt)$  to itself. Hence,  $LQ(\ker g) \to LQ(M)$ is not a monomorphism. This shows that  $L$  does not commute with kernels of deflations. In particular, L is not a conflation-exact functor

**Remark 5.20.** Although the functor  $L$  is not conflation-exact, it maps deflations to deflations (as it maps cokernels to cokernels and Preglid<sub>Λ</sub> FR is quasi-abelian). Thus, given a conflation  $X \rightarrow Y \rightarrow Z$ of glider representations, we obtain a sequence  $LX \to LY \to LZ$  of pregliders. Let K be the kernel of  $LY \rightarrow LZ$ . As the natural map  $LX \rightarrow K$  becomes an isomorphism under the functor Q (this uses that Q is conflation-exact and that  $Q \circ L \cong 1$ , the morphism  $LX \to K$  is a weak isomorphism. So, for every  $\lambda \in \Lambda$ , the map  $LX(\lambda) \to K(\lambda)$  is an isomorphism.

5.4.3. *The category* Glid<sub>Λ</sub> FR need not be (2-sided) exact. We now give two examples to show that Glid<sub>Λ</sub> FR is not a (2-sided) exact structure. In particular, Glid<sub>Λ</sub> FR need not be a quasi-abelian (or abelian) category.

**Example 5.21.** Let  $R = k[t]$  be a polynomial ring in one variable t over some field k. Let  $\Gamma = \mathbb{Z}$  and let  $\Lambda = \{-1, 0\}$ . Let FR be the  $\mathbb{Z}^+$ -filtration

$$
k \subseteq k[t]^{\leq 1} \subseteq k[t]^{\leq 2} \subseteq k[t]^{\leq 3} \subseteq \dots
$$

The following commutative diagram defines morphisms f, g and h in Preglid<sub>Λ</sub> FR.



Clearly f and h are inflations in Preglid<sub>Λ</sub> FR. Note that ker(g), coker(g)  $\in$  Mod(k[t]) and hence g is a weak  $Mod(k[t])$ -isomorphism by proposition 5.5. By definition, the maps  $Q(g \circ f)$  and  $Q(h)$  are inflations in Glid<sub>Λ</sub> FR. We claim that the composition  $Q(h) \circ Q(g \circ f)$  is not an inflation in Glid<sub>Λ</sub> FR. It follows that  $G\text{lid}_{\Lambda} FR$  does not satisfy axiom L1 and hence  $G\text{lid}_{\Lambda} FR$  is not an exact category.

Assume that  $Q(h) \circ Q(g \circ f) = Q(h \circ g \circ f)$  is an inflation in Glid<sub>A</sub> FR. By proposition 5.17, the cokernel of  $hgf$  in Preglid<sub>A</sub> FR is given by  $k \hookrightarrow k \hookrightarrow k$  and ker(coker( $hgf$ )) is given by  $0 \hookrightarrow kt \oplus kt^2 \hookrightarrow tk[t]$ . It follows that  $hqf$  is not the kernel of its cokernel in Preglid<sub>Λ</sub> FR and thus that  $hqf$  is not an inflation in Glid<sub>Λ</sub>  $FR$ .

**Example 5.22.** In this example, we provide another illustration of the failure of axiom L1 in Glid<sub>Λ</sub> FR. Using the same notation as in example 5.18, consider the following morphisms of prefragments:



where  $\Delta: V \to V \oplus V$  is the diagonal map. It is easy to see that f and g can be extended to inflations in Glid<sub>Λ</sub> FR but, as in example 5.18, the composition  $g \circ f$  is not an inflation in Glid<sub>Λ</sub> FR. Hence, the composition of inflations need not be an inflation in  $\text{Glid}_{\Lambda} FR$ .

5.4.4. *The functors*  $j_*, j_!$  *do not restrict to functors*  $\text{Prefix}_{\Lambda} FR \to \text{Prefix}_{\Lambda} FR$ . It is shown in proposition 4.1 that the top row in figure 1 is a recollement. The functors  $j_*, j_!,$  in general, fail to restrict to functors Prefrag<sub>Λ</sub>  $FR \rightarrow$  Preglid<sub>Λ</sub>  $FR$ . The following example is based on [14, example 1.3.3].

**Example 5.23.** Set  $\Gamma = \mathbb{Z}, \Lambda = -\mathbb{N}, R = \mathbb{Q}$ , and set

$$
FR = \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{Q} \subseteq \ldots
$$

Let  $M \in \text{Prefix}_{\Lambda} FR$  be the cokernel of the obvious map  $\text{Hom}_{\mathcal{F}_{\Lambda}R}(0,-) \xrightarrow{\times 2} \text{Hom}_{\mathcal{F}_{\Lambda}R}(0,-)$ , explicitly, M is determined by

$$
M(\lambda) = \begin{cases} 0 & \text{if } \lambda \neq 0, \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \lambda = 0. \end{cases}
$$

Note that  $j_!(M)(\infty) = 0$  and thus  $j_!(M) \notin \text{Preglid}_{\Lambda} FR$ . This shows that  $j_!$  does not restrict to a functor  $\text{Prefix}_{\Lambda} FR \rightarrow \text{Prefix}_{\Lambda} FR.$ 

Note that  $j_*(M)(\infty) = 0$  but  $j_*(M)(0) = \mathbb{Z}_2$ , it follows that  $j_*(M) \notin \text{Preglid}_{\Lambda} FR$ . Hence, the functor  $j_*$  does not restrict to a functor Prefrag<sub>Λ</sub>  $FR \rightarrow$  Preglid<sub>Λ</sub>  $FR$ .

5.4.5. Prefrag<sub>A</sub> FR is not the exact hull of Glid<sub>A</sub> FR. It is shown in [35] (see also [23]) that a deflationexact category C can be embedded in an exact category  $\overline{C}$  in a 2-universal way: there is a conflation-exact embedding  $i: \mathcal{C} \to \overline{\mathcal{C}}$  such that, for each exact category  $\mathcal{E}$ , the natural functor

$$
- \circ i \colon \operatorname{Hom}_{\operatorname{exact}}(\overline{\mathcal{C}}, \mathcal{E}) \to \operatorname{Hom}_{\operatorname{exact}}(\mathcal{C}, \mathcal{E})
$$

is an equivalence. In particular, every conflation-exact functor  $C \to \mathcal{E}$  factors essentially uniquely through  $i\colon \mathcal{C} \to \overline{\mathcal{C}}$ .

It might now be tempting to assume that  $\text{Prefix}_{\Delta}FR$  is the exact hull of Glid<sub>Λ</sub> FR (especially in light of proposition 8.6 below). However, in the notation of example 5.23, the only sub-prefragments of M are 0 and M itself. As M is not a glider, we see that M cannot occur as an extension of gliders. Hence, Prefrag<sub>Λ</sub> FR is not the exact hull of Glid<sub>Λ</sub> FR.

### 6. Noetherian objects

So far, we have considered the category of all glider representations of a filtered ring  $FR$ . In this section, we look at Noetherian objects in deflation quasi-abelian categories. Our first result is theorem 6.2, stating that the subcategory of Noetherian objects is a Serre subcategory (see definition 2.23), and hence itself quasi-abelian (see [36, lemma 4]). We then provide a number of equivalent formulations of when an object of Glid<sub>Λ</sub> FR is Noetherian in proposition 6.5.

We will use the following proposition, which is a straightforward adaptation of a similar statement in for quasi-abelian categories ([38, proposition 1.1.4], see also [36, proposition 1]).

**Proposition 6.1.** In a deflation quasi-abelian category, a morphism  $f: X \to Y$  factors as  $X \stackrel{d}{\to} \text{coim } f \stackrel{i}{\to}$ Y *where* d *is a deflation and* i *is a monomorphism.*

6.1. Noetherian objects in deflation quasi-abelian categories. We now look at the subcategory of Noetherian objects in a deflation quasi-abelian category. Recall that an object  $X$  in a category is called *Noetherian* if any ascending sequence of subobjects of X is stationary. The following theorem is a straightforward adaptation of a similar result for abelian categories.

**Theorem 6.2.** Let C be a deflation quasi-abelian category. The full subcategory  $\mathcal{N} \subseteq \mathcal{C}$  of Noetherian *objects is a Serre subcategory. In particular,* N *is deflation quasi-abelian.*

*Proof.* Consider a conflation  $X \rightarrow Y \rightarrow Z$  in C with  $Y \in \mathcal{N}$ . As the composition of monomorphisms is a monomorphism, we find that  $X \in \mathcal{N}$ . To show that  $Z \in \mathcal{N}$ , consider an ascending sequence  $Z_0 \hookrightarrow Z_1 \hookrightarrow$  $\cdots$  of subobjects of Z. Taking the pullback along  $Y \rightarrow Z$  (and using that pullbacks of monomorphisms are again monomorphisms), we find an ascending sequence  $Y_0 \hookrightarrow Y_1 \hookrightarrow \cdots$  of subobjects of Y:



As  $Y \in \mathcal{N}$ , we know that  $Y_i \hookrightarrow Y$  is an isomorphism for  $i \gg 0$ . It follows that  $Z_i \hookrightarrow Z$  is an isomorphism as well.

Assume now that  $X, Z \in \mathcal{N}$ . We show that  $Y \in \mathcal{N}$ . Let  $Y_0 \hookrightarrow Y_1 \hookrightarrow \cdots$  be an ascending sequence of subobjects of Y. Taking pullbacks along  $X \rightarrowtail Y$  yields a diagram

$$
X \cap_{Y} Y_{i} \rightarrowtail Y_{i} \longrightarrow Y/(X \cap_{Y} Y_{i})
$$
  
\n
$$
\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow
$$
  
\n
$$
X \rightarrowtail Y \longrightarrow Z
$$

where the downward arrows are monomorphisms. Indeed, the left-most arrow is a monomorphism as it is the pullback of a monomorphism. For the right-most arrow, it follows from the pullback property that the map  $X \cap_Y Y_i \to Y_i$  is the kernel of the composition  $Y_i \to Y \to Z$ , so that  $Y_i \to Y/(X \cap_Y Y_i)$  is the coimage of the composition  $Y_i \to Y \to Z$ . Proposition 6.1 now shows that the right-most map in the above diagram is a monomorphism.

As  $X, Z \in \mathcal{N}$ , we may assume the outer monomorphisms are isomorphisms for  $i \geq 0$ . It now follows from the short five lemma ([4, lemma 5.3] or [36, lemma 3]) that  $Y_i \hookrightarrow Y$  is an isomorphism as well. Hence,  $Y \in \mathcal{N}$ .

6.2. Noetherian glider representations. For later use, we introduce the category of Noetherian prefragments and glider representations.

**Definition 6.3.** We write prefrag<sub>Λ</sub> FR, preglid<sub>Λ</sub> FR, and glid<sub>Λ</sub> FR for the full subcategories of Prefrag<sub>Λ</sub> FR, Preglid<sub>Λ</sub> FR, and Glid<sub>Λ</sub> FR, respectively, consisting of all Noetherian objects.

**Corollary 6.4.** *The categories* prefrag<sub>Λ</sub> FR and preglid<sub>Λ</sub> FR are quasi-abelian. The category glid<sub>Λ</sub> FR *is deflation quasi-abelian.*

*Proof.* This follows from proposition 5.1 and theorem 5.12, together with theorem 6.2.

The following proposition helps recognizing Noetherian glider representations.

**Proposition 6.5.** Let  $\Gamma = \mathbb{Z}$  and  $\Lambda = \mathbb{Z}^{\leq 0}$ . Let FR be a nonnegative filtration of a ring R. Let  $M \in$  $Gli d<sub>\Lambda</sub> FR$  *be a glider representation. The following are equivalent.* 

- *(1) M is Noetherian in* Glid<sub>Λ</sub>  $FR$ ,
- *(2) M is Noetherian in* Prefrag<sub>Λ</sub>  $FR$ ,
- *(3) M is Noetherian in* Mod  $\mathcal{F}_{\Lambda}R$ ,
- *(4) the*  $F_0R$ *-module*  $\bigoplus_{\lambda \in \Lambda} M(\lambda)$  *is Noetherian,*
- *(5) the*  $F_0R$ *-module*  $M(0)$  *is Noetherian and*  $M(\lambda) = 0$  *for*  $\lambda \ll 0$ *.*

*Proof.* As each of the subcategories Glid<sub>Λ</sub>  $FR \subseteq \text{Prefix}_{A} FR \subseteq \text{Mod } \mathcal{F}_{A} R$  is closed under subobjects (see proposition 4.2 and corollary 5.11), it is clear that the first three statements are equivalent. The other statements are easy.

**Example 6.6.** Let  $\Gamma$ ,  $\Lambda$  be as in proposition 6.5. Let FR be the given by

$$
F_i R = \begin{cases} 0 & i < 0, \\ \mathbb{C} & i = 0, \\ \mathbb{C}[t] & i > 0. \end{cases}
$$

The projective prefragment  $\mathcal{F}_{\Lambda}R(0,-)$  is Noetherian, but the projective fragment  $\mathcal{F}_{\Lambda}R(-1,-)$  is not Noetherian (as  $\mathcal{F}_{\Lambda}R(-1,0) \cong \mathbb{C}[t]$  is not a finitely generated C-vector space).

The preglider  $L(\mathcal{F}_{\Lambda}R(0,-)) = \overline{\mathcal{F}}_{\Lambda}R(0,-)$  is not Noetherian.

# 7. Natural gliders and pregliders

Let Γ be an ordered group and let  $\Lambda \subseteq \Gamma$  be any subset. Let FR be a Γ-filtered ring. In this section, we assume that  $\Lambda$  has a maximal element which we denote by 0. In order to discuss the concept of a natural glider from [14], we introduce the category  $NPreglid_A FR$  of natural pregliders over FR.

7.1. **Natural pregliders.** As in §2.4, the inclusion  $n: \{0, \infty\} \to \Lambda \coprod \{\infty\}$  induces a restriction functor  $n^* \colon \text{Mod}\overline{\mathcal{F}}_{\Lambda}R \to \text{Mod}\overline{\mathcal{F}}_{\{0\}}R$ , which has a left adjoint  $n_1$  and a right adjoint  $n_*$ . These adjoints are fully faithful (see [33, remark 2.3 and example 2.13]). The adjoint triple  $(n_1, n^*, n_*)$  restricts to the corresponding subcategories of pregliders, as can be seen from the following explicit formulation.

**Proposition 7.1.** *The natural functor*  $n^*$ :  $\text{Preglid}_{\Lambda} FR \to \text{Preglid}_{\{0\}} FR$  *has a left adjoint given by* 

$$
n_!(M)(\lambda) = \begin{cases} M(\lambda) & \lambda \in \{0, \infty\} \\ 0 & otherwise, \end{cases}
$$

*and a right adjoint given by*

$$
n_*(M)(\lambda) = \{ m \in M(\infty) \mid F_{\lambda^{-1}}R \cdot m \subseteq \text{im } M(1_{0,\infty}) \}.
$$

**Definition 7.2.** We write NPreglid<sub>Λ</sub> FR for the essential image of the functor  $n_*$ : Preglid<sub>{0}</sub> FR  $\rightarrow$ Preglid<sub>Λ</sub> FR. An object of NPreglid<sub>Λ</sub> FR is called a *natural preglider*. We write  $\nu$ : NPreglid<sub>Λ</sub> FR  $\rightarrow$ Preglid<sub>Λ</sub> FR for the embedding and  $\rho$ : Preglid<sub>Λ</sub> FR → NPreglid<sub>Λ</sub> FR for the corresponding left adjoint.

**Remark 7.3.** For each  $M \in \text{Preglid}_{\Lambda} FR$ , the unit of the adjunction  $M \to \nu \circ \rho(M)$  is a monomorphism (as, for every  $\lambda \in \Lambda$ , it is given by the inclusion  $M(1_{\lambda,\infty})$ :  $M(\lambda) \to M^*(\lambda)$ ) and an epimorphism (as the map  $M(\infty) \to (\nu \circ \rho(M))(\infty)$  is an isomorphism).

**Remark 7.4.** Since NPreglid<sub>Λ</sub>  $FR \simeq$  Preglid<sub>{0}</sub>  $FR$ , we know that NPreglid<sub>Λ</sub>  $FR$  is quasi-abelian. However, as the following example shows, the embedding  $NPreglid_{\Lambda} FR \rightarrow Preglid_{\Lambda} FR$  need not commute with colimits.

**Example 7.5.** We set  $\Gamma = \mathbb{Z}$  and  $\Lambda = \mathbb{Z}^{\leq 0}$ . Let  $R = \mathbb{C}[t]$  with filtration  $\mathbb{C} \subseteq \mathbb{C}[t] \subseteq \mathbb{C}[t] \subseteq \ldots$  where  $F_0R = \mathbb{C}$ . We consider the following map between natural pregliders:



given by multiplication by t. One readily verifies that f is an inflation in Preglid<sub>Λ</sub> FR, and that the cokernel (in Preglid<sub>Λ</sub>  $FR$ ) is not a natural preglider.

**Proposition 7.6.** *Let*  $s: K \rightarrow M$  *be an inflation in* Preglid $\Lambda$  FR. If M *is a natural preglider, then so is* K.

*Proof.* Using that M is a natural preglider, the adjunction gives the following commutative diagram:



As Preglid<sub>Λ</sub> FR is a quasi-abelian category, we know that the canonical morphism  $K \to \nu \circ \rho(K)$  is an inflation. As it is also an epimorphism (see remark 7.3), we find that it is an isomorphism. Hence,  $K$  is a natural preglider.

**Proposition 7.7.** *The subcategory* NPreglid<sub>Λ</sub> FR *lies extension-closed in* Preglid<sub>Λ</sub> FR.

*Proof.* Let  $X \rightarrow Y \rightarrow Z$  be a conflation in Preglid<sub>A</sub> FR. Assume that X and Z are natural pregliders. Applying  $\nu \circ \rho$  yields the commutative diagram



As  $\nu \circ \rho$  is left exact, we know that  $i = \text{ker}(\rho)$ . Furthermore, as the composition  $Y \to \nu \circ \rho(Y) \to Z$  is a deflation, and Preglid<sub>Λ</sub> FR is quasi-abelian, we find that  $p: \nu \circ \rho(Y) \to Z$  is a deflation. Hence, the bottom row is a conflation. The short five lemma now shows that the canonical morphism  $Y \to \nu \circ \rho(Y)$ is an isomorphism. Hence, Y is a natural preglider.

**Remark 7.8.** The above puts forward two different conflation structures on NPreglid<sub>Λ</sub> FR.

- (1) As NPreglid<sub>Λ</sub> FR  $\simeq$  Preglid<sub>{0}</sub> FR, we know that NPreglid<sub>Λ</sub> FR is a quasi-abelian category. The conflation structure is induced by Preglid<sub>Λ</sub> FR via the embedding  $n_!$ : Preglid<sub>{0}</sub> FR  $\rightarrow$  $Preglid<sub>Λ</sub> FR.$
- (2) By proposition 7.7, we know that the exact structure of  $\text{Preglid}_{\Lambda} FR$  induces an exact structure on NPreglid<sub>Λ</sub> FR: the conflations in NPreglid<sub>Λ</sub> FR are those sequences which are conflations in  $Preglid<sub>Λ</sub> FR.$

In general, these two conflation structures need not coincide. As an example, the morphism  $f$  in example 7.5 is an inflation in the first exact structure (as it is the kernel of its cokernel), but not in the second exact structure.

7.2. Natural gliders. Having discussed the category of natural pregliders, we now turn to the category of natural gliders.

**Definition 7.9.** A glider  $M \in \text{Glid}_{\Lambda} FR$  is called a *natural glider* if it is isomorphic to  $Q(\overline{M})$  for a natural preglider  $\overline{M}$ . We write NGlid<sub>Λ</sub> FR for the full subcategory of Glid<sub>Λ</sub> FR given by the natural gliders.

**Remark 7.10.** The category NGlid<sub>Λ</sub> FR is the essential image of NPreglid<sub>Λ</sub> FR under the localization functor Q: Preglid<sub>Λ</sub>  $FR \rightarrow$  Glid<sub>Λ</sub>  $FR$ .

In proposition 7.12, we establish a recognition result for natural gliders. We start with the following lemma.

**Lemma 7.11.** Let  $s: M \overset{\sim}{\to} N$  be a weak isomorphism in  $\text{Preglid}_{\Lambda} FR$  (i.e.  $s \in \Sigma$ ). If N is a natural *preglider, then so is* M.

*Proof.* As  $s: M \to N$  is a weak isomorphism, it factors as  $M \stackrel{\sim}{\to} I \stackrel{\sim}{\to} N$ . It follows from proposition 7.6 that I is a natural preglider, and then from proposition 7.7 that M is a natural preglider (the last statement uses that ker( $M \stackrel{\sim}{\twoheadrightarrow} I$ ) is a natural preglider).

**Proposition 7.12.** A glider  $M \in \text{Glid}_{\Lambda} FR$  is natural if and only if  $L(M) \in \text{Preglid}_{\Lambda} FR$  is a natural *preglider.*

*Proof.* If  $L(M)$  is a natural preglider, then  $QL(M) \cong M$  is a natural glider. For the reverse implication, let  $M \in \text{Glid}_{\Lambda} FR$  be a natural glider. We know that there is a natural preglider  $\overline{M} \in \text{Preglid}_{\Lambda} FR$ such that  $Q(\bar{M}) \cong M$ . By adjointness, we find a weak isomorphism  $L(M) \stackrel{\sim}{\to} \bar{M}$ . Lemma 7.11 shows that  $L(M)$  is a natural preglider.

The following is an analogon of 7.6 (see also [14, lemma 1.6.2(3)]).

**Proposition 7.13.** *Let*  $s: K \rightarrow M$  *be an inflation in* Glid<sub>Λ</sub> FR. If M *is a natural glider, then so is* K.

*Proof.* Let  $K \nightharpoonup M \nightharpoonup N$  be a conflation in Glid<sub> $\Lambda$ </sub> FR with  $M \in \text{NGlid}_{\Lambda}$  FR. Applying the functor L: Glid<sub>A</sub>  $FR \to \text{Preglid}_{\Lambda} FR$ , we find a sequence  $LK \stackrel{Li}{\to} LM \stackrel{Lp}{\to} LN$ . The map Li factors as  $LK \stackrel{\sim}{\to}$ ker  $L_p \rightarrow LM$ , where the first map is a weak isomorphism. It now follows from propositions 7.6 and 7.12 that  $LK \in \text{NPreglid}_{\Lambda} FR$ , and hence  $K \in \text{NGlid}_{\Lambda} FR$ , as required.

**Corollary 7.14.** *The conflation structure on* NGlid<sub>Λ</sub> FR *induced by the embedding* NGlid<sub>Λ</sub> FR  $\rightarrow$  $Gli d<sub>Λ</sub> FR$  *gives*  $NGli d<sub>Λ</sub> FR$  *the structure of a deflation-exact category.* 

The following diagram can be appended to the diagram given in figure 1.

Theorem 7.15. *The following diagram commutes:*



*Moreover, the functors are fully faithful, the vertical functors have left adjoints, and the horizontal functors have right adjoints.*

*Proof.* That the functor L: Glid<sub>A</sub>  $FR \rightarrow$  Preglid<sub>A</sub>  $FR$  restricts to the natural gliders and pregliders, has been shown in proposition 7.12. This shows that the diagram commutes. By definition of the category  $NGlid_{\Lambda} FR$ , the functor  $Q$ : Preglid<sub> $\Lambda$ </sub>  $FR \to Glid_{\Lambda} FR$  also restricts to natural gliders and pregliders. This shows that the horizontal functors have right adjoints.

It follows from proposition 7.1 that  $\nu$  has a left adjoint  $\rho$ : Preglid<sub>Λ</sub> FR  $\rightarrow$  NPreglid<sub>Λ</sub> FR. It follows from proposition 2.11(1b) (with  $F = \rho \circ L$ ,  $H = \sigma$ , and  $G = Q$ ) that  $Q \circ \rho \circ L$  is left adjoint to  $\sigma$ .

Corollary 7.16. *The category* NGlid<sub>Λ</sub> FR *is complete and cocomplete.* 

*Proof.* It follows from theorem 7.15 that NGlid<sub>Λ</sub> FR is a reflective subcategory of Glid<sub>Λ</sub> FR, as well as a coreflective subcategory of  $NPreglid_A FR$ , both of which are complete and cocomplete.

**Remark 7.17.** As in proposition 7.1, the category  $\text{Preglid}_{\{0\}}FR$  admits two fully faithful embeddings into Preglid<sub>Λ</sub> FR: via n<sub>!</sub> and via n<sub>\*</sub>. Both essential images contain the subcategory  $i_*(\text{Mod } R)$ .

Since  $n_1(\text{Preglid}_{\{0\}} FR)$  is a Serre subcategory of Preglid<sub>A</sub> FR (and  $i_*(\text{Mod } R)$  is deflation-percolating in Preglid<sub>Λ</sub> FR, see proposition 5.5), we find that  $i_*(\text{Mod } R)$  is a deflation-percolating subcategory of  $n_1(\text{Preglid}_{0} FR)$ . The quotient  $n_1(\text{Preglid}_{0} FR)/i_*(\text{Mod } R)$  is equivalent to Glid<sub>{0}</sub>  $FR \simeq \text{Mod } S$ .

It follows from proposition 7.6 that  $i_*(\text{Mod } R)$  is a deflation-percolating subcategory of NPreglid<sub>Λ</sub> FR. It follows from lemma 7.11 that  $NPreglid_A FR/i_*(Mod R)$  is equivalent to  $NGlid_A FR$  as conflation categories.

Note that  $n_1(\text{Preglid}_{\text{f01}} FR) \simeq \text{NPreglid}_{\Lambda} FR$  as categories, but not as conflation categories.

8. The derived category of glider representations as a Verdier localization

In §3, we introduced the category of glider representations as a localization of the category of pregliders. In §5, we showed that Glid<sub>Λ</sub> FR can be seen as the quotient Preglid<sub>Λ</sub>  $FR/i_*(\text{Mod } R)$ , giving Glid<sub>Λ</sub> FR the structure of a deflation-exact category. In this section, we show that this quotient induces a Verdier localization sequence of the derived categories.

8.1. Projective gliders. We start by recalling the definition of a projective object in a deflation-exact category (see, for example, [4, 8]).

**Definition 8.1.** Let  $\mathcal{E}$  be a deflation-exact category.

(1) An object  $P \in \mathcal{E}$  is called *projective* if  $\text{Hom}(P, -)$ :  $\mathcal{E} \to \text{Ab}$  is an exact functor. We say that  $\mathcal{E}$  has *enough projectives* if for every object M of  $\mathcal E$  there is a deflation  $P \to M$  where P is projective.

(2) Dually, an object I is called *injective* if  $Hom(-, I)$ :  $\mathcal{E}^{\circ} \to Ab$  is an exact functor. We say that  $\mathcal{E}$ has *enough injectives* if for every object M of  $\mathcal E$  there is an inflation  $M \rightarrow I$  where I is injective.

We write Proj  $\mathcal{E}$  and Inj  $\mathcal{E}$  for the full subcategories of projectives and injectives, respectively.

The following proposition (see [23, proposition 3.22]) characterizes projective modules in a deflationexact category.

Proposition 8.2. Let  $\mathcal E$  be a deflation-exact category. The following are equivalent:

- *(1)* P *is projective.*
- *(2) For all deflations*  $f: X \rightarrow Y$  *and any map*  $g: P \rightarrow Y$  *there exists a map*  $h: P \rightarrow X$  *such that*  $g = f \circ h$ .
- *(3) Any deflation*  $f: X \rightarrow P$  *is a retraction, i.e. there exist a map*  $g: P \rightarrow X$  *such that*  $f \circ g = 1_P$ *.*
- **Proposition 8.3.** *(1) An object*  $M \in \text{Preglid}_{\Lambda} FR$  *is projective if and only if*  $\iota(M) \in \text{Mod }\overline{\mathcal{F}}_{\Lambda} R$  *is projective.*
	- (2) An object  $M \in \operatorname{Prefrag}_{\Lambda} FR$  is projective if and only if  $n(M) \in \operatorname{Mod} \mathcal{F}_{\Lambda} R$  is projective.

*Proof.* The functor  $\iota$ : Preglid<sub>A</sub>  $FR \to Mod \overline{\mathcal{F}}_hR$  reflect conflations. This shows that if an object  $\iota(M) \in$ Mod  $\overline{\mathcal{F}}_{\Lambda}R$  is projective, then  $M \in \text{Preglid}_{\Lambda} FR$  is projective. For the other direction, assume that  $M \in \text{Preglid}_{\Lambda} FR$  is projective. Let  $P \to \iota(M)$  be an epimorphism in Mod  $\overline{\mathcal{F}}_{\Lambda} R$  where P is projective. As all projective objects in Mod  $\overline{\mathcal{F}}_{\Lambda}R$  are pregliders (and projective in Preglid<sub> $\Lambda$ </sub> FR), and M is projective as a preglider, we find that M is a direct sumand of P. Hence,  $\iota(M)$  is a direct summand of P. We find that  $\iota(M)$  is projective.

The other statement can be proven in an analogous fashion.

**Proposition 8.4.** *(1) An object*  $M \in \text{Glid}_{\Lambda} FR$  *is projective if and only if*  $\phi(M) \in \text{Prefrag}_{\Lambda} FR$  *is projective.*

(2) An object  $M \in \text{Glid}_{\Lambda} FR$  *is projective if and only if*  $L(M) \in \text{Preglid}_{\Lambda} FR$  *is projective.* 

*Proof.* The proof of the first statement is analogous to the proof of proposition 8.3. For the second statement, as  $L$  has a conflation-exact right adjoint, we find that  $L$  maps projective objects to projective objects. Indeed, for every projective  $M \in \text{Glid}_{\Lambda} FR$ , we have that the functor  $\text{Hom}(LM, -) \cong$  $Hom(M, Q(-))$  is conflation-exact.

Assume now that LM is projective in Preglid<sub>Λ</sub> FR. Let  $K \rightarrow N \rightarrow M$  be any conflation in Glid<sub>Λ</sub> FR, ending in M. As L commutes with cokernels and Preglid<sub>A</sub> FR is quasi-abelian, we find that  $LN \rightarrow LM$ is a deflation. As  $LM$  is projective, this deflation splits. Applying the functor  $Q$ , we find that the given map  $N \rightarrow M$  is a split deflation. As any conflation ending in M splits, we find that M is projective.  $\square$ 

**Corollary 8.5.** *The categories*  $\text{Preglid}_{\Lambda} FR$ ,  $\text{Prefrag}_{\Lambda} FR$  and  $\text{Glid}_{\Lambda} FR$  have enough projectives.

*Proof.* This follows from propositions 8.3 and 8.4. □

8.2. A recollement on the derived level. We begin by observing that the upwards arrows in the diagram in figure 1 induce derived equivalences.

Proposition 8.6. *The following functors are triangle equivalences:*

- $(1)$  *ι*:  $\mathbf{D}^{\mathrm{b}}(\text{Preglid}_{\Lambda} FR) \to \mathbf{D}^{\mathrm{b}}(\text{Mod}\overline{\mathcal{F}}_{\Lambda} R)$ ,
- (2)  $\eta \colon \mathbf{D}^{\mathrm{b}}(\operatorname{Prefrag}_{\Lambda} FR) \to \mathbf{D}^{\mathrm{b}}(\operatorname{Mod} \mathcal{F}_{\Lambda} R)$ ,
- $(3) \phi: \mathbf{D}^{\mathrm{b}}(\text{Glid}_{\Lambda} FR) \to \mathbf{D}^{\mathrm{b}}(\text{Prefix}_{\Lambda} FR).$

*Proof.* It is shown in corollary 8.5 that Preglid<sub>Λ</sub> FR has enough projectives. It follows from proposition 8.3 that the functor  $\iota$ : Preglid<sub>A</sub>  $FR \to \text{Mod}(\overline{\mathcal{F}}_{\Lambda}R)$  induces an equivalence between the categories of projective objects, and hence between the homotopy categories of projectives. The result now follows from [23]. The other proofs are similar.  $\square$ 

Proposition 8.7. *The recollement*

$$
\operatorname{Mod} R \xleftarrow{i^* \atop i^!} \operatorname{Mod} \overline{\mathcal{F}}_{\Lambda} R \xleftarrow{j^* \atop j^* \atop j_*} \operatorname{Mod} \mathcal{F}_{\Lambda} R
$$

*lifts to a recollement on the bounded derived categories.*

*Proof.* As the categories  $Mod(R)$ ,  $Mod(\overline{\mathcal{F}}_{\Lambda}R)$  and  $Mod(\mathcal{F}_{\Lambda}R)$  have enough injective and projective objects, the six functors  $i^*, i_*, i^!, j_!, j^*$  and  $j_*$  lift to triangle functors on the bounded derived categories. It now follows from [33, theorem 7.2; lemma 7.3] that we obtain a recollement

$$
\mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(R)) \xleftarrow[\text{Ri}^{\mathrm{i}}]{i^*} \mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(\overline{\mathcal{F}}_{\Lambda}R)) \xleftarrow[\text{Ri}^{\mathrm{i}}]{Lj_!} \mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(\mathcal{F}_{\Lambda}R)).
$$

of triangulated categories.

Note that the sequence  $\mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(R)) \xrightarrow{i_*} \mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(\overline{\mathcal{F}}_{\Lambda}R)) \xrightarrow{j^*} \mathbf{D}^{\mathrm{b}}(\mathrm{Mod}(\mathcal{F}_{\Lambda}R))$  is a Verdier localization sequence. By proposition 5.5 and theorem 2.33, the sequence

$$
\mathbf{D}_{\mathrm{Mod}(R)}^{\mathrm{b}}(\mathrm{Preglid}_{\Lambda}FR) \to \mathbf{D}^{\mathrm{b}}(\mathrm{Preglid}_{\Lambda}FR) \xrightarrow{Q} \mathbf{D}^{\mathrm{b}}(\mathrm{Glid}_{\Lambda}FR)
$$

is a Verdier localization sequence as well.

**Proposition 8.8.** *The natural functor*  $\mathbf{D}^{\text{b}}(\text{Mod}(R)) \to \mathbf{D}_{\text{Mod}(R)}^{\text{b}}(\text{Preglid}_{\Lambda} FR)$  *is a triangle equivalence.* 

*Proof.* By [23, proposition 5.9] it suffices to show that the following property holds: for each conflation  $E' \rightarrow E \rightarrow A$  in Preglid<sub>A</sub> FR with  $A \in Mod(R) \subseteq Predid_A FR$  there exists a commutative diagram



where the rows are conflations and the top row belongs to  $Mod(R)$ .

Now let  $E' \rightarrow E \rightarrow A$  be a conflation with  $A \in Mod(R)$ . As  $Mod(R)$  has enough projectives, there exists a conflation  $K \rightarrow P \rightarrow A$  with P a projective in  $Mod(R)$ . Since P is also projective in Preglid<sub>Λ</sub> FR, we obtain a commutative diagram



where the map  $P \to E$  is obtained by the lifting property of the projective P. Clearly  $K \in Mod(R)$  as well. This shows the claim.

The following result summarizes the results of this section.

Proposition 8.9. *There is a Verdier localization sequence*

$$
\mathbf{D}^{\mathrm{b}}(\mathrm{Mod}\,R)\xrightarrow{i^{*}}\mathbf{D}^{\mathrm{b}}(\mathrm{Preglid}_{\Lambda}FR)\xrightarrow{Q}\mathbf{D}^{\mathrm{b}}(\mathrm{Glid}_{\Lambda}FR),
$$

*which can be completed to a recollement of triangulated categories.*

#### 9. Glider representations of bialgebra filtrations

In this section, we consider the category of glider representations associated to bialgebras. The definitions are based on the glider representations of groups [12, 14]. Our aims in this section are to related the monoidal category of glider representations to the notion of semi-Hopf categories (see §2.9), and to show that one can, in general, recover the original bialgebra from the monoidal category of glider representations.

Thus, let k be a field,  $\Gamma$  be an ordered group, B a k-bialgebra and FB a  $\Gamma$ -filtration of B by bialgebras, i.e. a collection of subbialgebras  $(FB_{\gamma})_{\gamma \in \Gamma^+}$  of B satisfying

- (1)  $FB_{\gamma} \cdot FB_{\gamma'} \subseteq FB_{\gamma\gamma'}$ ,
- (2)  $\Delta (FB_{\gamma}) \subseteq FB_{\gamma} \otimes FB_{\gamma}$ .

**Example 9.1.** Consider a group  $G$  and a finite group filtration

$$
1 \subseteq G_1 \subseteq G_2 \subseteq \cdots \subseteq G_n = G.
$$

From this filtration, one obtains the associated filtration of group algebras over some field  $k$ 

$$
k \subseteq k[G_1] \subseteq k[G_2] \subseteq \cdots \subseteq k[G_n] = k[G].
$$

This is a filtration of  $k[G]$  by bialgebras as above.

9.1. The monoidal structure and the connection with semi-Hopf categories. With notations as introduced before, we show that the filtered companion category  $\mathcal{F}_{A}B$  is a semi-Hopf category and that the category Glid<sub>Λ</sub> FB of glider representations is a full monoidal subcategory of Mod  $\mathcal{F}_{\Lambda}B$ . Moreover, the natural embedding Glid<sub>Λ</sub>  $FB \rightarrow$  Mod  $\mathcal{F}_{\Lambda}$  lifts to an equivalence on the level of derived categories.

**Proposition 9.2.** *The categories*  $\mathcal{F}_{\Lambda}B$  *and*  $\overline{\mathcal{F}}_{\Lambda}B$  *have the structure of k-linear semi-Hopf categories.* 

*Proof.* We only show that  $\mathcal{F}_{\Lambda}B$  is a k-linear semi-Hopf category, the case  $\overline{\mathcal{F}}_{\Lambda}B$  is similar. Let  $\lambda_1, \lambda_2, \lambda_3 \in$ Λ and set

$$
\circ_{\lambda_1, \lambda_2, \lambda_3}:FB_{\lambda_3\lambda_2^{-1}}\otimes FB_{\lambda_2\lambda_1^{-1}}\to FB_{\lambda_3\lambda_1^{-1}}: a\otimes b\mapsto ab.
$$

The map  $\circ_{\lambda_1,\lambda_2,\lambda_3}$  is well-defined as  $(FB_\gamma)_{\gamma \in \Gamma^+}$  is an algebra filtration. Moreover, as B is a bialgebra, the multiplication map  $\circ_{\lambda_1,\lambda_2,\lambda_3}$  is a coalgebra map. It follows that  $\mathcal{F}_{\Lambda}B$  is enriched over  $\underline{\mathcal{C}}(\mathcal{M}_k)$  and thus  $\mathcal{F}_{\Lambda}B$  is a k-linear semi-Hopf category.

Let  $\text{Mod}_k(\mathcal{F}_{\Lambda}B)$  and  $\text{Mod}_k(\overline{\mathcal{F}}_{\Lambda}B)$  be the categories of covariant additive functors to vec $(k)$ . Define a tensor product  $\otimes$  on  $Mod_k(\mathcal{F}_{\Lambda}B)$  by setting  $(M \otimes N)(\lambda) = M(\lambda) \otimes_k N(\lambda)$  for each  $\lambda \in \Lambda \coprod {\infty}$ .

**Proposition 9.3.** *The categories*  $\text{Mod}_k(\mathcal{F}_{\Lambda}B)$  *and*  $\text{Mod}_k(\overline{\mathcal{F}}_{\Lambda}B)$  *are* k-linear monoidal categories. More*over, the tensor product is exact.*

*Proof.* It follows from [3, proposition 3.2] that  $Mod_k(\mathcal{F}_{\Lambda}B)$  and  $Mod_k(\overline{\mathcal{F}}_{\Lambda}B)$  are k-linear tensor categories. The exactness of the tensor product is straightforward to verify.  $\hfill \Box$ 

The exactness of the tensor product over k immediately gives the following corollary.

**Corollary 9.4.** *The categories* Preglid<sub>Λ</sub> FB and Prefrag<sub>Λ</sub> FB inherit a monoidal structure from Mod<sub>k</sub>( $\overline{\mathcal{F}}$ <sub>Λ</sub>B) *and*  $\text{Mod}_k(\mathcal{F}_{\Lambda}B)$ *, respectively.* 

**Proposition 9.5.** *(1) The subcategory*  $i_*(\text{Mod}_k(B)) \subseteq \text{Mod}_k(\mathcal{F}_A B)$  *is a tensor ideal.* 

- *(2) The subcategory*  $i_*(\text{Mod}_k(B)) \subseteq \text{Preglid }_A FB$  *is a tensor ideal.*
- *(3) The category* Glid<sub>Λ</sub> FB *is a monoidal category, and the quotient functor* Q: Preglid<sub>Λ</sub> FR  $\rightarrow$  $Gli d<sub>A</sub> FR$  *is universal in the category of monoidal categories and functors.*

*Proof.* The first two statements are trivial. The last statement follows from the second one (see [17, corollary 1.4]).

**Remark 9.6.** The monoidal structure on Glid<sub>Λ</sub>  $FB = \Sigma^{-1}$  Preglid<sub>Λ</sub>  $FR$  (see definition 3.8) can be described as follows. As  $Ob(Glid_\Lambda FR) = Ob(Preglid_\Lambda FR)$ , the tensor product on gliders is the same as the tensor product on pregliders. Let now  $M \nightharpoonup M' \to X$  and  $N \nightharpoonup N' \to Y$  be morphisms in Glid<sub>A</sub> FR. The corresponding morphism from  $M \otimes N$  to  $X \otimes Y$  is given by:

$$
M \otimes N \xleftarrow{\sim} M' \otimes N' \to X \otimes Y.
$$

Here, we use that if  $s, t \in \Sigma$ , then  $s \otimes t \in \Sigma$ .

As Q: Glid<sub>A</sub>  $FB \rightarrow$  Preglid<sub>A</sub>  $FB$  is the identity on objects, we can give Q the structure of a monoidal functor by choosing the identity map for  $J_{M,N}$ :  $Q(M) \otimes Q(N) \rightarrow Q(M \otimes N)$ .

**Remark 9.7.** The tensor product on Glid<sub>Λ</sub> FB is conflation-exact; but need not commute with colimits. As an example, the cokernel of the morphism  $f: M \to N$  in Glid<sub>A</sub> FR from example 5.18 is zero. Let T be the prefragment given by  $0 \subseteq \mathbb{C}$ . Note that T is a glider with  $T(\infty) = \mathbb{C}S_3$ . The cokernel of the map  $f \otimes T : M \otimes T \to N \otimes T$  is isomorphic to T.

In particular, the monoidal structure on Glid<sub>Λ</sub> FR is not closed (i.e. the tensor product does not have a right adjoint).

**Proposition 9.8.** *The functor*  $\phi$ : Glid<sub>Λ</sub>  $FB \rightarrow$  Preglid<sub>Λ</sub>  $FB$  *is a monoidal functor.* 

*Proof.* Directly from proposition 9.5. □

**Remark 9.9.** The category Glid<sub>Λ</sub> FR is a full subcategory of Preglid<sub>Λ</sub> FR via the functor  $\phi$ . There is a unique structure of a monoidal functor on  $\phi$  such that  $\phi \circ Q = i_{*}$  (this follows from proposition 9.5); the morphisms  $J_{M,N}$ :  $\phi(M) \otimes \phi(N) \to \phi(M \otimes N)$  are given by the identities.

Combining the results of this section, we obtain the following theorem.

**Theorem 9.10.** Let B a k-bialgebra and FB a  $\Gamma$ -filtration of B by bialgebras. Let  $\Lambda \subseteq \Gamma$  be any subset. *The filtered companion category* FΛB *has the structure of a semi-Hopf category and there is a monoidal triangle equivalence*  $\mathbf{D}^{\mathrm{b}}(\text{Glid}_{\Lambda}FB) \to \mathbf{D}^{\mathrm{b}}(\text{Mod}\,\mathcal{F}_{\Lambda}B).$ 

*Proof.* It was verified in proposition 9.2 that  $\mathcal{F}_{\Lambda}B$  has the structure of a semi-Hopf category. The monoidal equivalence  $\mathbf{D}^{\rm b}(\text{Glid}_{\Lambda}FB) \to \mathbf{D}^{\rm b}(\text{Mod}\,\mathcal{F}_{\Lambda}B)$  is induced by the monoidal functor  $\eta \circ \phi$ : Glid $_{\Lambda}FB \to$  $\operatorname{Mod} \mathcal{F}_{\Lambda} B$ .

9.2. Glider representations and isocategorical groups. Let  $k$  be a field and let  $G$  be a group. It is known that one cannot recover the group G from the monoidal category  $\text{Mod } kG$  alone. Indeed, we say that the groups G and H are *isocategorical* over k if Mod kG and Mod kH are monoidally equivalent. Examples of nonisomorphic isocategorical groups are given in [19, Theorem 1.2].

In [13], the authors started from the Z-filtered algebra  $F(k)$  given by

$$
F_i(kG) = \begin{cases} 0 & i < 0, \\ k \cdot e_G & i = 0, \\ kG & i \ge 1, \end{cases}
$$

and let  $\Lambda = \{0, 1\}$ . Similar to the study of the characters of a group, one can look at various decategorifications of the category glid<sub>Λ</sub>  $F(k)$  of glider representations, such as the glider character ring or the (reduced) representation ring [10, 14]. Here, it can be shown that these invariants are sufficient to distinguish between the groups  $Q_8$  and  $D_8$  ([13, 14]) and even between some isocategorical groups ([10]). As of yet, there is no example known of groups  $G$  and  $H$  that cannot be distinguished using the glider character ring and the (reduced) representation ring.

In this section, we approach a similar question from a different perspective: can one recover the group G from the monoidal category Glid<sub>Λ</sub>  $F(kG)$ ? As the glider character ring and the (reduced) representation ring only depend on the monoidal category Glid<sub>Λ</sub>  $F(k)$ , the information one can recover from these rings is bounded above by the information one can recover from the category Glid<sub>Λ</sub>  $F(k)$ . In corollary 9.20 below, we show that the monoidal category  $\text{Glid}_{\Lambda} F(k)$  alone is sufficient to reconstruct the group G.

We will proceed with more generality. Let  $B$  be a finite-dimensional  $k$ -bialgebra, and consider the bialgebra  $\mathbb Z$ -filtered algebra  $FB$  given by

(1) 
$$
F_i B = \begin{cases} 0 & i < 0, \\ k \cdot 1_B & i = 0, \\ B & i \ge 1. \end{cases}
$$

We refer to such a filtration as the *standard one-step filtration* of B. We write mod B for the category of Noetherian (or, equivalently, finite-dimensional) modules.

For ease of notation, we consider the category glid<sub>Λ</sub> FR of Noetherian glider representations as a subcategory of Prefrag<sub>Λ</sub> FR via the embedding  $\phi$ : Glid<sub>Λ</sub> FR  $\rightarrow$  Prefrag<sub>Λ</sub> FR, i.e. we will describe an object of glid $\Lambda$  FR as an  $\mathcal{F}_{\Lambda}$ R-module.

As in definition 6.3, we write glid<sub>Λ</sub> FB for the subcategory of noetherian objects in Glid<sub>Λ</sub> FB. By proposition 6.5, these are exactly the glider representation with finite total dimension. In particular, as the bialgebra B is finite-dimensional,  $P_0 = \mathcal{F}_{\Lambda}R(0, -)$  and  $P_{-1} = \mathcal{F}_{\Lambda}R(-1, -)$  lie in glid<sub> $\Lambda$ </sub> FR.

Our goal is to recover the bialgebra B from the monoidal category glid<sub>Λ</sub> FB. We provide a short overview. First, we define full monoidal subcategories M and V of glid<sub>Λ</sub> FB. Here, the category M can be thought of as consisting of all gliders  $M \in \text{glid}_{\Lambda} FB$  for which  $M(1_{-1,0})$ :  $M(-1) \to M(0)$ is an isomorphism; the category V consists of all  $M \in \text{glid}_{\Lambda} FB$  for which  $M(-1) = 0$  (and hence,  $V \simeq \text{vec}_k$ ). The embedding  $V \to \text{glid}_{\Lambda} FB$  has a monoidal left adjoint. We then obtain a fiber functor  $\mathcal{M} \to \text{glid}_{\Lambda} FB \to \mathcal{V} \simeq \text{vec}_k$ . Finally, we show that the reconstruction theorem of finite-dimensional bialgebras applied to this fiber functor yields the bialgebra B.

We will be careful to construct the categories  $V, M$  and the fiber functor  $M \to V$  using only categorical properties of glid<sub>Λ</sub> FR (thus, without referring to the internal structure of the objects of glid<sub>Λ</sub> FR).

We start with specifying the full subcategories  $\mathcal M$  and  $\mathcal V$  of glid<sub> $\Lambda$ </sub> FB.

Notation 9.11. The category glid<sub>Λ</sub> FB has precisely two isomorphism classes of indecomposable projective objects:  $P_{-1}$  and  $P_0$ . They are distinguished by the property that  $Hom(P_{-1}, P_0) = 0$  while Hom( $P_0, P_{-1}$ )  $\neq 0$ .

We consider the subcategory M of glid<sub>Λ</sub> FB given by those objects M for which Hom( $P_0$ , M)  $\cong$  $Hom(P_{-1}, M)$  as vector spaces.

We write V for the full subcategory of glid<sub>Λ</sub> FB consisting of those objects M for which Hom( $P_{-1}$ , M) = 0.

**Remark 9.12.** (1) Note that  $P_0 \in \mathcal{V}$ .

(2) For any  $M \in \text{glid}_{\Lambda} FR$ , we have  $\dim_k \text{Hom}(P_0, M) < \infty$  (see proposition 6.5). As

$$
M(1_{-1,0})\colon M(-1)\to M(0)
$$

is a monomorphism, we find that  $M \in \mathcal{M}$  if and only if  $M(1_{-1,0})$  is an isomorphism.

**Proposition 9.13.** *The categories M and V are monoidal subcategories of* glid<sub>Λ</sub>  $FB$ .

*Proof.* Directly from the definitions. □

**Proposition 9.14.** *The monoidal embedding*  $V \to \text{glid}_{\Lambda} FB$  *has a monoidal right adjoint*  $R: \text{glid}_{\Lambda} FB \to$ V*, which is unique up to monoidal natural equivalence.*

*Proof.* The right adjoint is given by mapping an object  $M \in \text{glid}_{\Lambda} FB$  to the functor

$$
R(M)(i) = \begin{cases} 0 & i = -1, \\ M(0) & i = 0. \end{cases}
$$

It is clear that this is a monoidal functor. As  $R$  is an adjoint to the embedding in the 2-category of monoidal categories, monoidal functors, and monoidal natural transformations,  $R$  is unique up to monoidal natural equivalence.

**Lemma 9.15.** *(1) A monoidal equivalence*  $E: \text{vec}_k \to \text{vec}_k$  *is monoidally naturally isomorphic to the identity.*

(2) Let  $H_1, H_2$ :  $vec_k \rightarrow V$  *be monoidal functors. If*  $H_1$  *and*  $H_2$  *are equivalences, then there is a monoidal natural isomorphism*  $H_1 \rightarrow H_2$ .

*Proof.* The first statement follows from the reconstruction theorem of finite-dimensional bialgebras, cf. [20, Theorem 5.2.3]).

For the second part, let  $H_1, H_2$ :  $\mathrm{vec}_k \to \mathcal{V}$  be as in the statement of the proposition, and let  $H_1^{-1}$  be a monoidal quasi-inverse of  $H_1$ . It is established in the first part that  $H_1^{-1} \circ H_2 \cong 1_{\text{vec}}$ . This shows that  $H_1 \cong H_2$ , as required.

**Remark 9.16.** The composition  $\Xi: \mathcal{M} \to \text{glid}_{\Lambda} FB \to \mathcal{V} \to \text{vec}_k$  is a faithful monoidal functor. In theorem 9.19 below, we show that, by applying the reconstruction theorem of finite-dimensional bialgebras, we can recover the bialgebra B.

For the next lemma, recall that a finite-dimensional B-module is a pair  $(V, \rho_V)$  where V is a finitedimensional vector space and  $\rho_V : B \to \text{End}(V)$  is an algebra homomorphism.

**Lemma 9.17.** *There is a monoidal equivalence*  $\Phi$ : mod  $B \to M$ , *mapping a* B-module  $M = (V, \rho_V)$  *to the functor*  $\Phi(M)$ :  $\mathcal{F}_{\Lambda}B \to \text{vec}_k$  *given by*  $\Phi(M)(-1) = \Phi(M)(0) = V$  *and, for each*  $b \in B = \text{Hom}(-1, 0)$ *, the corresponding map*  $\Phi(M)(b)$ :  $\Phi(M)(-1) \to \Phi(M)(0)$  *is given by the action of* B *on* V *(i.e.*  $\Phi(M)(b) =$  $\rho(b)$ ).

*Proof.* It is straightforward to verify that this correspondence is a monoidal functor. To see that it is an equivalence, we will construct a quasi-inverse. Let  $N \in \mathcal{M}$ . As dim  $N(-1) = \dim N(0)$ , the monomorphism  $N(1_{-1,0})$ :  $N(-1) \rightarrow N(0)$  is an isomorphism. As N is a glider, there is a preglider  $N' \in \text{Preglid}_{\Lambda} FR$  such that  $Q(N') \cong N$ . It is now easy to see that  $N(-1) = N(0)$  is a B-submodule of  $N'(\infty)$ . The functor  $\Psi: \mathcal{M} \to \text{mod } B$ , given by mapping N to the B-submodule  $N(-1)$  is a quasi-inverse to  $\Phi$ .

**Remark 9.18.** Note that the construction of the functor  $\Phi$  in lemma 9.17 is based on the forgetful functor mod  $B \to \text{vec}_k$ .

We now come to the main result of this subsection.

**Theorem 9.19.** Let  $\Gamma = \mathbb{Z}$  and let  $\Lambda = \{-1,0\} \subset \mathbb{Z}$ . Let B and B' be finite-dimensional k-algebras and let FB and FB' be their standard one-step filtrations. The categories  $\text{glid}_{\Lambda}$  FB and  $\text{glid}_{\Lambda}$  FB' are *monoidally equivalent if and only if*  $B \cong B'$  *as bialgebras.* 

*Proof.* It is clear that an isomorphism  $B \cong B'$  induces a monoidal equivalence glid<sub>Λ</sub> FB and glid<sub>Λ</sub> FB'. For the other direction, consider the subcategory  $\mathcal V$  of glid<sub>Λ</sub> FB. There is a monoidal functor given by mapping  $V \in \text{vec}_k$  to the glider representation  $M_V$  given by  $M_V(0) = V$  and  $M_V(-1) = 0$ . It follows from lemma 9.15 that this monoidal functor is unique (up to monoidal natural isomorphism), and hence, does not depend on the bialgebra B.

The composition  $\Xi \colon \mathcal{M} \to \text{glid}_{\Lambda} FB \to \mathcal{V} \to \text{vec}_k$  is a faithful monoidal functor. We claim that the bialgebra obtained from the reconstruction theorem of finite-dimensional bialgebras ([20, Theorem 5.2.3]) is B. Consider the monoidal equivalence  $\Phi: \text{mod } B \to \mathcal{M}$  from lemma 9.17. It is straightforward to verify that the composition  $\Phi \circ \Xi$ : mod  $B \to \mathcal{M} \to \text{glid}_{\Lambda} FB \to \mathcal{V} \to \text{vec}_k$  is the usual fiber functor. It now follows from the reconstruction theorem of finite-dimensional bialgebras that one recovers the bialgebra B from End $(\Xi)$ .

As we have recovered the bialgebra B from only the monoidal category glid<sub>Λ</sub> FB, we infer that glid<sub>Λ</sub> FB and glid<sub>Λ</sub> FB' are monoidally equivalent if and only if  $B \cong B'$  as bialgebras.

Restricting to the case where the bialgebra  $B$  is a group algebra  $kG$ , we see that the monoidal category glid<sub>Λ</sub>  $F(kG)$  of glider representations, associated with the trivial group filtration  $\{e\} \subseteq G$  is sufficient to recover the group.

**Corollary 9.20.** *Let* G and H *be finite groups. There is a group isomorphism*  $G \cong H$  *if and only if there is a monoidal equivalence* glid<sub>Λ</sub>  $F(kG) \simeq$  glid<sub>Λ</sub>  $F(kH)$ .

**Remark 9.21.** The monoidal category glid<sub>Λ</sub>  $F(k)$  of Noetherian glider representations retains more information than the monoidal category mod  $kG$  of finite-dimensional modules. In contrast, the Grothendieck ring of glid<sub>Λ</sub>  $F(k)$  is isomorphic to the product  $\mathbb{Z} \times \mathbb{Z}$  (this follows from the derived equivalence  $\mathbf{D}^{\mathrm{b}}(\text{glid}_{\Lambda} F(kG)) \simeq \mathbf{D}^{\mathrm{b}}(\text{mod }\mathcal{F}_{\Lambda}(kG))).$  As such, the Grothendieck ring of glid $\Lambda F(kG)$  is disassociated from the group  $G$ .

As the following example illustrates, there is no direct generalization of corollary 9.20 to the category of prefragments.

**Example 9.22.** Let G and H be any groups with the same number of elements. Consider the standard one-step bialgebra filtrations,  $F(kG)$  and  $F(kH)$ , of the group algebras kG and kH. Note that there is a coalgebra isomorphism  $kG \to kH$ , mapping grouplike elements to grouplike elements (and hence mapping  $1 \in kG$  to  $1 \in kH$ ), inducing an equivalence  $\mathcal{F}_{\Lambda}(kH) \to \mathcal{F}_{\Lambda}(kG)$  of semi-Hopf categories. We obtain a monoidal equivalence  $\text{Prefix}_{\mathbb{A}} F(kH) \to \text{Prefix}_{\mathbb{A}} F(kG)$ , even though G and H need not be isomorphic groups.

9.3. Recovering an algebra from the category of glider representations. The proof of theorem 9.19 can easily be adapted to work in the setting of finite-dimensional algebras instead of bialgebras.

**Theorem 9.23.** Let  $\Gamma = \mathbb{Z}$  and let  $\Lambda = \{-1,0\} \subset \mathbb{Z}$ . Let FA and FA' be the standard one-step filtrations *of finite-dimensional* k*-algebras* A *and* A′ *. The categories* glid<sup>Λ</sup> F A *and* glid<sup>Λ</sup> F A′ *are equivalent if and only if*  $A \cong A'$  *as algebras.* 

*Proof.* The definition of the indecomposable projective objects,  $P_0$  and  $P_{-1}$ , does not use the monoidal structure. Similarly, the definition of the subcategories  $\mathcal M$  and  $\mathcal V$  from notation 9.11 carries over to this setting. As in lemma 9.17, we infer that  $\mathcal{M} \simeq \text{mod } A$ . Under this equivalence, the functor  $F =$  $\text{Hom}_{\text{grid}_A} F_A(P_0, -) : \mathcal{M} \to \text{vec}_k$  corresponds to the forgetful functor  $\text{Hom}(A, -)$ : mod  $A \to \text{vec}_k$ . We then recover A as the algebra  $\text{Hom}(F, F)$ .

As for bialgebras, there is no direct generalization of theorem 9.23 to the setting of prefragments.

**Example 9.24.** Assume that A and A' be finite-dimensional k-algebras of the same dimension. In this case, there is an equivalence of categories  $\mathcal{F}_{\Lambda}A \to \mathcal{F}_{\Lambda}A'$  mapping  $1 \in A$  to  $1 \in A'$ . The natural functor Mod  $\mathcal{F}_{\Lambda}A' \to \text{Mod }\mathcal{F}_{\Lambda}A$  induces an equivalence Prefrag<sub> $\Lambda$ </sub>  $FA' \to \text{Prefix}_{\Lambda}FA$ , and thus an equivalence prefrag<sub>Λ</sub>  $FA' \rightarrow$  prefrag<sub>Λ</sub>  $FA$ , even though A and A' need not be isomorphic.

## Appendix A. Comparison to earlier works

In this appendix, we provide a comparison of some notions in this paper to existing literature about glider representations. Our main reference for this appendix is [14].

Fragments over filtered rings were first introduced in [29, 30] as a generalization of modules; since then, the definition has been amended (see [15] or [14]). To avoid confusion, we opt to refer to the original concept (see [30]) as a prefragment, reserving the notion of a fragment for the one introduced in [14].

The definition of a glider as we use it, has been introduced in  $[15]$ , as a prefragment M whose partial actions  $\phi: F_i R \times M_{-i} \to M$  are induced by an ambient R-module  $\Omega_M$ . A morphism between glider representations (as given in [14]) is required to be compatible with a chosen ambient R-module. As expounded on in remark 2.6, this compatibility condition obscures whether the composition of glider

morphisms is well-defined. Indeed, in  $[14, \S1.7]$ , a sequence of morphisms is called a glider sequence if the sequence is composable.

As the functor embedding  $\phi$ : Glid<sub>Λ</sub> FR  $\rightarrow$  Prefrag<sub>Λ</sub> FR is fully faithful (see proposition 4.8), we can alternatively define the category of glider representations as follows.

Definition A.1. Let FR be a Z-filtered ring. A *glider representation* over FR consists of an R-module  $\Omega$  together with a  $\mathbb{Z}^{\leq 0}$ -indexed chain sequence of subgroups:

$$
\ldots \subseteq M_{-2} \subseteq M_{-1} \subseteq M_0 \subseteq \Omega
$$

such that the action  $R \otimes \Omega \to \Omega$  induces maps  $F_i R \otimes M_j \to M_{i+j}$ .

Let  $M_{\bullet} \subseteq \Omega_M$  and  $N_{\bullet} \subseteq \Omega_N$  be glider representations over FR. A *morphism of glider representations*  $f: (M_{\bullet} \subseteq \Omega_M) \to (N_{\bullet} \subseteq \Omega_N)$  is given by an additive map  $f: M_0 \to N_0$  satisfying the following conditions:

- (1) for all  $j \in \mathbb{Z}^{\leq 0}$ , we have  $f(M_j) \subseteq N_j$ , and
- (2)  $\forall i \in \mathbb{Z}$  and  $\forall j \leq -i$ , the following diagram commutes:

$$
F_i R \otimes M_j \longrightarrow M_{i+j}
$$

$$
\downarrow 1 \otimes f \qquad \qquad f
$$

$$
F_i R \otimes N_j \longrightarrow N_{i+j}
$$

or, equivalently, for all  $r \in F_i R$  and  $m \in M_j$ , we have  $f(rm) = rf(m)$ .

Following proposition 5.8, a sequence  $K \to M \to N$  is a *conflation* if and only each of the induced sequences  $0 \to K_j \to M_j \to N_j \to 0$  is exact in Mod  $F_0R$  (for all  $j \in \mathbb{Z}^{\leq 0}$ ).

- **Remark A.2.** (1) This definition is based on proposition 4.8. In this definition, the role of the  $R$ module  $\Omega_M$  is to determine whether the prefragment is, in fact, a glider representation; it plays no role in the definition of the morphisms, nor in determining whether a sequence of gliders is a conflation.
	- (2) As in [14], it suffices to require (2) for  $i + j = 0$ .
	- (3) By taking  $i = 0$  in the definition of a glider morphism, we see that  $f: M_i \to N_i$  is an  $F_0R$ -module homomorphism, for all  $j \leq 0$ .
	- (4) We do not claim that the map  $f: M_0 \to N_0$  can be extended to an R-morphism  $f_{\Omega}: \Omega_M \to \Omega_N$ (as is illustrated in example 3.16), but it follows from proposition 4.9 that this can be done after possibly re-choosing  $\Omega_M$ .

In fact, given the fragment part  $\dots \subseteq M_{-2} \subseteq M_{-1} \subseteq M_0$  of a glider, there is a canonical way of completing this to a glider in the sense of definition A.1 by applying the functor L: Glid<sub>Λ</sub>  $FR \rightarrow$ Preglid<sub>Λ</sub>  $FR$  (see §4.4).

(5) The category Frag<sub>Λ</sub> FR of fragments, considered in [14], is an additive subcategory of the category of prefragments, containing the category of glider representations. Thus, Glid<sub>Λ</sub> FR  $\subseteq$  $\text{Frag}_{\Lambda} FR \subseteq \text{Prefix}_{\Lambda} FR.$ 

*Limits and colimits.* Limits and colimits of fragments and gliders are discussed in [14, §1.8]. The categories of pregliders, prefragments, and (natural) gliders are complete and cocomplete. With the description of a glider from definition A.1, limits are taken pointwise. As is illustrated in example 5.18, colimits cannot be taken pointwise (see proposition 5.3).

*The conflation structure on the category of glider representations.* A morphism in a conflation category is called *admissible* if it admits a deflation-inflation factorization, i.e.  $f: X \rightarrow Y$  is admissible if it factors as  $X \rightarrow Z \rightarrow Y$ . Admissible morphisms have been called *strict* in [14, proposition 1.7.1]. A strict monomorphism is an inflation, and a strict epimorphism is a deflation.

In particular, a morphism  $f: M \to N$  of glider representations is a deflation if and only if, for all  $i \in \mathbb{Z}^{\leq 0}$ , the map  $f_i \colon M_i \to N_i$  is an epimorphism.

What is called the image of a morphism  $f: X \to Y$  in [14] is coker(ker f), and is often called the coimage of  $f$  (see, for example,  $[8, 28]$ ). As the category of gliders is not an abelian category, the natural morphism from the coimage of a morphism to its image need not be an isomorphism. However, as Glid<sub>A</sub> FR is deflation quasi-abelian, a morphism  $f: M \to N$  of gliders is the composition of a deflation d:  $M \rightarrow \text{coim } f$  and a monomorphism  $m: \text{coim } f \hookrightarrow N$  (see [36, corollary 1]).

*Noetherian glider representations.* Noetherian fragments were introduced in [14, §2.3]. It follows from proposition 6.5 that a glider is Noetherian if and only if it is Noetherian as a prefragment. Our theorem 6.2 and proposition 6.5 are analogues of [14, theorems 2.3.2 and 2.3.4].

*Projective glider representations.* A glider  $M \in \text{Glid}_{\Lambda} FR$  is projective if and only if the prefragment part  $\phi(M) \in \text{Prefix}_{\mathcal{A}} F M$  is projective. A glider representation is freely generated in the sense of [14, definition 2.1.3] if and only if it is a direct sum of standard projectives. A glider representation is projective if and only if it is a direct summand of a freely generated projective (a similar statement for fragments has been shown in [14, proposition 2.2.5]).

*Monoidal structure for glider representations of filtered groups.* Let k be a field. In [14, proposition 4.6.7 and definition 4.6.8], a tensor product of k-linear fragments and glider representations of filtered groups were defined: the product  $M \otimes N$  is defined via  $(M \otimes N)_{-i} \cong M_{-i} \otimes_k N_{-i}$ . This coincides with the product considered in §9. In particular, for a filtered group  $G_0 \subseteq G_1 \subseteq \ldots \subseteq G_n \subseteq \ldots$ , the category of glider representations is (monoidally) derived equivalent to the category of representations of a semi-Hopf category. Following  $\S9.2$ , one can recover the group G from the monoidal category of glider representations.

*Natural gliders.* Let  $\Omega$  be an R-module. Given an S-submodule  $M_0 \subseteq \Omega$ , we can build a glider representation by setting  $M_{-i}^* = \{m \in M_0 \mid F_iR \cdot m \subseteq M_0\}$ . Such glider representations (as well as those isomorphic) are called *natural gliders* in [14, definition 1.3.1]. In earlier work [29, 30], the same concept was called a natural fragment. The definitions given in [14] and in §7 coincide. Our proposition 7.13 recovers [14, lemma 1.6.2(3)]. It follows from theorem 7.15 that the category NGlid<sub>Λ</sub> FR of natural glider representations is a reflective subcategory of Glid<sub>Λ</sub> FR. As such, the category NGlid<sub>Λ</sub> FR is complete and cocomplete, and a limit of natural gliders is again a natural glider. This answers a question posed at the end of  $[14, §1.8]$ .

## **REFERENCES**

- 1. Jiří Adámek and Jiří Rosický, Locally presentable and accessible categories, London Mathematical Society Lecture Note Series, vol. 189, Cambridge University Press, Cambridge, 1994.
- 2. Maria Jesus Asensio, Michel Van den Bergh, and Freddy Van Oystaeyen, A new algebraic approach to microlocalization of filtered rings, Trans. Amer. Math. Soc. 316 (1989), no. 2, 537–553.
- 3. E. Batista, S. Caenepeel, and J. Vercruysse, Hopf categories, Algebr. Represent. Theory 19 (2016), no. 5, 1173–1216.
- 4. Silvana Bazzoni and Septimiu Crivei, One-sided exact categories, J. Pure Appl. Algebra 217 (2013), no. 2, 377–391.
- 5. A. A. Beïlinson, J. Bernstein, and P. Deligne, Faisceaux pervers, Analysis and topology on singular spaces, I (Luminy, 1981), Astérisque, vol. 100, Soc. Math. France, Paris, 1982, pp. 5-171.
- 6. A. Bondal and M. Van den Bergh, Generators and representability of functors in commutative and noncommutative geometry, Mosc. Math. J. 3 (2003), no. 1, 1–36, 258.
- 7. Mitchell Buckley, Timmy Fieremans, Christina Vasilakopoulou, and Joost Vercruysse, Oplax hopf algebras, arXiv preprint arXiv:1710.01465 (2017).
- 8. Theo Bühler, *Exact categories*, Expo. Math.  $28$  (2010), no. 1, 1–69.
- 9. Frederik Caenepeel, Glider representations of chains of semisimple Lie algebra, Comm. Algebra 46 (2018), no. 11, 4985–5005.
- 10. Frederik Caenepeel and Geoffrey Janssens, Glider representation rings with a view on distinguishing groups, arXiv preprint arXiv:1911.11733 (2019).
- 11. Frederik Caenepeel and Fred Van Oystaeyen, Clifford theory for glider representations, Algebr. Represent. Theory 19 (2016), no. 6, 1477–1493.
- 12. Glider representations of group algebra filtrations of nilpotent groups, Algebr. Represent. Theory 21 (2018), no. 3, 529–550.
- 13. Generalized characters for glider representations of groups, Algebr. Represent. Theory (2019), 1-24.
- 14. Glider representations, Monographs and Research Notes in Mathematics, Chapman & Hall/CRC, 2019.
- 15. Localizations and sheaves of glider representations, J. Pure Appl. Algebra 223 (2019), no. 6, 2635–2672.
- 16. S. Caenepeel and T. Fieremans, Descent and Galois theory for Hopf categories, J. Algebra Appl. 17 (2018), no. 7, 1850120, 39.
- 17. Brian Day, Note on monoidal localisation, Bull. Austral. Math. Soc. 8 (1973), 1–16.
- 18. Spencer E. Dickson, A torsion theory for Abelian categories, Trans. Amer. Math. Soc. 121 (1966), 223–235.
- 19. Pavel Etingof and Shlomo Gelaki, Isocategorical groups, Internat. Math. Res. Notices (2001), no. 2, 59–76.
- 20. Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik, Tensor categories, Mathematical Surveys and Monographs, vol. 205, American Mathematical Society, Providence, RI, 2015.
- 21. P. Gabriel and M. Zisman, Calculus of fractions and homotopy theory, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35, Springer-Verlag New York, Inc., New York, 1967.
- 22. Andreas Heider, Two results from Morita theory of stable model categories, 2007, p. 69.
- 23. Ruben Henrard and Adam-Christiaan van Roosmalen, Derived categories of one-sided exact categories and their localizations, arXiv preprint arXiv:1903.12647 (2019).
- 24. , Localizations of one-sided exact categories, arXiv preprint arXiv:1903.10861 (2019).
- 25. J. P. Jans, Some aspects of torsion, Pacific J. Math. 15 (1965), 1249–1259.
- 26. Masaki Kashiwara, D-modules and microlocal calculus, Translations of Mathematical Monographs, vol. 217, American Mathematical Society, Providence, RI, 2003, Translated from the 2000 Japanese original by Mutsumi Saito, Iwanami Series in Modern Mathematics.
- 27. J. C. McConnell, Quantum groups, filtered rings and Gelfand-Kirillov dimension, Noncommutative ring theory (Athens, OH, 1989), Lecture Notes in Math., vol. 1448, Springer, Berlin, 1990, pp. 139–147.
- 28. Barry Mitchell, Theory of categories, Pure and Applied Mathematics, Vol. XVII, Academic Press, New York-London, 1965.
- 29. Sayed Nawal and Freddy Van Oystaeyen, An introduction of fragmented structures over filtered rings, Comm. Algebra 23 (1995), no. 3, 975–993.
- 30. , Projective aspects of fragments over filtered rings, Comm. Algebra 24 (1996), no. 1, 165–185.
- 31. Amnon Neeman, The derived category of an exact category, J. Algebra 135 (1990), no. 2, 388–394.
- 32. Constantin Năstăsescu and F. Van Oystaeyen, Graded and filtered rings and modules, Lecture Notes in Mathematics, vol. 758, Springer, Berlin, 1979.
- 33. Chrysostomos Psaroudakis, Homological theory of recollements of abelian categories, Journal of Algebra 398 (2014), 63–110.
- 34. Chrysostomos Psaroudakis and Jorge Vitória, Recollements of module categories, Appl. Categ. Structures 22 (2014), no. 4, 579–593.
- 35. Alexander Rosenberg, Homological algebra of noncommutative spaces, I, preprint Max Planck, Bonn: MPIM 11-69 (2011).
- 36. Wolfgang Rump, Almost abelian categories, Cahiers Topologie Géom. Différentielle Catég. 42 (2001), no. 3, 163–225.
- 37. Pierre Schapira and Jean-Pierre Schneiders, Derived categories of filtered objects, Astérisque (2016), no. 383, 103-120.
- 38. Jean-Pierre Schneiders, Quasi-abelian categories and sheaves, Mém. Soc. Math. Fr. (N.S.) (1999), no. 76, vi+134.
- 39. Bo Stenström, Rings of quotients, Springer-Verlag, New York-Heidelberg, 1975, Die Grundlehren der Mathematischen Wissenschaften, Band 217, An introduction to methods of ring theory.

40. James Wallbridge, Homotopy theory in a quasi-abelian category, arXiv preprint arXiv:1510.04055 (2015).

Ruben Henrard, Universiteit Hasselt, Campus Diepenbeek, Departement WNI, 3590 Diepenbeek, Belgium E-mail address: ruben.henrard@uhasselt.be

Adam-Christiaan van Roosmalen, Universiteit Hasselt, Campus Diepenbeek, Departement WNI, 3590 Diepenbeek, Belgium

 $E\text{-}mail\;address:$  adamchristiaan.vanroosmalen@uhasselt.be