Made available by Hasselt University Library in https://documentserver.uhasselt.be

Association of greenspace exposure with telomere length in preschool children Peer-reviewed author version

Miri, Mohammad; De Prado-Bert, Paula; Alahabadi, Ahmad; Najafi, Moslem Lari; Rad, Abolfazl; Moslem, Alireza; Aval, Hamideh Ebrahimi; Ehrampoush, Mohammad Hassan; Bustamante, Mariona; Sakhvidi, Mohammad Javad Zare; NAWROT, Tim; Sunyer, Jordi & Dadvand, Payam (2020) Association of greenspace exposure with telomere length in preschool children. In: Environmental Pollution, 266 (Part 1) (Art N° 115228).

DOI: 10.1016/j.envpol.2020.115228 Handle: http://hdl.handle.net/1942/32869

Association of Greenspace Exposure with Telomere Length in Preschool Children

Mohammad Miri^{1, δ}, Paula de Prado-Bert^{2, 3,4, δ}, Ahmad Alahabadi¹, Moslem Lari Najafi⁵, Abolfazl Rad⁶, Alireza Moslem⁶, Hamideh Ebrahimi Aval⁶, Mohammad Hassan Ehrampoush⁷, Mariona Bustamante^{2,3,4}, Mohammad Javad Zare Sakhvidi^{8,9}, Tim Nawrot^{10,11}, Jordi Sunyer^{2,3,4}, Payam Dadvand^{2,3,4, *}

- Non-communicable Diseases Research Center, Department of Environmental Health, School of Public Health, Sabzevar University of Medical Sciences, Sabzevar, Iran.
- 2. ISGlobal, Barcelona, Spain.
- 3. Pompeu Fabra University, Barcelona, Catalonia, Spain.
- 4. Ciber on Epidemiology and Public Health (CIBERESP), Madrid, Spain.
- Pharmaceutical Sciences and Cosmetic Products Research Center, Kerman University of Medical Sciences, Kerman, Iran.
- Cellular and Molecular Research Center, Sabzevar University of Medical Sciences, Sabzevar, Iran.
- Department of Environmental Health, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- Occupational Health Research Center, Department of Occupational Health, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- Univ Rennes 1, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) – UMR_S 1085, F-35000 Rennes, France.
- 10. Centre for Environmental Sciences, Hasselt University, Hasselt, Belgium.

11. Department of Public Health & Primary Care, Leuven University, Leuven, Belgium δ these authors equally contributed.

* Corresponding authors

Payam Dadvand

Institute of Global Health of Barcelona. (ISGlobal) Carrer Dr. Aiguader 88. 08003-Barcelona, Spain. Phone: +34 93 214 73 29 E-mail: <u>payam.dadvand@isglobal.org</u>

Acknowledgments: This study was supported by Sabzevar University of medical sciences. Payam Dadvand is funded by a Ramón y Cajal fellowship (RYC-201210995) awarded by the Spanish Ministry of Economy and Competitiveness. The sponsor or funding organizations had no role in the design or conduct of this research.

Declarations of interest: All authors declare that they have no conflicting interests.

Abbreviations: Particulate Matter 2.5 with diameter less than 2.5 micrometer (PM2.5), Telomere Length (TL), Leukocyte Telomere Length (LTL), Normalized Difference Vegetation Index (NDVI), Quantitative Polymerase Chain Reaction (qPCR), Telomere amplification product (T), Single-copy gene (S), Telomere / Single copy gene ratio (T/S), Reactive Oxygen Species (ROS), Interquartile Range (IQR), Confidence Interval (CI).

1 Abstract

2

Exposure to greenspace has been associated with a wide range of health benefits; however, the 3 available evidence on the association of this exposure with telomere length (TL), an early 4 5 marker of ageing, is still scarce. We investigated the association of greenspace exposure with 6 TL in a sample of 200 preschool children (aged 5-7 years) residing in Sabzevar, Iran (2017). 7 We comprehensively characterized different aspects of greenspace exposure encompassing residential, kindergarten, and total (including both residential and kindergarten) surrounding 8 9 greenspace (using satellite-derived Normalized Difference Vegetation Index), residential and 10 kindergarten distance to green spaces, time spent in private gardens and public green spaces, 11 and the number of plant pots at home. Relative leukocyte TL (LTL) in blood samples of the study participants was measured using quantitative polymerase chain reaction (qPCR). We 12 13 applied mixed effects linear regression models with kindergarten and qPCR plate as random 14 effects, to estimate the association of indicators of greenspace exposure (one at a time) with LTL, controlled for relevant covariates. We observed an inverse association between distance 15 from home and kindergarten to green spaces larger than 5000m² and LTL. Moreover, higher 16 17 total surrounding greenspace at 300m and 500m buffers and higher surrounding greenspace at 300m buffer around kindergarten and home were associated with longer LTL. Furthermore, 18 19 longer time spent (h/week) in the public green spaces was associated with longer LTL. Our findings for residential and kindergarten distance to any green space (regardless of the size), 20 residential surrounding greenspace at 100m and 500m buffers, kindergarten surrounding 21 22 greenspace at 100m buffer, time spent in private gardens (h/week) and the number of plant pots at home were not conclusive. Our findings were generally suggestive for a positive 23

- 24 association between greenspace exposure and LTL in preschool children. More studies are
- 25 needed to confirm these findings in other settings with different climates and populations.

26

27 Keywords: Urbanization; Natural Environments; Ageing; Development; LMICs; Pediatric

28 **1. Introduction**

29 The on-going urbanization worldwide has led an increasing number of the global population, 30 including children, living in urban areas where residents often have less access to natural environments, including green spaces (UN-HABITAT 2016). Further to being more urbanized, 31 32 the global population is also ageing, particularly in urban areas. According to a recent United Nations report, between 2000 and 2015, the number of people aged 60 years or over increased by 33 69% in urban areas compared to 25% in rural areas (United Nations 2015). As a result, healthy 34 35 ageing is becoming a top public health priority worldwide. Early life, including pre- and postnatal periods and early childhood, is considered as an important window of susceptibility for 36 environmental exposures when the aging process has already started (Bijnens et al. 2017, Dadvand 37 et al. 2013). According to the Developmental Origins of Health and Disease (DOHaD) concept, 38 exposures during this window may permanently change the body's physiology, structure, and 39 40 metabolism and hence promote health or diseases (e.g., non-communicable diseases (NCDs)) in later stages of life (Barker 1997, Dietert et al. 2010, Gluckman and Hanson 2004). For example, 41 exposure to a hostile environment during this critical period of development and growth, has been 42 43 shown to induce a number of short- and long-term adaptation responses such as changes in endocrine or metabolic function, or in gene expression or differentiation that, in turn, could lead 44 45 to adverse health outcomes such as obesity, ischemic heart disease, hypertension, or diabetes 46 during the adult life (Mandy and Nyirenda, 2018). Accordingly, reducing exposure to 47 environmental risk factors and/or enhancing exposure to beneficial environmental factors during early life have been proposed to be one of the means to promote healthy ageing (United Nations 48 2015, Kuh et al. 2014, Martens and Nawrot 2018). Considering its beneficial health effects 49 50 (Nieuwenhuijsen et al. 2014, WHO Regional Office for Europe 2016), exposure to greenspace

during early life could therefore have important implications for promoting healthy ageing and in
the metropolitan area of Tehran (Iran) has already been considered as an alternative strategy for it
(Ahmadi et al., 2017).

54

Telomeres are specialized structures located at the ends of human chromosomes to protect their 55 56 integrity, block its degradation, provide genome stability, and avoid loss of genetic information (Lu et al. 2013). Telomere length (TL) is considered as a cellular marker of aging and its shortening 57 has been associated with a higher risk of developing different adverse health outcomes such as 58 59 cardiovascular diseases (Haycock et al., 2014), including myocardial infarction (Brouilette et al., 2003), hypertension (Benetos et al., 2001), atherosclerosis (Samani et al., 2001) or stroke (Ding et 60 al., 2012), and type 2 diabetes mellitus (Willeit et al., 2014). Various environmental and lifestyle 61 factors such as stress (Entringer, Epel, Kumsta, Lin, Hellhammer, Blackburn, Wüst, et al. 2011, 62 Mathur et al. 2016), physical inactivity (Mundstock et al. 2015, Arsenis et al. 2017, Tucker 2017), 63 64 social environment (Needham et al. 2012), air pollution (Miri et al. 2018), and noise have been associated with shorter TL (Martens and Nawrot 2018). In this context, exposure to greenspace, 65 could potentially result in less shortening of TL through reducing stress (Entringer, Epel, Kumsta, 66 67 Lin, Hellhammer, Blackburn, Wust, et al. 2011, Stigsdotter et al. 2010, Roe et al. 2013), fostering social interaction and social cohesion (Mitchell et al. 2014, Gascon et al. 2015), increasing physical 68 69 activity (Arsenis et al. 2017, Mundstock et al. 2015, Fanshawe et al. 2007), and reducing exposure 70 to urban-related environmental hazards such as air pollution (Dadvand et al. 2012, Dadvand et al. 71 2015a) and noise (Meillere et al. 2015, Dzhambov and Dimitrova 2014). However, the evidence 72 available on a potential impact of exposure to greenspace on TL is still very scarce (Bijnens et al. 73 2015), with no study available on such an impact in children. Moreover, to date, most of the

74	available studies on health effects of greenspace exposure have been conducted in high-income
75	countries and there is a paucity of such studies for low- and middle-income countries. This study
76	aimed to evaluate the association of exposure to greenspace with TL in preschool children in Iran.
77	
78	
79	2. Methods
80	2.1. Study setting
81	This study was conducted in Sabzevar, a town with a population of ~240,000 residents (2016)

located in the northeast of Iran (coordinates: 36°12′ N 57°35′, elevation: 977 m). Approximately
21,000 of the population are between 5 to 9 years old. Sabzevar has an arid climate with four
distinct seasons and an annual average rainfall of 176.8 mm. The highest precipitation is during
winter and most vegetation can be seen in spring. The annual mean relative humidity is 43%, and
the annual mean, minimum and maximum temperature in the city are 18, -2 and 45 °C respectively
(Khorasan Razavi Weather Center, http://www.razavimet.ir/fa/node/38). About 27% (7.1 km²) of
Sabzevar city is covered by green space resulting in a 29.4 m² of green space per capita.

89

90 **2.2.** Study population

The enrolment of the study participants (i.e., preschool children) was conducted through kindergartens during March 2017 to June 2017. Of over 80 kindergartens in Sabzevar, 60 kindergartens with about 900 children aged between 5 to 7 years were selected randomly, who accepted participation in our study. Briefing sessions were held with the children's parents in every kindergarten, and the research aims, steps and inclusion/exclusion criteria were fully explained to them. Finally, 200 participants from 27 kindergartens (seven children on average from each kindergarten) who met our inclusion criteria were enrolled in our study (participation rate of 22%)
(Figure 1). We included healthy children who did not constantly use supplements/vitamins and drugs, had no genetic disease, and were in kindergarten for at least one year and did not change their home since birth. Before the enrollment into the study, the children's parents/legal guardians signed the consent form approved by the Clinical Research Ethical Committee (IR.SSU.SPH.REC.1395.66) of the Shahid Sadoughi University of Medical Science, Yazd, Iran.

103

104

2.3. Exposure assessment

We carried out a comprehensive assessment of exposure by characterizing different aspects of exposure to greenspace including: a) residential as well as kindergarten distance from green spaces (any size as well as those larger than 5000 m²) as an indicator of *access to green spaces* (Ludlow, Mitchell, and Webster 2003), b) surrounding greenspace at residential as well as kindergarten address, c) time spent in private green spaces as well as public gardens (measured as hours per week), and d) the number of plant pots at participant's home. This strategy resulted in generating 16 exposure variables for each participant, as detailed below.

112

2.3.1. Access to green space

Distance from home address as well as kindergartens of participants to nearest green space (any green space) and distance to green space with area more than 5000 m² (major green space) (Ludlow, Mitchell, and Webster 2003) were calculated as surrogates of access to green space (Ludlow, Mitchell, and Webster 2003). Data needed for this purpose were calculated using land

use map of the study area (2015) prepared by the municipality of Sabzevar in ArcGIS v10.5software.

119

2.3.2. Surrounding greenspace

We applied satellite-derived Normalized Difference Vegetation Index (NDVI) to characterize surrounding greenspace separately for residential and kindergarten addresses. NDVI is commonly used as a numerical indicator of greenspace (i.e., photosynthetically active vegetation), which our previous study showed to be a valid measure to characterize green spaces (Gascon et al., 2016). The index depends on the visible and near-infrared light reflected by land surface and is calculated as (Weier and Herring 2011):

126
$$NDVI = (NIR - VIS)/(NIR + VIS)$$

where NIR and VIR are respectively land surface reflectance of near-infrared and red (visible)
parts of spectrum. It ranges from -1 to + 1, where 0 indicates no vegetation and + 1 means the
highest possible density of green leaves (Dadvand et al. 2017, Gascon et al. 2016). The NDVI for
the study area was calculated based on the Landsat images at 30m x 30m resolution obtained on
June 15, 2017 (Figure 1).

Greenspace surrounding residential and kindergarten address(es) of each participant was
abstracted as the average of NDVI in buffers of 100m, 300m and 500m around his/her geocoded
home address and kindergarten (Dadvand, de Nazelle, Figueras, et al. 2012).

Total surrounding greenspace index was calculated for each buffer (100, 300 and 500m) by
averaging NDVI values around home address and kindergartens separately for each buffer,
weighted by the daytime, which children spent at home (18 h) and kindergartens (6 h) (Dadvand
et al. 2015b).

139 2.3.3. Time spent in green spaces

Data on the amount of time spent in different types of green space was obtained through questionnaires answered by the parents (Dadvand, Poursafa, et al. 2018, Dadvand, Hariri, et al. 2018). The time spent in green space was asked separately for the time spent in private gardens such as yard garden, patio, etc (h/week), and the time spent in the public green spaces such as park, forest, etc (h/week)

145

2.3.4. Number of plant pots at home

146 The data on the number of plant pots at participant's home was collected through a questionnaire147 filled by parents.

148

149 **2.4.** Measurement of leukocyte telomere length

Two mL of blood were taken from each child in the morning at the reference lab of Sabzevar and transferred to laboratory in Vacutainer® Plus Plastic K2EDTA Tubes (USA) within less than two h and kept in -80 °C until analyses. The DNA was extracted from blood samples using DNA extraction kit (GeNet Bio, Korea) according to the manufacturer's instructions. The purity ratio (A260/230) and yield (A260/280) (ng/ μ L) of extracted DNA samples were determined using Nanodrop spectrophotometer (BIO INTELLECTICA Nano100, Canada). All DNA samples were stored at -80 °C until the time of analyses.

To measure LTL, we performed quantitative real-time polymerase chain reactions (qRT-PCR) based on the method by Cawthon which determines telomere copy number relative to single-gene (human beta-globin) copy number (T/S ratio) (Cawthon 2009, Cawthon 2002, Pieters, Janssen, and Dewitte 2016). This ratio is known as relative LTL. Q-RT-PCR was carried out using SYBR 161 green PCR master mix 2x (Amplicon, Denmark) on a CFX96 TouchTM Real-Time PCR Detection USA) with the following forward 5'-162 System (Bio-Rad, primers: telomere CGGTTTGTTTGGGTTTGGGTTTGGGTTTGGGTTTGGGTT-3', 5'-163 telomere reverse GGCTTGCCTTACCCTTACCCTTACCCTTACCCT-3', single-copy gene forward 5'-164 5'-GCTTCTGACACAACTGTGTTCACTAGC-3', 165 and single-copy gene reverse 166 CACCAACTTCATCCACGTTCACC-3'. Telomere and single-copy numbers were evaluated in triplicate (3 measured per individual) in a total of 10 plates. Each reaction for telomere evaluation 167 contained 5 µL of SYBR green PCR master mix 2x, 0.1 µL of forward primer (10 pmol/µL), 0.9 168 169 μ L of reverse primer (10 pmol/ μ L) and 1 μ L (25 ng/ μ L) of DNA. Single-copy gene (SCG) (SCG) was performed as the telomere reaction, with the exception that we added 0.3 μ L (10 pmol/ μ L) of 170 forward primer and 0.7 μ L (10 pmol/ μ L) of reverse primer. The cycling thermal profile was the 171 same for telomere and single-copy gene including one cycle at 95°C for 15 min, followed by 50 172 cycles of 95°C for 15 seconds, 60 °C for 20 seconds, and 72°C for 20 seconds. Each plate contained 173 174 5 serial concentrations of a standard human genomic DNA sample (150 ng, 50 ng, 16.7 ng, 5.55 ng and 1.85 ng per well), giving amplification efficiencies ranging from 95 to 105%. Melting curve 175 analyses were used at the end of each cycle to confirm amplification specificity and absence of 176 177 primer dimers. Finally, the relative T/S ratio was calculated as follow (Cawthon 2009, Cawthon 2002): 178

179
$$Relative \ T/S = \frac{2^{-(Ct_{TL}-Ct_{SCG})sample}}{2^{-(Ct_{TL}-Ct_{SCG})control}}$$

Where Ct_{TL} and Ct_{SCG} were the average Ct of each sample and single copy gene, respectively. In this study, the average value of T/S ratio of samples (the numerator) was used as control (the denominator).

183	To provide reproducibility, inter-assay variability was measured in three different days. The % of
184	coefficient of variation (%CV) was 0.74% for LTL and 47% for single-copy gene. LTL is
185	expressed in relative units as the ratio of telomere copy number proportional to single-copy gene
186	number (T/S) relative to the average telomere length in the entire sample set (T/S ratio).
187	
188	2.5. Statistical analyses
189	
190	2.5.1. Main analyses
191	Considering the multilevel nature of our data, we developed linear mixed effect models with
192	relative LTL as the outcome, indicators of greenspace exposure (one at a time) as fixed effect
193	predictor and the kindergartens and qPCR plates as random effects. The models were further
194	adjusted for a priori selected set of variables: age (continuous), sex (girl/boy), body mass index
195	(BMI, continuous) (Gielen et al. 2018), exposure to environmental tobacco smoke at home
196	(yes/no) (Theall et al. 2013) and two indicators of household socioeconomic status (SES) as well
197	as two indicators of the neighborhood SES (Needham et al. 2012). Parental education as the highest

education level between paternal and maternal education of the participant (no education/ primary school, secondary school, or university) and income (<15 million and >15 million Rials) were used as indicators of household SES, while the percentages of illiterate adults as well as unemployment in the census tract based on the last census in Iran (2016) were applied as indicators of the neighborhood SES. The associations were reported for each interquartile range (IQR) increase in NDVI, distance from green space areas, time spent in the green space area, and number of flowerpots based on all study participants. Consistent with previous studies of the environmental

205	determinants of TL (Miri et al., 2019; Zhao et al., 2018), relative LTL was transformed using its
206	natural logarithm to assure the normal distribution of the residuals of our models. Consequently,
207	to enhance the interpretation of the regression models we expressed all β coefficients and 95%
208	confidence intervals (CIs) as the percentage increase (or decrease) in LTL for each IQR increase
209	in the exposure variables. The percent change in LTL was calculated as (e ^(β) -1) × 100%, and 95%
210	CI were calculated as $(e^{[(\beta\pm 1.96\times SE)]}-1) \times 100\%$, in which SE is the estimated standard error and β is
211	the estimated regression coefficient. Stata 14 (StataCorp L.P., College Station, TX) has been used
212	for all statistical analyses.

213

2.5.2. Sensitivity analyses 214

215 The models were further adjusted for age squared (continuous), car ownership (yes/no) and home ownership (yes/no), and also for environmental tobacco smoke in other places than the home 216 (yes/no). 217

218

219 2.5.3. Stratified analyses

Main analyses were stratified by sex (girl/boy), parental education (no education/primary school, 220 221 secondary school, or university) and percentage of unemployment in the census tract in tertiles. The associations were reported per IQR increase in each greenspace indicator based on participants 222 223 in each stratum.

224

2.5.4. Mediation analyses 225

226 We explored the mediation role of air pollution in the association between total surrounding greenspace (including both residential and kindergarten surrounding greenspace) and telomere 227 length, for which we could hypothesize such a mediation. Our previous study based on the same 228 sample of schoolchildren demonstrated a negative association between exposure to PM_{2.5} and LTL 229 (Moslem et al., 2020). Estimated PM_{2.5} levels (for detailed information regarding exposure 230 231 assessment, see the Supporting information in page 41 Supplemental Materials) at home address 232 and kindergarten were used to develop total $PM_{2.5}$ exposure as the annual average of $PM_{2.5}$ levels 233 at home and kindergarten, weighted by the time the child would spend at home (18 h) and 234 kindergarten (6 h). Following the approach of Baron and Kenny (Baron and Kenny 1986) for mediation analyses, we fitted separately a model for the outcome given the mediator and the 235 exposure, and a model for the mediator given the exposure to compute the percentage of mediation. 236 Kindergartens and qPCR plates were considered as random effects in the model for the outcome. 237 Bootstrap was used to obtain percentile-based 95% confidence intervals for the different estimates. 238

239

3. Results

241

242 **3.1.** Study population

Descriptive statistics of the sociodemographic characteristics and exposure variables of the study participants are presented in Table 1. From 200 participants included in the study, 89 (44.5%) were girls and 111 (55.5%) were boys. The median (IQR) age of participants was 7 (1) years. Most of the participants had either the father or the mother with high school education (57%) and 13% of children had exposure to environmental tobacco smoke at home (Table 1). The median (IQR) T/S ratio of children was 0.83 (0.7) with larger T/S ratio for girls compared to boys (0.89 vs. 0.79)
(Supplemental materials, Table S1). A graphical display of the correlation matrix between
greenspace indicators was developed (Figure 2). In our study NDVI values of different buffers
around home and kindergarten as well as total surrounding greenspace were positively correlated.
As expected, distances to any green spaces and larger than 5000 m² were negatively correlated
with NDVI indicators and also with time spent mainly in public green spaces.

254

255 **3.2.** Main analyses

256 Percent changes (95% CIs) in LTL in relation to the indicators of greenspace exposure are presented in Table 2, in which the IQR for each exposure variable is also displayed. In fully 257 adjusted models, we observed that an IQR increase in distance from home and as well as 258 kindergarten to green spaces larger than 5000 m² was associated with a 20.5% (95% C = -29.9%, 259 -9.7%) and 21.8% (95% CI = -31.4%, -10.8%) decrease in LTL (Table 2). For the distance to any 260 green space regardless of its size we did not observe any association with LTL. As presented in 261 Table 2, we observed that an IQR increase in greenspace around home and kindergarten as well as 262 total surrounding greenspace in a 300m buffer, was associated with a 8.3% (95% CI = 1.2%, 263 264 15.9%), 13.3% (95% CI = 2.2%, 25.7%) and a 12.0% (95% CI = 3.1%, 21.7%) increase in LTL, respectively. We also observed that an IQR increase in greenspace around kindergarten and total 265 266 surrounding greenspace in a 500m buffer was related with a 14.5% (95% CI = 0.4%, 28.4%) and 267 an 8.4% (95% CI = 0.3%, 17.1%) increase in LTL, respectively. Similarly, we observed positive associations for surrounding greenspace in a 100m buffer around home and kindergarten and in a 268 500m buffer around home with LTL. Spending more time in the public green spaces (park, forest, 269

etc.) was associated with a 39.0% (95% CI = 22.8%, 57.3%) increase in LTL. Longer time spent
in private gardens and more plant pots at home did not show any association with LTL (Table 2)."

273

274

3.3. Sensitivity analyses

The results of the models further adjusted for age squared, car ownership and home ownership were generally similar to the main analyses in terms of strength of the association and direction (Supplemental materials, Table S2). Moreover, after further adjustment of analyses for the exposure to environmental tobacco smoke in other places than the home, we did not observe any notable change in our findings either in terms of direction and strength of the associations.

280

3.4. Stratified analyses

The results of the stratified analyses by sex and SES are shown in table S3 and S4 in Supplemental materials. In the sex-stratified analyses for the residential (100m buffer), kindergarten (300m buffer) and total surrounding (100m buffer) greenspace, we observed positive associations ranging from a 17% to a 26% increase in LTL among girls and null for boys. For the other indicators of greenspace exposure, the associations for boys were also null (Table S3) except the time spent in public greenspace and residential distance to greenspaces for which we observed similar associations for boys and girls.

288

In the stratified analyses based on socioeconomic status (see Supplemental Material, Table S4), we observed some suggestions for a potential trend across the strata of parental education with stronger associations for children with higher parental education. Considering the indicator of neighborhood SES (% of unemployment in the census tract), the stratified analyses in tertiles suggested that stronger associations could be shown in the 1st tertile, which corresponds to the lower percentage of unemployment. In addition, interaction terms suggested a stronger association in the 1st tertile of % unemployment for surrounding greenspace around kindergarten (300m), and for public green space they suggested a higher and positive association in children whose parents had higher education level; however, all p-values for the interaction terms were more than 0.01.

298

3.5. Mediation Analyses

Our mediation analysis showed that 30.0% (95% CIs: -1.71, 3.12) of the association between total
surrounding (300m buffer) telomere length could be mediated through exposure to PM_{2.5}.

301

302 **4. Discussion**

To our knowledge, this is the first study on the association between greenspace exposure and TL 303 in children. Moreover, our study provides new evidence regarding the health effects of green 304 spaces in children from low and middle-income countries. Our study benefitted from a 305 306 comprehensive assessment of greenspace exposure, including residential and kindergarten surrounding greenspace and distance to green spaces as well as time spent in private and public 307 308 green spaces and the number of plant pots at home. We found that more greenspace surrounding home (300m buffer), kindergarten (300m and 500m buffer), and combination of these two (i.e., 309 310 total surrounding greenspace at 300m and 500m buffer), less distance to green spaces larger than 311 5000 m^2 from home and kindergarten, and longer time spent in public green spaces were positively associated with LTL. For the residential and kindergarten surrounding greenspace in other buffers, 312

313 distance to any green space (regardless of its size) and the number of plant pots at home, we did not observe any association. We observed some suggestions for stronger associations for girls, for 314 315 those participants with parents having higher educational levels and living in neighborhoods with lower percentage of unemployment; however, these patterns were not consistent for all indicators 316 of greenspace exposure. Moreover, our stratified analyses were likely under-powered, and this 317 318 issue needs to be considered when interpreting our findings for these analyses. We also found that about one-third of our observed association between total surrounding greenspace and LTL could 319 320 be explained by reduction in exposure to air pollution ($PM_{2.5}$).

321

4.1. Interpretation of the findings

We are not aware of any study on the association between greenspace exposure and TL in children; 322 323 therefore, it is not possible to compare our findings with those of others. However, our findings 324 are in line with some previous observations and they might suggest that exposure to greenspace can potentially result in less shortening of LTL, which could promote healthy aging by reducing 325 326 the risk of developing different adverse health outcomes that have been already associated with 327 shorter telomeres. So far, only two studies have evaluated the association between greenspace exposure and TL (Woo et al. 2009, Bijnens et al. 2015). Bijnens et al. assessed TL from placental 328 329 tissues of 211 twins and showed that maternal residential surrounding greenspace (measured using NDVI) during pregnancy was positively associated with TL (Bijnens et al. 2015). In a cohort study 330 331 of 976 men aged 65 and over, Woo et al. found shorter TL in participants living in zones with 332 lower green spaces presence (Woo et al. 2009). It is worthy to mention that in our study, the 333 greenspace surrounding home and kindergarten for each participant were not strongly correlated, 334 suggesting that our observed associations between these indicators and TL were not dependent on each other. 335

We observed an inverse association between the distance to a green space larger than 5000m² and 337 LTL, but not for any green space regardless of its size. Previous studies have reported that larger 338 339 green spaces are more likely to be used for physical activity (McCormack et al. 2010) and mitigate air pollution, which in turn, have been associated with longer LTL (Arsenis et al. 2017, Martens 340 and Nawrot 2018). In this context, we also found some suggestions for stronger associations for 341 342 measures of surrounding greenspace in larger buffer sizes. This is consistent with previous evidence in which the associations between green spaces indicators in larger buffers and health 343 outcomes were stronger (Su et al. 2019) and more relevant to physical activity levels(McMorris 344 345 et al. 2015; Borwning and Lee 2017). Previous studies have reported that more time spent in green spaces is associated with lower stress, better mental health, and improved social contacts (Beyer 346 et al. 2014, Amoly et al. 2014, Dadvand, Hariri, et al. 2018), which, in turn, could be associated 347 348 with longer LTL (Mathur et al. 2016, Starkweather et al. 2014, Costa et al. 2015). These findings are in line with our results in which, more time spent in public green spaces was associated with 349 longer LTL. Even though our study is the first to evaluate the sex- and SES-stratified associations 350 between greenspace exposure and LTL, previous literature has shown that health effects of green 351 spaces could differ between sex and SES. Our stratified analyses did not show any effect 352 353 modifications by sex or SES, but the results were suggestive for stronger associations in girls compared to boys and in higher SES. A number of studies have observed that the relationships 354 between green spaces exposure and health outcomes could be stronger in girls (Annerstedt et al. 355 356 2012, Roe et al. 2013), which goes in line with our findings. However, the evidence is not consistent as some studies are not supportive for these associations or showed an inverse 357 relationship with TL, stronger in boys (Ruijsbroek et al. 2017, Richardson et al. 2017). Regarding 358

socioeconomic status, different studies have evaluated the modification of the health benefits of greenspace exposure by SES, however is still unclear its role. In some cases, stronger relationships have been found for participants with lower SES (Maas et al. 2006, McEachan et al. 2016), and in other studies non-statistically significant associations were observed (Casey et al. 2016). Therefore, it might be essential to continue considering the effect modification analyses by sex and SES when evaluating the health effects of greenspace exposure, as it seems that the available evidence is still inconsistent.

366

4.2. Biological plausibility

The mechanisms underlying health effects of green spaces are yet to be established. Some of the 367 possible mechanisms suggested are promoting social contacts, inducing psychological 368 restoration/stress reduction, improving functioning of the immune system due to microbial input, 369 promoting physical activity and reducing obesity, and mitigating urban-related environmental 370 371 hazards including air pollution, noise, and heat (Nieuwenhuijsen et al. 2014, WHO Regional Office for Europe 2016). Some of these mechanisms could also be relevant for TL. Higher levels 372 of physical activity and lower levels of obesity have been associated with reduced inflammation 373 and oxidative stress and enhanced restoration of TL (Mundstock et al. 2015). Conversely, 374 psychological and life stress (Law et al. 2016, Epel et al. 2004), noise exposure (Meillere et al. 375 2015), and air pollution (Martens and Nawrot 2018, Miri et al. 2018) have been associated with 376 shorter TL. The results of the mediation analyses showed that air pollution could explain about 377 one-third of our observed association between total surrounding greenspace and LTL. This finding 378 379 is in line with previous evidence that linked the existence of green spaces with a reduction of PM2.5 concentrations (Chen et al. 2016, Dadvand, de Nazelle, Triguero-Mas, et al. 2012, 380

381 Dadvand et al. 2015b). At the same time, a recent systematic review and meta-analysis of the 382 available evidence showed that higher long-term exposure to PM2.5 was associated with 383 shortened telomeres (Miri et al. 2018). Such an exposure has been hypothesized to induce TL shortening through reactive oxygen species (ROS) mediated telomeric DNA damage and 384 inflammation (Kordinas, Ioannidis, and Chatzipanagiotou 2016, Miri et al. 2018). Additionally, it 385 386 has been reported that accelerated telomere shortening could be associated with different clinical outcomes such as cancer, cardiovascular diseases or diabetes (Haycock et al., 2014; Willeit et al., 387 2014), which are also associated with exposure to high levels of PM2.5 over a long period of time 388 (Cao et al., 2018; Hayes et al., 2020; Lao et al., 2019). 389

390

4.3. Limitations

391 Our study should be interpreted in the context of its limitations. Firstly, our measures of proximity to green spaces did not address the quality of these spaces, while quality aspects such as safety, 392 393 aesthetics, biodiversity, facilities and amenities could have had implications for our analyses. 394 Additionally, to develop the total surrounding greenspace, we assumed that the children on 395 average spent 6 hours at kindergarten and 18 hours at home. Such an assumption was based on the average of working hours of kindergartens in our study area and not collected data from our 396 participants. Moreover, by only including greenspace surrounding home and kindergarten to 397 398 develop the total surrounding greenspace, we have effectively overlooked exposures that could have happened in other potentially important microenvironments such as homes of friends and 399 400 relatives, commuting routes to and from home, or public places that could have been visited 401 frequently by participants and their families. Furthermore, our exposure assessment through NDVI data did not differentiate the type of vegetation or land cover, which could have influenced 402 403 our observed associations for the greenspace surrounding homes and kindergartens. Additionally,

404 our satellite data for the greenspace indicators around home and kindergarten was at 30m x 30m resolution and by using images with higher resolution, we might have been able to obtain a more 405 406 precise assessment of exposure; however, such a higher resolution was less likely to result in a notable change in our observed associations (Su et al., 2019). Moreover, our data on time spent in 407 green spaces was obtained through questionnaires, which could have resulted in exposure 408 409 measurement error. Moreover, our measure of exposure to indoor plant was merely based on the number of plant pots home, while lacking information on important determinants of this exposure 410 411 such as visibility of the plants for the children (given their height), species of the plants (which 412 could have been relevant, among others, to their ability to reduce air pollution), or the placement of plants at home (e.g. whether they were located in places where the child spent much of her/his 413 time). The unavailability of such detailed information might have been the reason for our null 414 findings for the indoor plants. Additionally, in our study we did not cover greenspace exposure 415 during prenatal period as we did not have data on the maternal residential address during this 416 417 period or their potential mobility during pregnancy. The potentially important impact of prenatal greenspace exposure on TL remains as an open question for the future studies. Lastly, our study 418 419 did not cover exposure to other air pollutants such as ozone, that could have a key mediator role 420 in the pathway between greenspace exposure and LTL as the observed with PM2.5. In addition, we did not collect data on physical activity, which, could act as another mediator, given that green 421 422 spaces have been reported, although inconsistently, to enhance physical activity (McCrorie et al., 423 2014) and, at the same time, physical activity has been suggested to induce potential protection 424 against the shortening of TL (Arsenis et al., 2017)."

426 **5.** Conclusions

We observed a positive association of residential/kindergarten greenspace, as well as time spent 427 428 in public green spaces and inverse association of distance to major green spaces with LTL in a sample of Iranian preschool children. Given the rapid urbanization coupled with ageing population 429 worldwide, healthy ageing has become a top public health priority worldwide. Our findings, if 430 confirmed by future studies, could provide policymakers with evidence base to implement targeted 431 interventions to decelerate ageing process early in life. Further research is warranted to evaluate 432 433 these associations using longitudinal design based on repeated measurements of TL in other 434 settings with different climates and genetic backgrounds. These studies are also recommended to disentangle the effects of pre- and postnatal greenspace exposures and the role of air pollution. 435

437 **References**

439 Amoly, E, P Dadvand, J Forns, M López-Vicente, X Basagaña, J Julvez, M Alvarez-Pedrerol, MJ Nieuwenhuijsen, and J Sunyer. 2014. "Green and blue spaces and Behavioural 440 Development in Barcelona Schoolchildren: The BREATHE Project." Environ Health 441 Perspect 122:1351-8. 442 Annerstedt, Matilda, Per-Olof Östergren, Jonas Björk, Patrik Grahn, Erik Skärbäck, and Peter 443 Währborg. 2012. "Green qualities in the neighbourhood and mental health – results from 444 a longitudinal cohort study in Southern Sweden." BMC Public Health 12 (1):337. doi: 445 10.1186/1471-2458-12-337. 446 Arsenis, N. C., T. You, E. F. Ogawa, G. M. Tinsley, and L. Zuo. 2017. "Physical activity and 447 448 telomere length: Impact of aging and potential mechanisms of action." Oncotarget 8 (27):45008-45019. doi: 10.18632/oncotarget.16726. 449 Barker, D. J. 1997. "The fetal origins of coronary heart disease." Acta Paediatr Suppl 422:78-450 451 82. 452 Baron, Reuben M., and David A. Kenny. 1986. "The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations." 453 454 Journal of Personality and Social Psychology 51 (6):1173-1182. doi: 10.1037/0022-3514.51.6.1173. 455 Benetos, A., K. Okuda, M. Lajemi, M. Kimura, F. Thomas, J. Skurnick, C. Labat, K. Bean, and 456 A. Aviv. 2001. "Telomere length as an indicator of biological aging: the gender effect 457 and relation with pulse pressure and pulse wave velocity." Hypertension 37 (2 Pt 2):381-458 5. 459 460 Beyer, Kirsten M. M., Andrea Kaltenbach, Aniko Szabo, Sandra Bogar, F. Javier Nieto, and Kristen M. Malecki. 2014. "Exposure to neighborhood green space and mental health: 461 evidence from the survey of the health of Wisconsin." International journal of 462 environmental research and public health 11 (3):3453-3472. doi: 463 464 10.3390/ijerph110303453. Bijnens, E. M., M. P. Zeegers, C. Derom, D. S. Martens, M. Gielen, G. J. Hageman, M. 465 Plusquin, E. Thiery, R. Vlietinck, and T. S. Nawrot. 2017. "Telomere tracking from birth 466

to adulthood and residential traffic exposure." *BMC Med* 15 (1):205. doi:

468 10.1186/s12916-017-0964-8.

- Bijnens, E., M. P. Zeegers, M. Gielen, M. Kicinski, G. J. Hageman, D. Pachen, C. Derom, R.
- Vlietinck, and T. S. Nawrot. 2015. "Lower placental telomere length may be attributed to
 maternal residential traffic exposure; a twin study." *Environ Int* 79:1-7. doi:
- 472 10.1016/j.envint.2015.02.008.
- Brouilette, S., R. K. Singh, J. R. Thompson, A. H. Goodall, and N. J. Samani. 2003. "White cell
 telomere length and risk of premature myocardial infarction." *Arterioscler Thromb Vasc Biol* 23 (5):842-6. doi: 10.1161/01.atv.0000067426.96344.32.
- 476 Casey, Joan A., Peter James, Kara E. Rudolph, Chih-Da Wu, and Brian S. Schwartz. 2016.

477 "Greenness and Birth Outcomes in a Range of Pennsylvania Communities."

- 478 International journal of environmental research and public health 13 (3):311. doi:
- 479 10.3390/ijerph13030311.
- Cawthon, R. M. 2009. "Telomere length measurement by a novel monochrome multiplex
 quantitative PCR method." *Nucleic Acids Res* 37 (3):e21. doi: 10.1093/nar/gkn1027.
- 482 Cawthon, Richard M. 2002. "Telomere measurement by quantitative PCR." *Nucleic Acids Res*483 30 (10):e47-e47.
- Chen, Jiquan, Liuyan Zhu, Peilei Fan, Li Tian, and Raffaele Lafortezza. 2016. "Do green spaces
 affect the spatiotemporal changes of PM(2.5) in Nanjing?" *Ecological processes* 5:7-7.
 doi: 10.1186/s13717-016-0052-6.
- 487 Costa, Danielle de Souza, Daniela Valadão Freitas Rosa, Alexandre Guimarães Almeida Barros,
 488 Marco Aurélio Romano-Silva, Leandro Fernandes Malloy-Diniz, Paulo Mattos, and
- 489 Débora Marques de Miranda. 2015. "Telomere length is highly inherited and associated
- 490 with hyperactivity-impulsivity in children with attention deficit/hyperactivity disorder."
- 491 *Front Mol Neurosci* 8:28. doi: 10.3389/fnmol.2015.00028.
- 492 Dadvand , P. , A. de Nazelle, Margarita Triguero-Mas, Anna Schembari, Marta Cirach, Elmira
 493 Amoly, Francesc Figueras, Xavier Basagaña, Bart Ostro, and Mark Nieuwenhuijsen.
- 494 2012. "Surrounding Greenness and Exposure to Air Pollution During Pregnancy: An
- 495 Analysis of Personal Monitoring Data " *Environ Health Perspect* 120 (9):1286-90.

- 496 Dadvand, P., A. de Nazelle, F. Figueras, X. Basagana, J. Su, E. Amoly, M. Jerrett, M. Vrijheid,
- J. Sunyer, and M. J. Nieuwenhuijsen. 2012. "Green space, health inequality and
 pregnancy." *Environ Int* 40:110-115. doi: 10.1016/j.envint.2011.07.004.
- Dadvand, P., S. Hariri, B. Abbasi, R. Heshmat, M. Qorbani, M. E. Motlagh, X. Basagana, and R.
 Kelishadi. 2018. "Use of green spaces, self-satisfaction and social contacts in
- adolescents: A population-based CASPIAN-V study." *Environ Res* 168:171-177. doi:
 10.1016/j.envres.2018.09.033.
- Dadvand, P., J. Parker, M. L. Bell, M. Bonzini, M. Brauer, L. A. Darrow, U. Gehring, S. V.
 Glinianaia, N. Gouveia, E. H. Ha, J. H. Leem, E. H. van den Hooven, B. Jalaludin, B. M.
- Jesdale, J. Lepeule, R. Morello-Frosch, G. G. Morgan, A. C. Pesatori, F. H. Pierik, T.
- 506 Pless-Mulloli, D. Q. Rich, S. Sathyanarayana, J. Seo, R. Slama, M. Strickland, L.
- 507 Tamburic, D. Wartenberg, M. J. Nieuwenhuijsen, and T. J. Woodruff. 2013. "Maternal
- 508 exposure to particulate air pollution and term birth weight: a multi-country evaluation of
- 509 effect and heterogeneity." *Environ Health Perspect* 121 (3):267-373. doi:
- 510 10.1289/ehp.1205575.
- 511 Dadvand, P., P. Poursafa, R. Heshmat, M. E. Motlagh, M. Qorbani, X. Basagana, and R.
- 512 Kelishadi. 2018. "Use of green spaces and blood glucose in children; a population-based
- 513 CASPIAN-V study." *Environ Pollut* 243 (Pt B):1134-1140. doi:
- 514 10.1016/j.envpol.2018.09.094.
- 515 Dadvand, P., I. Rivas, X. Basagaña, M. Alvarez-Pedrerol, J. Su, M. De Castro Pascual, F.
- 516 Amato, M. Jerret, X. Querol, J. Sunyer, and MJ. Nieuwenhuijsen. 2015a. "The
- Association between Greenness and Traffic-related Air Pollution at Schools." *Sci Total Environ* 523:59-63.
- 519 Dadvand, P., C. Tischer, M. Estarlich, S. Llop, A. Dalmau-Bueno, M. Lopez-Vicente, A.
- 520 Valentin, C. de Keijzer, A. Fernandez-Somoano, N. Lertxundi, C. Rodriguez-Dehli, M.
- 521 Gascon, M. Guxens, D. Zugna, X. Basagana, M. J. Nieuwenhuijsen, J. Ibarluzea, F.
- 522 Ballester, and J. Sunyer. 2017. "Lifelong Residential Exposure to Green Space and
- 523 Attention: A Population-based Prospective Study." *Environ Health Perspect* 125
- 524 (9):097016. doi: 10.1289/ehp694.
- 525 Dadvand, Payam, Audrey de Nazelle, Margarita Triguero-Mas, Anna Schembari, Marta Cirach,
 526 Elmira Amoly, Francesc Figueras, Xavier Basagaña, Bart Ostro, and Mark

- 527 Nieuwenhuijsen. 2012. "Surrounding greenness and exposure to air pollution during
- pregnancy: an analysis of personal monitoring data." *Environmental health perspectives*120 (9):1286-1290. doi: 10.1289/ehp.1104609.
- 530 Dadvand, Payam, Ioar Rivas, Xavier Basagaña, Mar Alvarez-Pedrerol, Jason Su, Montserrat De
- 531 Castro Pascual, Fulvio Amato, Michael Jerret, Xavier Querol, Jordi Sunyer, and Mark J.
- 532 Nieuwenhuijsen. 2015b. "The association between greenness and traffic-related air
- pollution at schools." *Science of The Total Environment* 523 (Supplement C):59-63. doi:
 https://doi.org/10.1016/j.scitotenv.2015.03.103.
- Dietert, R. R., J. C. DeWitt, D. R. Germolec, and J. T. Zelikoff. 2010. "Breaking patterns of
 environmentally influenced disease for health risk reduction: immune perspectives."
 Environ Health Perspect 118 (8):1091-9. doi: 10.1289/ehp.1001971.
- 537 *Environ Health Perspect* 118 (8):1091-9. doi: 10.1289/ehp.1001971.
 538 Ding, H., C. Chen, J. R. Shaffer, L. Liu, Y. Xu, X. Wang, R. Hui, and D. W. Wang. 2012.
- 539 "Telomere length and risk of stroke in Chinese." *Stroke* 43 (3):658-63. doi:
- 540 10.1161/strokeaha.111.637207.
- 541 Dzhambov, A. M., and D. D. Dimitrova. 2014. "Urban green spaces' effectiveness as a
 542 psychological buffer for the negative health impact of noise pollution: a systematic
 543 review." *Noise Health* 16 (70):157-65. doi: 10.4103/1463-1741.134916.
- 544 Entringer, S., E. S. Epel, R. Kumsta, J. Lin, D. H. Hellhammer, E. H. Blackburn, S. Wust, and P.
- 545 D. Wadhwa. 2011. "Stress exposure in intrauterine life is associated with shorter telomere 546 length in young adulthood." *Proc Natl Acad Sci U S A* 108 (33):E513-8. doi:
- 547 10.1073/pnas.1107759108.
- Entringer, Sonja, Elissa S. Epel, Robert Kumsta, Jue Lin, Dirk H. Hellhammer, Elizabeth H.
 Blackburn, Stefan Wüst, and Pathik D. Wadhwa. 2011. "Stress exposure in intrauterine
 life is associated with shorter telomere length in young adulthood." *Proc Natl Acad Sci USA* 108 (33):E513-E518. doi: 10.1073/pnas.1107759108.
- 552 Epel, E. S., E. H. Blackburn, J. Lin, F. S. Dhabhar, N. E. Adler, J. D. Morrow, and R. M.
- Cawthon. 2004. "Accelerated telomere shortening in response to life stress." *Proc Natl Acad Sci U S A* 101 (49):17312-5. doi: 10.1073/pnas.0407162101.
- Fanshawe, T. R., P. J. Diggle, S. Rushton, R. Sanderson, P. W. W. Lurz, S. V. Glinianaia, M. S.
 Pearce, L. Parker, M. Charlton, and T. Pless-Mulloli. 2007. "Modelling spatio-temporal

- variation in exposure to particulate matter: a two-stage approach." *Environmetrics* 19
 (6):549-566.
- Gascon, Mireia, Marta Cirach, David Martínez, Payam Dadvand, Antònia Valentín, Antoni
 Plasència, and Mark J Nieuwenhuijsen. 2016. "Normalized difference vegetation index
 (NDVI) as a marker of surrounding greenness in epidemiological studies: The case of
 Barcelona city." Urban Forestry & Urban Greening 19:88-94.
- Gascon, Mireia, Margarita Triguero-Mas, David Martínez, Payam Dadvand, Joan Forns, Antoni
 Plasància, and Mark Nieuwenhuijsen. 2015. "Mental Health Benefits of Long-Term
 Exposure to Residential Green and Blue Spaces: A Systematic Review." *Int J Environ Res Public Health* 12 (4):4354-4379.
- Gielen, Marij, Geja J. Hageman, Evangelia E. Antoniou, Katarina Nordfjall, Massimo Mangino, 567 Muthuswamy Balasubramanyam, Tim de Meyer, Audrey E. Hendricks, Erik J. Giltay, 568 Steven C. Hunt, Jennifer A. Nettleton, Klelia D. Salpea, Vanessa A. Diaz, Ramin 569 Farzaneh-Far, Gil Atzmon, Sarah E. Harris, Lifang Hou, David Gilley, Iiris Hovatta, 570 Jeremy D. Kark, Hisham Nassar, David J. Kurz, Karen A. Mather, Peter Willeit, Yun-571 572 Ling Zheng, Sofia Pavanello, Ellen W. Demerath, Line Rode, Daniel Bunout, Andrew Steptoe, Lisa Boardman, Amelia Marti, Belinda Needham, Wei Zheng, Rosalind 573 574 Ramsey-Goldman, Andrew J. Pellatt, Jaakko Kaprio, Jonathan N. Hofmann, Christian Gieger, Giuseppe Paolisso, Jacob B. H. Hjelmborg, Lisa Mirabello, Teresa Seeman, 575 576 Jason Wong, Pim van der Harst, Linda Broer, Florian Kronenberg, Barbara Kollerits, Timo Strandberg, Dan T. A. Eisenberg, Catherine Duggan, Josine E. Verhoeven, 577 578 Roxanne Schaakxs, Raffaela Zannolli, Rosana M. R. dos Reis, Fadi J. Charchar, Maciej Tomaszewski, Ute Mons, Ilja Demuth, Andrea Elena Iglesias Molli, Guo Cheng, Dmytro 579 580 Krasnienkov, Bianca D'Antono, Marek Kasielski, Barry J. McDonnell, Richard Paul Ebstein, Kristina Sundquist, Guillaume Pare, Michael Chong, and Maurice P. Zeegers. 581 582 2018. "Body mass index is negatively associated with telomere length: a collaborative cross-sectional meta-analysis of 87 observational studies." The American Journal of 583 584 Clinical Nutrition 108 (3):453-475. doi: 10.1093/ajcn/nqy107. 585 Gluckman, Peter D., and Mark A. Hanson. 2004. "Living with the past: evolution, development,
 - and patterns of disease." Science 305 (5691):1733-1736.

- 587 Haycock, P. C., E. E. Heydon, S. Kaptoge, A. S. Butterworth, A. Thompson, and P. Willeit.
- 2014. "Leucocyte telomere length and risk of cardiovascular disease: systematic review
 and meta-analysis." *Bmj* 349:g4227. doi: 10.1136/bmj.g4227.
- 590 Kordinas, Vasileios, Anastasios Ioannidis, and Stylianos Chatzipanagiotou. 2016. "The
- 591 Telomere/Telomerase System in Chronic Inflammatory Diseases. Cause or Effect?"
 592 *Genes* 7 (9):60. doi: 10.3390/genes7090060.
- Kuh, Diana, Sathya Karunananthan, Howard Bergman, and Rachel Cooper. 2014. "A life-course
 approach to healthy ageing: maintaining physical capability." *The Proceedings of the Nutrition Society* 73 (2):237-248. doi: 10.1017/S0029665113003923.
- Law, E., A. Girgis, L. Sylvie, J. Levesque, and H. Pickett. 2016. "Telomeres and Stress:
- 597 Promising Avenues for Research in Psycho-Oncology." *Asia Pac J Oncol Nurs* 3
 598 (2):137-147. doi: 10.4103/2347-5625.182931.
- Lu, W., Y. Zhang, D. Liu, Z. Songyang, and M. Wan. 2013. "Telomeres-structure, function, and
 regulation." *Exp Cell Res* 319 (2):133-41. doi: 10.1016/j.yexcr.2012.09.005.
- Ludlow, David, Clare Mitchell, and M Webster. 2003. "European common indicators: towards a
 local sustainability profile."
- Maas, Jolanda, Robert A. Verheij, Peter P. Groenewegen, Sjerp De Vries, and Peter
- Spreeuwenberg. 2006. "Green space, urbanity, and health: how strong is the relation?" J
 Epidemiol Community Health 60 (7):587-592.
- Martens, D. S., and T. S. Nawrot. 2018. "Ageing at the level of telomeres in association to
 residential landscape and air pollution at home and work: a review of the current
 evidence." *Toxicol Lett.* doi: 10.1016/j.toxlet.2018.06.1213.
- Mathur, Maya B., Elissa Epel, Shelley Kind, Manisha Desai, Christine G. Parks, Dale P. Sandler,
 and Nayer Khazeni. 2016. "Perceived stress and telomere length: A systematic review,
- 611 meta-analysis, and methodologic considerations for advancing the field." *Brain Behave*
- 612 *Immune* 54 (Supplement C):158-169. doi: <u>https://doi.org/10.1016/j.bbi.2016.02.002</u>.
- 613 McCormack, Gavin R., Melanie Rock, Ann M. Toohey, and Danica Hignell. 2010.
- 614 "Characteristics of urban parks associated with park use and physical activity: A review
 615 of qualitative research." *Health Place* 16 (4):712-726.
- McEachan, R. R. C., S. L. Prady, G. Smith, L. Fairley, B. Cabieses, C. Gidlow, J. Wright, P.
 Dadvand, D. van Gent, and M. J. Nieuwenhuijsen. 2016. "The association between green

- 618 space and depressive symptoms in pregnant women: moderating roles of socioeconomic
- 619 status and physical activity." *J Epidemiol Community Health* 70 (3):253-259. doi:

620 10.1136/jech-2015-205954.

- Meillere, Alizée, François Brischoux, Cécile Ribout, and Frédéric Angelier. 2015. "Traffic noise
 exposure affects telomere length in nestling house sparrows." *Biology letters* 11
 (9):20150559.
- Miri, M., M. Nazarzadeh, A. Alahabadi, M. H. Ehrampoush, A. Rad, M. H. Lotfi, M. H.
 Sheikhha, M. J. Z. Sakhvidi, T. S. Nawrot, and P. Dadvand. 2018. "Air pollution and
 telomere length in adults: A systematic review and meta-analysis of observational
 studies." *Environ Pollut* 244:636-647. doi: 10.1016/j.envpol.2018.09.130.
- Mitchell, C., J. Hobcraft, S. S. McLanahan, S. R. Siegel, A. Berg, J. Brooks-Gunn, I. Garfinkel,
 and D. Notterman. 2014. "Social disadvantage, genetic sensitivity, and children's
- 630 telomere length." *Proc Natl Acad Sci U S A* 111 (16):5944-9. doi:
- 631 10.1073/pnas.1404293111.
- Mundstock, E., H. Zatti, F. M. Louzada, S. G. Oliveira, F. T. Guma, M. M. Paris, A. B. Rueda,
 D. G. Machado, R. T. Stein, M. H. Jones, E. E. Sarria, F. M. Barbe-Tuana, and R.
- Mattiello. 2015. "Effects of physical activity in telomere length: Systematic review and
 meta-analysis." *Ageing Res Rev* 22:72-80. doi: 10.1016/j.arr.2015.02.004.
- Needham, B. L., J. R. Fernandez, J. Lin, E. S. Epel, and E. H. Blackburn. 2012. "Socioeconomic
 status and cell aging in children." *Soc Sci Med* 74 (12):1948-51. doi:

638 10.1016/j.socscimed.2012.02.019.

- 639 Nieuwenhuijsen, M. J., H. Kruize, C. Gidlow, S. Andrusaityte, J. M. Anto, X. Basagana, M.
- 640 Cirach, P. Dadvand, A. Danileviciute, D. Donaire-Gonzalez, J. Garcia, M. Jerrett, M.
- Jones, J. Julvez, E. van Kempen, I. van Kamp, J. Maas, E. Seto, G. Smith, M. Triguero,
- 642 W. Wendel-Vos, J. Wright, J. Zufferey, P. J. van den Hazel, R. Lawrence, and R.
- 643 Grazuleviciene. 2014. "Positive health effects of the natural outdoor environment in
- typical populations in different regions in Europe (PHENOTYPE): a study programme
 protocol." *BMJ Open* 4 (4):e004951. doi: 10.1136/bmjopen-2014-004951.
- Pieters, N., B. G. Janssen, and H. Dewitte. 2016. "Biomolecular Markers within the Core Axis of
 Aging and Particulate Air Pollution Exposure in the Elderly: A Cross-Sectional Study."
 Environ Health Perspect 124 (7):943-950. doi: 10.1289/ehp.1509728.

- Richardson, Elizabeth A., Jamie Pearce, Niamh K. Shortt, and Richard Mitchell. 2017. "The role
 of public and private natural space in children's social, emotional and behavioural
 development in Scotland: A longitudinal study." *Environ Res* 158:729-736.
- Roe, Jenny J., Catharine Ward Thompson, Peter A. Aspinall, Mark J. Brewer, Elizabeth I. Duff,
- David Miller, Richard Mitchell, and Angela Clow. 2013. "Green space and stress:
- Evidence from cortisol measures in deprived urban communities." *Int J Environ Res Public Health* 10 (9):4086-4103.
- Ruijsbroek, A, S. M Mohnen, M Droomers, H Kruize, C Gidlow, R Grazuleviciene, S
 Andrusaityte, and et al. 2017. "Neighbourhood green space, social environment and
 mental health: an examination in four European cities." *International Journal of Public Health*, 62 (6):657–667.
- Samani, N. J., R. Boultby, R. Butler, J. R. Thompson, and A. H. Goodall. 2001. "Telomere
 shortening in atherosclerosis." *Lancet* 358 (9280):472-3. doi: 10.1016/s01406736(01)05633-1.
- Starkweather, Angela R., Areej A. Alhaeeri, Alison Montpetit, Jenni Brumelle, Kristin Filler,
 Marty Montpetit, Lathika Mohanraj, Debra E. Lyon, and Colleen K. Jackson-Cook. 2014.
 "An integrative review of factors associated with telomere length and implications for
- biobehavioral research." *Nurs Res* 63 (1):36-50. doi: 10.1097/nnr.0000000000000000.
- 667 Stigsdotter, U. K., O. Ekholm, J. Schipperijn, M. Toftager, F. Kamper-Jorgensen, and T. B.
 668 Randrup. 2010. "Health promoting outdoor environments--associations between green
- 669 space, and health, health-related quality of life and stress based on a Danish national
- 670 representative survey." *Scand J Public Health* 38 (4):411-7. doi:
- **671** 10.1177/1403494810367468.
- Su, Jason G., Payam Dadvand, Mark J. Nieuwenhuijsen, Xavier Bartoll, and Michael Jerrett.
- 673 2019. "Associations of green space metrics with health and behavior outcomes at
 674 different buffer sizes and remote sensing sensor resolutions." *Environment International*
- 675 126:162-170. doi: https://doi.org/10.1016/j.envint.2019.02.008.
- Theall, Katherine P, Sarah McKasson, Emily Mabile, Lauren F Dunaway, and Stacy S Drury.
- 677 2013. "Early hits and long-term consequences: tracking the lasting impact of prenatal
 678 smoke exposure on telomere length in children." *American Journal of Public Health* 103
 679 (S1):S133-S135.

- 680 Tucker, Larry A. 2017. "Physical activity and telomere length in U.S. men and women: An
- 681 NHANES investigation." *Prev Med* 100 (Supplement C):145-151. doi:
 682 https://doi.org/10.1016/j.ypmed.2017.04.027.
- 683 UN-HABITAT. 2016. Urbanization and Development: Emerging futures. World Cities Report.
 684 New York United Nations: UN HABITAT.
- 685 United Nations, Department of Economic. 2015. ""World
- 686 UrbanizationProspects:The2014Revision."
- 687 <u>https://esa.un.org/unpd/wup/publications/files/wup2014-report.pdf</u>."
- Weier, J., and D. Herring. 2011. *Measuring Vegetation (NDVI & EVI)*. Available:
 http://earthobservatory.nasa.gov/Features/MeasuringVegetation.
- 690 WHO regional Office for Europe. 2016. "Urban green spaces and health a review of evidence."
- WHO regional Office for Europe,, accessed July 31. <u>http://www.euro.who.int/en/health-</u>
 <u>topics/environment-and-health/urban-health/publications/2016/urban-green-spaces-and-</u>
 health-a-review-of-evidence-2016.
- Willeit, P., J. Raschenberger, E. E. Heydon, S. Tsimikas, M. Haun, A. Mayr, S. Weger, J. L.
 Witztum, A. S. Butterworth, J. Willeit, F. Kronenberg, and S. Kiechl. 2014. "Leucocyte
- telomere length and risk of type 2 diabetes mellitus: new prospective cohort study and
- 697 literature-based meta-analysis." *PLoS One* 9 (11):e112483. doi:
- 698 10.1371/journal.pone.0112483.
- Woo, J., N. Tang, E. Suen, J. Leung, and M. Wong. 2009. "Green space, psychological
 restoration, and telomere length." *Lancet* 373 (9660):299-300. doi: 10.1016/s01406736(09)60094-5.
- Zhao, Bing, Ha Q. Vo, Fay H. Johnston, and Kazuaki Negishi. 2018. "Air pollution and telomere
 length: a systematic review of 12,058 subjects." *Cardiovascular diagnosis and therapy* 8
 (4):480-492. doi: 10.21037/cdt.2018.06.05.

Figure and Table captions

Figure 1. Map of Sabzevar, Normalized Difference Vegetation Index (NDVI,) and location of the kindergartens.

Figure 2. Graphical representation of the spearman's correlations between indicators of greenspace exposure.

Table 1. Baseline and exposure characteristics of the study population.

Table 2. Adjusted mixed effects regression models of the association between Log LTL and measures of green spaces. Percentage changes in LTL are presented per one interquartile range (IQR) change in green space indicator.

Baseline and exposure characteristics	Sample (N=200)
LTL, T:S ratio	0.83 (0.7)
Age, years	7 (1)
Missings, n	1
Sex, %	
Male	89 (44.5)
Female	111 (55.5)
Missings, n	0
Environmental tobacco smoke at home, %	
No	169 (84.5)
Yes	26 (13.0)
Missings, n	5
BMI, kg/m2	14.72 (2.6)
Missings, n	17
Parental education in 3 cat, %	
Elementary	52 (26.0)
High School	114 (57.0)
University	32 (16.0)
Missings, n	2
Income, %	
≤ 15 million Rials	173 (86.5)
>15 million Rials	22 (11.0)
Missings, n	5
Illiterate adults, % per census tract	25.2 (28.0)
Missings, n	0
Unemployment, % per census tract	7 (7.9)
Missings, n	0
Distance to any green spaces (in meters)	
Home	106.8 (118.5)
Kindergarten	88.1 (109.2)
Distance to green space \geq 5000 m2 (in meters)	
Home	269.7 (335.4)
Kindergarten	328.5 (497.6)
Surrounding green space index (NDVI)	
Home – 100m	0.067 (0.015)
Home – 300m	0.072 (0.012)
Home – 500m	0.073 (0.012)
Kindergarten – 100m	0.068 (0.011)
Kindergarten – 500m	0.075 (0.019)
Kindergarten – 500m	0.075 (0.016)
Total surrounding green space index (NDVI)	0.060 (0.015)
Total surrounding green space -100m	0.009(0.013) 0.072(0.012)
Total surrounding green space -500m	0.075(0.013)
Time spart in the group space	0.073 (0.011)
Private garden (h/waek)	6 (12)
Public green space (h/week)	187(21.6)
r uone green space (II/ week)	10.7 (21.0)
Plant pots at home (number)	3 (6)

Table 1. Baseline and exposure characteristics of the study population.

Values are mean (SD) for continuous normal distributed variables, median (IQR=interquartile range) for continuous non-normal distributed variables, and frequency (percentage) for categorical variables. LTL = Leukocyte Telomere Length; BMI = Body Mass Index; NDVI = Normalized Difference Vegetation Index.

	Percentage change (95% CI)	p-value ¹
Distance to any green space (in meters)		
Home (IQR=118.5)	0.02 (-9.3, 10.0)	0.996
Kindergarten (IQR=109.23)	8.6 (-4.6, 23.6)	0.213
Distance to green space \geq 5000 m ² (in meters)		
Home (IQR=335.42)	-20.5 (-29.9, -9.7)	<0.001
Kindergarten (IQR=497.64)	-21.8 (-31.4, -10.8)	<0.001
Surrounding greenspace		
Home-100m (IQR=0.016)	5.4 (-2.6, 14.0)	0.191
Home-300m (IQR=0.012)	8.3 (1.2, 15.9)	0.021
Home-500m (IQR=0.012)	6.4 (-0.9, 14.2)	0.085
Kindergarten-100m (IQR=0.011)	3.5 (-2.9, 10.4)	0.289
Kindergarten-300m (IQR=0.019)	13.3 (2.2, 25.7)	0.018
Kindergarten-500m(IQR=0.016)	14.5 (0.4, 28.4)	0.043
Total surrounding greenspace index		
Total surrounding greenspace-100 m (IQR=0.011)	7.6 (-2.0, 18.1)	0.125
Total surrounding greenspace-300 m (IQR=0.019)	12.0 (3.1, 21.7)	0.007
Total surrounding greenspace-500m (IQR=0.016)	8.4 (0.3, 17.1)	0.041
Time spent in the green space (h/week)		
Private garden (IOR=12)	4.9 (-9.1, 21.1)	0.511
Public green space (IOR=21.62)	39.0 (22.8, 57.3)	<0.001
Plant pots at home (IOR=6)	4.1 (-4.0. 13.0)	0.330

Table 2. Adjusted mixed effects regression models of the association between Log LTL and measures of green spaces. Percentage changes in LTL are presented per one interquartile range (IQR) change in green space indicator.

Each cell represents an independent model: 95% CI = 95% confidence interval; LTL = Leukocyte Telomere Length. ¹p: wald test p values. Mixed effects regression models for greenspace exposure were adjusted for age, sex, body mass index (BMI), parental education, income, environmental tobacco smoke at home, illiterate percent per census tract and unemployed percent per census tract. Note: Kindergartens and qRT-PCR plates id were used as random effects in all models. The percentage changes (95% CIs) reported based on 1 IQR increase in distance from home and kindergartens to the closest green space, 1 IQR increase in NDVI, 1 IQR increase for spent time in the public or private green spaces and 1 IQR increase in indoor plant pots at home.