

## Tilting Bundles on Hypertoric Varieties

Non Peer-reviewed author version

VAN DEN BERGH, Michel & SPENKO, Spela (2021) Tilting Bundles on Hypertoric Varieties. In: INTERNATIONAL MATHEMATICS RESEARCH NOTICES, 2021 (2) , p. 1034 -1042.

DOI: 10.1093/imrn/rnz218

Handle: <http://hdl.handle.net/1942/34163>

# TILTING BUNDLES ON HYPERTORIC VARIETIES

ŠPELA ŠPENKO AND MICHEL VAN DEN BERGH

ABSTRACT. Recently McBreen and Webster constructed a tilting bundle on a smooth hypertoric variety (using reduction to finite characteristic) and showed that its endomorphism ring is Koszul.

In this short note we present alternative proofs for these results. We simply observe that the tilting bundle constructed by Halpern-Leistner and Sam on a generic open GIT substack of the ambient linear space restricts to a tilting bundle on the hypertoric variety. The fact that the hypertoric variety is defined by a quadratic regular sequence then also yields an easy proof of Koszulity.

## 1. INTRODUCTION

Below  $k$  is an algebraically closed ground field of characteristic 0. Let  $T$  be torus and let  $W$  be a symplectic representation of  $T$ . Then  $W$  is equipped with a canonical moment map<sup>1</sup>  $\mu : W \rightarrow \mathfrak{t}^*$  with  $\mathfrak{t} = \text{Lie}(T)$ . Throughout we assume that the action of  $T$  is faithful which implies that  $\mu$  is surjective and flat.

Let  $X(T)$  be the characters of  $T$ . For  $\chi \in X(T)$  let  $W^{ss,\chi}$  be the semi-stable part of  $W$  with respect to the linearization  $\chi \otimes \mathcal{O}_W$ . Recall that  $X(T)_{\mathbb{R}}$  is equipped with a so-called secondary fan such that  $\chi$  is in the interior of a maximal cone if and only if  $W^{ss,\chi}/T$  is a (smooth) Deligne-Mumford stack [CLS11, Theorem 14.3.14]. We will call such  $\chi$  *generic*. For generic  $\chi$ , the DM stack  $W^{ss,\chi}/T$  is a crepant resolution of the GIT quotient  $W//T$  [ŠVdB17c, Lemma A.2 and its proof]. Building on the methods developed in [ŠVdB17a] Halpern-Leistner and Sam constructed a tilting bundle on  $W^{ss,\chi}/T$  [HLS16]. See Theorem 3.2 below.

For  $\xi \in \mathfrak{t}^*$  the *hypertoric variety* associated to the data  $(\chi, \xi)$  is the GIT quotient  $\mu^{-1}(\xi)^{ss,\chi}/T$ . For  $\chi$  generic,  $\mu^{-1}(\xi)^{ss,\chi}/T$  is also a smooth Deligne-Mumford stack which is a crepant resolution of the hypertoric variety  $\mu^{-1}(\xi)//T$  (see §4).

The following is our main result.

**Theorem 1.1.** Let  $\chi$  be generic and let  $\mathcal{T}$  be the tilting bundle on  $W^{ss,\chi}/T$  constructed in [HLS16] (see Theorem 3.2).

---

1991 *Mathematics Subject Classification.* 13A50, 14M15, 32S45.

*Key words and phrases.* Invariant theory, tilting bundle, noncommutative resolution.

The first author is a FWO [PEGASUS]<sup>2</sup> Marie Skłodowska-Curie fellow at the Free University of Brussels (funded by the European Union Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 665501 with the Research Foundation Flanders (FWO)). During part of this work she was also a postdoc with Sue Sierra at the University of Edinburgh.

The second author is a senior researcher at the Research Foundation Flanders (FWO). While working on this project he was supported by the FWO grant G0D8616N: “Hochschild cohomology and deformation theory of triangulated categories”.

<sup>1</sup>If  $\langle -, - \rangle$  is the symplectic bilinear form on  $W$ , then one has  $\mu(w)(v) = (1/2)\langle vw, w \rangle$  for  $w \in W, v \in \mathfrak{t}$ .

- (1) For  $\xi \in \mathfrak{t}^*$  the restriction  $\mathcal{T}_\xi$  of  $\mathcal{T}$  to  $\mu^{-1}(\xi)^{ss, \chi}/T$  is a tilting bundle.
- (2) Put  $\Lambda = \text{End}_{W^{ss, \chi}/T}(\mathcal{T})$ ,  $\Lambda_\xi = \text{End}_{\mu^{-1}(\xi)^{ss, \chi}/T}(\mathcal{T}_\xi)$ . Then  $\Lambda_\xi$  is the quotient of  $\Lambda$  by the defining relations of  $\mu^{-1}(\xi)$  which form a regular sequence.
- (3)  $\Lambda_\xi$  is a “non-commutative crepant resolution” [VdB04] of  $k[\mu^{-1}(\xi)//T]$ .
- (4)  $\Lambda_0$  is Koszul when graded using the dilation  $G_m$ -action on  $W$ .

The fact that  $\mu^{-1}(\xi)^{ss, \chi}/T$  admits a tilting bundle with Koszul endomorphism ring when  $\xi = 0$  was proved in [MW18] using the Bezrukavnikov-Kaledin method based on reduction mod  $p$ .

**Remark 1.2.**  $\Lambda$  is an NCCR for  $W//T$  (see Remark 3.3) but it is not Koszul, except in trivial cases. This follows from Proposition 4.1 applied in the same way as in the proof of Theorem 1.1(4). For example in the case of the conifold ( $G_m$  acting with weights  $1, 1, -1, -1$  on a 4-dimensional representation)  $\Lambda$  has cubic relations. This is in fact expected as explained in the next remark.

**Remark 1.3.** From the fact that  $\Lambda_0$  is Koszul one obtains in particular that it is quadratic. Using the explicit form of  $\mathcal{T}$  (see Theorem 3.2) one then quite easily obtains a quiver presentation of  $\Lambda_0$ . See [MW18, Corollary 3.18]. This presentation of  $\Lambda_0$  may be lifted to a quadratic non-homogeneous presentation of  $\Lambda$  as  $\text{Sym}(\mathfrak{t})$ -algebra (with  $|\mathfrak{t}| = 2$ ). This then yields a more complicated presentation of  $\Lambda$  as  $k$ -algebra, possibly involving extra quadratic generators and cubic and even quartic relations.

Similar results as those of McBreen and Webster have been announced by Tatsuyuki Hikita [Hik17].

## 2. ACKNOWLEDGEMENT

This note is mostly the result of a discussion with Travis Schedler during the conference “Quantum geometric and algebraic representation theory” at the Hausdorff Institute in October 2017, which was attended by both authors. We are grateful to Travis for his interest in this work. We would also like to thank Theo Raedschelders for drawing our attention to the work of McBreen and Webster.

## 3. THE TILTING BUNDLE ON $W^{ss, \chi}/T$

Recall that a tilting bundle on an algebraic stack  $\mathcal{Y}$  is a vector bundle  $\mathcal{T}$  such that  $\text{Ext}_{\mathcal{Y}}^{>0}(\mathcal{T}, \mathcal{T}) = 0$  and such that  $\mathcal{T}$  generates  $D_{\text{Qch}}(\mathcal{Y})$  in the sense that  $\mathcal{T}^\perp = 0$ .

For benefit of the reader we now describe explicitly the tilting bundle on  $W^{ss, \chi}/T$  constructed in [HLS16]. The construction in fact only requires that  $W$  is *quasi-symmetric* [ŠVdB17a], i.e. the sum of weights of  $W$  on each line through the origin is zero.

Let  $(\beta_i)_{i=1}^{2\epsilon}$  be the  $T$ -weights of  $W$ . Let  $\varepsilon \in X(T)$ . Put

$$\Sigma = \left\{ \sum_i a_i \beta_i \mid a_i \in ]-1, 0] \right\} \subset X(T)_{\mathbb{R}}, \quad \bar{\Sigma}_\varepsilon = \bigcup_{r>0} \bar{\Sigma} \cap (r\varepsilon + \bar{\Sigma}).$$

The following result indicates the combinatorial significance of the zonotope  $\bar{\Sigma}$ .

**Proposition 3.1** ([HLS16, Proposition 2.1]). Assume that  $W$  is quasi-symmetric. A character  $\chi \in X(T)$  is generic if and only if it is not parallel to any face of  $\bar{\Sigma}$ .

For  $\varepsilon \in X(T)$  put  $\mathcal{L}_\varepsilon = X(T) \cap (1/2)\bar{\Sigma}_\varepsilon$ . The following is one of the main results of [HLS16].

**Theorem 3.2** ([HLS16, Corollary 4.2]). Assume that  $W$  is quasi-symmetric and that  $\chi \in X(T)$  is generic. Then for any generic  $\varepsilon \in X(T)$

$$\mathcal{T} = \bigoplus_{\mu \in \mathcal{L}_\varepsilon} \mathcal{O}_{W^{ss,\chi}} \otimes_k \mu$$

defines a tilting bundle on  $W^{ss,\chi}/T$ .

**Remark 3.3.** If  $W$  is “generic” in the sense of [ŠVdB17a, Definition 1.3.4], i.e. the complement of  $\{x \in W \mid Tx \text{ is closed and } \text{Stab } x = \{e\}\}$  has codimension  $\geq 2$  then  $\Lambda = \text{End}_{W^{ss,\chi}/T}(\mathcal{T}) = (\text{End}_k(\bigoplus_{\mu \in \mathcal{L}_\varepsilon} \mu) \otimes_k k[W])^T$  is the NCCR of  $W//T$  constructed in [ŠVdB17a]. Assuming that  $T$  acts faithfully and that  $W$  is symplectic, one can always reduce to the case of generic  $W$ .

If  $W$  is not generic then all except for two  $T$ -weights of  $W$  (whose sum is 0) lie in a hyperplane. Let  $W'$  be a vector subspace of  $W$  spanned by the weight vectors whose weights lie in this hyperplane. Then  $T = T' \times G_m$ , where  $G_m$  acts trivially on  $W'$  and the action of  $T'$  is faithful on  $W'$ . This induces the decomposition of  $X(T)$  and  $\mathfrak{t}$ . The corresponding projections onto  $X(T')$  and  $\mathfrak{t}'$  will be denoted by  $'$ . We have  $W//T = W'//T' \times_k \mathbb{A}^1$ , and for a generic  $\chi$  also  $W^{ss,\chi}/T = W^{ss,\chi'}/T' \times_k \mathbb{A}^1$ . Moreover,  $\mathcal{L}_\varepsilon = \mathcal{L}'_{\varepsilon'} \times \{m\}$  ( $\mathcal{L}'_{\varepsilon'}$ , with respect to  $(W', T')$ ) for some  $m \in \mathbb{Z}$ . Repeating if necessary we reduce to the case that  $W$  is generic.

We also note that the corresponding hypertoric varieties do not change. Let  $\mu, \mu'$  correspond to the moment maps associated to  $W, W'$ , resp. Then  $\mu^{-1}(\xi)//T = \mu'^{-1}(\xi')//T', \mu^{-1}(\xi)^{ss,\chi}/T = \mu'^{-1}(\xi')^{ss,\chi'}/T'$  (for a generic  $\chi$ ).

#### 4. PROOFS

We revert to the setting of the introduction, i.e.  $W$  is a symplectic representation of the torus  $T$ . Assume  $\chi \in X(T)$  is a generic character. As explained in the introduction  $W^{ss,\chi}/T$  is a DM stack which yields a (stacky) crepant resolution  $\pi : W^{ss,\chi}/T \rightarrow W//T$  of the Gorenstein variety [Sta83, Corollary 13.3]  $W//T$ . The fact that the stabilizers of the  $T$ -action on  $W^{ss,\chi}$  are finite yields by the defining property of the moment map that  $\mu|_{W^{ss,\chi}}$  is smooth. Hence for every  $\xi \in \mathfrak{t}^*$ ,  $\mu^{-1}(\xi)^{ss,\chi}/T$  is also a smooth Deligne-Mumford stack.

Since  $T$  is commutative,  $\mu^{-1}(\xi) \subset W$  is cut out by a  $T$ -invariant regular sequence of global sections (see [Vin86, Theorem 2] and [Sch95, Proposition 9.4]). In particular  $\mu^{-1}(\xi)//T$  is cut out by a regular sequence in  $W//T$  and the same regular sequence cuts out  $\mu^{-1}(\xi)^{ss,\chi}/T$  in  $W^{ss,\chi}/T$ . It follows from this that  $\mu^{-1}(\xi)//T$  is Gorenstein and that  $\pi$  restricts to a crepant resolution  $\pi_\xi : \mu^{-1}(\xi)^{ss,\chi}/T \rightarrow \mu^{-1}(\xi)//T$ .

Finally note that  $\mu^{-1}(\xi)//T$  is normal. For  $\xi = 0$  this is shown in the proof of [BK12, Proposition 4.11]. For the benefit of the reader we give an elementary proof valid for general  $\xi$ . Using Remark 3.3 we first reduce to the generic case and in that case we will show that  $\mu^{-1}(\xi)$  is normal (the quotient of a normal variety is normal). Since  $\mu^{-1}(\xi)$  is Cohen-Macaulay, it suffices by Serre’s normality criterion to prove that  $\mu^{-1}(\xi)$  is regular in codimension 1. Thus, it is sufficient to prove that codimension in  $\mu^{-1}(\xi)$  of the intersection of  $\mu^{-1}(\xi)$  with the non-smooth locus of  $\mu$  is  $\geq 2$ . We will in fact verify that this codimension is  $\geq 3$ !

By the defining property of the moment map, the non-smooth locus of  $\mu$  is the locus of points in  $W$  whose stabilizer has dimension  $> 0$ . This is a union of  $T$ -invariant subspaces of  $W$ . Each such subspace  $W'$  is again a symplectic vector space with a faithful action of a quotient torus  $T/T'$  of  $T$ , where  $T'$  denotes the generic stabilizer of  $W'$  (hence by hypotheses  $\dim T' > 0$ ) and moreover  $\mu$  restricts to a moment map for the  $T/T'$ -action. Thus, as above,  $W' \cap \mu^{-1}(\xi)$  is cut out by a regular sequence of length  $\dim T/T'$  in  $W'$ . Hence the dimension of  $W' \cap \mu^{-1}(\xi)$  is equal to  $c := \dim W' - \dim T/T'$ .

Choose a filtration  $W = W_0 \supseteq W_1 \supseteq \cdots \supseteq W_t = W'$  of  $T$ -invariant symplectic subspaces such that  $\dim W_{i+1} = \dim W_i - 2$  and let  $T_i$  be the generic stabilizer of  $W_i$ . Write  $c_i = \dim W_i - \dim T/T_i = \dim W_i \cap \mu^{-1}(\xi)$ .

The dimension of  $T/T_i$  is equal to the rank of the sublattice of  $X(T)$  spanned by the weights of  $W_i$ . Hence by genericity we have  $\dim T/T_1 = \dim T$ , and thus  $c_1 = c_0 - 2$ . Since  $\dim T' > 0$  we have  $t \geq 2$ . In addition for  $i \geq 1$  we have  $c_{i+1} \leq c_i - 1$  ( $\dim W_i$  goes down by 2 but  $\dim T/T_i$  goes at most down by 1). So we get  $c = c_t \leq c_0 - 3$  and we are done.

*Proofs of Theorem 1.1(1) and 1.1(2).* Let  $i : \mu^{-1}(\xi)^{ss, \chi}/T \rightarrow W^{ss, \chi}/T$  be the inclusion. We have to prove that  $\mathcal{T}_\xi = i^* \mathcal{T}$  is a tilting bundle on  $\mu^{-1}(\xi)^{ss, \chi}/T$ . We have

$$\mathrm{Ext}_{\mu^{-1}(\xi)^{ss, \chi}/T}^*(i^* \mathcal{T}, i^* \mathcal{T}) = \mathrm{Ext}_{W^{ss, \chi}/T}^*(\mathcal{T}, i_* i^* \mathcal{T}).$$

Now  $\mu^{-1}(\xi)^{ss, \chi}$  is cut out in  $W^{ss, \chi}$  by an invariant regular sequence. Tensoring the corresponding Koszul resolution of  $i_* \mathcal{O}_{\mu^{-1}(\xi)^{ss, \chi}}$  with  $\mathcal{T}$  we obtain a left resolution  $K_\bullet$  of  $i_* i^* \mathcal{T}$  which consists of direct sums of  $\mathcal{T}$ . Using the fact that  $\mathrm{Ext}_{W^{ss, \chi}/T}^{>0}(\mathcal{T}, \mathcal{T}) = 0$  we obtain

$$(4.1) \quad \mathrm{Ext}_{W^{ss, \chi}/T}^*(\mathcal{T}, i_* i^* \mathcal{T}) = H^*(\mathrm{Hom}_{W^{ss, \chi}/T}(\mathcal{T}, K_\bullet)).$$

Since  $K_\bullet$  lives in degree  $\leq 0$  this implies  $\mathrm{Ext}_{\mu^{-1}(\xi)^{ss, \chi}/T}^{>0}(i^* \mathcal{T}, i^* \mathcal{T}) = 0$ . Since also tautologically  $\mathrm{Ext}_{\mu^{-1}(\xi)^{ss, \chi}/T}^{<0}(i^* \mathcal{T}, i^* \mathcal{T}) = 0$  we obtain that the right-hand side of (4.1) is in fact a resolution of  $\Lambda_\xi$ , proving Theorem 1.1(2).

To finish the proof of Theorem 1.1(1) we have to check the generation property. Assume that  $\mathrm{RHom}_{\mu^{-1}(\xi)^{ss, \chi}/T}(i^* \mathcal{T}, \mathcal{G}) = 0$  for  $\mathcal{G} \in D_{\mathrm{Qch}}(\mu^{-1}(\xi)^{ss, \chi}/T)$ . Then by adjunction  $i_* \mathcal{G} \in \mathcal{T}^\perp$  and hence  $i_* \mathcal{G} = 0$ . It follows that  $\mathcal{G} = 0$ .  $\square$

*Proof of Theorem 1.1(3).* Since  $\pi_\xi$  is a crepant resolution, it follows that  $\Lambda_\xi = \pi_* \mathcal{E}nd_{\mu^{-1}(\xi)^{ss, \chi}/T}(\mathcal{T}_\xi)$  is an NCCR of  $\mu^{-1}(\xi)//T$  (see e.g. [ŠVdB17b, Corollary 4.3] and Corollary 4.7 in loc. cit. with its proof, together with Remark 3.3).  $\square$

The proof of Theorem 1.1(4) will be based on the following more general criterion.

**Proposition 4.1.** Let  $\Delta$  be a  $\mathbb{N}$ -graded homologically homogeneous  $k$ -algebra [BH84] (see also [VdB04, §3], [SVdB08, §2]) such that  $\Delta_0$  is semi-simple. Assume that  $\Delta$  is finite as a module over a central subring  $R$  of Krull-dimension  $d$ . Then the following are equivalent.

- (1)  $\Delta$  is Koszul.
- (2) The invertible  $\Delta$ -bimodule  $\omega_\Delta := \mathrm{Hom}_R(\Delta, \omega_R)$  (see [SVdB08, Proposition 2.6]) is generated as right module in degree  $d$ .

Moreover in (2) we may replace “right” by “left”.

*Proof.* Note that by [SVdB08, Lemma 2.4(2), Proposition 2.6]  $\omega_\Delta[d]$  is a “rigid dualizing complex” for  $\Delta$ . Hence by the uniqueness of rigid dualizing complexes together with the proof of<sup>2</sup> [VdB97, Proposition 8.2(2)] we obtain as  $\Delta$ -bimodules:  $\omega_\Delta[d] = R\Gamma_{\Delta_{\geq 1}}(\Delta)^\vee$  where  $(-)^\vee$  denotes the graded  $k$ -dual and  $R\Gamma_{\Delta_{\geq 1}}$  is the derived functor of  $\varinjlim_n R\mathrm{Hom}_\Delta(\Delta/\Delta_{\geq n}, -)$ . Moreover by [VdB97, Theorem 5.1] for  $M \in D(\mathrm{Gr}(\Delta))$  we obtain as objects in  $D(\mathrm{Gr}(\Delta^\circ))$

$$R\mathrm{Hom}_\Delta(M, \omega_\Delta[d]) \cong R\Gamma_{\Delta_{\geq 1}}(M)^\vee.$$

Applying this with a graded simple  $\Delta$ -module  $\mathbb{S}$ , concentrated in degree zero we find

$$R\mathrm{Hom}_\Delta(\mathbb{S}, \omega_\Delta[d]) = \mathbb{S}^*;$$

i.e.  $R\mathrm{Hom}_\Delta(-, \omega_\Delta[d])$  restricts to the canonical bijection between the graded simple left and right  $\Delta$ -modules, concentrated in degree zero.

We will now prove (2)  $\Rightarrow$  (1). Since  $\omega_\Delta$  is invertible and generated in degree  $d$  we have that  $\omega_\Delta \otimes_\Delta \mathbb{S} = \mathbb{S}^\dagger(-d)$  where  $\mathbb{S}^\dagger$  is a graded simple  $\Delta$ -module concentrated in degree zero. Hence we obtain

$$(4.2) \quad R\mathrm{Hom}_\Delta(\mathbb{S}, \Delta) = R\mathrm{Hom}_\Delta(\omega_\Delta[d] \otimes_\Delta \mathbb{S}, \omega_\Delta[d]) = \mathbb{S}^{\dagger*}(d)[-d]$$

Let  $P_\bullet = \cdots \rightarrow P_1 \rightarrow P_0$  be the minimal graded projective resolution of  $\mathbb{S}$  over  $\Delta$ . As  $\Delta$  is homologically homogeneous,  $P_\bullet$  is of length  $d$ , and by (4.2)  $\mathrm{Hom}_\Delta(P_\bullet, \Delta)[d]$  is a projective resolution of  $\mathbb{S}^{\dagger*}(d)$ . Hence  $P_d = P(\mathbb{S}^\dagger)(-d)$ , where  $P(\mathbb{S}^\dagger)$  is the graded projective cover of  $\mathbb{S}^\dagger$ .

Let  $f_i$  be the minimal degree of an element in  $P_i$ . We claim  $f_{i-1} < f_i$ . To see this note that the graded Jacobson radical of  $\Delta$  is equal to  $\Delta_{\geq 1}$  (as  $\Delta_0$  is semi-simple). Hence the image of  $d_i : P_i \rightarrow P_{i-1}$  is in  $\Delta_{\geq 1}P_{i-1}$ .

An element  $x$  of degree  $f_i$  in  $P_i$  is not in  $\Delta_{\geq 1}P_i$  and therefore it is not in the image of  $d_{i+1}$ . As  $P_\bullet$  is acyclic it follows that  $d_i(x) \neq 0$ . Writing  $d_i(x) = \sum_j l_j y_j$  with homogeneous  $l_j \in \Delta_{\geq 1}$ ,  $y_j \in P_{i-1}$ , we see that some  $y_j$  must be non-zero. The fact  $l_j$  has strictly positive degree implies  $\deg(y_j) < \deg(x)$  which proves our claim.

From the inequalities

$$d = f_d > f_{d-1} > \cdots > f_1 > f_0 = 0$$

we obtain  $f_i = i$ .

We now use a dual argument. Let  $f_i^*$  be the maximal degree of a generator of  $P_i$ . Using the dual resolution  $\mathrm{Hom}_\Delta(P_\bullet, \Delta)(-d)[d]$  of  $\mathbb{S}^\dagger$  (which is also minimal) we get that  $-d + f_{d-i}^* = -i$ , i.e.  $f_i^* = i$ . Therefore  $P_i$  is purely generated in degree  $i$  and so the resolution  $P_\bullet$  is linear.

The implication (1)  $\Rightarrow$  (2) is similar. From the nature of the minimal resolutions of the simples we get  $\omega_\Delta \otimes_\Delta \mathbb{S} = \mathbb{S}^\dagger(d)$  for all  $\mathbb{S}$ . This implies that  $\omega_\Delta$  is right generated in degree  $d$ .

Since Koszulity is left right symmetric, the hypothesis of right generation in (2) may indeed be replaced by left generation.  $\square$

<sup>2</sup>The reference [VdB97] is concerned with the case  $\Delta_0 = k$  but this hypothesis is not used in an essential way.

*Proof of Theorem 1.1(4).* Let  $s = \dim T$ ,  $2e = \dim W$  and  $d = \dim \mu^{-1}(0)//T$ . Thus  $d = 2e - 2s$ .

By Theorem 3.2 have  $\Lambda_0 = (\text{End}_k(\bigoplus_{\mu \in \mathcal{L}_\xi} \mu) \otimes_k k[\mu^{-1}(0)])^T$ . Hence  $\Lambda_0$  is  $\mathbb{N}$ -graded and its part of degree zero is semi-simple.<sup>3</sup> Therefore by Proposition 4.1 we have to prove that  $\omega_{\Lambda_0}$  is generated in degree  $d$ .

Put  $R = k[W//T]$ ,  $R_0 = k[\mu^{-1}(0)//T]$ . We have

$$\omega_{R_0} = \text{Ext}_R^s(R_0, \omega_R) = \text{Ext}_R^s(R_0, R(-2e)) = R_0(2s - 2e) = R_0(-d)$$

where the first equality is the adjunction formula, the second equality follows from [Sta83, Corollary 13.3] and the third equality follows from the fact that  $R_0$  is cut out from  $R$  by a quadratic regular sequence of length  $s$ . As  $\Lambda_0$  is an NCCR by Theorem 1.1(3) we have  $\text{Hom}_{R_0}(\Lambda_0, R_0) = \Lambda_0$ . It follows that  $\omega_{\Lambda_0} = \text{Hom}_{R_0}(\Lambda_0, \omega_{R_0}) = \Lambda_0(-d)$ , finishing the proof.  $\square$

#### REFERENCES

- [BH84] K. A. Brown and C. R. Hajarnavis, *Homologically homogeneous rings*, Trans. Amer. Math. Soc. **281** (1984), 197–208.
- [BK12] G. Bellamy and T. Kuwabara, *On deformation quantizations of hypertoric varieties*, Pacific J. Math. **260** (2012), no. 1, 89–127.
- [CLS11] D. A. Cox, J. B. Little, and H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011.
- [Hik17] T. Hikita, *Tilting generators for hypertoric varieties*, <https://www.math.kyoto-u.ac.jp/~ivan.ip/KTGU/Workshop2017/program.html>, Feb. 20 2017, lecture at “The 2nd KTGU Mathematics Workshop for Young Researchers”, Kyoto University.
- [HLS16] D. Halpern-Leistner and S. Sam, *Combinatorial constructions of derived equivalences*, arXiv:1601.02030 [math.AG], 2016.
- [MW18] M. McBreen and B. Webster, *Homological Mirror Symmetry for Hypertoric Varieties I*, arXiv:1804.10646 [math.AG], 2018.
- [Sch95] G. W. Schwarz, *Lifting differential operators from orbit spaces*, Ann. scient. Éc. Norm. Sup. **28** (1995), no. 3, 253–305.
- [Sta83] R. P. Stanley, *Combinatorics and commutative algebra*, Progress in Mathematics, vol. 41, Birkhäuser Boston Inc., Boston, MA, 1983.
- [SVdB08] J. T. Stafford and M. Van den Bergh, *Noncommutative resolutions and rational singularities*, Michigan Math. J. **57** (2008), 659–674.
- [ŠVdB17a] Š. Špenko and M. Van den Bergh, *Non-commutative resolutions of quotient singularities*, Invent. Math. **210** (2017), no. 1, 3–67.
- [ŠVdB17b] ———, *Noncommutative crepant resolutions for some toric singularities I*, arXiv:1701.0525 [math.AG], 2017.
- [ŠVdB17c] ———, *Noncommutative crepant resolutions for some toric singularities II*, arXiv:1707.08245 [math.AG], 2017.
- [VdB97] M. Van den Bergh, *Existence theorems for dualizing complexes over non-commutative graded and filtered rings*, J. Algebra **195** (1997), no. 2, 662–679.
- [VdB04] ———, *Non-commutative crepant resolutions*, The legacy of Niels Henrik Abel, Springer, Berlin, 2004, pp. 749–770.
- [Vin86] É. B. Vinberg, *Complexity of action of reductive groups*, Functional Anal. Appl. **20** (1986), no. 1, 1–11.

DEPARTEMENT WISKUNDE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, B-1050 ELSENE  
E-mail address: [spela.spenko@vub.be](mailto:spela.spenko@vub.be)

DEPARTEMENT WNI, UNIVERSITEIT HASSELT, UNIVERSITAIRE CAMPUS, B-3590 DIEPENBEEK  
E-mail address: [michel.vandenbergh@uhasselt.be](mailto:michel.vandenbergh@uhasselt.be)

<sup>3</sup>This is the only place where we use the specific form of  $\mathcal{T}$ .