Made available by Hasselt University Library in https://documentserver.uhasselt.be

Evaluation of Processing Speed of Different Cognitive Functions Across the Life Span Using Cognitive Mobile Games Peer-reviewed author version

BONNECHERE, Bruno (2022) Evaluation of Processing Speed of Different Cognitive Functions Across the Life Span Using Cognitive Mobile Games. In: Games for health journal, 11 (2), p. 132 -140.

DOI: 10.1089/g4h.2021.0144 Handle: http://hdl.handle.net/1942/36798

EVALUATION OF PROCESSING SPEED OF DIFFERENT COGNITIVE FUNCTIONS ACROSS THE LIFESPAN USING COGNITIVE MOBILE GAMES

Bonnechère B.^{1,2*}

¹ REVAL Rehabilitation Research Center, Faculty of Rehabilitation Sciences, Hasselt University, Diepenbeek, Belgium

² Department of Psychiatry and Behavioural and Clinical Neurosciences, University of Cambridge, Cambridge, United Kingdom CB2 0SZ

Corresponding author: Prof. dr. Bonnechère Bruno: bruno.bonnechere@uhasselt.be

Agoralaan Building A 3590 Diepenbeek - Belgium

Words count: 2994

Figures: 3

Tables: 2

References: 62

ABSTRACT

Objective: Processing speed are important indicators of cognitive functioning and normal aging. However, the tools used to evaluate these are often rather simplistic and only assess one cognitive component. The aim of this study was to use cognitive mobile games (CMG) to evaluate the evolution of reaction times over the lifespan during different cognitive tasks.

Methodology: We carried out a retrospective observational study in which we obtained anonymized results of 15,000 subjects. Scores of five CMG that train arithmetic, vocabulary, response control, visual attention and recognition, and working memory were analysed.

Results: Overall, we observed a highly statistically significant decrease (p < .001) in processing speed and a decrease of accuracy (p < .001) with increasing participant age, indicating that for each cognitive function tested, older participants performed cognitive tasks more slowly than younger participants. We also observed an interaction between the age of the participants and the number of errors. These results are consistent with physiological data with respect to aging and cognition.

Conclusion: Owing to their wide availability and ease of use, CMG could be used as a simple tool to monitor cognitive function such as processing speed. Further studies are needed to study the influence of the pathologies on those variables.

INTRODUCTION

The World Health Organization estimated that the population aged above 60 years old will double in size by 2050.¹ In the context of the aging of the population, the accurate and accessible assessment of cognitive function is thus of increasingly high importance for both public and individual health. Clinically, normal aging is associated with progressive decline in cognitive functions.² Usually, cognition is divided into several sub-functions such as attention, memory, language and visuospatial abilities, these sub-functions are each impacted differently by the process of aging.^{3,4} Indeed, following childhood's fast cognitive growth, early adulthood (i.e., the ages of 20–39 years) is marked by relative stability and peak cognitive function, then many declines occur in the cognitive system.⁵ The functions most affected by age are the processing speed (PS), executive functions,⁶ inhibitory function,⁷ and episodic memory⁸ while verbal abilities⁹ and implicit memory seems to be preserved.¹⁰ As the name indicates, PS relates to the rate at which information is detected, interpreted, comprehended, and reacted to. PS is often regarded a critical component of attention; certainly, the majority of tests of attention are speedsensitive, if not speed-dependent, however some specialized neuropsychological testing disentangle attentional accuracy and response speed.¹¹ PS deficiencies may impair cognitive processing in various cognitive areas (e.g., placing a limit on the amount of information that can be attended to or encoded at one time). Processing speed, on the other hand, may be separated from other cognitive domains and subdomains.¹² From a clinical point of view, PS is an important factor to evaluate since it is a good indicator of cognitive function and thus of declining cognitive performance in healthy aging¹³ and in dementia.¹⁴ Furthermore, it has been highlighted that PS has an effect on other cognitive skills; PS is responsible for a significant amount of the decreases in cognitive ability associated with aging. Additionally, studies suggest that reductions in PS have an effect on daily functioning and driving behavior.¹⁵ Clinically, cognition is evaluated using different scores and scales that can assess global cognition or more specifically different cognitive subfunctions.¹⁶ So far, measures of reaction time are still performed using so-called simple reaction time tasks or more complex methods such as the flanker reaction time task.¹³ The mean speed of responding is the most commonly used measure in the assessment of reaction time.¹⁷ Currently, there is still a lack of information about the evolution of PS during various cognitive tasks across the lifespan using automated computerized solutions (e.g., digital biomarkers).¹⁸ One study previously investigated the PS in different activities using tablets and apps.¹⁹ The authors assessed cognitive function of a large sample size of more than 15,000 participants using a computerized cognitive task battery composed of 13 tasks that were categorized into three cognitive constructs of memory, attention and processing speed. The authors found that PS mediated the relationship between age and other cognitive performance scores. However, despite the large sample size, only a limited number of participants (n = 837) aged above 60 years old were included in this study. There is thus a lack of information about the evolution of PS at old and very old age. Since PS is a good indicator of healthy aging, it is therefore important to study how PS evolves with ages and to analyze if there is any difference between the different cognitive functions. Today, the development and use of Cognitive Mobile Games (CMG) is becoming more and more popular in a clinical context and in research thanks to the worldwide success of cognitive video games such as 'How old is your Brain' by Dr. Kawashima,²⁰ and other popular apps. The games have been developed to train and challenge the brain in an attempt to preserve or improve cognitive functions.^{21–23} Besides the brain training aspects provided, CMG also offer other interesting possibilities, such as allowing simultaneous cognitive evaluation while performing the exercises.^{24–26} Such assessments can be done using directly the scores provided by the CMG ^{27,28} or by developing new applications to collect more psychological measurements such as the reaction time and PS.²⁹ Remote health assessments that collect real-world data outside of clinic settings need a thorough grasp of the appropriate data collection, quality assurance, analysis, and interpretation methodologies.³⁰ Therefore, the main aim of this study was to establish the validity of the outcomes obtained from the CMG in the study of human cognition. The specific aim of this study was to determine if the PS in different CMG could be used as an indicator of cognitive aging. To do so we examined the PS age-related change of differences and hypothesized that, as for the cognitive functions, the change in PS in the different cognitive functions should not be equally affected by the aging process and that these changes may affect the strategies of the participants (e.g., focus on correct answers rather than speed).

MATERIAL AND METHODS

We carried out a retrospective observational study in which we obtained anonymized results of 15,000 subjects ranging from 18 to 83 years old (43±10 years old). This study was approved by the Cambridge Psychology Research Ethics Committee (Pre.2020.28), and all participants agreed that their data could be used for research purposes when installing the app. To have the same number of participants in every category of age (i.e., balance design), subjects were randomly selected (using simple random sampling) from a list of participants that had played five CMG provided by Peak brain training (www.peak.net, London - UK). The 5 CMG were selected based on a previous study that computed correlations between scores obtained for these 5 particular CMG and scores in two clinically-established cognitive assessments (the Mini-Mental State Examination and the Addenbrooke's Cognitive Evaluation) in elderly subjects with and without cognitive impairments.²⁷ To avoid risk of bias induced by the training or familiarization of the CMG we only analysed the first session of play. Screenshots, descriptions, main objectives and how the PS are computed for the five CMG are presented in Table 1. For the different CMG the main cognitive abilities trained were defined based on previous research.³¹ The CMG were played on a smartphones or tablets. The CMG response time data are automatically recorded by the application.

The main outcomes were the PS evaluated by the processing time (PT defined as the time required to perform each task in the CMG, see Table 1), we separately analysed PT for correct and incorrect answers. As secondary outcomes, we also computed the accuracy (number of correct trials divided by the total number of trials) as well as the number of correct trials as an indicator of the efficacy of the training.

We analysed the different outcomes of each CMG using mixed models³² and tested the interactions between the age of the participants and the PT for correct and incorrect responses. Statistical analyses were performed at an overall significance level of 0.05. Statistical analyses were conducted in RStudio (version 1.2.5042) with R version 3.6.3, using the LME4 package to run the mixed effect models.³³

RESULTS

Results of the mixed models analysis for the 5 different CMG are presented in Table 2. The evolution of the PT, the number of success and the accuracy over age is presented in Figure 1. First concerning the accuracy we observed that, except for arithmetic (measured by *Square Numbers*, $\beta = 0.003$ [-0.004 ; 0.011], p = .345), there is statistically significant linear decrease of accuracy with age. The most important decline is for task shifting (measured by *Must Sort*, $\beta = -0.102$ [-0.112 ; -0.091], p < .001) and the less important, yet highly significant for working memory (measured by *Rush Back*, $\beta = -0.039$ [-0.046 ; -0.032], p < .001).

For the PT, first concerning the correct response there is a significant effect of age, regardless of the cognitive functions. The effect size (estimated by the beta's)³⁴ is more important for the vocabulary (measured by *Word Pairs*, $\beta = 44$ ms [42; 46]) and less important for task shifting (measured by *Must Sort*, $\beta = 6.2$ ms [5.9; 6.7]), see Figure 2 for complete results. Similar results were obtained for the PT during incorrect responses: the change in time to give incorrect

responses is always more important that the change for correct ones. To visualize the evolution of the PT across lifespan between the different cognitive abilities we first centred the PT by removing the average results of the younger participants (18-21 years old) for the different CMG and then plot the evolution of PT through ages in Figure 3. To detect change in slope across aging we applied a Continuous-piecewise-linear Pruned Optimal Partitioning for the different CMG³⁵ We observed three different patterns. For task shifting and working memory a small and linear evolution of the PT. For the visual attention the evolution is also linear but more marked ($\beta = 6.2$ for Must Sort, $\beta = 9.5$ for Rush Back and $\beta = 20$ for Unique). Finally, for vocabulary and arithmetic abilities the PT increased linearly until approximately 50-55 years old, after this first phase we observed and important change in the increase of PT. For the vocabulary, this change occurs at 52 years old and for arithmetic at 56 years old. Next, we compared the slope before and after this change. For vocabulary the slope is 17 [14; 21] before 52 years old and 73 [64; 82], the difference is significant (p < .001); for the arithmetic abilities the slope is 8 [6; 10] before 56 years old and 36 [31; 41] after, yielding to a significant difference (p < .001).

Finally, we assessed the interaction between correct and incorrect responses and found significant interactions for all assessed cognitive functions with ages indicating that the older the subjects are the more important the difference in PT between correct and incorrect response is.

DISCUSSION

The main results of this study if that CMG can be used as an indicator of cognitive function to evaluate the PS of participants within different cognitive tasks. The observed increase in PT for the five CMG are coherent with physiological data and previous studies.^{6,13,29} Analyses showed that the smallest effect was observed for fluency (*Word Pairs*), consistent with the notion that

vocabulary and fluency are the most preserved cognitive function with age.^{4,36,37} Indeed fluency task seems to be only reduced at old or very old age,^{38,39} in our study we observed an important increase in the PT around 50 years old but both the accuracy and the number of responses decreased linearly. On the other hand, task shifting (Must Sort) is the most affected with a large effect size, which is also in accordance with the literature since this task is the closest one to a simple reaction time tests, and previous studies highlighted that this function is particularly strongly affected by aging.^{3,6} It has been previously shown that older adults experience more difficulties in tasks shifting compared to younger individuals due to highest cognitive cost (i.e., change in performance on no-switch trials in dual-task blocks compared to no-switch trials in single-task blocks).⁴⁰ The difficulties are more marked when the tasks also require inhibition skills,⁴¹ which was not the case in this study. Concerning the other functions, for visual attention (Unique) the linear increased in PT, more important for incorrect responses, is also consistent with neurophysiological knowledge on aging where a decrease of selective attention is observed ⁴² that may be due to the deterioration of the field of view and vision.⁴³. For working memory (Rush Back) the same trend was observed which is also consistent with the literature were it has been showed that older adults tended to show less improvement in scores after *n*-back training (similar task than the Rush Back) than younger adults.⁴⁴ Finally for the arithmetic ability (Square Numbers), the time required to perform the computation stay stable until 50 years old and then there is an important increase, the accuracy is also decreased. A recent study show that arithmetic abilities skills are preserved in healthy elderly adults and that older adults could even outperform young adults because they more often retrieve arithmetic facts from long-term memory.⁴⁵ Our results did not support those findings as we observed a decreases of the accuracy, number of response and increase of PT. This may be due to the educational background of the participants. The high variability and the different patterns of PT changes, evaluated through the CMG and presented in Figures 1 and 3, are in line with previous studies. A recent study highlighted the fact that no single measure of cognitive performance and performance variability produces the same findings with respect to age related change.⁴⁶ The authors observed that the age of peak performance varied significantly across metrics, with young adults performing best on measures of median PT, middle-aged adults performing best on certain measures of PT variability, and older adults performing best on accuracy.

Another important finding of this study is the interaction between the age of the participants and the difference between PT for correct and incorrect measurements. While a slight decrease in accuracy was also found with age, this interaction indicates that older participants tend to have a higher PT for incorrect responses compared to younger participants. Several mechanisms could explain these differences. The reduced response inhibition is reflected by poorer performance in incongruent trials where prepotent responses can interfere with other correct responses.⁴⁷ Older adults also demonstrate difficulty forming and retrieving episodic memories. One proposed mechanism is that older adults are impaired at binding information into nonoverlapping representations, which is a key function of the hippocampus.⁴⁸

The results of this study must be interpreted with caution and three main limitations should be borne in mind. Due to the study design, the first - and probably the most important one - is the selection bias of the participants. Since all the participants were users of the apps, it implies that they are familiar with smartphone and mobile apps. Despite the fact that we analysed only the first session of training to avoid training of familiarization effects, there may have been a transfer of the abilities trained in other apps into this one as a study highlighted that owner and regular owner of smartphones have a reduced risk of dementia compared to people that do not own one mobile device.⁴⁹ This effect may be more important in younger participants, who are more familiar with smartphones and the use of apps or games,⁵⁰ therefore the observed difference in PT may be overestimated. The second limitation is that we do not have information about the participants but it is well documented that several factors, not only age, influence

cognitive function and abilities such as education,⁵¹ lifestyle-related factors,⁵² genetics⁵³ or comorbidities (i.e., diabetes,⁵⁴ chronic respiratory diseases,⁵⁵ cardiovascular diseases⁵⁶ or stroke⁵⁷). Since we do not have access to these information we can not speculate the influence of those parameters on the results. The third limitation is related to the use of different technologies (hardware and software). Here again we do not have information on the devices used. Since the PT are recorded in ms, the accuracy may vary depending on the types of devices used. However, given the large sample size we can assume that the vast majority of the participants playing with this kind of apps are cognitively healthy and that the effects of these factors, if any, must be mitigate by the large number of participants.

Despite the above-mentioned limitation, the results of this study are in accordance with current neurophysiological knowledge. Remote digital studies have the potential to alter the user experience in comparison to conventional clinical trials and provide additional issues to the researcher that must be addressed in order to conduct a successful study.⁵⁸ The proposed method to assess PS has many advantages: largely available, affordable, efficient administration, automated scoring, fun and evaluation not requiring the presence of a healthcare professional even at old age (above 80 years old). Another advantage is that, with the exception of Square Numbers and Word Pairs, the CMG are culturally and educationally unbiased which is particularly important for cognitive evaluation, since it is widely known that education has a confounding effect on cognitive assessment.⁵⁹ The above aspects are particularly interesting in the context of low- and mid-income countries: owed to the global increase in life expectancy, these countries are faced with the high price of the disease of the elderly with limited human resources.⁶⁰ CMG could be used as practical tool for the routine screening of PS and therefore as a method of longitudinal assessment and follow-up. However, before being used at-large scale and in routine, further studies must focus on the changes in PT in various pathologies affecting cognitive function such as dementia, Mild Cognitive Impairment, stroke, multiple sclerosis of Parkinson's disease. Important efforts must also be done to link the results with neurophysiological ⁶¹ and imaging data.⁶² These efforts should help determine the best criteria for assessing and classifying participants or patients, as well as determining thresholds and values to be reached during regular monitoring.

CONCLUSION

This study highlights the potential use of CMG as an indicator of the different cognitive functions. Beside the positive effect of brain training using CMG, these types of apps could also be used to performed regular follow-up of cognitive functions or as new outcomes for interventional or physiological studies. Performing an application-based study also enable researchers to do cross-sectional studies at a cheap cost. Due of the device's portability, it is an ideal tool for cross-cultural studies. Additionally, the device's mobility enables data collection in places or populations that may not be readily accessible or equipped with equipment capable of conducting lengthy testing. Additionally, mobile apps might be beneficial for cognitive and psychological testing. The evaluation tools may be used to evaluate and define persons who are at an increased risk of cognitive impairment or illness. Further research should focus on the use of CMG in older adults with and without risk of cognitive impairment and in neurological patients suffering from cognitive impairment.

ACKNOWLEDGMENTS

The author would like to thank Peak (www.peak.net) for providing access to the results of the CMG.

AUTHOR CONTRIBUTION

The author confirms sole responsibility for the following: study conception and design, analysis and interpretation of results, and manuscript preparation

CONFLICT OF INTEREST

The author declare no conflict of intersest

FUNDINGS

None

REFERENCES

- WHO. Ageing and health. https://www.who.int/news-room/fact-sheets/detail/ageing-andhealth (2018).
- Eshkoor, S. A., Hamid, T. A., Mun, C. Y. & Ng, C. K. Mild cognitive impairment and its management in older people. *Clin Interv Aging* 10, 687–693 (2015).
- Vogel, A., Salem, L. C., Andersen, B. B. & Waldemar, G. Differences in quantitative methods for measuring subjective cognitive decline - results from a prospective memory clinic study. *Int Psychogeriatr* 28, 1513–1520 (2016).
- Ofen, N. & Shing, Y. L. From perception to memory: changes in memory systems across the lifespan. *Neurosci Biobehav Rev* 37, 2258–2267 (2013).
- Ballesteros, S., Mayas, J. & Reales, J. M. Cognitive function in normal aging and in older adults with mild cognitive impairment. *Psicothema* 25, 18–24 (2013).
- Tam, A., Luedke, A. C., Walsh, J. J., Fernandez-Ruiz, J. & Garcia, A. Effects of reaction time variability and age on brain activity during Stroop task performance. *Brain Imaging Behav* 9, 609–618 (2015).
- Baltes, P. B. & Lindenberger, U. Emergence of a powerful connection between sensory and cognitive functions across the adult life span: a new window to the study of cognitive aging? *Psychol Aging* 12, 12–21 (1997).
- Gutchess, A. H. & Park, D. C. Effects of Aging on Associative Memory for Related and Unrelated Pictures. *Eur J Cogn Psychol* 21, 235–254 (2009).
- 9. The handbook of aging and cognition. (Psychology Press, 2008).
- Ballesteros, S., Bischof, G. N., Goh, J. O. & Park, D. C. Neural correlates of conceptual object priming in young and older adults: an event-related functional magnetic resonance imaging study. *Neurobiology of Aging* 34, 1254–1264 (2013).

- Silva, M. A. & Lee, J. M. Neurocognitive testing. in *Reference Module in Neuroscience and Biobehavioral Psychology* B9780128229637001000 (Elsevier, 2021). doi:10.1016/B978-0-12-822963-7.00047-5.
- Agelink van Rentergem, J. A. *et al.* The Factor Structure of Cognitive Functioning in Cognitively Healthy Participants: a Meta-Analysis and Meta-Analysis of Individual Participant Data. *Neuropsychol Rev* 30, 51–96 (2020).
- Chen, K.-C., Weng, C.-Y., Hsiao, S., Tsao, W.-L. & Koo, M. Cognitive decline and slower reaction time in elderly individuals with mild cognitive impairment. *Psychogeriatrics* 17, 364–370 (2017).
- Prado, C. E., Watt, S., Treeby, M. S. & Crowe, S. F. Performance on neuropsychological assessment and progression to dementia: A meta-analysis. *Psychol Aging* 34, 954–977 (2019).
- Vance, D. E. Speed of processing in older adults: a cognitive overview for nursing. J Neurosci Nurs 41, 290–297 (2009).
- Harvey, P. D. Domains of cognition and their assessment. *Dialogues Clin Neurosci* 21, 227–237 (2019).
- Booth, T. *et al.* Reaction time variability and brain white matter integrity. *Neuropsychology* 33, 642–657 (2019).
- Piau, A., Wild, K., Mattek, N. & Kaye, J. Current State of Digital Biomarker Technologies for Real-Life, Home-Based Monitoring of Cognitive Function for Mild Cognitive Impairment to Mild Alzheimer Disease and Implications for Clinical Care: Systematic Review. *J Med Internet Res* 21, e12785 (2019).
- Lee, H. *et al.* Examining cognitive function across the lifespan using a mobile application. *Computers in Human Behavior* 28, 1934–1946 (2012).

- Nouchi, R. *et al.* Brain training game boosts executive functions, working memory and processing speed in the young adults: a randomized controlled trial. *PLoS ONE* 8, e55518 (2013).
- 21. Gates, N. J. *et al.* Computerised cognitive training for maintaining cognitive function in cognitively healthy people in midlife. *Cochrane Database Syst Rev* **3**, CD012278 (2019).
- 22. Gates, N. J. *et al.* Computerised cognitive training for preventing dementia in people with mild cognitive impairment. *Cochrane Database Syst Rev* **3**, CD012279 (2019).
- 23. Gates, N. J. *et al.* Computerised cognitive training for 12 or more weeks for maintaining cognitive function in cognitively healthy people in late life. *Cochrane Database Syst Rev* CD012277 (2020) doi:10.1002/14651858.CD012277.pub3.
- Koo, B. M. & Vizer, L. M. Mobile Technology for Cognitive Assessment of Older Adults: A Scoping Review. *Innov Aging* 3, igy038 (2019).
- 25. Bonnechère, B. *et al.* The Use of Mobile Games to Assess Cognitive Function of Elderly with and without Cognitive Impairment. *J. Alzheimers Dis.* **64**, 1285–1293 (2018).
- Bonnechère, B. *et al.* Age-Associated Capacity to Progress When Playing Cognitive Mobile Games: Ecological Retrospective Observational Study. *JMIR Serious Games* 8, e17121 (2020).
- 27. Bonnechère, B. *et al.* The Use of Mobile Games to Assess Cognitive Function of Elderly with and without Cognitive Impairment. *J. Alzheimers Dis.* **64**, 1285–1293 (2018).
- 28. Van Hove, O. *et al.* The use of cognitive mobile games to assess cognitive function of healthy subjects under various inspiratory loads. *Medicine in Novel Technology and Devices* 1, 100005 (2019).
- 29. Khaligh-Razavi, S.-M. *et al.* Integrated Cognitive Assessment: Speed and Accuracy of Visual Processing as a Reliable Proxy to Cognitive Performance. *Sci Rep* **9**, 1102 (2019).

- Warmerdam, E. *et al.* Long-term unsupervised mobility assessment in movement disorders.
 Lancet Neurol 19, 462–470 (2020).
- Bonnechère, B., Klass, M., Langley, C. & Sahakian, B. J. Brain training using cognitive apps can improve cognitive performance and processing speed in older adults. *Sci Rep* 11, 12313 (2021).
- Di Brisco, A. M. & Migliorati, S. A new mixed-effects mixture model for constrained longitudinal data. *Stat Med* 39, 129–145 (2020).
- Bates, D., M\u00e4chler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using Ime4. J. Stat. Soft. 67, (2015).
- 34. Nieminen, P., Lehtiniemi, H., Vähäkangas, K., Huusko, A. & Rautio, A. Standardised regression coefficient as an effect size index in summarising the reported findings between quantitative exposure and response variables in epidemiological studies. *Epidemiology, Biostatistics and Public Health* (2013) doi:10.2427/8854.
- 35. Fearnhead, P., Maidstone, R. & Letchford, A. Detecting Changes in Slope With an L₀ Penalty. *Journal of Computational and Graphical Statistics* **28**, 265–275 (2019).
- Verhaeghen, P. Aging and vocabulary scores: a meta-analysis. *Psychol Aging* 18, 332–339 (2003).
- 37. Methqal, I. *et al.* Age-Related Shift in Neuro-Activation during a Word-Matching Task. *Front Aging Neurosci* **9**, 265 (2017).
- 38. Murphy, D. H. & Castel, A. D. Age-related similarities and differences in the components of semantic fluency: analyzing the originality and organization of retrieval from long-term memory. *Aging, Neuropsychology, and Cognition* 1–14 (2020) doi:10.1080/13825585.2020.1817844.
- 39. Steen-Baker, A. A. *et al.* The effects of context on processing words during sentence reading among adults varying in age and literacy skill. *Psychol Aging* **32**, 460–472 (2017).

- 40. Eich, T. S. *et al.* Functional brain and age-related changes associated with congruency in task switching. *Neuropsychologia* **91**, 211–221 (2016).
- Eich, T. S., MacKay-Brandt, A., Stern, Y. & Gopher, D. Age-Based Differences in Task Switching Are Moderated by Executive Control Demands. *GERONB* gbw117 (2016) doi:10.1093/geronb/gbw117.
- 42. Malavita, M. S., Vidyasagar, T. R. & McKendrick, A. M. The Effect of Aging and Attention on Visual Crowding and Surround Suppression of Perceived Contrast Threshold. *Invest Ophthalmol Vis Sci* 58, 860–867 (2017).
- 43. Matthews, K., Nazroo, J. & Whillans, J. The consequences of self-reported vision change in later-life: evidence from the English Longitudinal Study of Ageing. *Public Health* 142, 7–14 (2017).
- 44. Rhodes, R. E. & Katz, B. Working memory plasticity and aging. *Psychol Aging* 32, 51–59 (2017).
- 45. Thevenot, C. *et al.* The use of automated procedures by older adults with high arithmetic skills during addition problem solving. *Psychol Aging* **35**, 411–420 (2020).
- 46. Rutter, L. A., Vahia, I. V., Forester, B. P., Ressler, K. J. & Germine, L. Heterogeneous Indicators of Cognitive Performance and Performance Variability Across the Lifespan. *Front Aging Neurosci* 12, 62 (2020).
- 47. Yano, M. Aging effects in response inhibition: general slowing without decline in inhibitory functioning. *J Hum Ergol (Tokyo)* **40**, 129–139 (2011).
- 48. Clark, R., Hazeltine, E., Freedberg, M. & Voss, M. W. Age differences in episodic associative learning. *Psychol Aging* **33**, 144–157 (2018).
- Jin, Y., Jing, M. & Ma, X. Effects of Digital Device Ownership on Cognitive Decline in a Middle-Aged and Elderly Population: Longitudinal Observational Study. *J. Med. Internet Res.* 21, e14210 (2019).

- 50. Granic, I., Lobel, A. & Engels, R. C. M. E. The benefits of playing video games. *Am Psychol* 69, 66–78 (2014).
- Xu, W. *et al.* Education and Risk of Dementia: Dose-Response Meta-Analysis of Prospective Cohort Studies. *Mol. Neurobiol.* 53, 3113–3123 (2016).
- 52. Livingston, G. *et al.* Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet* **396**, 413–446 (2020).
- 53. van der Lee, S. J. *et al.* The effect of APOE and other common genetic variants on the onset of Alzheimer's disease and dementia: a community-based cohort study. *Lancet Neurol* 17, 434–444 (2018).
- 54. Li, C.-I. *et al.* Risk score prediction model for dementia in patients with type 2 diabetes. *Eur J Neurol* **25**, 976–983 (2018).
- 55. Zhang, X. *et al.* Chronic Obstructive Pulmonary Disease as a Risk Factor for Cognitive Dysfunction: A Meta-Analysis of Current Studies. *J. Alzheimers Dis.* **52**, 101–111 (2016).
- 56. Stefanidis, K. B., Askew, C. D., Greaves, K. & Summers, M. J. The Effect of Non-Stroke Cardiovascular Disease States on Risk for Cognitive Decline and Dementia: A Systematic and Meta-Analytic Review. *Neuropsychol Rev* 28, 1–15 (2018).
- Zheng, F., Yan, L., Zhong, B., Yang, Z. & Xie, W. Progression of cognitive decline before and after incident stroke. *Neurology* 93, e20–e28 (2019).
- 58. Omberg, L. *et al.* Remote smartphone monitoring of Parkinson's disease and individual response to therapy. *Nat Biotechnol* (2021) doi:10.1038/s41587-021-00974-9.
- Ogunniyi, A., Lekwauwa, U. G. & Osuntokun, B. O. Influence of education on aspects of cognitive functions in non-demented elderly Nigerians. *Neuroepidemiology* 10, 246–250 (1991).
- 60. Bonnechère, B. & Sahakian, B. J. Can Mobile Technology Help Prevent the Burden of Dementia in Low- and Mid-Income Countries? *Front Public Health* **8**, 554938 (2020).

- Stojan, R. & Voelcker-Rehage, C. A Systematic Review on the Cognitive Benefits and Neurophysiological Correlates of Exergaming in Healthy Older Adults. *J Clin Med* 8, E734 (2019).
- 62. Kraft, J. N. *et al.* Structural Neural Correlates of Double Decision Performance in Older Adults. *Front Aging Neurosci* **12**, 278 (2020).

FIGURES CAPTION

Figure 1: Evolution of the processing time according to age for the different CMG. Green color indicates the PT for correct responses, red color for incorrect responses, black dashed lines represent the accuracy. The small boxes (blue lines) represent the number of correct answers per CMG session.

Figure 2: Yearly changes in processing time across the different cognitive abilities for correct and incorrect responses.

Figure 3: Evolution of processing time for the different cognitive functions across lifespan. To ease the comparison of the different functions and interpretation data were centered for each different CMG by removing the average values of younger participants (18-21 years old).