

Prevalence and biopsychosocial factors associated with chronic low back pain in urban and rural communities in Western Africa: a population-based door-to-door survey in Benin

Peer-reviewed author version

KOSSI, Oyene; Yamadjako, Deneuve; TIMMERMANS, Annick; MICHIELS, Sarah; Adoukonou, Thierry & JANSSENS, Lotte (2022) Prevalence and biopsychosocial factors associated with chronic low back pain in urban and rural communities in Western Africa: a population-based door-to-door survey in Benin. In: *European spine journal* (Print), 31 (8).

DOI: 10.1007/s00586-022-07345-1

Handle: <http://hdl.handle.net/1942/37851>

1 **Title :** Prevalence and biopsychosocial factors associated with chronic low back pain in urban
2 and rural communities in Western Africa: A population-based door-to-door survey in Benin

3

4 Oyéné Kossi^{1,2}, Deneuve Yamadjako¹, Annick Timmermans², Sarah Michiels², Thierry
5 Adoukonou^{1,3}, Lotte Janssens²

6

7 ¹ ENATSE (Ecole Nationale de Santé Publique et de Surveillance Epidémiologique),
8 Université de Parakou, Parakou, Benin.

9 ² REVAL, Rehabilitation Research Center, Hasselt University, Belgium.

10 ³ Department of Neurology, Faculty of Medicine, University of Parakou, Parakou, Benin.

11

12

13 **Corresponding Author:**

14 Oyene Kossi

15 ENATSE (Ecole Nationale de Santé Publique et de Surveillance Epidémiologique)

16 Université de Parakou, Parakou,

17 03 BP 10 Parakou, Benin

18 Phone: (+229) 95 01 66 05

19 Email: oyene.kossi@gmail.com

20

1 **ACKNOWLEDGMENTS**

2 The authors are grateful to Bakita Houeze, Mariam Salami, Edith Daga, Joanita Djohou,
3 Cyriaque Guedenon, Rivalin Gilardi Aho-Glele, and to Fidèle Agossou, from the ‘ENATSE
4 (Ecole Nationale de Santé Publique et de Surveillance Epidémiologique)’, University of
5 Parakou, for their helpful contribution to the data collection.

6

7 **STUDY FUNDING**

8 None

9

10 **DISCLOSURES**

11 The authors declare no potential conflicts of interest concerning the research, authorship, and
12 publication of this article.

13

1 **ABSTRACT**

2 **Purpose:** This study aimed to assess the prevalence of chronic low back pain (CLBP) and
3 related biopsychosocial factors in urban and rural communities in Benin.

4 **Methods:** This is a population-based observational cross-sectional survey. An interviewer-
5 administered electronic questionnaire was used to collect information on demographic, socio-
6 economic, behavioral, and psychological factors relating to CLBP risk factors and medical
7 history of participants. The numeric pain rating scale and the Beck Depression Inventory were
8 used to assess pain intensity and the level of depression, respectively. Bivariate analyses were
9 performed to investigate the association between sociodemographic, behavioral, and
10 psychological factors and CLBP. Sequential multiple regression analyses were subsequently
11 performed to predict the occurrence of CLBP.

12 **Results:** A total of 4320 participants, with a mean age±SD of 32.9±13.1 years, of which
13 40.7% were females and 50.1% from an urban area, were enrolled in the study. We found a
14 global prevalence rate of CLBP of 35.5% [95% CI: 34.1%–36.9%]. The prevalence in urban
15 areas was 30.68% [95%CI: 28.9%–32.8%]) whilst 40.2% was found in rural areas [95%CI:
16 38.1%–42.2%]. Age ($p<0.001$), level of education ($p=0.046$), marital status ($p<0.001$),
17 working status ($p<0.003$), tobacco use ($p<0.016$) and regular physical activity ($p<0.011$) were
18 associated with CLBP. In urban areas, only the level of education was able to predict the
19 prevalence of CLBP ($R^2=61\%$). In rural areas, CLBP was predicted by age, marital and
20 working status ($R^2=89\%$).

21 **Conclusions:** This study showed a high prevalence of CLBP among urban and rural
22 communities in Benin. Age, level of education, marital status, and working status were
23 significantly associated with CLBP in Benin.

24

25 **Keywords:** Chronic low back pain, prevalence, biopsychosocial factors.

26

1 INTRODUCTION

2 Chronic low back pain (CLBP) is defined as “low back pain lasting for more than three
3 months, or as episodic low back pain within 6 months” [1]. CLBP is known as one of the
4 most common and disabling chronic pain conditions, affecting up to 19.6% of individuals
5 aged 20 to 59 years in high-income countries [2]. CLBP is among the most prevalent causes
6 of work absence and healthcare consumption worldwide [3]. Despite these facts, CLBP has
7 rarely been a focus of public health programs, especially in low- and middle-income countries
8 [4]. This was recently confirmed by a call for action initiative [5]. As a result, the socio-
9 economic problem of CLBP is currently underestimated and has even been ignored for a long
10 time, mainly due to its low mortality rate and because of considered often as being
11 irreversible or simply part of the ageing process.

12 To date, very few population-based studies investigated the prevalence of CLBP and its
13 associated factors in Africa [6–8]. Twenty years ago, Omokhodion assessed the prevalence of
14 low back pain in a rural community in South West Nigeria and found a 44% prevalence rate
15 of low back pain [9]. Risk factors were male gender and farming as an occupation. More
16 recently, Igwesi-Chidobe et al.,[7] investigated which biopsychosocial factors associate with
17 CLBP disability in rural Enugu State, South-eastern Nigeria. These authors did not report the
18 prevalence rate of CLBP but found that illness perceptions, pain intensity, catastrophizing,
19 fear-avoidance beliefs, lack of social support, and female gender were significant predictors
20 of self-reported and performance-based disability amongst people with CLBP in these areas.
21 Overall, an issue that has become important in the modern healthcare system is the rural
22 health. Rural and urban social environments differ so much that studies should not generalize
23 findings across these populations [10,11]. Specifically, some previous research indicated that
24 rural populations are unique in culture, economics, lifestyle, values, population mix, social
25 organization, and behaviors relating to illness and healthcare [10,12]. A study in India

1 determined that more people with a rural background reported severe chronic pain than those
2 with an urban background [13]. The authors explained these findings by the lack of social
3 support as well as living or working in socially isolated environments.

4 To the best of our knowledge, no studies have been carried out in West African countries
5 investigating CLBP prevalence in rural compared to urban areas. Therefore, this study aimed
6 to assess the prevalence of CLBP in Benin, and subsequently investigate the associated
7 biopsychosocial factors in urban and rural communities.

8

9 **METHODS**

10 **Study design and setting**

11 This is a population-based cross-sectional survey conducted from April to June 2021 in six
12 cities in the Republic of Benin: three urban cities (Cotonou, Abomey-Calavi, and Parakou)
13 and three rural cities (Ketou, Dassa-Zoumè, and Pèrèrè). Rural cities are characterized as
14 small cities composed of mainly rural areas with suburbs and villages, as opposed to large
15 cities with a mainly urban character. Figure 1 shows the selected cities on a map of the
16 Republic of Benin.

17

18 **Ethics considerations**

19 This study received approval from the biomedical ethics committee of the University of
20 Parakou (certificate number: 0429/CLERB-UP/P/SP/R/SA). Informed consent and agreement
21 to participate in the study were obtained via signature or thumbprint.

22

23 **Study size**

24 A minimal sample size of 689 participants per city was estimated according to the Schwartz
25 formula, $N = (Z_{\alpha/2} * p * q) / i^2$, where N = minimal sample size, p = 25.80% (prevalence)[14]; q =

1 (1-p), $Z\alpha = 1,96$ (for $\alpha = 5\%$), and $i = 4\%$ (accuracy). A margin of 10% was applied to cover
2 potential refusals to participate.

3

4 **Participants**

5 Figure 2 describes the selection flow chart of the participants. The identification and
6 recruitment of the participants were conducted using a three-stage sampling technique. The
7 National Institute of Statistics and Economic Analysis (INSAE) provided the initial frame
8 based on data from the last general census (2013) of population and housing (RGPH4) in
9 Benin. The first stage consisted of a simple random sampling technique to select six cities:
10 one rural city and one urban city in the north (Departments of Borgou and Alibori), two rural
11 cities in the center (Departments of Collines and Plateaux), and two urban cities in the south
12 (Departments of Atlantique and Littoral). The second stage comprised a selection of 50% of
13 neighborhoods in the selected cities through a simple random sampling approach. The number
14 of households to be surveyed in each neighborhood was obtained by dividing the sample size
15 by the number of neighborhoods. Participating households were identified by a systematic
16 sampling approach. The first household to be surveyed in each neighborhood was randomly
17 identified from the middle of the neighborhood by throwing a pen and by subsequently
18 following the direction of the pen direction. The third and last stage consisted of a random
19 selection of one individual per household according to the Kish method as recommended by
20 the WHO [15].

21

22 **Inclusion criteria**

23 To be included in the study, participants had to meet the following criteria: aged ≥ 18 years,
24 resident of the city for ≥ 6 months, and absence of any major cognitive impairments that could
25 interfere with the survey response.

1

2 **Exclusion criteria**

3 Exclusion criteria were pregnancy, history of spinal surgery, and red flags indicative of
4 serious spinal pathologies like cancer, traumas, or infection.

5

6 **Variables and measurement**

7 Sociodemographic information, as well as behavioral data, were collected using a general
8 questionnaire. Participants' body height was obtained from their ID card information while
9 their body weight was obtained using a mechanical scale, type SECA. Those with $18.5 \leq$
10 Body Mass Index (BMI) ≤ 25 were classified as having a normal body weight while those
11 with BMI > 25 were classified as being overweight or those with BMI > 30 as obese [16].
12 Those who smoked occasionally as well as those who smoked regularly were all considered
13 as being tobacco users. We also recorded peoples' self perspectives on being regularly
14 physically active or not. Those who were physically active were consecutively questioned
15 about the weekly frequency and duration of their physical activity

16

17 ***Numeric pain rating scale (NPRS)***

18 The Numerical Pain Rating Scale (NPRS) was used to evaluate pain severity and has been
19 reported to have good psychometric properties and clinical applicability [17]. The scores
20 range from 0 with 'no pain' to 10 with 'unacceptable pain'.

21

22 ***Presence of chronic low back pain***

23 The presence of CLBP was defined as having pain (NPRS score > 0) between the 12th rib and
24 the gluteal cleft, with or without radiation to the legs [18], lasting at least 12 weeks without a
25 specific underlying pathology or occurring episodically within 6 months [18].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Beck Depression Inventory (BDI)

We used the BDI to assess the severity of depression.[19] The BDI is a widely used tool to discriminate between chronic pain patients with and without major depression [20]. The total score of the BDI ranges from 0 to 63 with higher scores corresponding to a higher level of depression. Specifically, a score of 0-9 corresponds to the absence of depression while 10-15 corresponds to mild depression, 16-23 to moderate depression, and 24-63 to severe depression. The reliability of the BDI is high (Cronbach's $\alpha = 0.88$) [21].

Study procedures

An interviewer-administered electronic questionnaire was used to collect information on sociodemographic, behavioral, and psychological factors relating to CLBP risk factors, and medical history including past diagnosis of CLBP. All researchers involved in data collection were briefed extensively on the study methodology and underwent rigorous training to ensure consistency and compliance with the study procedures.

Statistical analyses

Data were analyzed using Epi Info 7.2.2.6 software. Data were assessed for normality using the graphical method of Q-Q plot. Nominal and ordinal variables were presented as proportions while quantitative variables with normal distribution were presented as means with standard deviation (SD). Bivariate analyses with Fisher's exact tests were used to investigate associations between sociodemographic, behavioral, psychological data, and geographical areas while the associations with CLBP were analyzed through the odds ratio (OR) with a 95% confidence interval. Confidence intervals could not overlap nor include 0, and in case they did, the difference was deemed not to be significant. Variables with a

1 significant association with CLBP were entered into a sequential multiple logistic regression
2 analysis to determine factors predicting CLBP. An adjusted coefficient of determination (R^2),
3 which considers the number of selected variables, was used to avoid overestimating the
4 predictive variance of the regression equation. Interactions between variables were examined
5 by testing the significance of the model and individual variables without each interaction and
6 after adding it. The association of each variable with CLBP was tested for inclusion in the
7 model using a significance level that was higher than the one set to define a cofactor.[22] This
8 is usually set at $P < 0.2$ for bivariate analysis and then set at $P < 0.05$ for retaining a factor in the
9 final model [22].

10

11

12 **RESULTS**

13 **Sociodemographic characteristics**

14 Table 1 describes the sociodemographic characteristics of the sample. Of the 4320
15 respondents, 50.09% were enrolled in an urban area, and 40.67% were females. The
16 mean \pm SD age of respondents was 32.85 \pm 13.08 years. Most of the participants were living
17 together as a couple (62.22%) and more than three quarters were independent workers
18 (56.64%) or were unemployed (26.44%). All the examined sociodemographic characteristics
19 were significantly different between urban and rural cities. Specifically, in urban cities, there
20 were significantly more young people ($p=0.001$), more males ($p=0.001$), more people with a
21 high education level ($p < 10^{-6}$), fewer self-employed people ($p < 10^{-4}$), and more paid workers
22 ($p < 10^{-4}$) compared to rural areas.

23

24 **Behavioral and psychological factors**

1 Table 2 presents the distribution of the behavioral and psychological factors in the sample. All
2 the examined behavioral and psychological factors were significantly different between urban
3 and rural cities, except for tobacco use ($p=0.368$). Specifically, significantly more people in
4 urban areas declared that they practice regular physical activity compared to rural areas ($p<10^{-6}$).
5 However, in urban areas, most of the people practice 1–2 times a week whilst those in rural
6 areas practice at least 3 times per week ($p<10^{-6}$). Overall the time spent on physical activity
7 was balanced between the two areas. Regarding BMI and depression, there were more
8 overweight or obese people ($p<10^{-3}$), and more people with depression ($p<10^{-6}$) in urban
9 compared to rural cities. Specifically, the mean \pm SD BMI of the sample was 22.75 ± 3.94
10 kg.m^{-2} . A total of 567 people, 13.13% [95%CI: 12.15%–14.16%] were overweight while 245
11 people, 5.67% [95%CI: 5.02%–6.40%] were obese. There were significantly more overweight
12 people in urban areas compared to rural areas (14.56% [95%CI: 13,13%–16,11%] versus
13 11.69% [95%CI: 10,40%–13,11%] respectively, $p=0.022$). The prevalence of people with
14 obesity was similar in both areas ($p=0.74$).

15

16 **Overall and area-specific prevalence of CLBP**

17 Table 3 shows the global prevalence and the region-specific prevalence of CLBP in this study.
18 We found a global prevalence rate of 35.49% [95%CI: 34.07% – 36.93%] of CLBP. This
19 prevalence varied between cities. Overall, rural areas showed a significantly higher
20 prevalence rate (40.17% [95%CI: 38.12% – 42.25%]) compared to urban areas (30.68%
21 [95%CI: 28.91% – 32.80%]) ($p<10^{-6}$).

22

23 **The association between sociodemographic factors and CLBP**

24 Table 4 shows the results of the association between sociodemographic factors and CLBP in
25 urban and rural areas. In urban areas, education level and work status were significantly

1 associated with CLBP. More specifically, bivariate analyses showed that a higher education
2 level was associated with a lower likelihood of having CLBP. Also, unemployed people and
3 students had 1.77 higher odds of having CLBP compared to self-employed people.
4 Furthermore, in rural areas, age, marital status, and working status were associated with
5 CLBP. Also, those who were living together as a couple had 1.48 higher odds of CLBP
6 compared to those who were living alone. In addition, retirees and those who have unpaid
7 work showed 1.53 higher odds compared to self-employed people.

8

9 **The association between behavioral/psychological factors and CLBP**

10 Table 5 presents the results of the association between behavioral and psychological factors
11 on the one hand and CLBP on the other hand, in urban and rural areas. No association was
12 found.

13

14 **Factors predicting CLBP**

15 Table 6 presents the final model predicting CLBP in both urban and rural areas. In urban
16 areas, only education level was predictive with a lower education level being a predicting
17 factor for having CLBP. This model explains about 61% of the total variance of the odds of
18 having CLBP in urban areas. On the other hand, in rural areas, CLBP was predicted by age,
19 marital status, and working state. This model explains about 89% of the total observed
20 variance of the odds of having CLBP in rural areas.

21

1 **DISCUSSION**

2 This study aimed to assess the prevalence of CLPB in Benin and subsequently investigate the
3 associated biopsychosocial factors in urban and rural communities in Benin. Overall, the
4 results showed high prevalence rates of CLBP up to 35%. Several factors were associated
5 with having CLBP such as age, level of education, marital status, working status, tobacco use,
6 and regular physical activity. However, in urban areas, only the level of education
7 significantly predicted CLBP while in rural areas age, marital status, and working status were
8 the significant predictors.

9 The prevalence rate of CLBP found in Benin is far higher than that reported by several
10 population-based studies in developed countries such as the USA (8.1%)[23], Spain
11 (11.12%)[24], and Canada (15.7% – 23.3%)[25]. This is in line with a previous review[26]
12 which estimated the point prevalence of low back pain among Africans at 39% which is
13 considerably higher than the global low back pain prevalence estimate (18.3%) reported by
14 Hoy et al.(2012)[27]. Overall, estimates from the global burden of disease study in 2017
15 found that globally low back pain continued to be the leading cause of years lived with
16 disability[28]. Nevertheless, countries and health-related organizations continue to prioritize
17 communicable diseases over non-communicable diseases such as low back pain. This is
18 especially true in the current Covid-19 pandemic context. In 2018, The Lancet Low Back Pain
19 Series made a call for action on the management of low back pain burden from governments,
20 policymakers, and the broader society [29]. Authors suggested establishing integrated and
21 collaborative approaches built upon affordable solutions to the growing burden of low back
22 pain in low- and middle-income countries such as Benin [30,31]. This is especially relevant
23 given that many of the risk factors for CLBP are shared by other non-communicable chronic
24 diseases.

1 The findings of this study showed that people aged above 60 had two times more odds of
2 having CLBP compared to 18-29-year-old people, specifically in rural areas. Earlier research
3 confirms that the incidence and prevalence of CLBP increase with older age [32,33]. Various
4 age-related physical and psychological changes (e.g., degenerative changes, physical
5 inactivity, slower reaction time, and changes in central pain processing), as well as multiple
6 risk factors (e.g., genetic, gender, and ethnicity), may affect the incidence, prognosis and
7 management of CLBP in older adults [34]. In the context of this study, specifically in rural
8 areas, agriculture is the predominant socio-economic occupation. Unfortunately, most of the
9 farmers still practice agriculture with rudimentary means due to their low accessibility to
10 mechanization. It is well established that physical and psychosocial work stressors relate to
11 employees' work-related musculoskeletal symptoms [38]. Then, the accumulation of
12 physically-demanding working hours over the years in conjunction with other risk factors
13 would explain the high exposure to CLBP of older adults compared to young. As a
14 consequence, it is reasonable that work status is associated with CLBP in both rural and urban
15 areas and that it is among the predictors of CLBP in rural areas. Moreover, biophysical factors
16 including working conditions and psychosocial factors including back pain beliefs may also
17 contribute to the significant discrepancy found in the prevalence of CLBP concerning
18 geographic residence. Negative beliefs about back pain are described as a signal of an
19 impending threat, which may lead to fear of movement, decreased function and activity, and
20 consequently persistent chronic disability [39].

21 In accordance with the above, the results of the present study showed that education level is
22 significantly lower in rural compared to urban areas with about one-third of people in rural
23 areas having no formal education. In addition, this study revealed that people with higher
24 education levels presented lower odds of developing CLBP compared with illiterates and this
25 factor was the only one that significantly explained the occurrence of CLBP in people living

1 in urban areas. These findings are consistent with those of prior studies identifying lower
2 education level as being associated with an increased risk of low back pain and associated
3 disability [40,41]. A recent study from KwaZulu-Natal (South Africa) also found that people
4 with no formal education had about 6 times more risk of having CLBP [42]. This association
5 could reflect variations in behavioral and environmental risk factors as well as variations in
6 living and work conditions. People with higher education levels are more likely to be in
7 professional, managerial, or other skilled occupations that are generally less physically
8 demanding and where there is more flexibility to eliminate pain-provoking job situations
9 [41,43]. In addition, people with higher education levels are more likely to have adequate
10 access to health services and to develop adaptive stress coping strategies [41].

11

12 **STUDY STRENGTH AND LIMITATIONS**

13 The findings of this study involving a representative sample of 4320 participants are a
14 valuable contribution to the evidence on CLBP and its associated factors in Benin. Our results
15 are consistent with models corrected for potential selection bias meaning that our findings
16 may be generalized to the population. In addition, the multiple logistic regression model used
17 to test for associations and predict the occurrence of CLBP is appropriate and easy to interpret
18 for a large audience. It not only provides a measure of how appropriate a predictor
19 (coefficient size) is but also its direction of association (positive or negative).

20 Limitations of our study include recall bias that may have affected the declarative reports of
21 participants about their CLBP. This includes the estimation of the duration of pain. Therefore,
22 chances of under- or over-estimating the complaint cannot be overcome with certainty, but we
23 expect this influence to be minimal. Another limitation of this study is that it did not include
24 the burden or consequences of CLBP such as days of sick leave, visits to physicians or
25 physiotherapists, length of hospitalization, functional limitations, and quality of life. This

1 information is crucial to establishing a more comprehensive view of whether or not CLBP is a
2 major health problem in Benin.

3

4

5 **CONCLUSION**

6 This study showed a high prevalence of CLBP among urban and rural communities in Benin.

7 Age, level of education, marital status, and working status were significantly associated with

8 CLBP. We suggest that future studies should examine the burden of CLBP in Benin. We also

9 suggest that the health authorities pay more attention to primary prevention and effective

10 management of CLBP by addressing the modifiable risk factors.

11

1 **STUDY FUNDING**

2 None

3

4 **DISCLOSURES**

5 The authors declare no potential conflicts of interest concerning the research, authorship, and
6 publication of this article.

7

8

1 REFERENCES

- 2 1. Balagué F, Mannion AF, Pellisé F, Cedraschi C. Non-specific low back pain.
3 Lancet. 4 févr 2012;379(9814):482-91.
- 4 2. Meucci RD, Fassa AG, Faria NMX. Prevalence of chronic low back pain: a
5 systematic review. Rev Saude Publica. 2015;49.
- 6 3. GBD 2013 DALYs and HALE Collaborators, Murray CJL, Barber RM, Foreman
7 KJ, Abbasoglu Ozgoren A, Abd-Allah F, et al. Global, regional, and national
8 disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy
9 life expectancy (HALE) for 188 countries, 1990-2013: quantifying the
10 epidemiological transition. Lancet. 28 nov 2015;386(10009):2145-91.
- 11 4. Hoy D, Geere JA, Davatchi F, Meggitt B, Barrero LH. A time for action:
12 Opportunities for preventing the growing burden and disability from
13 musculoskeletal conditions in low- and middle-income countries. Best Pract Res
14 Clin Rheumatol. juin 2014;28(3):377-93.
- 15 5. Sharma S, McAuley JH. Low Back Pain in Low- and Middle-Income Countries,
16 Part 1: The Problem. J Orthop Sports Phys Ther. mai 2022;52(5):233-5.
- 17 6. Igwesi-Chidobe CN, Coker B, Onwasigwe CN, Sorinola IO, Godfrey EL.
18 Biopsychosocial factors associated with chronic low back pain disability in rural
19 Nigeria: a population-based cross-sectional study. BMJ Glob Health.
20 2017;2(3):e000284.
- 21 7. Igwesi-Chidobe CN, Kitchen S, Sorinola IO, Godfrey EL. « A life of living
22 death »: the experiences of people living with chronic low back pain in rural
23 Nigeria. Disabil Rehabil. avr 2017;39(8):779-90.
- 24 8. Tarimo N, Diener I. Knowledge, attitudes and beliefs on contributing factors
25 among low back pain patients attending outpatient physiotherapy treatment in
26 Malawi. S Afr J Physiother. 2017;73(1):395.
- 27 9. Omokhodion FO. Low back pain in a rural community in South West Nigeria.
28 West Afr J Med. juin 2002;21(2):87-90.
- 29 10. Goode AP, Freburger JK, Carey TS. The influence of rural versus urban residence
30 on utilization and receipt of care for chronic low back pain. J Rural Health.
31 2013;29(2):205-14.
- 32 11. Rafferty AP, Luo H, Egan KL, Bell RA, Gaskins Little NR, Imai S. Rural,
33 Suburban, and Urban Differences in Chronic Pain and Coping Among Adults in
34 North Carolina: 2018 Behavioral Risk Factor Surveillance System. Prev Chronic
35 Dis. 18 févr 2021;18:E13.

- 1 12. Probst JC, Laditka SB, Wang JY, Johnson AO. Effects of residence and race on
2 burden of travel for care: cross sectional analysis of the 2001 US National
3 Household Travel Survey. *BMC Health Serv Res.* 9 mars 2007;7:40.
- 4 13. Varma VK, Malhotra A, Chaturvedi SK, Chari P. Sociodemographic study of
5 patients with chronic pain. *Indian J Psychiatry.* avr 1986;28(2):119-25.
- 6 14. Zomalhèto Z, Mikponhoué RCN, Wanvoègbe A, Adikpéto I, Ayélo P. Prévalence
7 et facteurs associés à la lombalgie chez les conducteurs de taxi moto à Porto-
8 Novo (Bénin). *The Pan African Medical Journal [Internet].* 2019 [cité 24 oct
9 2021];32. Disponible sur:
10 <https://www.ncbi.nlm.nih.gov/sites/ppmc/articles/PMC6560951/>
- 11 15. Kish L. A Procedure for Objective Respondent Selection within the Household.
12 *Journal of the American Statistical Association.* sept 1949;44(247):380-7.
- 13 16. CDC. Defining Adult Overweight and Obesity [Internet]. Centers for Disease
14 Control and Prevention. 2021 [cité 9 déc 2021]. Disponible sur:
15 <https://www.cdc.gov/obesity/adult/defining.html>
- 16 17. Farrar JT, Young JP, LaMoreaux L, Werth JL, Poole MR. Clinical importance of
17 changes in chronic pain intensity measured on an 11-point numerical pain rating
18 scale. *Pain.* nov 2001;94(2):149-58.
- 19 18. Rozenberg S, Foltz V, Fautrel B. Treatment strategy for chronic low back pain.
20 *Joint Bone Spine.* déc 2012;79(6):555-9.
- 21 19. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for
22 measuring depression. *Arch Gen Psychiatry.* juin 1961;4:561-71.
- 23 20. Geisser ME, Roth RS, Robinson ME. Assessing depression among persons with
24 chronic pain using the Center for Epidemiological Studies-Depression Scale and
25 the Beck Depression Inventory: a comparative analysis. *Clin J Pain.* juin
26 1997;13(2):163-70.
- 27 21. Wang YP, Gorenstein C. Psychometric properties of the Beck Depression
28 Inventory-II: a comprehensive review. *Braz J Psychiatry.* déc 2013;35(4):416-31.
- 29 22. Zhang Z. Variable selection with stepwise and best subset approaches. *Ann*
30 *Transl Med.* avr 2016;4(7):136.
- 31 23. Johannes CB, Le TK, Zhou X, Johnston JA, Dworkin RH. The prevalence of
32 chronic pain in United States adults: results of an Internet-based survey. *J Pain.*
33 nov 2010;11(11):1230-9.
- 34 24. Palacios-Ceña D, Alonso-Blanco C, Hernández-Barrera V, Carrasco-Garrido P,
35 Jiménez-García R, Fernández-de-las-Peñas C. Prevalence of neck and low back

- 1 pain in community-dwelling adults in Spain: an updated population-based
2 national study (2009/10-2011/12). *Eur Spine J.* mars 2015;24(3):482-92.
- 3 25. Alkherayf F, Agbi C. Cigarette smoking and chronic low back pain in the adult
4 population. *Clin Invest Med.* 1 oct 2009;32(5):E360-367.
- 5 26. Morris LD, Daniels KJ, Ganguli B, Louw QA. An update on the prevalence of
6 low back pain in Africa: a systematic review and meta-analyses. *BMC*
7 *Musculoskelet Disord.* 21 juin 2018;19(1):196.
- 8 27. Hoy D, Bain C, Williams G, March L, Brooks P, Blyth F, et al. A systematic
9 review of the global prevalence of low back pain. *Arthritis Rheum.* juin
10 2012;64(6):2028-37.
- 11 28. Wu A, March L, Zheng X, Huang J, Wang X, Zhao J, et al. Global low back pain
12 prevalence and years lived with disability from 1990 to 2017: estimates from the
13 Global Burden of Disease Study 2017. *Ann Transl Med.* mars 2020;8(6):299.
- 14 29. Buchbinder R, van Tulder M, Öberg B, Costa LM, Woolf A, Schoene M, et al.
15 Low back pain: a call for action. *Lancet.* 9 juin 2018;391(10137):2384-8.
- 16 30. Vlaeyen JWS, Maher CG, Wiech K, Van Zundert J, Meloto CB, Diatchenko L, et
17 al. Low back pain. *Nat Rev Dis Primers.* 13 déc 2018;4(1):52.
- 18 31. Nduwimana I, Nindorera F, Thonnard JL, Kossi O. Effectiveness of walking
19 versus mind-body therapies in chronic low back pain: A systematic review and
20 meta-analysis of recent randomized controlled trials. *Medicine (Baltimore).* 28
21 août 2020;99(35):e21969.
- 22 32. Dionne CE, Dunn KM, Croft PR. Does back pain prevalence really decrease with
23 increasing age? A systematic review. *Age Ageing.* mai 2006;35(3):229-34.
- 24 33. Thomas E, Peat G, Harris L, Wilkie R, Croft PR. The prevalence of pain and pain
25 interference in a general population of older adults: cross-sectional findings from
26 the North Staffordshire Osteoarthritis Project (NorStOP). *Pain.* juill
27 2004;110(1-2):361-8.
- 28 34. Wong AY, Karppinen J, Samartzis D. Low back pain in older adults: risk factors,
29 management options and future directions. *Scoliosis Spinal Disord.* 18 avr
30 2017;12:14.
- 31 38. Eatough EM, Way JD, Chang CH. Understanding the link between psychosocial
32 work stressors and work-related musculoskeletal complaints. *Applied*
33 *Ergonomics.* 1 mai 2012;43(3):554-63.
- 34 39. Liu H, Huang L, Yang Z, Li H, Wang Z, Peng L. Fear of Movement/(Re)Injury:
35 An Update to Descriptive Review of the Related Measures. *Front Psychol.*
36 2021;12:696762.

- 1 40. Dionne CE, Von Korff M, Koepsell TD, Deyo RA, Barlow WE, Checkoway H.
2 Formal education and back pain: a review. *J Epidemiol Community Health*. juill
3 2001;55(7):455-68.
- 4 41. Reisbord LS, Greenland S. Factors associated with self-reported back-pain
5 prevalence: a population-based study. *J Chronic Dis*. 1985;38(8):691-702.
- 6 42. Kahere M, Ginindza T. The prevalence and risk factors of chronic low back pain
7 among adults in KwaZulu-Natal, South Africa: an observational cross-sectional
8 hospital-based study. *BMC Musculoskelet Disord*. 15 nov 2021;22:955.
- 9 43. Haber LD. Disabling effects of chronic disease and impairment. II. Functional
10 capacity limitations. *J Chronic Dis*. mars 1973;26(3):127-51.
- 11