



# BMJ Open Association between incident falls and subsequent fractures in patients attending the fracture liaison service after an index fracture: a 3-year prospective observational cohort study

Lisanne Vranken <sup>1,2</sup>, Caroline E Wyers <sup>1,2</sup>, Robert Y Van der Velde<sup>1,2</sup>, Heinrich M J Janzing,<sup>3</sup> Sjoerd Kaarsemakers,<sup>4</sup> Johanna Driessen,<sup>5,6</sup> John Eisman,<sup>7,8</sup> Jacqueline R Center,<sup>7,9</sup> Tuan V Nguyen,<sup>7,10</sup> Thach Tran,<sup>7</sup> Dana Bliuc,<sup>7</sup> Piet Geusens,<sup>11,12</sup> Joop P van den Bergh<sup>1,2</sup>

**To cite:** Vranken L, Wyers CE, Van der Velde RY, *et al.* Association between incident falls and subsequent fractures in patients attending the fracture liaison service after an index fracture: a 3-year prospective observational cohort study. *BMJ Open* 2022;**12**:e058983. doi:10.1136/bmjopen-2021-058983

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-058983>).

Received 27 November 2021  
Accepted 04 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Joop P van den Bergh;  
jvdbergh@viecuri.nl

## ABSTRACT

**Objectives** To evaluate the risk of subsequent fractures in patients who attended the Fracture Liaison Service (FLS), with and without incident falls after the index fracture.

**Design** A 3-year prospective observational cohort study.

**Setting** An outpatient FLS in the Netherlands.

**Participants** Patients aged 50+ years with a recent clinical fracture.

**Outcome measures** Incident falls and subsequent fractures.

**Results** The study included 488 patients (71.9% women, mean age: 64.6±8.6 years). During the 3-year follow-up, 959 falls had been ascertained in 296 patients (60.7% (ie, fallers), and 60 subsequent fractures were ascertained in 53 patients (10.9%). Of the fractures, 47 (78.3%) were fall related, of which 25 (53.2%) were sustained at the first fall incident at a median of 34 weeks. An incident fall was associated with an approximately 9-fold (HR: 8.6, 95% CI 3.1 to 23.8) increase in the risk of subsequent fractures.

**Conclusion** These data suggest that subsequent fractures among patients on treatment prescribed in an FLS setting are common, and that an incident fall is a strong predictor of subsequent fracture risk. Immediate attention for fall risk could be beneficial in an FLS model of care.

**Trial registration number** NL45707.072.13.

## INTRODUCTION

Patients with a recent fracture have a high imminent risk of subsequent fractures as shown after most fractures,<sup>1–6</sup> and a high risk of subsequent falls, as shown after a recent hip fracture.<sup>7–11</sup> The Fracture Liaison Service (FLS) is considered the most effective organisational approach for secondary fracture prevention in patients after the age of 50 years with a recent fracture.

Most fractures are caused by a fall, but most falls do not result in a fracture.<sup>12 13</sup> Falls are a major contributing factor to the occurrence of fractures, independent and additive to the risk attributable to age and bone mineral

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Although this is one of the largest prospective studies in a Fracture Liaison Service population focusing on the incidence of falls after an index fracture, the number of patients is modest and the number of subsequent fractures relatively small.
- ⇒ Data on falls were collected prospectively using fall diaries, but no procedures were in place to validate self-reported falls.
- ⇒ No information was available on falls between the index fracture and enrolment in the study.
- ⇒ Relatively healthy patients participated in the study, which may have resulted in an underestimation of incident falls and subsequent fractures.

density (BMD).<sup>14–17</sup> Guidelines on the FLS, therefore, recommend fall prevention and prescription of anti-osteoporosis medication (AOM) in patients at a high risk.<sup>18–22</sup>

However, it is not well known to what extent the imminent risk of subsequent fractures after an index fracture can be attributed to incident falls. We hypothesised that the risk of subsequent fractures would be substantially higher in patients with falls after a recent fracture than in those without falls. The aim of this study was, therefore, to evaluate the incidence of falls and subsequent fractures, and the risk of subsequent fractures in those with and without falls after a recent index fracture in patients who attend the FLS.

## METHODS

### Study population and design

A 3-year prospective observational cohort study was conducted, including 500 consecutive patients aged between 50 years and

90 years, with a recent, radiologically confirmed clinical vertebral or non-vertebral low-trauma fracture, and who were willing and able to participate. Patients were recruited at the FLS in VieCuri Medical Center, Venlo, the Netherlands.

Low-trauma fractures were defined as fractures that resulted from a fall from standing height or less. Excluded were non-Caucasian patients, patients with bone metastasis, failure of prosthesis or osteomyelitis and patients with cognitive impairment.

According to standard care, a nurse specialised in osteoporosis invited all patients aged 50 years and older, who visited the emergency department because of a recent clinical vertebral or non-vertebral fracture, to the FLS. All patients who responded and agreed to be evaluated scheduled an appointment for fracture risk evaluation. Fracture risk evaluation included a detailed questionnaire for evaluation of risk factors for fractures and falls, including medical history and medication use. This questionnaire was based on the Dutch guidelines on osteoporosis and fracture prevention, and prevention of falls in the elderly.<sup>23 24</sup> Also, height and weight were measured, a BMD measurement with dual-energy X-ray absorptiometry of the lumbar spine, total hip and femoral neck, with vertebral fracture assessment was performed, and a blood sample was collected to detect contributors to secondary osteoporosis and metabolic bone disease.<sup>25</sup> According to the Dutch osteoporosis guideline,<sup>23</sup> AOM was started in patients with osteoporosis or having at least one moderate to severe prevalent vertebral fracture according to Genant *et al.*<sup>26</sup> Bisphosphonates and denosumab were first-choice treatments. Teriparatide was restricted to patients already on another AOM with at least three fractures, of which two were vertebral fractures.

### Falls and subsequent fractures

During the 3-year follow-up, patients were requested to record falls weekly in a fall diary. Fall registration started at the beginning of the study, mean 3.5±1.0 months after the index fracture. A fall was defined as an unintentional change in position resulting in coming to rest on the ground or other lower level.<sup>27</sup> Patients were asked to return their fall diaries by mail at 3 months and 6 months, and during the study visit at 1 year, 2 years and 3 years of follow-up. They were contacted by telephone if the fall diary was not received or incomplete. Patients were categorised as those with at least one incident fall (ie, faller) or without an incident fall (ie, non-faller) during follow-up.

When patients recorded a fall in their diary, they were also asked to record whether or not they sustained a subsequent clinical fracture as a direct result of the fall. Additionally, at 1-year, 2-year and 3-year follow-ups, patients had to complete a detailed questionnaire, including a question on whether they sustained a fracture due to another trauma than a fall or without an overt trauma. All subsequent fractures were radiologically confirmed according to radiology reports in the electronic patient

records. Since no imaging of the spine was performed at the end of the study, all reported vertebral fractures were symptomatic, clinical vertebral fractures. A distinction was made between subsequent fractures that were directly caused by a fall (ie, fall-related fractures) and those that occurred without an overt trauma or were the result of another trauma than a fall (ie, non-fall-related fractures).

### Data analysis

Baseline characteristics were compared between fallers and non-fallers, and between patients with and without subsequent fractures using the Student's t-test or Wilcoxon test for continuous variables, and Chi-squared or Fisher's exact test for categorical variables where appropriate. The incidence rate of falls and subsequent fractures per 100 person-years was estimated at 3 and 6 months and 1, 2 and 3 year follow-up, assuming a Poisson distribution. Kaplan Meier curves were made for incident falls and subsequent fractures, in which patients were included once, and only the first incident fall or subsequent fracture was included. Cox proportional hazards regression was used to determine the association between incident falls and subsequent fractures, yielding HRs and 95% CIs. Proportional hazard assumptions were not violated. Follow-up time was determined by the first subsequent fracture, lost to follow-up or the end of the study, whatever occurred first. All analyses were adjusted for the predefined covariates, including age, gender, index fracture type (major or hip vs any other fracture), BMD (lowest measured at lumbar spine, total hip and femoral neck) and prevalent vertebral fractures (moderate or severe vs mild or no prevalent vertebral fractures). A p value of <0.05 was considered statistically significant.

Two sensitivity analyses were planned: (1) excluding patients with index and subsequent finger or toe fractures and (2) by classifying patients with a non-fall-related subsequent fracture as non-faller, even if they fell at another time during follow-up.

### Patient and public involvement

Patients or members of the public were not involved in the design, or conduct, or reporting, or dissemination plans of the research.

## RESULTS

### Study population

Among 1220 patients who approached from the FLS, 1011 patients met the study criteria. Of the 1011 patients, 511 were not willing or able to participate in the study, and after excluding 12 patients with missing fall data, ultimately 488 patients were available for analysis (online supplemental figure 1) of whom 34 patients (7.0%) had incomplete follow-up data on incident falls (5 patients died, 8 withdrew consent and 21 had incomplete fall registration).

The mean time between the index fracture and FLS visit at which patients were included for this study was

**Table 1** Baseline characteristics of 488 participants stratified by incident fall and subsequent fracture status

	Total population (n=488)	Non-fallers (n=192)	Fallers (n=296)	P value	No subsequent fracture (n=435)	Subsequent fracture (n=53)	P value
Age (years)	64.6±8.6	64.4±8.0	64.8±9.0	0.608	64.5±8.8	65.3±7.1	0.488
Female gender	351 (71.9)	130 (67.7)	221 (74.7)	0.095	308 (70.8)	43 (81.1)	0.114
Baseline fracture							
Finger or toe	55 (11.3)	30 (15.6)	25 (8.4)	0.060	49 (11.3)	6 (11.3)	0.460
Minor	303 (62.1)	109 (56.8)	194 (65.5)		270 (62.1)	33 (62.3)	
Major	104 (21.3)	44 (22.9)	60 (20.3)		95 (21.8)	9 (17.0)	
Hip	26 (5.3)	9 (4.7)	17 (5.7)		21 (4.8)	5 (9.4)	
Fall related*	422 (86.5)	164 (85.4)	258 (87.2)	0.582	378 (86.9)	44 (83.0)	0.436
Fall previous year†							
0	349 (71.5)	155 (80.7)	194 (65.5)	<0.001	315 (72.4)	34 (64.2)	0.208
≥1	139 (28.5)	37 (19.3)	102 (34.5)		120 (27.6)	19 (35.8)	
BMI (kg/m <sup>2</sup> )	27.7±4.4	27.7±4.4	27.7±4.4	0.961	27.8±4.4	26.9±4.8	0.154
BMD							
Normal BMD	132 (27.1)	54 (28.1)	78 (26.4)	0.906	123 (28.3)	9 (17.0)	0.081
Osteopenia	249 (51.0)	97 (50.5)	152 (51.4)		222 (51.0)	27 (50.9)	
Osteoporosis	107 (21.9)	41 (21.4)	66 (22.3)		90 (20.7)	17 (32.1)	
Prevalent vertebral fracture‡§							
None	356 (73.0)	139 (72.4)	217 (73.3)	0.572	328 (75.4)	28 (52.8)	<0.001
Grade 1	62 (12.7)	22 (11.5)	40 (13.5)		54 (12.4)	8 (15.1)	
Grade 2–3	70 (14.3)	31 (16.1)	39 (13.2)		53 (12.2)	17 (32.1)	
Anti-osteoporosis treatment	167 (34.2)	70 (36.5)	97 (32.8)	0.402	142 (32.6)	25 (47.2)	0.035

Continuous variables are shown in mean±SD (SD) and categorical variables are shown as number of patients (%).  
 \*Signifying that fracture was caused by a fall.  
 †Fall resulting in baseline fracture not included.  
 ‡According to Genant *et al.*<sup>26</sup>  
 §According to most severe prevalent vertebral fracture.  
 BMD, bone mineral density; BMI, body mass index.

3.9±1.1 months for patients with a hip fracture and 3.5±1.0 months for patients with other fractures. Baseline characteristics of the 488 study participants are presented in table 1. Mean age was 64.6±8.6 years and 71.9% of the patients were women. In 86.5% of patients, the index fracture was caused by a fall, and 28.5% of patients had at least one other fall in the year before the start of the study. At baseline, 21.9% of patients were diagnosed with osteoporosis, 51.1% with osteopenia and 27.1% had a normal BMD. Lowest BMD was measured at the femoral neck in 470 participants, at the total hip in 3 participants and at the lumbar spine in 15 participants. Moderate to severe (ie, grade 2–3) prevalent vertebral fractures were present in 14.3% of patients. AOM was prescribed in 34.2% of patients (8 (1.6%) were already using AOM and 159 (32.6%) started using AOM at baseline visit).


Compared with eligible FLS attenders, who were not willing or able to participate in our study, patients included in our study were younger, had fewer major or hip fractures and had a higher BMD, and a lower

proportion had prevalent vertebral fractures (see online supplemental table 1).

### Falls

During a median follow-up of 3 years (range: 0.1–3.0 years), 296 patients (60.7%) recorded 959 falls, corresponding to 68.6 falls per 100 person-years. The cumulative fall incidences and incidence rates per 100 person-years at 3 months and 6 months, and at 1-year, 2-year and 3-year follow-ups are presented in figure 1. Of the 296 patients with at least one fall, 115 (38.9%) had one fall and 181 (61.1%) had two or more falls (up to 39 falls in one patient).

A first fall was recorded by 189/488 patients (38.7%) during the 1st year of follow-up, by 56/299 (18.7%) during the 2nd, and by 51/243 (21.0%) during the 3rd year of follow-up. The median time to the first fall was 34 weeks (range: 1–156 weeks). Of the 959 falls, 47 (4.9%) resulted in a subsequent fall-related fracture.



**Figure 1** Cumulative incidence of falls stratified by gender.

There were no significant differences in baseline characteristics between patients with and without a fall during the 3-year follow-up, except for that a higher proportion of patients with incident falls reported at least one fall in the year before the start of the study (34.5% vs 19.3%,  $p < 0.001$ ) (see [table 1](#)). There were no significant differences in baseline characteristics between patients with one fall and those with multiple falls (data not shown).

### Subsequent fractures


In total, 53 patients (10.9%) recorded 60 subsequent fractures, corresponding to 4.29 subsequent fractures per 100 person-years. The cumulative subsequent fracture incidences and incidence rates (per 100-person years) at 3 months and 6 months, and at 1-year, 2-year and 3-year follow-ups are presented in [figure 2](#). Of all subsequent fractures, 47 (78.3%) were fall-related and 13 (21.7%) were non-fall-related. Fall-related subsequent fracture sites were: radius and ulna (n=9), tibia and fibula (n=8), proximal femur (n=4), metatarsal (n=4), hand phalanx (n=4), symptomatic vertebra (n=3), proximal humerus (n=3), clavicle (n=3), costal bones (n=2), scapula (n=2), pelvic bone (n=1), metacarpal (n=1), tarsal (n=1), patella (n=1) and foot phalanx (n=1), whereas subsequent non-fall-related fractures sites were symptomatic vertebral

(n=5), metatarsal (n=2), foot phalanx (n=5) and hand phalanx (n=1). Half (53.2%) of all fall-related subsequent fractures were sustained at the first fall.

Baseline characteristics for patients with and without subsequent fractures are presented in [table 1](#).

Of the 296 patients with at least one fall, 41 (13.9%) had 46 fall-related subsequent fractures, 7 (2.4%) had seven non-fall-related subsequent fractures and 1 (0.3%) had 1 fall-related and 1 non-fall-related subsequent fractures. Of the 192 patients without a fall, 4 (2.1%) had 5 non-fall-related subsequent fractures. Of note, the risk of subsequent fractures was higher in patients with at least one fall than in those without a fall (adjusted HR: 8.6; 95% CI 3.1 to 23.8; cumulative incidence: 16.6% vs 2.1%) ([figure 3](#) and [table 2](#)). Results were similar when femoral neck BMD instead of the lowest BMD was used for adjustments (adjusted HR: 8.3; 95% CI 3.0 to 23.0). Additionally, subsequent fracture risk was higher in patients with moderate or severe prevalent vertebral fractures than in those with no or mild prevalent vertebral fractures (adjusted HR: 3.9; 95% CI 2.1 to 7.3; cumulative incidence: 24.3% vs 8.6%) ([table 2](#)).

The association between falls and subsequent fractures remained significant in sensitivity analyses (1) excluding



**Figure 2** Cumulative incidence of subsequent fractures stratified by gender.

patients with index and subsequent finger and toe fractures (adjusted HR: 8.2; 95% CI 2.5 to 26.6), and (2) by classifying patients with a non-fall-related subsequent fracture as non-faller (adjusted HR: 2.9; 95% CI 1.5 to 5.6).

## DISCUSSION


In this 3-year prospective observational cohort study in patients aged 50+ years with a recent clinical fracture, treated according to current Dutch osteoporosis guidelines at an FLS, 60.7% of patients had at least one fall and 10.9% had at least one subsequent fracture. The majority (78.3%) of subsequent fractures was caused by a fall, and of all fall-related subsequent fractures, 53.2% occurred at the first fall. Subsequent fracture risk was ninefold higher in fallers than in non-fallers.

Literature reporting fall incidence in patients with a fracture is limited. Comparable to our results, van Helden *et al*<sup>28</sup> reported a 3-month fall incidence of 15% in patients with a recent fracture at an FLS, and Matsumoto *et al*<sup>29</sup> reported a 1-year fall incidence of 40% in ambulatory patients with a recent fracture. Various other studies included older, hip fracture patients and reported higher 1 year fall incidences up to 55%,<sup>7–11</sup> except for the study from Yeh *et al* that reported a lower 1-year fall incidence (31%).<sup>30</sup> Higher fall incidences in hip fracture studies can partially be explained by the older study population. Unfortunately, other fall risk factors cannot be compared.

An explanation for the lower fall incidence in the study by Yeh *et al* may be that information on the occurrence of falls was provided by patients and family caregivers, which may have resulted in under registration of falls.

A comparison between the fall incidence in our study and that in the general population is difficult to make, because population-based studies were conducted in a 65+ years old, community-dwelling population, whereas approximately 50% of our study population was <65 years old. The proportion of community-dwelling people aged 65+ years sustaining at least 1 fall over a 1-year period ranged from 28% to 35%,<sup>31–33</sup> with an increasing incidence with increasing age.<sup>34</sup> The 1-year fall incidence reported in our study is comparable to that in an older (65+ years aged) population, and, therefore, relatively high. However, in contrast to what has been reported in the literature, we found no higher 3-year fall incidence with increasing age. An explanation for this could be that, especially in the older age group, relatively more healthy patients participated in our study, resulting in a lower fall incidence in older age group. Another explanation could be that patients aged 50–65 years are more physically active, and, therefore, fall more often.

Compared with our results, previously published FLS studies reported lower,<sup>35 36</sup> similar<sup>28 37 38</sup> and higher<sup>39 40</sup> subsequent fracture rates. Differences can be explained by differences in patient selection. Studies that included



**Figure 3** Cumulative incidence of subsequent fractures stratified by fall status.

older patients<sup>39</sup> and patients with more severe fractures<sup>40</sup> reported higher subsequent fracture rates, whereas studies that excluded hand and foot index and subsequent fractures<sup>35</sup> or frail patients reported lower rates.<sup>36</sup>

In 2010, the Dutch population consisted of approximately 6000000 people aged 50+ years, of whom 119419 sustained a fracture that year,<sup>41</sup> corresponding to a calculated annual fracture incidence of 2.0% in the general Dutch population of 50+ years. Compared with the general Dutch population of 50+ years, the fracture incidence was more than 2 times higher in our study, even in

the 3rd year of follow-up. In our study, fracture incidence remained high despite treatment according to the current osteoporosis guideline, raising the question of what more can be done to prevent subsequent fractures. Even though conflicting results have been published about the effect of fall prevention strategies on subsequent fracture,<sup>42</sup> we hypothesise that fall interventions could be effective in patients at highest risk, namely those with a recent fracture at risk of falling. Furthermore, according to literature, recurrent fallers have an almost fourfold increased odds of sustaining a fall-related fracture compared with individuals with a single fall.<sup>43</sup> However, we found that the majority of subsequent fall-related fractures occur at the first fall after the index fracture, with a median time to the first fall of 34 weeks. Interestingly, fall incidence was higher in the first year of follow-up compared with the second and third years. This may indicate an imminent fall risk, which may attribute to the imminent subsequent fracture risk after an index fracture.<sup>1-6</sup> This implies that the FLS patients with a high fall risk should be identified immediately, because there is a small window of opportunity to prevent falls and fall-related subsequent fractures.

Remarkably, in contrast to previous studies indicating that imminent fracture risk that was highest in the 1st year after an index fracture,<sup>44 45</sup> there was a linear subsequent fracture incidence during a 3-year follow-up in this study. An explanation for the linear subsequent fracture incidence may be the relatively healthy patients who

**Table 2** Predictors of refracture: results of the Cox's proportional hazard model

Predictor	Unit of comparison	HR and 95% CI	P value
Gender	Women vs men	1.39 (0.68 to 2.83)	0.362
Age	+5 years	0.97 (0.82 to 1.13)	0.662
Index fracture	Major or hip vs all other	0.68 (0.35 to 1.33)	0.263
BMD	-0.12 g/cm <sup>2</sup>	1.30 (0.95 to 1.78)	0.101
Prevalent vertebral fracture	Yes vs no	3.88 (2.07 to 7.27)	<0.0001
Fall	Yes vs no	8.58 (3.09 to 23.8)	<0.0001

BMD, bone mineral density.

agreed to participate in our study. Compared with non-attenders, they were younger, and a lower proportion had a major baseline fracture, a prevalent vertebral fracture and osteoporosis, and if indicated, were more likely to receive AOM. Importantly, in addition to falls, moderate to severe prevalent vertebral fractures at baseline were associated with subsequent fractures, even though anti-osteoporosis medication had been prescribed to these patients according to the current Dutch osteoporosis guideline.

This study has several limitations. Although, this is one of the largest prospective studies in an FLS population focusing on the incidence of falls after an index fracture, the number of patients is modest and the number of subsequent fractures relatively low. Therefore, the association between falls and fall-related, and non-fall-related subsequent fractures could not be analysed separately. A fall 'not resulting in a subsequent fracture' might indicate frailty of patients, and might be different from those falls that directly resulted in a subsequent fracture. Future studies are needed to investigate this difference. Finally, because of small numbers, subgroup analyses should not be performed. Furthermore, data on falls were collected prospectively using fall diaries that had to be returned at 3 months and 6 months, and 1 year, 2 years and 3 years. However, no procedures were in place to validate self-reported falls, and it is possible that recall bias could have led to underregistration of falls. Moreover, no information was available on falls between the index fracture and enrolment in the study. Finally, relatively healthy patients participated in the study. Compared with non-attenders, they were younger, a lower proportion had a major baseline fracture, a prevalent vertebral fracture and osteoporosis. The proportion of patients with a fall and subsequent fractures could be expected to be even higher in the total FLS population.

In conclusion, in this 3-year prospective observational cohort study in FLS patients, subsequent fracture incidence was high despite being prescribed anti-osteoporosis medications according to the current Dutch osteoporosis guidelines. Subsequent fracture risk was 9-fold higher in fallers than in non-fallers, and the majority of fall-related subsequent fractures occurred at the first fall at a median time of 34 weeks. These findings emphasise that immediate attention for fall risk reduction could be beneficial in FLS care. Various risk factors, including comorbidities, medication use, polypharmacy and alcohol use among others, contribute to patient's fall risk and further research is needed to determine predictors for falls to identify patients at highest risk of falling.

#### Author affiliations

<sup>1</sup>Department of Internal Medicine, VieCuri Medical Centre, Venlo, The Netherlands

<sup>2</sup>Department of Internal Medicine, Research School NUTRIM, Maastricht University Medical Centre+, Maastricht, The Netherlands

<sup>3</sup>Department of Surgery, VieCuri Medical Centre, Venlo, The Netherlands

<sup>4</sup>Department of Orthopedic Surgery, VieCuri Medical Centre, Venlo, The Netherlands

<sup>5</sup>Department of Clinical Pharmacy and Toxicology, Maastricht University Medical Centre+, Maastricht, The Netherlands

<sup>6</sup>Department of Clinical Pharmacy and Toxicology, Maastricht University, Maastricht, The Netherlands

<sup>7</sup>Department of Osteoporosis and Bone Biology, Garvan Institute of Medical Research, Sydney, New South Wales, Australia

<sup>8</sup>School of Medicine, The University of Notre Dame, Sydney, New South Wales, Australia

<sup>9</sup>School of Population Health, UNSW, Sydney, New South Wales, Australia

<sup>10</sup>School of Biomedical Engineering, University of Technology Sydney, Sydney, New South Wales, Australia

<sup>11</sup>Department of Internal Medicine, Subdivision Rheumatology, Research School CAPHRI, Maastricht University Medical Centre, Maastricht, The Netherlands

<sup>12</sup>Faculty of Medicine and Life Sciences, Hasselt University, Diepenbeek, Belgium

**Contributors** LV collected data, carried out data analysis and drafted the manuscript. CEW and JPvdB developed the study design, wrote the research protocol, collected data and critically reviewed the manuscript. PG developed the study design, wrote the research protocol and critically reviewed the manuscript. RYVdV collected data and critically reviewed the manuscript. TVN and TT assisted with data analysis and critically reviewed the manuscript. HMJJ, SK, JD, JE, JRC and DB critically reviewed the manuscript. JPvdB was responsible for the overall content as the guarantor. All authors approved the final version of the manuscript.

**Funding** This work was supported by the Weijerhorst Foundation, grant number not applicable. The Weijerhorst Foundation was not involved in conducting the study, analyzing the data or writing the manuscript.

**Competing interests** LV, CEW, RYVdV, HMJJ, SK, JD, JE, TT and DB have nothing to disclose. JRC reports honoraria for educational talks and part support for educational meetings from Amgen, and advisory board participation for Amgen and Bayer, outside the submitted work. TVN reports honoraria for lectures sponsored from Merck and participation as executive member Asia Pacific Consortium on Osteoporosis, outside the submitted work. PG reports grants from Amgen, Pfizer, MSD, UCB, Abbott, Lilly, BMS, Novartis, Roche and Will Pharma, and honoraria for lectures from Amgen and Lilly, outside the submitted work. JPvdB reports grants for lectures from UCB and Amgen, outside the submitted work.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Ethics approval** This study has been approved by the Independent Review Board Nijmegen and complied with the Declaration of Helsinki. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** No data are available.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Lisanne Vranken <http://orcid.org/0000-0002-6925-4397>

Caroline E Wyers <http://orcid.org/0000-0001-7662-3990>

#### REFERENCES

- Johnell O, Kanis JA, Odén A, *et al*. Fracture risk following an osteoporotic fracture. *Osteoporos Int* 2004;15:175–9.
- Kanis JA, Johnell O, De Laet C, *et al*. A meta-analysis of previous fracture and subsequent fracture risk. *Bone* 2004;35:375–82.

- 3 van Helden S, Cals J, Kessels F, *et al.* Risk of new clinical fractures within 2 years following a fracture. *Osteoporos Int* 2006;17:348–54.
- 4 Bliuc D, Nguyen TV, Eisman JA, Center JR., Risk of subsequent fracture after Low-Trauma fracture in men and women. *JAMA. American Medical Association* 2007;297:387–94.
- 5 Briggs AM, Sun W, Miller LJ, *et al.* Hospitalisations, admission costs and re-fracture risk related to osteoporosis in Western Australia are substantial: a 10-year review. *Aust N Z J Public Health* 2015;39:557–62.
- 6 Balasubramanian A, Zhang J, Chen L, *et al.* Risk of subsequent fracture after prior fracture among older women. *Osteoporos Int* 2019;30:79–92.
- 7 McKee KJ, Orbell S, Austin CA, *et al.* Fear of falling, falls efficacy, and health outcomes in older people following hip fracture. *Disabil Rehabil* 2002;24:327–33.
- 8 Shumway-Cook A, Ciol MA, Gruber W, *et al.* Incidence of and risk factors for falls following hip fracture in community-dwelling older adults. *Phys Ther* 2005;85:648–55.
- 9 Kristensen MT, Foss NB, Kehlet H. Timed "up & go" test as a predictor of falls within 6 months after hip fracture surgery. *Phys Ther* 2007;87:24–30.
- 10 Berggren M, Stenvall M, Olofsson B, *et al.* Evaluation of a fall-prevention program in older people after femoral neck fracture: a one-year follow-up. *Osteoporos Int* 2008;19:801–9.
- 11 Lloyd BD, Williamson DA, Singh NA, *et al.* Recurrent and injurious falls in the year following hip fracture: a prospective study of incidence and risk factors from the sarcopenia and hip fracture study. *J Gerontol A Biol Sci Med Sci* 2009;64:599–609.
- 12 Masud T, Morris RO. Epidemiology of falls. *Age Ageing* 2001;30 Suppl 4:3–7.
- 13 Morrison A, Fan T, Sen SS, *et al.* Epidemiology of falls and osteoporotic fractures: a systematic review. *Clinicoecon Outcomes Res* 2013;5:9–18.
- 14 Geusens P, Autier P, Boonen S, *et al.* The relationship among history of falls, osteoporosis, and fractures in postmenopausal women. *Arch Phys Med Rehabil* 2002;83:903–6.
- 15 Kaptoge S, Benevolenskaya LI, Bhalla AK, *et al.* Low BMD is less predictive than reported falls for future limb fractures in women across Europe: results from the European prospective osteoporosis study. *Bone* 2005;36:387–98.
- 16 Harvey NC, Odén A, Orwoll E, *et al.* Falls predict fractures independently of FRAX probability: a meta-analysis of the osteoporotic fractures in men (MROS) study. *J Bone Miner Res* 2018;33:510–6.
- 17 Leslie WD, Morin SN, Lix LM, *et al.* Fracture prediction from self-reported falls in routine clinical practice: a registry-based cohort study. *Osteoporos Int* 2019;30:2195–203.
- 18 Eisman JA, Bogoch ER, Dell R, *et al.* Making the first fracture the last fracture: ASBMR Task force report on secondary fracture prevention. *J Bone Miner Res* 2012;27:2039–46.
- 19 Lems WF, Dreinhöfer KE, Bischoff-Ferrari H. EULAR/EFORT recommendations for management of patients older than 50 years with a fragility fracture and prevention of subsequent fractures. *Ann Rheum Dis* 2016;annrheumdis-2016-210289–10.
- 20 Åkesson K, Marsh D, Mitchell PJ, *et al.* Capture the fracture: a best practice framework and global campaign to break the fragility fracture cycle. *Osteoporosis International* 2013;24:2135–52.
- 21 Dreinhöfer KE, Mitchell PJ, Bégué T, *et al.* A global call to action to improve the care of people with fragility fractures. *Injury* 2018;49:1393–7.
- 22 Javaid MK, Sami A, Lems W, *et al.* A patient-level key performance indicator set to measure the effectiveness of fracture liaison services and guide quality improvement: a position paper of the IOF capture the fracture Working group, National osteoporosis Foundation and fragility fracture network. *Osteoporos Int* 2020;4:e001806.
- 23 Werkgroep CBO. *Richtlijn Osteoporose en Fractuurpreventie, derde herziening [Dutch]*. Utrecht: CBO, 2011.
- 24 Richtlijn Preventie van valincidenten bij ouderen [Dutch], 2017. Available: [https://richtlijnen database.nl/richtlijn/preventie\\_van\\_valincidenten\\_bij\\_ouderen/startpagina\\_-\\_preventie\\_van\\_valincidenten.html](https://richtlijnen database.nl/richtlijn/preventie_van_valincidenten_bij_ouderen/startpagina_-_preventie_van_valincidenten.html)
- 25 Bours SPG, van Geel TACM, Geusens PPMM, *et al.* Contributors to secondary osteoporosis and metabolic bone diseases in patients presenting with a clinical fracture. *J Clin Endocrinol Metab* 2011;96:1360–7.
- 26 Genant HK, CY W, van Kuijk C. Vertebral fracture assessment using a semiquantitative technique. *Journal of bone and mineral research. John Wiley and Sons and The American Society for Bone and Mineral Research* 1993;8:1137–48.
- 27 The prevention of falls in later life. A report of the Kellogg international work group on the prevention of falls by the elderly. *Dan Med Bull* 1987;34:1–24.
- 28 van Helden S, Wyers CE, Dagnelie PC, *et al.* Risk of falling in patients with a recent fracture. *BMC Musculoskelet Disord* 2007;8:348.
- 29 Matsumoto H, Makabe T, Morita T, *et al.* Accelerometry-based gait analysis predicts falls among patients with a recent fracture who are ambulatory: a 1-year prospective study. *Int J Rehabil Res* 2015;38:131–6.
- 30 Yeh H-F, Shao J-H, Li C-L, *et al.* Predictors of postoperative falls in the first and second postoperative years among older hip fracture patients. *J Clin Nurs*. 2017;26:3710–23. 3rd ed..
- 31 Prudham D, Evans JG. Factors associated with falls in the elderly: a community study. *Age Ageing* 1981;10:141–6.
- 32 Campbell AJ, Reinken J, Allan BC, *et al.* Falls in old age: a study of frequency and related clinical factors. *Age Ageing* 1981;10:264–70.
- 33 Blake AJ, Morgan K, Bendall MJ, *et al.* Falls by elderly people at home: prevalence and associated factors. *Age Ageing* 1988;17:365–72.
- 34 World Health Organization WHO. *Who global report on falls prevention in older age*, 2017: 1–53.
- 35 Eekman DA, van Helden SH, Huisman AM, *et al.* Optimizing fracture prevention: the fracture liaison service, an observational study. *Osteoporos Int* 2014;25:701–9.
- 36 Lih A, Nandapalan H, Kim M, *et al.* Targeted intervention reduces refracture rates in patients with incident non-vertebral osteoporotic fractures: a 4-year prospective controlled study. *Osteoporos Int* 2011;22:849–58.
- 37 Van der Kallen J, Giles M, Cooper K, *et al.* A fracture prevention service reduces further fractures two years after incident minimal trauma fracture. *Int J Rheum Dis* 2014;17:195–203.
- 38 Huntjens KMB, van Geel TACM, van Helden S, *et al.* The role of the combination of bone and fall related risk factors on short-term subsequent fracture risk and mortality. *BMC Musculoskelet Disord* 2013;14:721.
- 39 Sanli I, van Helden SH, Ten Broeke RHM, *et al.* The role of the fracture liaison service (FLS) in subsequent fracture prevention in the extreme elderly. *Aging Clin Exp Res* 2019;31:1105–11.
- 40 Deloumeau A, Moltó A, Roux C, *et al.* Determinants of short term fracture risk in patients with a recent history of low-trauma non-vertebral fracture. *Bone* 2017;105:287–91.
- 41 Lötters FJB, van den Bergh JP, de Vries F. *Current and future incidence and costs of Osteoporosis-Related fractures in the Netherlands: combining claims data with BMD measurements*. Springer US: Calcified Tissue International, 2016: 235–43.
- 42 Wang Q, Jiang X, Shen Y, *et al.* Effectiveness of exercise intervention on fall-related fractures in older adults: a systematic review and meta-analysis of randomized controlled trials. *BMC Geriatr* 2020;20:322–11.
- 43 Pluijm SMF, Smit JH, Tromp EAM, *et al.* A risk profile for identifying community-dwelling elderly with a high risk of recurrent falling: results of a 3-year prospective study. *Osteoporos Int* 2006;17:417–25.
- 44 van Geel TACM, Huntjens KMB, van den Bergh JPW, *et al.* Timing of subsequent fractures after an initial fracture. *Curr Osteoporos Rep* 2010;8:118–22.
- 45 van Geel TACM, van Helden S, Geusens PP, *et al.* Clinical subsequent fractures cluster in time after first fractures. *Ann Rheum Dis* 2009;68:99–102.