

Characterization and diversity of seed endophytic bacteria of the endemic holoparasitic plant *Cistanche armena* (Orobanchaceae) from a semi-desert area in Armenia

Peer-reviewed author version

PETROSYAN, Kristine; THIJS, Sofie; Piwowarczyk, Renata; Ruraz, Karolina; VANGRONSVELD, Jaco & Kaca, Wiesław (2022) Characterization and diversity of seed endophytic bacteria of the endemic holoparasitic plant *Cistanche armena* (Orobanchaceae) from a semi-desert area in Armenia. In: SEED SCIENCE RESEARCH, 32 (4), p. 264-273.

DOI: 10.1017/S0960258522000204

Handle: <http://hdl.handle.net/1942/38848>

1                    **Bacterial seed endophytes of the holoparasitic endemic**  
2                    ***Cistanche armena* (Orobanchaceae) from a semi-desert area in Armenia**

3  
4                    **Kristine Petrosyan<sup>1,3\*</sup>, Sofie Thijs<sup>3</sup>, Renata Piwowarczyk<sup>2</sup>, Karolina Ruraż<sup>2</sup>, Jaco**  
5                    **Vangronsveld<sup>3,4</sup> and Wiesław Kaca<sup>1</sup>**

6                    <sup>1</sup>Department of Microbiology, Institute of Biology, Jan Kochanowski University,  
7                    Uniwersytecka 7, 25-406 Kielce, Poland

8                    <sup>2</sup>Center for Research and Conservation of Biodiversity, Department of Environmental  
9                    Biology, Institute of Biology, Jan Kochanowski University, Uniwersytecka 7, 25-406 Kielce,  
10                    Poland

11                    <sup>3</sup>Centre for Environmental Sciences, Environmental Biology Research Group, Hasselt  
12                    University, Agoralaan building D, 3590 Diepenbeek, Belgium

13                    <sup>4</sup>Institute of Biological Sciences, Department of Plant Physiology and Biophysics, Faculty of  
14                    Biology and Biotechnology, Maria Curie-Skłodowska University, 19 Akademicka, 20–033  
15                    Lublin, Poland

16                    **\* Correspondence:**

17                    Corresponding Author

18                    Department of Microbiology, Institute of Biology, Jan Kochanowski University,  
19                    Uniwersytecka 7, 25-406 Kielce, Poland,

20                    Tel: (48) 735046235

21                    e-mail: kristine.petrosyan@phd.ujk.edu.pl

22                    kristine.petrosyan@uhasselt.be

23  
24                    **Keywords:** abiotic stress, *Bacillus*, dust-seeds, *Pantoea*, parasitic plants, PGP traits, seeds  
25                    microbiome

26  
27                    **Declaration of all sources of financial support**

28                    The author acknowledges financial support through the project “Development Accelerator of  
29                    the Jan Kochanowski University of Kielce,” co-financed by the European Union under the  
30                    European Social Fund, (K.P., POWR.03.05.00-00-Z212/18, 2019-2023). This study was  
31                    supported by grants from the Jan Kochanowski University (K.R., 666 065, 2019), (W.K.;  
32                    K.P., SUPB.RN. 21.235, 2021-2022). The field research in this study in Armenia was  
33                    partially financed by the National Geographic grant (R.P., GEFNE 192-16, 2017). This study  
34                    was also supported by a BOF-BILA grant from Hasselt University Belgium BOF21BL12  
35                    (K.P.; J.V., 2021-2022) and the Hasselt University Methusalem project (J.V., 08M03VGRJ).

## 45 Abstract

46 We explored the seed-associated bacterial endophytic microbiome in seeds of the endemic  
47 holoparasitic species *Cistanche armena* from a saline and arid habitat in Armenia. A  
48 combination of culture-dependent and molecular techniques was employed for identifying the  
49 seed endomicrobiome (culturable and unculturable). From surface sterilized seeds, 10 phyla,  
50 256 endophytic bacterial genera were identified. Of the culturable strains, we also  
51 investigated the plant growth-promoting (PGP) traits. Most of the isolates were spore  
52 forming, halotolerant, and alkaliphile *Bacillus* spp., indicating that the endophytic bacteria of  
53 *C. armena* seeds own traits related to the natural habitat of their host plant. Our results  
54 confirm that *Bacillus* species are common and dominated endophytes from plants growing on  
55 saline and arid soils. *Pantoea* spp. and *Stenotrophomonas* are more favourable PGP  
56 endophytes in seeds of *C. armena*. The PGP traits of these bacteria, such as production of  
57 auxins, ACC-deaminase and organic acids have the potential to improve the tolerance of their  
58 host plants against the abiotic stresses present in their natural habitat. To the best of our  
59 knowledge, that is the first report concerning bacterial seed endophytes of the *C. armena*.

60

## 61 Introduction

62 With approximately 4,750 species, parasitic plants constitute 1.6% of the angiosperms  
63 (Nickrent, 2020). Parasitism, especially holoparasitism, represents the most extreme  
64 interaction between plants, with strong associations between host and parasite biogeography,  
65 ecology, and probably with diversification (Schneider and Moore, 2017). Orobanchaceae is  
66 the largest parasitic plant family with 102 genera and over 2100 species (Nickrent, 2020). One  
67 of the most peculiar in this family is the genus *Cistanche* Hoffmanns. & Link, which includes  
68 approximately 25 species, and is found mainly in arid, semiarid and halophytic habitats across  
69 Eurasia and North Africa. These magnificent, achlorophyllous species, with fleshy stems,  
70 long underground stolons and intensely colored inflorescences grow as obligate parasite  
71 (holoparasite) on the roots of host-plant species mainly belonging to the Chenopodiaceae,  
72 Zygophyllaceae, Tamaricaceae, and Plumbaginaceae (Piwowarczyk et al., 2019). Species  
73 belonging to this genus have been widely used in traditional Chinese medicine for centuries  
74 (Li et al., 2016; Piwowarczyk et al., 2020a).

75 A particularity of parasitic plants is their production of huge numbers of seeds, which are  
76 also among the smallest of all seed plants. With a length of less than 1 mm they are often  
77 called ‘dust seeds’ (Yoneyama et al., 2008; Eriksson and Kainulainen, 2011; Piwowarczyk,  
78 2013). The seeds possess a unique simple structure, contain only a reduced embryo, as a  
79 spherical body without a plumule, and radicle or cotyledons. The reticulated testa of these  
80 seeds with polygonal and sometimes deeply submerged walls might enhance the contact of  
81 the seed surface with water or facilitate the seed dispersal by wind. The endothelium (inner  
82 testa layer) containing mucilage and labyrinthine walls, allows rapid absorption of water,  
83 which is crucial for imbibition and subsequent germination (Piwowarczyk et al., 2020b). The  
84 cutinized endothelium has a protective role in the underground part of the plant life cycle  
85 (Dinesh et al., 2015; Piwowarczyk et al., 2019). Lipids are the main storage material in the  
86 seeds of Orobanchaceae (Ruraż et al., 2020). For germination, *Cistanche* seeds need to be  
87 very nearby their preferred host. Germination depends on hormones-strigolactones exuded  
88 from the host root (Yoneyama et al., 2008). Seeds of *Cistanche*, like related *Orobanche* s.l.  
89 species, seem to be resistant to harsh environmental conditions and stay viable in the soil for  
90 several decades (Joel et al., 2007). Among the wide range of plant protection mechanisms, the  
91 endophytic microbes have a specific role for improving the plant tolerance against different  
92 biotic and abiotic stresses (Shrivastava and Kumar, 2015).

93 Recently, the interest in plant endophytes from ecosystems with harsh environmental  
94 conditions, especially saline soils has increased (Hryniewicz et al., 2019; Manasa et al.,  
95 2020). Such endophytes can have the potential to mitigate the impacts of adverse conditions  
96 such as soil salinization, high concentrations of metals and climate change (Hallmann et al.,  
97 1997; Truyens et al., 2016; Manjunatha et al., 2017; Hemida and Reyad, 2019). Most of the  
98 seed associated bacteria are considered to have an environmental origin and to be important  
99 for the adaptation of their host to harsh environmental conditions (Frank et al., 2017).  
100 Therefore, tissues of halotolerant plants also contain halophilic bacterial communities  
101 (Etesami and Beattie, 2018) and the composition of seed-associated bacterial communities  
102 should be closely related to the soil bacterial communities. Besides of the obligate  
103 endophytes, plant tissues can be colonized by soil bacteria as well. This is explained by  
104 possible migration of bacteria from the soil to the seeds (Frank et al., 2017). According to  
105 Barret et al. (2016), the endophytes reach the seeds by: internal transmission through the  
106 vascular system and floral transmission (external transmission) through the stigma, fruits, or  
107 flowers. Indeed, during the early stages of seed development, the endophytes reach the seeds  
108 via the xylem and nonvascular plant tissues. Bacteria can also use the floral pathway to reach  
109 the seeds. Though, the floral route has a selective function, and only endophytes with  
110 biocontrol ability and nonhost pathogens can reach the seeds.

111 So far, ample of endophytes have been isolated from different seeds of many wild and  
112 agricultural/sylvicultural herbaceous and woody plant species (e.g., Ulrich et al., 2008;  
113 Truyens et al., 2013, 2014, 2016; Asaf et al., 2017; Glassner et al., 2018; Sánchez-López et  
114 al., 2018; Compant et al., 2019), including some holoparasitic species (tissue and seeds) such  
115 as *Phelipanche aegyptiaca*, *P. ramosa*, and *Orobancha hederæ* (Iasur Kruh et al., 2017;  
116 Fitzpatrick and Schneider, 2020; Huet et al., 2020; Durlík et al., 2021). The microbiome of *P.*  
117 *aegyptiaca* in different developmental stages was investigated by Iasur Kruh et al. (2017).  
118 Surface-sterilized tissues of roots, haustoria and shoots harbored bacteria belonging to the  
119 Proteobacteria (*Rhizobium*, *Pseudomonas*, *Comamonadaceae* sp., *Sphingomonas* and  
120 *Burkholderia*, *Actinobacter* sp., *Bacillus* sp.). In addition, *Novosphingobium* and  
121 *Methylophilus* were reported as specific endophytes for this plant species (Iasur Kruh et al.,  
122 2017). A study of the endophytic microbiome of *O. hederæ* reported that *Orobancha* leaves  
123 (scales) contain Acidobacteria, Proteobacteria, Verrucomicrobia and bacteria belonging to the  
124 *Enterobacteriaceae*, *Pseudomonadaceae*, and *Rhizobiaceae* (Fitzpatrick and Schneider,  
125 2020). The first report about seed endophytes of the holoparasitic *P. ramosa* reported a  
126 dominance of four bacterial phyla, i.e., Proteobacteria, Bacteroidetes, Actinobacteria,  
127 Firmicutes (Huet et al., 2020). In another study on surface sterilized seeds of *P. ramosa*,  
128 culturable *Brevibacterium frigoritolerans* and *Bacillus simplex* were isolated (Durlík et al.,  
129 2021) (Table 1). Different bacterial phyla also have been isolated from plants growing in arid  
130 and semiarid regions, like *Larrea tridentata*, from the desert plant *Salsola* (Soussi et al., 2016)  
131 and the saline wetland species *Salicornia* (Szymańska et al., 2018). Furthermore, some  
132 authors argue that the bacterial phyla Proteobacteria, Bacteroidetes, Firmicutes,  
133 Planctomycetes, Actinobacteria, Fibrobacteres are common for halotolerant plants from arid  
134 and wetland soils (Soussi et al., 2016; Asaf et al., 2017; Szymańska et al., 2018).

135 Although many investigations highlight the importance of endophytes in plant health, the  
136 knowledge concerning communities of bacterial seed endophytes, especially about the  
137 microbiome of seeds of holoparasitic plant species is still limited (Iasur Kruh et al., 2017;  
138 Fitzpatrick and Schneider, 2020; Huet et al., 2020; Durlík et al., 2021). Therefore, the major  
139 objective of our study was to explore the bacterial endophytes (culturable and unculturable)  
140 from seeds of the holoparasitic endemic plant *Cistanche armena* (K. Koch) M.V. Agab.  
141 (Orobanchaceae) from a saline and semi-desert habitat of Armenia. The other aim was to

142 investigate the potential plant growth-promoting (PGP) traits of the culturable seed  
143 endophytes that might have a role in plant responses and tolerance to abiotic stresses.

144 The present study combined culture-dependent and molecular approaches. Moreover, the  
145 effectivity of the sterilization method is a crucial step to isolate just the seed endophytes. For  
146 this purpose, the micromorphology of the seeds was studied to help us to select the  
147 appropriate method of surface sterilization, due to the unique structure of the reticulated testa  
148 and the endothelium of the seed coat. Molecular techniques were used to identify the  
149 culturable bacteria and to describe the diversity of the microbial communities in seeds of the  
150 examined plant species. PGP traits such as the ability to produce Indole-3-acetic acid (IAA),  
151 ACC-deaminase, siderophores and organic acids of the culturable endophytic bacterial strains  
152 were also investigated.

153 To the best of our knowledge this is the first report about bacterial seed endophytes of the  
154 holoparasitic endemic plant species *C. armena*.

155

156

### 157 **Materials and methods**

#### 158 ***Species natural habitat and plant material***

159 Mature seeds of *Cistanche armena* (Orobanchaceae) were used. *C. armena* (K. Koch) M.V.  
160 Agab. is an endemic, critically endangered species. It is known only from the Ararat and  
161 Armavir provinces in central Armenia, in the Arax River valley and at the foot of Mount  
162 Ararat, NW of the village Lusarat, near the Khor Virap monastery (39°53'01' N, 44°34'49' E)  
163 at about 820–840 m above sea level (Piwowarczyk et al., 2017, 2019). This locality is one of  
164 the hottest and extremely arid regions of Armenia. The mean daily air temperature ranges  
165 from a maximum of 42°C in July to a minimum of -33°C in January. The average annual  
166 rainfall is 300 mm, while the annual evaporation reaches up to 1,000 mm. The area is  
167 characterized by strong salinity (total salt content of the soil 1–3%) with considerable  
168 carbonization (Panosyan et al., 2018). It is a semi-desert, with sandy, saline soils and a  
169 halophytic vegetation. *C. armena* parasitizes *Alhagi maurorum* (Fabaceae) and *Salsola*  
170 *dendroides* (Chenopodiaceae) (Fig. 1A, B).

171 The mature seeds were collected in June 2017. Seeds from at least 10 plant individuals of  
172 the total population from the region were collected. Mature and dry seeds were collected from  
173 dry fruits and used for further experiments. The seeds were collected and identified by Renata  
174 Piwowarczyk, and herbarium materials were deposited in the Herbarium of the Jan  
175 Kochanowski University in Kielce (KTC), Poland. The seeds were dried under natural  
176 conditions. Field studies, including the collection of plant and seed material complied with  
177 relevant local, institutional, national, and international guidelines, permissions, and  
178 legislation.

179

#### 180 ***Microscopic observation and morphometric analysis of seeds***

181 General seed morphology was studied using an Axio Zoom.V16 Stereo Zoom system (Carl  
182 Zeiss, Germany) in bright-field illumination (objective lenses PlanApo Z 1.5×, FWD = 30  
183 mm) and processed in ImageJ software using Fiji macros. The terminology of seed surfaces  
184 was taken from Barthlott (1981), and Piwowarczyk et al. (2020b). At least 30 seeds were  
185 examined, and quantitative and qualitative morphological characteristics were determined  
186 several times for each seed (Fig. 2).

#### 187 ***Method for seed surface sterilization and cultivation conditions of culturable seed*** 188 ***endophytic bacteria***

189 The aim of seed surface sterilization was to obtain only the endophytic bacterial communities  
190 of the seeds. For this purpose, 50 mg seeds were transferred into 1.5-mL Eppendorf tubes,  
191 submersed in 70% ethanol for 60 s, then 1 mL of 0.85% sterile NaCl solution was added,  
192 followed by shaking on a vortex (8,000 rpm) at 21°C for 2.5 h. Subsequently, the washed  
193 seeds were kept at 4°C for 15 min. Before rinsing with sterile double distilled water, the seeds  
194 were centrifuged for 30 s at 12,000 rpm (13,400 × g). The washing process was repeated five  
195 times with a decreasing time of shaking from 2 h to 30 min (2 h, 1.5 h, 60 min, 45 min, and 30  
196 min). Each time samples were centrifuged for 30 s, rinsed with sterile double distilled water,  
197 and kept at 4°C for 15 min. The rinsing procedure was repeated three times. For proving the  
198 effectiveness of the sterilization procedure, the last rinsing water was plated on previously  
199 prepared Petri dishes with LB medium. The surface sterilized seeds were mechanically  
200 homogenized using a sterile pellet pestle (Kimble®) in 0.5ml 10mM MgSO<sub>4</sub>. Part of the  
201 homogenous seed suspension was used for DNA extraction, another part for isolation of  
202 culturable bacteria.

### 203 *Total DNA extraction from seeds, library preparation, and Illumina sequencing*

204 For identification of the total (cultivable and uncultivable) bacterial community the  
205 homogenized suspension of the surface sterilized seeds was used. The DNA isolation was  
206 performed using the Mobio Power Plant protocol. The isolation of total bacterial DNA was  
207 conducted in 4 replicates.

208 All DNA samples were subjected to bacterial 16S rRNA gene amplicon PCR. In the first  
209 round of 16S rRNA gene PCR, an amplicon of 291 bp was generated, using primers 515F-  
210 GTGYCAGCMGCCGCGGTAA and 806R- GGACTACNVGGGTWTCTAAT (Walters et  
211 al., 2016), with an Illumina adapter overhang nucleotide sequence, resulting in the following  
212 sequences, 515F-adaptor: 5'- TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG-  
213 3' and 806R-adaptor: 5'- GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA G-3'.  
214 For the first round of PCR the Q5 High-Fidelity DNA Polymerase system (M0491, NEB), a  
215 reaction volume of 25 µl per sample was prepared containing 1 µl of extracted DNA (final  
216 DNA-concentration per reaction 1-10 ng), 1x Q5 Reaction Buffer with 2 mM MgCl<sub>2</sub>, 200 µM  
217 dNTP mix, 1x Q5 High GC Enhancer (for the seed and bacterial samples), 0.25 µM forward  
218 or reverse primer, and 0.02 U µl<sup>-1</sup> Q5 High-Fidelity DNA polymerase, and for the seed  
219 endophytic extracts, additionally 0,5 µL mitoPNA blocker (2 µM final concentration added  
220 from a 50 µM stock), 0,5 µL (seeds) plastidPNA blocker (2 µM final concentration from 50  
221 µM stock) (Kusstatscher et al., 2021) were using. The PCR program started with an initial  
222 denaturation for 3 min at 98 °C, followed by a 10 sec denaturation at 98°C, a 30 sec annealing  
223 at 56°C for V3V4 (58°C for ITS) and a 30 sec extension at 72 °C, all three steps were repeated  
224 for a total of 30 cycles. The reaction was ended by a final 7 min extension at 72 °C. The  
225 amplified DNA was purified using the AMPure XP beads (Beckman Coulter) and the  
226 MagMax magnetic particle processor (ThermoFisher, Leuven, Belgium). Subsequently, 5 µl  
227 of the cleaned PCR product was used for the second PCR attaching the Nextera indices  
228 (Nextera XT Index Kit v2 Set A(FC-131-2001), and D (FC-131-2004), Illumina, Belgium).  
229 For these PCR reactions, 5 µl of the purified PCR product was used in a 25 µl reaction  
230 volume and prepared following the 16S Metagenomic Sequencing Library Preparation Guide.  
231 PCR conditions were the same as described above, but the number of cycles reduced to 20,  
232 and 55°C annealing temperature. PCR products were cleaned with the Agencourt AMPure XP  
233 kit, and then quantified using the Qubit dsDNA HS assay kit (Invitrogen) and the Qubit 2.0  
234 Fluorometer (Invitrogen). Once the molarity of the sample was determined, the samples were  
235 diluted down to 4 nM using 10 mM Tris pH 8.5 prior to sequencing on the Illumina MiSeq.  
236 Samples were sequenced using the MiSeq Reagent Kit v3 (600 cycle) (MS-102-3003) and  
237 15% PhiX Control v3 (FC-110-3001). For quality control, a DNA-extraction blank and PCR

238 blank were included throughout the process, and also the ZymoBIOMICS Microbial Mock  
239 Community Standard (D6300) to test efficiency of DNA extraction (Zymo Research).

240

### 241 *Bioinformatic processing of reads*

242 Sequences were demultiplexed using the Illumina Miseq software, and subsequently quality  
243 trimmed and primers removed using DADA2 1.10.1 (Callahan et al., 2016) in R version 3.5.1.  
244 Parameters for length trimming were set to keep the first 290 bases of the forward read and  
245 200 bases of the reverse read, maxN=0, MaxEE=(2,5) and PhiX removal. Error rates were  
246 inferred, and the filtered reads were dereplicated and denoised using the DADA2 default  
247 parameters. After merging paired reads and removal of chimeras via the  
248 removeBimeraDenovo function, an amplicon sequence variant (ASV) table was built and  
249 taxonomy assigned using the SILVA v138 training set (Quast et al., 2013; Yilmaz et al.,  
250 2014). The resulting ASVs and taxonomy tables were combined with the metadata file into a  
251 phyloseq object (Phyloseq, version 1.26.1) (McMurdie and Holmes, 2013). Contaminants  
252 were removed from the dataset using the package Decontam (version 1.2.1) applying the  
253 prevalence method with a 0.5 threshold value (Davis et al., 2018). A phylogenetic tree was  
254 constructed using a DECIPHER/Phangorn pipeline as described before (Murali et al., 2018).

### 255 *Data visualization and statistical analyses*

256 The ASV table was further processed removing organelles (chloroplast, mitochondria), and  
257 prevalence filtered using a 2% inclusion threshold (unsupervised filtering) as described by  
258 Callahan et al. (2016). Alpha-diversity metrics such as Chao1, Simpson's and Shannon's  
259 diversity indexes were calculated on unfiltered data using scripts from the MicrobiomeSeq  
260 package. Hypothesis testing was done using analysis of variance (ANOVA) and the Tukey  
261 Honest Significant Differences method (Tukey HSD). When assumptions of normality and  
262 homoscedasticity were not met, a Kruskal-Wallis Rank Sum test and a Wilcoxon Rank Sum  
263 test was performed. The results were summarized in boxplots. Relative abundances were  
264 calculated and visualized in bar charts using Phyloseq. All performed statistical tests were  
265 corrected for multiple testing and  $\alpha < 0.05$  was considered as statistically significant. All  
266 graphs were generated in R version 4.0.4.

### 267 *Isolation of culturable endophytes*

268 The first part of the suspension obtained after crushing the seeds (see above) was used for  
269 DNA extraction, the second part for isolation of culturable bacteria. Serial dilutions were  
270 made  $10^6$  cfu ml<sup>-1</sup> and then 100  $\mu$ l was plated onto 1/869 rich medium with composition:  
271 0.035 g L<sup>-1</sup> CaCl<sub>2</sub> x 2H<sub>2</sub>O, Glucose D 0.1 g L<sup>-1</sup>, NaCl 0.5 g L<sup>-1</sup>, Trypton 1 g L<sup>-1</sup>, Yeast Extract  
272 0.5 g L<sup>-1</sup>, Agar 15 g L<sup>-1</sup> (Eevers et al., 2015) and incubated at 30°C for 7 days. For further  
273 experiments, single, morphological diverse colonies were picked and purified. Subsequently,  
274 they were grown in 96-well master blocks and triplicated: one block was used for DNA-  
275 extraction, the second one was used for PGP tests and the third was stored at -45°C in 15%  
276 glycerol (75 g glycerol, 4.25 g NaCl, 425 ml dH<sub>2</sub>O).

### 277 *Genomic DNA extraction and taxonomic identification of the culturable endophytic 278 bacterial strains*

279 The DNA isolation was performed using standard procedure for DNA isolation from bacterial  
280 pellets with MagMAX. DNA was quantified with a Qubit® 2.0 Fluorometer  
281 (ThermoScientific, US) and checked for purity on a Nanodrop spectrophotometer  
282 (ThermoScientific, US) with an A260/A280 ratio of 1.7–2.0. The near full-length sequences

283 of the 16S rRNA gene were amplified with the primers 27f (5-  
284 AGAGTTTGATCMTGGCTCAG-3) and 1492r (5-GGTTACCTTGTTACGACTT-3). The  
285 products were checked on agarose gel and then shipped to Macrogen for 16S rRNA Sanger  
286 sequencing. Sequencing results were quality filtered using Geneious v4.8, were analyzed over  
287 the ribosomal database SILVA (<https://www.arb-silva.de/aligner/>) and NCBI GenBank  
288 databases using the program Standard Nucleotide BLAST and database RDP  
289 ([https://rdp.cme.msu.edu/seqmatch/seqmatch\\_intro.jsp](https://rdp.cme.msu.edu/seqmatch/seqmatch_intro.jsp)).

### 290 *Plant growth promoting (PGP) characteristics*

291 In order to evaluate the ability of the isolated strains to induce plant growth promotion, *in*  
292 *vitro* PGP tests were performed. All tests were performed at least two times.

293 The IAA production ability was tested using the Salkowski test. Bacteria were grown in a  
294 1/10 869 medium containing tryptophan (Patten and Glick, 2002). 25  $\mu$ l of bacterial  
295 suspension with 0.7 ml IAA medium were incubated for 4 days at 30°C and shaken at 150  
296 rpm in the dark. Thereafter, the suspension was centrifuged for 10 min at 4000 rpm. 1 ml  
297 Salkowski reagent was added to 0.5 ml supernatant. After 20 min reaction time colored pink  
298 means positive for IAA production.

299 To check for organic acid production the method of Cunningham & Kuiack was used. The  
300 bacteria were cultivated in a Sucrose Tryptone (ST) medium with composition: sucrose 20 g  
301 L<sup>-1</sup>, tryptone 5 g L<sup>-1</sup>, 10 ml trace element solution SET (Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O 20 mg L<sup>-1</sup>, H<sub>3</sub>BO<sub>3</sub>  
302 200 mg L<sup>-1</sup>, CuSO<sub>4</sub>·5H<sub>2</sub>O 20 mg L<sup>-1</sup>, FeCl<sub>3</sub> 100 mg L<sup>-1</sup>, MnCl<sub>2</sub>·4H<sub>2</sub>O 20 mg L<sup>-1</sup>, ZnCl<sub>2</sub> 280  
303 mg L<sup>-1</sup>). The bacterial suspension was incubated for 5 days at 30°C and 200 rpm, after which  
304 the pH-sensitive color indicator 100  $\mu$ L Alizarine Red S 0,1% was added (Cunningham and  
305 Kuiack, 1992). The organic acid production was checked after 15 min reaction time: yellow =  
306 positive, pink = negative.

307 ACC-deaminase activity was tested in SMN medium with 5 mM ACC as N-source with  
308 HCl and autoclaved (Belimov et al., 2005). SMN medium composition: 970mL: 0,4g L<sup>-1</sup>  
309 KH<sub>2</sub>PO<sub>4</sub>, 2 g L<sup>-1</sup> K<sub>2</sub>HPO<sub>4</sub> (pH 6,6), 10 mL MgSO<sub>4</sub> solution, 10 mL CaCl<sub>2</sub> solution and 10mL  
310 micronutrient stock were added after filter sterilization. 50 mL C-mix stock with 2 g L<sup>-1</sup>  
311 glucose, 2 g L<sup>-1</sup> sucrose, 2 g L<sup>-1</sup> Na-acetate, 2 g L<sup>-1</sup> Na-citrate, 2 g L<sup>-1</sup> Malic acid and 2 g L<sup>-1</sup>  
312 Mannitol and 10 mL ACC-stock were added. 250  $\mu$ L of the bacterial suspension added to 1.2  
313 mL SMN medium with 5 mM ACC as N-source were incubated for 3 days at 30°C and  
314 centrifuged at 4000 rpm for 15 min. The pellet was resuspended in 100  $\mu$ L 0,1M Tris-HCl  
315 buffer (pH 8,5) and 3  $\mu$ L toluene was added for cell lysis, and vortexed for 5 min. In next step  
316 10  $\mu$ L 0,5 M ACC and 100  $\mu$ L 0,1M Tris-HCl buffer (pH 8,5), vortexed and incubated for 30  
317 min at 30°C and 150 rpm. 690  $\mu$ L 0,56N HCl and 150  $\mu$ L 0,2% 2,4-dinitrophenylhydrazine in  
318 2N HCl and 1 mL 2N NaOH were added. The obtained results were evaluated: brown =  
319 positive, yellow = negative.

320 Production of siderophores was studied by using the 284 medium with 0.25  $\mu$ l optimal iron  
321 concentration with CAS solution (Schwyn and Neilands, 1987). Tris 6.06 g L<sup>-1</sup>, NaCl 4.68 g  
322 L<sup>-1</sup>, KCl 1.49 g L<sup>-1</sup>, NH<sub>4</sub>Cl 1.07 g L<sup>-1</sup>, Na<sub>2</sub>SO<sub>4</sub> 0.43g L<sup>-1</sup>, MgCl<sub>2</sub>·6H<sub>2</sub>O 0.2 g L<sup>-1</sup>, CaCl<sub>2</sub>·2H<sub>2</sub>O  
323 0.03 g L<sup>-1</sup>, Na<sub>2</sub>HPO<sub>4</sub>·2H<sub>2</sub>O 0.04 g L<sup>-1</sup>, S17 trace elements 1 ml, 0.25 mM Fe(III)Citrate  
324 solution, Sodium lactate (sol. 50%) 0.7 ml, D-(+)-glucose 0.52 g L<sup>-1</sup>, D-gluconic acid sodium  
325 salt 0.66 g L<sup>-1</sup>, D-(+) fructose 0.54 g L<sup>-1</sup>, Sodium succinate·6H<sub>2</sub>O 0.81 g L<sup>-1</sup>. The 284 medium  
326 with 0  $\mu$ l and 3  $\mu$ l were used as control. 800  $\mu$ L 284 medium (0  $\mu$ M, 0,25  $\mu$ M and 3  $\mu$ M Fe)  
327 with 20  $\mu$ L of the bacterial suspension were incubated for 5 days at 30°C and 200 rpm. 100  
328  $\mu$ L Chrom-Azuroil S Solution (CAS-Solution) were added. After 4 h reaction time, orange =  
329 positive, blue = negative.

330



331 **Results**332 ***Seed micromorphology***

333 *C. armena* seeds are dark brown, 541–1003  $\mu\text{m}$  long, 347–631  $\mu\text{m}$  wide with a 1.1–2.3  
 334 length-to-width ratio and 164333–445987  $\mu\text{m}^2$  area. The shape was oblongoid to ovoid, rarely  
 335 subrectangular. The seed ornamentation was constantly alveolate. The testa of the seeds had  
 336 smooth, thin outer periclinal walls adjacent to the inner periclinal wall with perforated (pitted)  
 337 sculpture. The seed coat surface was formed by polygonal and isodiametric cells with  
 338 different sizes, 41–159  $\mu\text{m}$  long and 33–96  $\mu\text{m}$  wide with a 1.0–3.1 length-to-width ratio. The  
 339 number of cells along the seed longitudinal axis was 7–13; in the lateral view; it varied from  
 340 34 to 79. The anticlinal walls were of slight depth with a width of 7.7–14.6  $\mu\text{m}$  (Fig. 2).

341 ***Seed endophytic bacterial community composition***

342 The number of paired raw Illumina reads after filtering low quality reads, adapters, barcodes  
 343 and primers, there were about 2300 effective read for the 4 replicates of *C. armena* seeds. The  
 344 Shannon-Wiener biodiversity index, Chao1 and Simpson indexes for the seed endophytes of  
 345 *C. armena* were 2.82, 27, 13.9 respectively (Supplementary Figure S1) with P-value 0.05. A  
 346 total of 75 different Operational Taxonomic Unit (OTU)s on genus level was found from 10  
 347 phyla. The relative abundance of the dominant bacteria comprising the seed endophytic  
 348 community at different taxonomic levels is presented in Supplementary Figure S2.

349 From the surface sterilized seeds, 10 phyla and 256 bacterial genera were identified. The  
 350 taxonomy of the sequences was described primarily at the phylum level. For the *C. armena*  
 351 seeds, we determined Proteobacteria, Firmicutes and Actinobacteriota, whereas the  
 352 Bacteroidetes, Acidobacteria, Verrucomicrobia, Mixococcota, Planctomycetes,  
 353 Patescibacteria and Chloroflexi were less abundant (Supplementary Figure S2). Firmicutes  
 354 were the predominating phylum in the seeds of the examined plant population, followed by  
 355 Proteobacteria and Actinobacteriota. The phylum Actinobacteriota was classified only in 3  
 356 biological replicates. Only Bacilli, Gammaproteobacteria and Actinobacteria dominated at the  
 357 class level (Table 2). Indeed, Bacilli were the most abundant class (Supplementary Figure  
 358 S2). The majority of endophytic bacterial community of seeds of *C. armena* belonged to the  
 359 order Bacillales that at genus level was represented by *Psychrobacillus*, *Bacillus* and  
 360 *Domibacillus*. The most abundant family of Firmicutes identified in examined seeds was  
 361 Planococcaceae with *Paenisporosarcina* as a predominant genus.

362 The Gammaproteobacteria were identified as another abundant class, that at the order level  
 363 was represented by Xanthomonadales, Pseudomonadales and Enterobacterales. At genus level  
 364 *Pseudomonas*, *Stenotrophomonas* and *Serratia* dominated (Table 2). Finally, *Microbacterium*  
 365 and *Curtobacterium* were the dominating genera of the phylum Actinobacteriota. Unclassified  
 366 groups were found also at different taxonomic levels. The results are presented based on the  
 367 most representative and dominating OTUs (identified at genus level with a relative abundance  
 368 higher than 1%).

369  
 370 ***Diversity of cultivable endophytes from surface-sterile seeds and in vitro characterization of***  
 371 ***PGP bacteria***

372 43 bacterial strains were picked up from the 1/869 medium. Using 16S rRNA gene Sanger  
 373 sequencing we found that 35 bacteria (81.4%) of the total isolates were Firmicutes and only  
 374 18.6% were Proteobacteria with *Stenotrophomonas maltophilia* and different strains of  
 375 *Pantoea*. The majority of Firmicutes isolates belonged to the genera *Bacillus* and  
 376 *Paenibacillus*.

377 36 strains scored positive for IAA production and only 3 strains of *Bacillus* spp. tested  
378 positive for siderophore production. Relatively similar outcomes were obtained for production  
379 of ACC-deaminase and organic acids: 26 and 27 strains respectively showed positive (Fig. 3).  
380 In the *in vitro* tests *Pantoea* spp. and *Stenotrophomonas maltophilia* demonstrated higher  
381 growth promoting capacities compared to *Bacillus* spp. and other isolates (Fig. 3).

### 382 Discussion

383 The seed surfaces of holoparasitic *C. armena* possess an alveolate ornamentation with  
384 perforated (pitted) sculpture formed by polygonal and isodiametric cells with different sizes.  
385 The quite coarse structure of the seed coat (Fig. 2) can complicate the surface sterilization of  
386 the seeds. The preliminary results obtained by applying the generally used sterilization  
387 protocols (Watts et al., 1993; Metwaly et al., 2018) showed to be inadequate. We assumed  
388 that the sterilizing agents could not always sufficiently reach the deepest zones of the coarse  
389 seed surface. Due to this, not all bacteria residing on the surface of the seeds could be  
390 eliminated. Finally, the combination of 70% ethanol and 0.85% NaCl sterile solution together  
391 with intense shaking showed to be adequate to remove all bacteria from the surfaces of *C.*  
392 *armena* seeds. This allowed us to isolate only the bacteria present inside the seeds. It is known  
393 that the majority of plant associated bacteria are unculturable, and it is often assumed that  
394 only 0.001-1% can be grown in laboratory conditions (Eevers et al., 2015). Consequently, in  
395 order to obtain more information about the composition of the total endophytic bacterial  
396 communities of the seeds (culturable and unculturable) of *C. armena*, molecular techniques  
397 were used. The Illumina MiSeq data showed that the seeds of *C. armena* were mainly  
398 inhabited by Gram-positive, spore forming *Bacilli* (36.8%) (Supplementary Figure S2). In  
399 case of a holoparasitic plant, like *C. armena*, this is very plausible because these seeds,  
400 similarly to *Orobanch* s.l., have to stay viable in the soil for several decades (Joel et al.,  
401 2007). Plant colonization by spore forming *Bacillus* spp. that possess potential to mitigate  
402 environmental stress can help plants to survive in harsh environmental conditions. *C. armena*  
403 adapted to the arid and saline environment of specific areas in Armenia (Piwowarczyk et al.,  
404 2017, 2019). We demonstrated that *C. armena* was colonized by halotolerant, alkalophilic,  
405 spore forming, motile *Bacillus* spp. strains (Petrosyan et al., 2022). Some isolated strains  
406 were also thermophilic. They are able to produce one or more hydrolytic enzymes, especially  
407 cellulase and protease. Some strains also produced amylase and pectinase too. Production of  
408 auxins (IAA) and gibberellins (GA) and phosphate solubilization was also characteristic for  
409 the *Bacillus* spp. isolated from the seeds of *C. armena*.

410 Our results demonstrated that at genus level *Paenibacillus*, *Bacillus*, *Psychrobacillus*,  
411 *Domibacillus* and *Paenisporosarcina* were well represented in the seeds of the investigated  
412 population of *C. armena* (Table 2). The dominating *Paenisporosarcina* have been described  
413 as *gen. nov.* and not sufficiently investigated (Parte, 2018). However, some members of the  
414 family Planococcaceae were isolated from a semi-arid tropical soil from India (Raj et al.,  
415 2013). Thus, their presence in the examined seeds is not surprising because of the natural  
416 habitats of *C. armena* (Fig. 1B).

417 Forty-three isolated strains were well adapted to the growing conditions of their host plant  
418 and showed potential PGP traits (production of organic acids, ACC-deaminase, IAA and  
419 siderophores). Most of the isolated strains (83.7%) were positive for IAA production (Fig. 3).  
420 Endophytic bacteria can increase plant growth through their ability to produce plant growth  
421 hormones, particularly auxins. Auxin producing PGP endophytes improve plant growth even  
422 under stress by effectively mitigating the effects of all the growth inhibiting conditions  
423 (Grobela et al., 2018). Respectively 26 and 27 of the isolates produced ACC-deaminase and  
424 organic acids, and only 3 *Bacillus* spp. could produce siderophores (Fig. 3). All these traits  
425 have potential to improve plant growth also under stress conditions (Grobela et al., 2018;

426 Shameer and Prasad, 2018). Hassan and Bano (2016) explored the IAA production of  
427 *Stenotrophomonas maltophilia* strains isolated from a halophytic herb *Cenchrus ciliaris* and  
428 mentioned that bacterial IAA production played a positive role in the salt tolerance of their  
429 host plant.

430 Compared to *Bacillus* spp. and *Paenibacillus* spp. strains that demonstrated relatively low  
431 levels of production of PGP compounds, *Pantoea* spp. and *Stenotrophomonas maltophilia*  
432 demonstrated a high production of IAA (100%), ACC-deaminase (100%) and organic acids  
433 (96.3%) (Fig. 3), which is in agreement with earlier reports (Singh and Jha, 2017; Lumactud  
434 and Fulthorpe, 2018). The production of various organic acids by seed endophytic  
435 *Paenibacillus* sp., *Pantoea* sp., and *Bacillus* sp. inhibits the growth of pathogens and can  
436 significantly enhance plant growth and resistance against plant pathogens (Herrera et al.,  
437 2016; Shahzad et al., 2017). The high levels of IAA production among *P. agglomerans* and *S.*  
438 *maltophilia* strains correspond with findings of other authors (Ambawade and Pathade 2015;  
439 Luziatelli et al., 2020).

440

### 441 Conclusion

442 We explored the endophytic bacterial community of the seeds of the endemic holoparasite *C.*  
443 *armena*. The sterilization procedure for the seed surface was optimized. Ten phyla and 256  
444 bacterial genera were identified. However, also some unclassified and unexplored taxonomic  
445 groups were found in the seeds.

446 Our results confirm that spore forming *Bacillus* spp. are common and dominated  
447 endophytes from seeds of plants growing in harsh environmental conditions, especially from  
448 arid saline soils. *Pantoea* spp. and *Stenotrophomonas* seem the most favourable PGP  
449 endophytes in seeds of *C. armena*. The PGP traits of these bacteria, such as production of  
450 IAA, ACC-deaminase and organic acids seem correlated with the natural habitat of their hosts  
451 and have the potential to improve plant tolerance against abiotic stresses. To elucidate the  
452 effective benefits of these endophytic bacteria for their host plants, particularly for the seeds,  
453 seed germination and development of the seedling, more research is required.

454

455 **Data availability.** The sequence data available in the NCBI Genbank  
456 (<https://www.ncbi.nlm.nih.gov/>) Sequence Read Archive with accession number  
457 PRJNA819412.

458

### 459 Supplementary material.

460 **Supplementary Figure S1.** The values of reads of *Cistanche armena* seed-endophytes after  
461 filtering (a) Chao1 index was 27, (b) Shannon-Wiener biodiversity index was 2.82, (c)  
462 Simpson index 13.9 for total 75 different OTUs. P-value: 0.05.

463 **Supplementary Figure S2.** The relative abundances of the dominated bacteria comprising  
464 the seed associated endophytic community of *Cistanche armena* species at different  
465 taxonomic levels at a) Phylum, b) Class, c) Order, d) Genera, e) Families.

466 **Acknowledgments.** The manuscript was prepared under “Partnership agreement governing  
467 the joint supervision and awarding of a doctorate diploma between Jan Kochanowski  
468 University in Kielce (Poland) and Hasselt University (Belgium)” (K.P.). We thank Dr. Yuliya  
469 Krasnylenko for taking photographs under a zoom microscope.

470

471 **Author contributions.** Conceptualization, K.P.; originator of the research topic, R.P.; field  
472 research, R.P.; methodology, K.P., W.K., J.V., K.R., S.T. and R.P.; Bioinformatic and  
473 statistical analysis, S.T.; resources, R.P., W.K., K.R., K.P. and J.V.; writing the original draft

474 preparation, K.P., R.P and K.R.; writing the review and editing, R.P., W.K., J.V.;  
475 visualization, K.P., R.P, S.T. and K.R. All authors read and approved the final manuscript.

476  
477 **Financial support.** The author acknowledges financial support through the project  
478 “Development Accelerator of the Jan Kochanowski University of Kielce,” co-financed by the  
479 European Union under the European Social Fund, (K.P., POWR.03.05.00-00-Z212/18, 2019-  
480 2023). This study was supported by grants from the Jan Kochanowski University (K.R.,  
481 666 065, 2019), (W.K.; K.P., SUPB.RN. 21.235, 2021-2022). The field research in this study  
482 in Armenia was partially financed by the National Geographic grant (R.P., GEFNE 192-16,  
483 2017). This study was also supported by a BOF-BILA grant from Hasselt University Belgium  
484 BOF21BL12 (K.P.; J.V., 2021-2022) and the Hasselt University Methusalem project (J.V.,  
485 08M03VGRJ).

486  
487 **Conflicts of interest.** The authors declare that the research was conducted in the absence of  
488 any commercial or financial relationships that could be construed as a potential conflict of  
489 interest.

490  
491  
492 **References**

- 493  
494 **Ambawade MS and Pathade GR.** (2015) Production of indole acetic acid (IAA) by  
495 *Stenotrophomonas maltophilia* BE25 isolated from roots of banana (*Musa* spp).  
496 *International Journal of Science and Research* 4(1), 2644-2650.
- 497 **Asaf S, Aaqil Khan M, Latif Khan A, Waqas M, Shahzad R, Kim A-Y, Kang S-M and**  
498 **Lee I-J.** (2017) Bacterial endophytes from arid land plants regulate endogenous hormone  
499 content and promote growth in crop plants: an example of *Sphingomonas sp.* and *Serratia*  
500 *marcescens*. *Journal of Plant Interactions* 12(1), 31–38.  
501 <https://doi.org/10.1080/17429145.2016.1274060>
- 502 **Barret M, Guimbaud J-F, Darrasse A and Jacques M-A.** (2016) Plant microbiota affects  
503 seed transmission of phytopathogenic microorganisms. *Molecular Plant Pathology* 17(6),  
504 791-795. <https://doi.org/10.1111/mpp.12382>
- 505 **Barthlott W.** (1981) Epidermal and seed surface characters of plants: systematic applicability  
506 and some evolutionary aspects. *Nordic Journal of Botany* 1(3), 345–355.  
507 <https://doi.org/10.1111/j.1756-1051.1981.tb00704.x>
- 508 **Belimov AA, Hontzeas N, Safronova VI, Demchinskaya SV, Piluzza G, Bullitta S and**  
509 **Glick BR.** (2005) Cadmium-tolerant plant growth-promoting bacteria associated with the  
510 roots of Indian mustard (*Brassica juncea* L. Czern.). *Soil Biology and Biochemistry* 37(2),  
511 241-250. <https://doi.org/10.1016/j.soilbio.2004.07.033>
- 512 **Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ and Holmes SP.** (2016)  
513 DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*  
514 13(7), 581–583. <https://doi.org/10.1038/nmeth.3869>
- 515 **Compant S, Samad A, Faist H and Sessitsch A.** (2019) A review on the plant microbiome:  
516 Ecology, functions, and emerging trends in microbial application. *Journal of Advanced*  
517 *Research* 19, 29–37. <https://doi.org/10.1016/j.jare.2019.03.004>
- 518 **Cunningham JE, Kuiack C.** (1992) Production of citric and oxalic acids and solubilization  
519 of calcium phosphate by *Penicillium bilaii*. *Applied Environmental Microbiology* 58(5),  
520 1451-1458. <https://journals.asm.org/doi/10.1128/aem.58.5.1451-1458.1992>
- 521 **Davis NM, Proctor DM, Holmes SP, Relman DA and Callahan BJ.** (2018) Simple  
522 statistical identification and removal of contaminant sequences in marker-gene and  
523 metagenomics data. *Microbiome* 6(1), 226. <https://doi.org/10.1186/s40168-018-0605-2>

- 524 **Dinesh R, Anandaraj M, Kumar A, Bini YK, Subila KP and Aravind R.** (2015) Isolation,  
 525 characterization, and evaluation of multi-trait plant growth promoting rhizobacteria for  
 526 their growth promoting and disease suppressing effects on ginger. *Microbiological*  
 527 *Research* **173**, 34–43. <https://doi.org/10.1016/j.micres.2015.01.014>
- 528 **Durlík K, Żarnowiec P, Piwowarczyk R. and Kaca W.** (2021) Culturable endophytic  
 529 bacteria from *Phelipanche ramosa* (Orobanchaceae) seeds. *Seed Science Research* **31**(1),  
 530 69–75. <https://doi.org/10.1017/S0960258520000343>
- 531 **Eevers N, Gielen M, Sánchez-López A, Jaspers S, White JC, Vangronsveld J. and**  
 532 **Weyens N.** (2015) Optimization of isolation and cultivation of bacterial endophytes  
 533 through addition of plant extract to nutrient media. *Microbial Biotechnology* **8**(4), 707–  
 534 715. <https://doi.org/10.1111/1751-7915.12291>
- 535 **Eriksson O. and Kainulainen K.** (2011) The evolutionary ecology of dust seeds.  
 536 *Perspectives in Plant Ecology, Evolution and Systematics* **13**(2), 73–87.  
 537 <https://doi.org/10.1016/j.ppees.2011.02.002>
- 538 **Etesami H and Beattie GA.** (2018) Mining halophytes for plant growth-promoting  
 539 halotolerant bacteria to enhance the salinity tolerance of non-halophytic crops. *Frontiers in*  
 540 *Microbiology* **9**, 148. <https://doi.org/10.3389/fmicb.2018.00148>
- 541 **Fitzpatrick CR and Schneider AC.** (2020) Unique bacterial assembly, composition, and  
 542 interactions in a parasitic plant and its host. *Journal of Experimental Botany* **71**(6), 2198–  
 543 2209. <https://doi.org/10.1093/jxb/erz572>
- 544 **Frank AC, Saldierna Guzmán JP and Shay JE.** (2017) Transmission of bacterial  
 545 endophytes. *Microorganisms* **5**(4), 70. <https://doi.org/10.3390/microorganisms5040070>
- 546 **Glassner H, Zchori-Fein E, Yaron S, Sessitsch A, Sauer U and Compant S.** (2018)  
 547 Bacterial niches inside seeds of *Cucumis melo* L. *Plant and Soil* **422**, 101–113.  
 548 <https://doi.org/10.1007/s11104-017-3175-3>
- 549 **Grobelak A, Kokot P, Świątek J, Jaskulak M, Rorat A.** (2018) Bacterial ACC deaminase  
 550 activity in promoting plant growth on areas contaminated with heavy metals. *Journal of*  
 551 *Ecological Engineering* **19**(5), 150–157. <https://doi.org/10.12911/22998993/89818>
- 552 **Hallmann J, Quadt-Hallmann A, Mahaffee WF and Kloepper JW.** (1997) Bacterial  
 553 endophytes in agricultural crops. *Canadian Journal of Microbiology* **43**(10), 895–914.  
 554 <https://doi.org/10.1139/m97-131>
- 555 **Hassan TU and Bano A.** (2016) Comparative effects of wild type *Stenotrophomonas*  
 556 *maltophilia* and its indole acetic acid-deficient mutants on wheat. *Plant Biology (Stuttgart,*  
 557 *Germany)* **18**(5), 835–841. <https://doi.org/10.1111/plb.12477>
- 558 **Hemida KA and Reyad AM.** (2019) Improvement salt tolerance of safflower plants by  
 559 endophytic bacteria. *Journal of Horticulture and Plant Research* **5**, 38–56.  
 560 <https://doi.org/10.18052/www.scipress.com/JHPR.5.38>
- 561 **Herrera SD, Grossi C, Zawoznik M and Groppa MD.** (2016) Wheat seeds harbour  
 562 bacterial endophytes with potential as plant growth promoters and biocontrol agents of  
 563 *Fusarium graminearum*. *Microbiological Research* **186-187**, 37–43.  
 564 <https://doi.org/10.1016/j.micres.2016.03.002>
- 565 **Hryniewicz K, Patz S and Ruppel S.** (2019) *Salicornia europaea* L. as an underutilized  
 566 saline-tolerant plant inhabited by endophytic diazotrophs. *Journal of Advanced Research*  
 567 **19**, 49–56. <https://doi.org/10.1016/j.jare.2019.05.002>
- 568 **Huet S, Pouvreau J-B, Delage E, Delgrange S, Marais C, Bahut M, Delavault P, Simier P**  
 569 **and Poulin L.** (2020) Populations of the parasitic plant *Phelipanche ramosa* influence  
 570 their seed microbiota. *Frontiers in Plant Science* **11**, 1075.  
 571 <https://doi.org/10.3389/fpls.2020.01075>
- 572 **Iasur Kruh L, Lahav T, Abu-Nassar J, Achdari G, Salami R, Freilich S and Aly R.**  
 573 (2017) Host-parasite-bacteria triangle: the microbiome of the parasitic weed *Phelipanche*

- 574 *aegyptiaca* and tomato-*Solanum lycopersicum* (Mill.) as a host. *Frontiers in Plant Science*  
 575 **8**, 269. <https://doi.org/10.3389/fpls.2017.00269>
- 576 **Joel DM, Hershenhorn Y, Eizenberg H, Aly R, Ejeta G, Rich PJ, Ransom JK,**  
 577 **Sauerborn J. and Rubiales D.** (2007). Biology and management of weedy root parasites.  
 578 In: J. Janick (ed.) *Horticultural Reviews*. John Wiley & Sons **33**, 267-349.  
 579 <https://doi.org/10.1002/9780470168011.ch4>
- 580 **Kusstatscher P, Adam E, Wicaksono WA, Bernhart M, Olimi E, Müller H and Berg G.**  
 581 (2021) Microbiome-assisted breeding to understand cultivar-dependent assembly in  
 582 *Cucurbita pepo*. *Frontiers in Plant Science* **12**, 642027.  
 583 <https://doi.org/10.3389/fpls.2021.642027>
- 584 **Li Z, Lin H, Gu L, Gao J and Tzeng C-M.** (2016) Herba *Cistanche* (Rou Cong-Rong): One  
 585 of the best pharmaceutical gifts of traditional chinese medicine. *Frontiers in Pharmacology*  
 586 **7**(41). <https://doi.org/10.3389/fphar.2016.00041>
- 587 **Lumactud R and Fulthorpe RR.** (2018) Endophytic bacterial community structure and  
 588 function of herbaceous plants from petroleum hydrocarbon contaminated and non-  
 589 contaminated sites. *Frontiers in Microbiology* **9**, 1926.  
 590 <https://doi.org/10.3389/fmicb.2018.01926>
- 591 **Luziatelli F, Ficca AG, Bonini P, Muleo R, Gatti L, Meneghini M, Tronati M, Melini F**  
 592 **and Ruzzi M** (2020) A genetic and metabolomic perspective on the production of indole-  
 593 3-acetic acid by *Pantoea agglomerans* and use of their metabolites as biostimulants in  
 594 plant nurseries. *Frontiers in Microbiology* **11**, 1475.  
 595 <https://doi.org/10.3389/fmicb.2020.01475>
- 596 **Manasa KM, Vasanthakumari MM, Nataraja KN and Uma Shaanker R.** (2020)  
 597 Endophytic fungi of salt adapted *Ipomea pes-caprae* L. R. Br: their possible role in  
 598 inducing salinity tolerance in paddy (*Oryza sativa* L.). *Current Science* **118**(9), 1448–1453.  
 599 <https://doi.org/10.18520/cs/v118/i9/1448-1453>
- 600 **Manjunatha BS, Asha AD, Nivetha N, Bandeppa, Govindasamy V, Rathi MS and**  
 601 **Sangeeta P.** (2017) Evaluation of endophytic bacteria for their influence on plant growth  
 602 and seed germination under water stress conditions. *International Journal of Current*  
 603 *Microbiology and Applied Sciences* **6**(11), 4061–4067.  
 604 <https://doi.org/10.20546/ijemas.2017.611.475>
- 605 **McMurdie PJ and Holmes S.** (2013) phyloseq: An R package for reproducible interactive  
 606 analysis and graphics of microbiome census data. *PLoS One* **8**(4), e61217.  
 607 <https://doi.org/10.1371/journal.pone.0061217>
- 608 **Metwaly A, Salama GMY and Ali GA.** (2018) Using hydrogen peroxide for reducing  
 609 bacterial contamination in date palm tissue culture. *International Journal of Advances in*  
 610 *Agricultural Science and Technology* **5**(4), 25–33.
- 611 **Murali A, Bhargava A, and Wright ES** (2018) IDTAXA: a novel approach for accurate  
 612 taxonomic classification of microbiome sequences. *Microbiome* **6**(1), 140.  
 613 <https://doi.org/10.1186/s40168-018-0521-5>
- 614 **Nickrent DL.** (2020) Parasitic angiosperms: How often and how many? *Taxon* **69**(1), 5–27.  
 615 <https://doi.org/10.1002/tax.12195>
- 616 **Panosyan H, Hakobyan A, Birkeland N-K and Trchounian A.** (2018) Bacilli community  
 617 of saline-alkaline soils from the Ararat Plain (Armenia) assessed by molecular and culture-  
 618 based methods. *Systematic and Applied Microbiology* **41**(3), 232–240.  
 619 <https://doi.org/10.1016/j.syapm.2017.12.002>
- 620 **Parte AC.** (2018) LPSN — List of Prokaryotic names with Standing in Nomenclature  
 621 (bacterio.net), 20 years on. *International Journal of Systematic and Evolutionary*  
 622 *Microbiology* **68**(6), 1825-1829. <https://doi.org/10.1099/ijsem.0.002786>

- 623 **Patten CL and Glick BR.** (2002) Role of *Pseudomonas putida* indoleacetic acid in  
 624 development of the host plant root system. *Applied and Environmental Microbiology*  
 625 **68**(8), 3795–3801. <https://doi.org/10.1128/AEM.68.8.3795-3801.2002>
- 626 **Petrosyan K, Piwowarczyk R, Ruraż K, Thijs S, Vangronsveld J and Kaca W** (2022)  
 627 Seed associated microbial communities of holoparasitic *Cistanche* species from Armenia  
 628 and Portugal p.125 in Proceedings from XVI International Conference on Plant Physiology  
 629 and Plant Science, January 2022, Zurich, Switzerland.
- 630 **Piwowarczyk R.** (2013) Seed productivity in relation to other shoot features for endangered  
 631 parasitic plant *Orobanche picridis* F.W. Schultz (Orobanchaceae). *Polish Journal of*  
 632 *Ecology* **61**(1), 55–64.
- 633 **Piwowarczyk R, Kwolek D, Góralski G, Denysenko M, Joachimiak AJ and Aleksanyan**  
 634 **A.** (2017) First report of the holoparasitic flowering plant *Cistanche armena* on Caspian  
 635 Manna (*Alhagi maurorum*) in Armenia. *Plant Disease* **101**(3), 512–512.  
 636 <https://doi.org/10.1094/PDIS-10-16-1469-PDN>
- 637 **Piwowarczyk R, Ochmian I, Lachowicz S, Kapusta I, Sotek Z and Błaszczak M.** (2020a)  
 638 Phytochemical parasite-host relations and interactions: A *Cistanche armena* case study.  
 639 *Science of The Total Environment* **716**, 137071.  
 640 <https://doi.org/10.1016/j.scitotenv.2020.137071>
- 641 **Piwowarczyk R, Ruraż K, Krasylenko Y, Kasińska J and Sánchez-Pedraja Ó.** (2020b)  
 642 Seed micromorphology of representatives of holoparasitic Orobanchaceae genera from the  
 643 Caucasus region and its taxonomic significance. *Phytotaxa* **432**(3), 223–251.  
 644 <https://doi.org/10.11646/phytotaxa.432.3.1>
- 645 **Piwowarczyk R, Sánchez Pedraja Ó, Moreno Moral G, Fayvush G, Zakaryan N,**  
 646 **Kartashyan N and Aleksanyan A.** (2019) Holoparasitic Orobanchaceae (*Cistanche*,  
 647 *Diphelypaea*, *Orobanche*, *Phelipanche*) in Armenia: distribution, habitats, host range and  
 648 taxonomic problems. *Phytotaxa* **386**(1), 001–106.  
 649 <https://doi.org/10.11646/phytotaxa.386.1.1>
- 650 **PNA Bio PCR Blockers.** mPNA & pPNA. Available at:  
 651 [https://www.pnabio.com/products/PCR\\_blocker.htm](https://www.pnabio.com/products/PCR_blocker.htm).
- 652 **Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J and Glöckner**  
 653 **FO.** (2013) The SILVA ribosomal RNA gene database project: improved data processing  
 654 and web-based tools. *Nucleic Acids Research* **41**(D1), D590–D596.  
 655 <https://doi.org/10.1093/nar/gks1219>
- 656 **Raj PS, Sasikala Ch, Ramaprasad EVV, Subhash Y, Busse H-J, Schumann P and**  
 657 **Ramana ChV.** (2013) *Chryseomicrobium amylolyticum* sp. nov., isolated from a semi-arid  
 658 tropical soil, and emended descriptions of the genus *Chryseomicrobium* and  
 659 *Chryseomicrobium imtechense*. *International Journal of Systematic and Evolutionary*  
 660 *Microbiology* **63**(Pt7), 2612–2617. <https://doi.org/10.1099/ijs.0.044552-0>
- 661 **Ruraż K, Piwowarczyk R, Gajdoš P, Krasylenko Y and Čertík M.** (2020) Fatty acid  
 662 composition in seeds of holoparasitic Orobanchaceae from the Caucasus region: Relation  
 663 to species, climatic conditions and nutritional value. *Phytochemistry* **179**, 112510.  
 664 <https://doi.org/10.1016/j.phytochem.2020.112510>
- 665 **Sánchez-López AS, Pintelon I, Stevens V, Imperato V, Timmermans J-P, González-**  
 666 **Chávez C, Carrillo-González R, Van Hamme J, Vangronsveld J and Thijs S.** (2018)  
 667 Seed endophyte microbiome of *Crotalaria pumila* unpeeled: identification of plant-  
 668 beneficial *Methylobacteria*. *International Journal of Molecular Sciences* **19**(1), 291.  
 669 <https://doi.org/10.3390/ijms19010291>
- 670 **Schneider AC and Moore AJ.** (2017) Parallel Pleistocene amphitropical disjunctions of a  
 671 parasitic plant and its host. *American Journal of Botany* **104**(11), 1745–1755.  
 672 <https://doi.org/10.3732/ajb.1700181>

- 673 **Schwyn B and Neilands JB.** (1987) Universal chemical assay for the detection and  
 674 determination of siderophores. *Analytical Biochemistry* **160**(1), 47-56.  
 675 [https://doi.org/10.1016/0003-2697\(87\)90612-9](https://doi.org/10.1016/0003-2697(87)90612-9)
- 676 **Shahzad R, Khan AL, Bilal S, Asaf S and Lee I-J.** (2017) Plant growth-promoting  
 677 endophytic bacteria versus pathogenic infections: an example of *Bacillus*  
 678 *amyloliquefaciens* RWL-1 and *Fusarium oxysporum* f. sp. *lycopersici* in tomato. *PeerJ* **5**,  
 679 e3107. <https://doi.org/10.7717/peerj.3107>
- 680 **Shameer S and Prasad TNVKV.** (2018) Plant growth promoting rhizobacteria for  
 681 sustainable agricultural practices with special reference to biotic and abiotic stresses. *Plant*  
 682 *Growth Regulation* **84**(3), 603–615. <https://doi.org/10.1007/s10725-017-0365-1>
- 683 **Shrivastava P and Kumar R.** (2015) Soil salinity: A serious environmental issue and plant  
 684 growth promoting bacteria as one of the tools for its alleviation. *Saudi Journal of*  
 685 *Biological Sciences* **22**(2), 123–131. <https://doi.org/10.1016/j.sjbs.2014.12.001>
- 686 **Singh RP and Jha PN.** (2017) The PGPR *Stenotrophomonas maltophilia* SBP-9 augments  
 687 resistance against biotic and abiotic stress in wheat plants. *Frontiers in Microbiology* **8**,  
 688 1945. <https://doi.org/10.3389/fmicb.2017.01945>
- 689 **Soussi A, Ferjani R, Marasco R, Guesmi A, Cherif H, Rolli E, Mapelli F, Ouzari HI,**  
 690 **Daffonchio D and Cherif A.** (2016) Plant-associated microbiomes in arid lands: diversity,  
 691 ecology and biotechnological potential. *Plant and Soil* **405**(1-2), 357–370.  
 692 <https://doi.org/10.1007/s11104-015-2650-y>
- 693 **Szymańska S, Borruso L, Brusetti L, Hulisz P, Furtado B and Hryniewicz K.** (2018)  
 694 Bacterial microbiome of root-associated endophytes of *Salicornia europaea* in  
 695 correspondence to different levels of salinity. *Environmental Science and Pollution*  
 696 *Research* **25**, 25420–25431. <https://doi.org/10.1007/s11356-018-2530-0>
- 697 **Truyens S, Beckers B, Thijs S, Weyens N, Cuypers A and Vangronsveld J.** (2016) The  
 698 effects of the growth substrate on cultivable and total endophytic assemblages of  
 699 *Arabidopsis thaliana*. *Plant and Soil* **405**(1-2), 325–336. [https://doi.org/10.1007/s11104-](https://doi.org/10.1007/s11104-015-2761-5)  
 700 [015-2761-5](https://doi.org/10.1007/s11104-015-2761-5)
- 701 **Truyens S, Jambon I, Croes S, Janssen J, Weyens N, Mench M, Carleer R, Cuypers A**  
 702 **and Vangronsveld J.** (2014) The effect of long-term Cd and Ni exposure on seed  
 703 endophytes of *Agrostis capillaris* and their potential application in phytoremediation of  
 704 metal-contaminated soils. *International Journal of Phytoremediation* **16**(7–8), 643–659.  
 705 <https://doi.org/10.1080/15226514.2013.837027>
- 706 **Truyens S, Weyens N, Cuypers A and Vangronsveld J.** (2013) Changes in the population  
 707 of seed bacteria of transgenerationally Cd-exposed *Arabidopsis thaliana*. *Plant Biology*  
 708 **15**(6), 971–981. <https://doi.org/10.1111/j.1438-8677.2012.00711.x>
- 709 **Ulrich K, Ulrich A and Ewald D.** (2008) Diversity of endophytic bacterial communities in  
 710 poplar grown under field conditions. *FEMS Microbiology Ecology* **63**(2), 169–180.  
 711 <https://doi.org/10.1111/j.1574-6941.2007.00419.x>
- 712 **Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, Gilbert**  
 713 **JA, Jansson JK, Caporaso JG, Fuhrman JA, Apprill A and Knight R.** (2016)  
 714 Improved bacterial 16S rRNA gene (V4 and V4-5) and fungal internal transcribed spacer  
 715 marker gene primers for microbial community surveys. *mSystems* **1**(1), e00009-15.  
 716 <https://doi.org/10.1128/mSystems.00009-15>
- 717 **Watts JE, de Villiers OT and Watts L.** (1993) Sterilization of wheat seeds for tissue culture  
 718 purposes. *South African Journal of Botany* **59**(6), 641–642. [https://doi.org/10.1016/s0254-](https://doi.org/10.1016/s0254-6299(16)30683-4)  
 719 [6299\(16\)30683-4](https://doi.org/10.1016/s0254-6299(16)30683-4)
- 720 **Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J,**  
 721 **Ludwig W and Glöckner FO.** (2014) The SILVA and “All-species Living Tree Project



722 (LTP)” taxonomic frameworks. *Nucleic Acids Research* **42**(D1), D643–D648.  
723 <https://doi.org/10.1093/nar/gkt1209>

724 **Yoneyama K, Xiaonan X, Sekimoto H, Takeuchi Y, Ogasawara S, Akiyama K, Hayashi**  
725 **H and Yoneyama K.** (2008) Strigolactones, host recognition signals for root parasitic  
726 plants and arbuscular mycorrhizal fungi, from Fabaceae plants. *New Phytologist* **179**(2),  
727 484-494. <https://doi.org/10.1111/j.1469-8137.2008.02462.x>

728  
729  
730

731 **Figures and tables captions:**

732

733 **Figure 1.** General habit of the studied species and its habitats: (A) parasitic plant *Cistanche*  
734 *armena*, (B) semi-deserts with halophytic vegetation - the natural habitat of *C. armena*.  
735 Photos by R. Piwowarczyk.

736

737 **Figure 2.** ZOOM microscopy micrographs of seeds of *Cistanche armena*.

738

739 **Figure 3.** PGP activity of tested bacteria and relative PGP traits between isolated bacterial  
740 species (%). The left figure presents the PGP activity for all tested isolates. The figure on  
741 right shows the relative IAA (blue), ACCD (violet), siderophore (green) and organic acids  
742 (red) production ability among the isolated bacterial genera.

743

744 **Table 1.** Endophytic bacterial taxa isolated from different tissues of holoparasitic plant  
745 species

746

747 **Table 2.** Cumulative list of dominating endophytic bacteria in the seeds of *Cistanche armena*  
748 and their taxonomic information

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772 **Table 1.** Endophytic bacterial taxa isolated from different tissues of holoparasitic plant species

Holoparasitic plant	Endophytic bacteria
<i>Phelipanche aegyptiaca</i> , host plant: tomato ( <i>Lycopersicum esculentum</i> )	Pre-haustorium stage $\alpha, \beta, \gamma, \delta$ Proteobacteria, Actinobacteria, Flavobacteria, Sphingobacteria
	Spider stage $\alpha, \beta, \gamma, \delta$ Proteobacteria, Flavobacteria, Sphingobacteria, Firmicutes
Iasur Kruh et al., 2017	Shoots $\alpha, \beta, \gamma$ Proteobacteria, Actinobacteria, Sphingobacteria, Clostridia, Flavobacteria, Firmicutes
<i>Orobanche hederæ</i> , host plant: ivy ( <i>Hedera</i> sp.)	Roots Armatimonadetes, Bacteroidetes, Proteobacteria, Actinobacteria, Acidobacteria, Verrucomicrobia
	Leaves Bacteroidetes, Actinobacteria, Proteobacteria
<i>Phelipanche ramosa</i> , host plants: oilseed rape ( <i>Brassica napus</i> ), hemp ( <i>Cannabis sativa</i> ), tomato ( <i>Solanum lycopersicum</i> ), tobacco ( <i>Nicotiana tabacum</i> ), sunflower ( <i>Helianthus annuus</i> ), melon ( <i>Cucumis melo</i> )	Seeds Proteobacteria, Actinobacteria, Bacteroidetes, Firmicutes
	Huet et al., 2020; Durlík et al., 2021

773  
774  
775  
776  
777  
778  
779  
780

**Table 2.** Cumulative list of dominating endophytic bacteria in the seeds of *Cistanche armena* and their taxonomic information

Phyla	Classes	Orders	Families	Genera
Firmicutes	Bacilli	Paenibacillales	Paenibacillaceae	<i>Paenibacillus</i>
		Bacillales	Bacillaceae	<i>Psychrobacillus</i> <i>Bacillus</i> <i>Domibacillus</i>
			Planococcaceae	<i>Paenisporosarcina</i>
Proteobacteria	$\gamma$ Proteobacteria	Xanthomonadales	Xanthomonadaceae	<i>Stenotrophomonas</i>
		Pseudomonadales	Pseudomonadaceae	<i>Pseudomonas</i>
		Enterobacteriales	Yersiniaceae	<i>Serratia</i>
Actinobacteriota	Actinobacteria	Micrococcales	Microbacteriaceae	<i>Microbacterium</i> <i>Curtobacterium</i>

781