

Comparison of the Quantity of Calcium in the Aortic Valve and the Coronary Arteries in Men Versus Women Who Underwent Transcatheter Aortic Valve Implantation



Gurpreet K. Singh, MD^a, E. Mara Vollema, MD^a, Jan Stassen, MD^{a,b}, Alexander van Rosendaal, MD^a, Tea Gegenava, MD^a, Frank van der Kley, MD^a, Juhani Knuuti, MD, PhD^d, Nina Ajmone Marsan, MD, PhD^a, Victoria Delgado, MD, PhD^{a,c}, and Jeroen J. Bax, MD, PhD^{a,d,*}

Several studies have shown an association between aortic stenosis (AS) and coronary atherosclerosis. This study aimed to evaluate the gender differences in aortic valve calcium (AVC) and coronary artery calcium (CAC) and the association between CAC and all-cause mortality in patients with severe AS. A total of 260 patients (80 ± 7 years, 39% men) with severe AS who were scheduled for transcatheter aortic valve implantation (TAVI) were included. AVC and CAC before TAVI were assessed by noncontrast cardiac computed tomography. Patients with coronary intervention or aortic valve replacement before cardiac computed tomography were excluded. Standard reference values of CAC score were used to classify the percentile groups and the distribution of AVC was assessed. The primary end point was all-cause mortality. In men, the AVC score was 3,911 Hounsfield units (HUs) (interquartile range [IQR] 2,525 to 5,259) and in women, 2,409 HU (IQR 1,588 to 3,359) ($p < 0.001$). CAC score in men was 824 HU (IQR 328 to 1,855) and in women, 478 HU (IQR 136 to 962) ($p < 0.001$). In men, the AVC score increased along with the CAC score, whereas in women, the AVC score was similar across the CAC percentile groups. During a median follow-up of 1,095 days, 59 patients (23%) died. No significant gender-difference was seen in all-cause mortality for CAC score ($p = 0.187$). Men with severe AS show higher AVC and CAC scores than women. Although the pattern of CAC distribution was similar between men and women, the AVC score increased along with the CAC score in men; whereas, in women, the AVC score remained similar across the various percentiles. CAC score was not associated with cumulative mortality in patients with severe AS who underwent TAVI. © 2022 Published by Elsevier Inc. (Am J Cardiol 2022;182:83–88)

Calcific aortic stenosis (AS) valve is the most common valvular heart disease in the aging population.^{1,2} Although the underlying pathophysiologic process of calcification of the aortic valve remains incompletely understood, several studies have shown an association between calcific AS, atherosclerosis, and the presence of cardiovascular risk factors.^{1–4} Therefore, it has been hypothesized that calcific AS and coronary atherosclerosis share similar pathophysiologic mechanisms. Interestingly, various studies evaluating the aortic valve and coronary artery calcium (CAC) score with cardiac computed tomography (CT) have shown important gender differences.^{5–7} Men tend to have higher loads of aortic valve calcium (AVC) and CAC than women. CAC measured on CT has also been considered as a risk marker

for cardiovascular events, with patients showing a CAC score >400 Hounsfield units (HUs) having a significantly increased risk for an adverse event.^{8,9} However, there are limited studies that explored the gender differences of the interaction between AVC and CAC and the association between CAC and all-cause mortality in patients with severe AS. Therefore, the present study evaluated the distribution of AVC and CAC in men and women with severe AS. In addition, the prognostic value of CAC in this population was evaluated.

Methods

A total of 260 patients with severe AS who underwent transcatheter aortic valve implantation (TAVI) were included. Patients with coronary artery bypass grafting, percutaneous coronary intervention, or aortic valve replacement before cardiac CT were excluded. The AVC and CAC scores were assessed using noncontrast-enhanced CT. The standard reference values of the CAC score for men and women were used to divide the population in percentile groups, and the distribution of AVC scores across those percentiles of CAC score was assessed.¹⁰ Clinical and demographic data were collected from the electronic patient

^aDepartment of Cardiology, Leiden University Medical Center, Leiden, The Netherlands; ^bDepartment of Cardiology, Jessa Hospital, Hasselt, Belgium; ^cHeart Institute, Hospital University Germans Trias i Pujol, Badalona, Spain; and ^dTurku Heart Center, University of Turku and Turku University Hospital, Turku, Finland. Manuscript received April 1, 2022; revised manuscript received and accepted July 5, 2022.

See page 87 for disclosure information.

*Corresponding author: Tel: +31 31 71 526 2020; fax: +31 31 71 526 6809

E-mail address: j.j.bax@lumc.nl (J.J. Bax).

records (EPD Vision, version 12.3.5.0, Leiden, The Netherlands). For this retrospective analysis, the institutional review board waived the need for patient written informed consent.

Transthoracic echocardiography was performed in patients at rest, before TAVI. With the patient in the left lateral decubitus position, 2-dimensional, color, and continuous- and pulsed wave Doppler data were obtained in the parasternal and apical views, using commercially available ultrasound systems (E9 and E95, GE Healthcare, Horten, Norway), equipped with M5S transducers. Data were stored digitally for offline analysis using the EchoPac software (version BT13; GE Medical Systems, Horten, Norway). Left ventricular (LV) volumes were measured on apical 2- and 4-chamber views and indexed to body surface area (BSA).¹¹ LV ejection fraction was calculated using the Simpson method.¹¹ LV dimensions were measured on the parasternal long-axis view and LV mass was calculated according to Devereux formula.¹¹ The aortic valve morphology was assessed on the parasternal short-axis images to identify the number of cusps (tricuspid vs bicuspid).¹² On the 3- and 5-chamber apical views, the peak aortic jet velocity was measured using continuous-wave Doppler and, according to the Bernoulli equation, the aortic valve mean and peak gradients were calculated. The pulsed wave Doppler images of the LV outflow tract were obtained from the same apical views and the aortic valve area (AVA) was calculated using the continuity equation. AVA was also indexed to BSA. Severe AS was defined as an AVA <1.0 cm² or indexed AVA <0.6 cm²/m².¹²

Multidetector row CT scans were performed using a 320-slice CT scanner (Aquilion ONE, Toshiba Medical Systems, Tochigi-ken, Japan) or with a 64-slice CT scanner (Aquilion 64, Toshiba Medical Systems).¹³ The multidetector CT acquisition protocol started with a prospective calcium scan (collimation 4 × 3.0 mm, tube voltage and current of 120 kV and 200 mA), followed by the contrast-enhanced CT.¹³ From the noncontrast-enhanced images, the Agatston AVC and CAC scores were assessed. Offline analysis of the data was performed in a remote workstation with dedicated CT software (Vitrea, Vital Images, Minnetonka, Minnesota).

The primary end point was all-cause mortality. Survival data were collected from the departmental Cardiology Information System (EPD version 12.6.2.0) and were complete for all patients.

Continuous variables are presented as mean ± SD if normally distributed and were analyzed using the unpaired Student's *t* test. Continuous variables that were not normally distributed are presented as median and interquartile range (IQR) and were analyzed using the Mann-Whitney *U* test. Categorical data are presented as frequencies and percentages and were analyzed using the chi-square test. Kaplan–Meier curves were performed for survival analysis, and differences between groups were analyzed using the log-rank test. A Cox proportional hazards model was conducted with gender and CAC >400 HU as co-variables. The groups were divided according to gender and CAC score < or >400 HU.^{8,9} A 2-sided *p* <0.05 was considered statistically significant. Statistical analysis was performed using the SPSS software (version 25.0; IBM, Armonk, New York).

Results

Clinical characteristics of the patient population are shown in [Table 1](#). A total of 260 patients, 102 men (mean age 79 ± 7 years) and 158 women (mean age 81 ± 7 years), were included. Women were significantly older, had a lower BSA, and less impaired renal function than men. Except for smoking, cardiovascular risk factors were not significantly different between both genders. There was no significant difference in medication use.

Echocardiographic and CT characteristics of the patient population are shown in [Table 2](#). Women had significantly smaller LV end-systolic and end-diastolic volumes indexed than men. There was no significant difference between men and women regarding the morphology of the aortic valve. The AVC score was significantly higher in men than women (2,409 HU [IQR 1,588 to 3,359], *p* <0.001). Similarly, the CAC score was significantly higher in men (824 HU [IQR 328 to 1,855]) than in women (478 HU [IQR 136 to 962], *p* <0.001). The distribution of AVC and CAC scores in the patient population are shown in [Fig. 1](#). Men had higher AVC and CAC scores compared with women. In men, the AVC score increased along with the CAC score, whereas in women the AVC scores were similar across the various percentiles of CAC score.

The Kaplan–Meier curve for all-cause mortality is shown in [Fig. 2](#). During a median follow-up of 1,095 (IQR 581 to 1,095) days, 59 patients (23%) died. There was no significant difference in mortality rates between men and women for each of the CAC score groups (*p* = 0.187). The hazard ratio was increased for mortality in men and CAC score >400 HU; although, this did not reach statistical significance ([Supplementary Table 1](#)).

Discussion

The present study showed that there is a discordance between the interaction of CAC and AVC between gender, whereas the cumulative mortality was similar in men and women.

In both men and women, AVC and CAC scores have been associated with an increased risk of atherosclerotic cardiovascular events and all-cause mortality.^{9,14–17} However, studies investigating the interaction between AVC and CAC scores in patients with severe AS are limited. Patients with severe AS tend to be older than the participants included in population-based studies, from which the categories of CAC score have been derived.¹⁸ In addition, they frequently present with concomitant coronary artery disease.^{19,20} Therefore, we investigated the interaction between these 2 CT-derived scores in a selected population, excluding patients with previous coronary artery or aortic valve intervention. Similar to previous studies, the present study demonstrated that men have higher AVC and CAC scores than women.^{5–7} Importantly, men also showed an association between the CAC score and the AVC score (with both scores increasing in parallel), whereas women did not show an increase in the AVC score along with the CAC score. The pathophysiologic mechanisms underlying this observation merit further investigation. A possible explanation could be the hormonal differences between

Table 1
Clinical characteristics

Variable	Total population(n = 260)	Men(n = 102)	Women(n = 158)	p-Value
Age (years)	80 ± 7	79 ± 7	81 ± 7	0.030
Body mass index (kg/m ²)	26 ± 5	27 ± 4	26 ± 6	0.637
Creatinine (μmol/liter)	88 (70-109)	100 (82-129)	80 (65-99)	<0.001
Systolic blood pressure (mmHg)	138 ± 23	134 ± 24	141 ± 23	0.024
Diastolic blood pressure (mmHg)	69 ± 13	70 ± 13	69 ± 13	0.425
CAD	46 (18%)	19 (19%)	27 (17%)	0.751
Diabetes mellitus	74 (28%)	35 (34%)	39 (25%)	0.093
Hypertension	192 (74%)	73 (72%)	119 (75%)	0.502
Dyslipidemia	135 (52%)	54 (53%)	81 (51%)	0.792
Smoker	69 (27%)	36 (35%)	33 (21%)	0.010
Family history of CAD	32 (23%)	10 (19%)	22 (26%)	0.372
β-blockers	138 (53%)	50 (49%)	88 (56%)	0.292
ACE inhibitors or ARB	129 (50%)	49 (48%)	80 (51%)	0.683
Statins	119 (46%)	54 (53%)	65 (41%)	0.062
Diuretics	157 (60%)	62 (61%)	95 (60%)	0.916
Aspirin	95 (37%)	39 (38%)	56 (35%)	0.648
Oral anticoagulants	102 (39%)	46 (45%)	56 (35%)	0.120

Values are mean ± SD, median [IQR] or n (%). Diabetes mellitus was defined as having a history of diabetes mellitus and medical therapy with insulin, oral glucose-lowering drugs or diet; Hypertension was defined as a systolic blood pressure of >140 mmHg and/or a diastolic blood pressure of >90 mmHg or prior use of antihypertensive medication; dyslipidemia was defined as previous statin use and/or having a documented history of dyslipidemia.

ACE = angiotensin converting enzyme; ARB = angiotensin receptor blocker; CAD = coronary artery disease.

men and women. It has been postulated that the female gender hormone (estrogen) plays a role in preventing the inflammatory process, which is associated with arterial and valvular calcification.²¹ However, even in postmenopausal women (in whom estrogen concentrations are declining), gender differences in the degree of arterial and valvular calcification are still observed.²² This might suggest that the protective effects of estrogen in younger female patients still remain in postmenopausal women. Nonetheless, irrespective of the hormonal status, studies have shown differences in vascular calcification in men and women despite the same severity of cardiovascular risk factors. These gender differences in calcification are also seen in the aortic valve. Simard et al²³ reported that for the same degree of AS severity, women had less valvular calcification but more fibrosis than men. Differences in coronary plaque

composition between genders have also been described. Women are more likely to have less calcified plaques and more noncalcified plaques than men.^{24,25} Moreover, the impact of various cardiovascular risk factors on coronary artery disease is different between genders, with diabetes mellitus and smoking being a stronger risk factor for coronary artery disease in women than men.^{26,27} As such, underlying cardiovascular risk factors could perhaps also have a different impact on the development of AVC in men compared with women. Previous studies have suggested a possible correlation between AVC and CAC scores.^{22,28} In a healthy population, Nasir et al²⁸ reported that the presence of AVC was independently associated with an increasing severity of coronary atherosclerosis. In contrast, Liyanage et al²² reported no correlation between AVC and CAC in elderly patients with severe AS. The present study provides

Table 2
Echocardiography and computed tomography characteristics

Variable	Total population(n=260)	Men(n=102)	Women(n=158)	p-Value
Echocardiography				
Tricuspid AV morphology	254 (98%)	101 (99%)	153 (97%)	0.252
LVOT diameter (mm)	20.2 ± 2.8	21.2 ± 3.6	19.6 ± 1.9	<0.001
LV end systolic volume indexed (ml/m ²)	23 (15-42)	27 (18-49)	21 (14-37)	0.004
LV end diastolic volume indexed (ml/m ²)	51 ± 22	56 ± 23	48 ± 20	0.005
LV ejection fraction (%)	54 ± 14	53 ± 14	55 ± 14	0.291
Stroke volume index (ml/m ²)	37 ± 13	36 ± 11	36 ± 14	0.354
Mean aortic pressure gradient (mmHg)	45 ± 16	44 ± 15	46 ± 17	0.433
Peak aortic pressure gradient (mmHg)	72 ± 26	70 ± 25	73 ± 27	0.255
Aortic valve area (cm ²)	0.71 ± 0.20	0.76 ± 0.19	0.68 ± 0.19	0.001
Computed tomography				
Aortic valve calcium score (HU)	2776 (1786-4168)	3911 (2525-5259)	2409 (1588-3359)	<0.001
Coronary artery calcium score (HU)	617 (172-1392)	824 (328-1855)	478 (136-962)	<0.001

Values are mean ± SD, median [IQR] or n (%).

AV = aortic valve; LV = left ventricular; LVOT = left ventricular outflow tract; HU = Hounsfield unit.




Figure 1. AVC and CAC in men (A) and women (B) categorized by percentiles using the standard reference of CAC score.

more insights in this topic and shows that the AVC score increases in parallel with the CAC score in men but not in women.

Several studies have reported that patients with an increased CAC score have an increased risk of cardiovascular events and mortality.^{9,14,17} A study in 13,092 asymptomatic participants that was performed by Nakanishi et al⁹ reported a strong association between the CAC score and long-term mortality, with the highest risk being observed in participants with a CAC score ≥ 400 . This finding was confirmed by Miedema et al¹⁷ in a much larger study (22,346 patients) and showed that patients with a CAC score >100 already had a fivefold increased risk for mortality because of coronary heart disease and a threefold increased risk for mortality related to cardiovascular disease, compared with patients with a CAC score of 0. When specifically looking at gender differences, Nakanishi et al⁹ did not find any significant difference in the risk of all-cause mortality between men and women at any level of CAC. Similarly, in our study with patients with severe AS who underwent TAVI, we did not detect any significant difference between men and women with a CAC score >400 compared with a CAC score

<400 . These findings suggest that although most patients with severe AS who underwent TAVI receive a preprocedural cardiac CT, CAC score cannot be used to risk stratify these patients or assess their risk for future events. Larger prospective trials are needed to confirm these results.

The present study has limitations. First, this is a single-center, retrospective study with inherent limitations related to its retrospective design. Second, this study only included patients who underwent TAVI and not surgical aortic valve replacement. Third, only patients with noncontrast-enhanced CT to assess the coronary and AVC were included, which could have resulted in selection bias. Fourth, different CT scans were used (64-slice CT vs 320-CT). Fifth, the aortic valve morphology was only assessed by transthoracic echocardiography.

In conclusion, men with severe AS have higher AVC and CAC scores than women. Although the pattern of CAC distribution is similar between men and women, the AVC score increases along with the CAC score in men; whereas in women, the AVC scores remained similar across the various percentiles. CAC score was not associated with cumulative mortality in patients with severe AS who underwent TAVI.




Figure 2. Kaplan–Meier analysis of all-cause mortality according to gender and CAC score < or >400 Hounsfield units.

Disclosures

The department of Cardiology of the Leiden University Medical Center received research grants from Abbott Vascular, Bayer, Bioventrix, Biotronik, Boston Scientific, Edwards Lifesciences, GE Healthcare, and Medtronic. Dr. Delgado received speaker fees from Abbott Vascular, Edward Lifesciences, GE Healthcare, MSD, and Medtronic. Dr. Marsan received speaker fees from Abbott Vascular and GE Healthcare. Dr. Bax received speaker fees from Abbott Vascular. The remaining authors have no conflicts of interest to declare.

Supplementary materials

Supplementary material associated with this article can be found in the online version at <https://doi.org/10.1016/j.amjcard.2022.07.008>.

- Baumgartner H, Falk V, Bax JJ, De Bonis M, Hamm C, Holm PJ, Jung B, Lancellotti P, Lansac E, Rodriguez Muñoz D, Rosenhek R, Sjögren J, Tornos Mas P, Vahanian A, Walther T, Wendler O, Windecker S, Zamorano JL. ESC Scientific Document Group. 2017 ESC/EACTS guidelines for the management of valvular heart disease. *Eur Heart J* 2017;38:2739–2791.
- Carabello BA, Paulus WJ. Aortic stenosis. *Lancet* 2009;373:956–966.
- Stewart BF, Siscovick D, Lind BK, Gardin JM, Gottdiener JS, Smith VE, Kitzman DW, Otto CM. Clinical factors associated with calcific aortic valve disease. Cardiovascular health study. *J Am Coll Cardiol* 1997;29:630–634.
- Yan AT, Koh M, Chan KK, Guo H, Alter DA, Austin PC, Tu JV, Wijeyesundera HC, Ko DT. Association between cardiovascular risk factors and aortic stenosis: the CANHEART Aortic Stenosis Study. *J Am Coll Cardiol* 2017;69:1523–1532.
- Aggarwal SR, Clavel MA, Messika-Zeitoun D, Cuffe C, Malouf J, Araoz PA, Mankad R, Michelena H, Vahanian A, Enriquez-Sarano M. Sex differences in aortic valve calcification measured by multidetector computed tomography in aortic stenosis. *Circ Cardiovasc Imaging* 2013;6:40–47.
- Nakao YM, Miyamoto Y, Higashi M, Noguchi T, Ohishi M, Kubota I, Tsutsui H, Kawasaki T, Furukawa Y, Yoshimura M, Morita H, Nishimura K, Kada A, Goto Y, Okamura T, Tei C, Tomoike H, Naito H, Yasuda S. Sex differences in impact of coronary artery calcification to predict coronary artery disease. *Heart* 2018;104:1118–1124.
- Pawade T, Clavel MA, Tribouilloy C, Dreyfus J, Mathieu T, Tastet L, Renard C, Gun M, Jenkins WSA, Macron L, Sechrist JW, Lacomis JM, Nguyen V, Galian Gay L, Cuéllar Calabria H, Ntalas I, Carlidge TRG, Prendergast B, Rajani R, Evangelista A, Cavalcante JL, Newby DE, Pibarot P, Messika Zeitoun D, Dweck MR. Computed tomography aortic valve calcium scoring in patients with aortic stenosis. *Circ Cardiovasc Imaging* 2018;11:e007146.
- Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, Himmelfarb CD, Khera A, Lloyd-Jones D, McEvoy JW, Michos ED, Miedema MD, Muñoz D, Smith SC Jr, Virani SS, Williams KA Sr, Yeboah J, Ziaeian B. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines [published correction appears in *Circulation* 2019;140:e649–e650] [published correction appears in *Circulation* 2020;141:e60] [published correction appears in *Circulation* 2020;141:e774] *Circulation* 2019;140:e596–e646.
- Nakanishi R, Li D, Blaha MJ, Whelton SP, Darabian S, Flores FR, Dailing C, Blumenthal RS, Nasir K, Berman DS, Budoff MJ. All-cause mortality by age and gender based on coronary artery calcium scores. *Eur Heart J Cardiovasc Imaging* 2016;17:1305–1314.
- Hoff JA, Chomka EV, Krainik AJ, Daviglius M, Rich S, Kondos GT. Age and gender distributions of coronary artery calcium detected by

- electron beam tomography in 35,246 adults. *Am J Cardiol* 2001;87:1335–1339.
11. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, Flachskampf FA, Foster E, Goldstein SA, Kuznetsova T, Lancellotti P, Muraru D, Picard MH, Rietzschel ER, Rudski L, Spencer KT, Tsang W, Voigt JU. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging [published correction appears in *Eur Heart J Cardiovasc Imaging* 2016;17:412] [published correction appears in *Eur Heart J Cardiovasc Imaging* 2016;17:969]. *Eur Heart J Cardiovasc Imaging* 2015;16:233–270.
 12. Baumgartner H, Hung J, Bermejo J, Chambers JB, Edvardsen T, Goldstein S, Lancellotti P, LeFevre M, Miller F Jr, Otto CM. Recommendations on the echocardiographic assessment of aortic valve stenosis: a focused update from the European Association of Cardiovascular Imaging and the American Society of Echocardiography. *J Am Soc Echocardiogr* 2017;30:372–392.
 13. van Rosendaal PJ, Kamperidis V, Kong WK, van Rosendaal AR, Marsan NA, Bax JJ, Delgado V. Comparison of quantity of calcific deposits by multidetector computed tomography in the aortic valve and coronary arteries. *Am J Cardiol* 2016;118:1533–1538.
 14. Lahti SJ, Feldman DI, Dardari Z, Mirbolouk M, Orimoloye OA, Osei AD, Graham G, Rumberger J, Shaw L, Budoff MJ, Rozanski A, Miedema MD, Al-Mallah MH, Berman D, Nasir K, Blaha MJ. The association between left main coronary artery calcium and cardiovascular-specific and total mortality: the Coronary Artery Calcium Consortium. *Atherosclerosis* 2019;286:172–178.
 15. Otto CM, Lind BK, Kitzman DW, Gersh BJ, Siscovick DS. Association of aortic-valve sclerosis with cardiovascular mortality and morbidity in the elderly. *N Engl J Med* 1999;341:142–147.
 16. Thomassen HK, Cioffi G, Gerdt E, Einarsen E, Midtbø HB, Mancusi C, Cramariuc D. Echocardiographic aortic valve calcification and outcomes in women and men with aortic stenosis. *Heart* 2017;103:1619–1624.
 17. Miedema MD, Dardari ZA, Nasir K, Blankstein R, Knickelbine T, Oberemba S, Shaw L, Rumberger J, Michos ED, Rozanski A, Berman DS, Budoff MJ, Blaha MJ. Association of coronary artery calcium with long-term, cause-specific mortality among young adults. *JAMA Netw Open* 2019;2:e197440.
 18. McClelland RL, Chung H, Detrano R, Post W, Kronmal RA. Distribution of coronary artery calcium by race, gender, and age: results from the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation* 2006;113:30–37.
 19. Vandeplas A, Willems JL, Piessens J, De Geest H. Frequency of angina pectoris and coronary artery disease in severe isolated valvular aortic stenosis. *Am J Cardiol* 1988;62:117–120.
 20. Rapp AH, Hillis LD, Lange RA, Cigarroa JE. Prevalence of coronary artery disease in patients with aortic stenosis with and without angina pectoris. *Am J Cardiol* 2001;87:1216–1217. A7.
 21. Zhang B, Miller VM, Miller JD. Influences of sex and estrogen in arterial and valvular calcification. *Front Endocrinol (Lausanne)* 2019;10:622.
 22. Liyanage L, Lee NJ, Cook T, Herrmann HC, Jagasia D, Litt H, Han Y. The impact of gender on cardiovascular system calcification in very elderly patients with severe aortic stenosis. *Int J Cardiovasc Imaging* 2016;32:173–179.
 23. Simard L, Côté N, Dagenais F, Mathieu P, Couture C, Trahan S, Bossé Y, Mohammadi S, Pagé S, Joubert P, Clavel MA. Sex-related discordance between aortic valve calcification and hemodynamic severity of aortic stenosis: is valvular fibrosis the explanation? *Circ Res* 2017;120:681–691.
 24. Shaw LJ, Min JK, Nasir K, Xie JX, Berman DS, Miedema MD, Whelton SP, Dardari ZA, Rozanski A, Rumberger J, Bairey Merz CN, Al-Mallah MH, Budoff MJ, Blaha MJ. Sex differences in calcified plaque and long-term cardiovascular mortality: observations from the CAC Consortium. *Eur Heart J* 2018;39:3727–3735.
 25. Bigeh A, Shekar C, Gulati M. Sex differences in coronary artery calcium and long-term CV mortality. *Curr Cardiol Rep* 2020;22:21.
 26. Njølstad I, Arnesen E, Lund-Larsen PG. Smoking, serum lipids, blood pressure, and sex differences in myocardial infarction. A 12-year follow-up of the Finnmark study. *Circulation* 1996;93:450–456.
 27. Huxley R, Barzi F, Woodward M. Excess risk of fatal coronary heart disease associated with diabetes in men and women: meta-analysis of 37 prospective cohort studies. *BMJ* 2006;332:73–78.
 28. Nasir K, Katz R, Al-Mallah M, Takasu J, Shavelle DM, Carr JJ, Kronmal R, Blumenthal RS, O'Brien K, Budoff MJ. Relationship of aortic valve calcification with coronary artery calcium severity: the Multi-Ethnic Study of Atherosclerosis (MESA). *J Cardiovasc Comput Tomogr* 2010;4:41–46.