

Cognitive Computerized Training for Older Adults and Patients with
Neurological Disorders: Do the Amount and Training Modality Count? An
Umbrella Meta-Regression Analysis

Peer-reviewed author version

BONNECHERE, Bruno & Klass, Malgorzata (2023) Cognitive Computerized Training
for Older Adults and Patients with Neurological Disorders: Do the Amount and
Training Modality Count? An Umbrella Meta-Regression Analysis. In: Games for
Health Journal, 12 (2) , p. 100-117.

DOI: 10.1089/g4h.2022.0120

Handle: <http://hdl.handle.net/1942/39776>

Abstract

Numerous applications have been created to train cognition and challenge the brain; a process known as computerized cognitive training (CCT). Despite potential positive results important questions remain unresolved: the appropriate training duration, the efficacy of CCT depending on its type (commercial or developed in-house for the rehabilitation of specific patients) and delivery mode (at-home or on-site), and the patients most likely to benefit such intervention. This study aims to perform an umbrella meta-analysis and meta-regression to determine if the type of CCT, the delivery mode, the amount of training, and participants' age at inclusion influence the improvement of the cognitive function. To do so we performed a umbrella meta-analysis. One hundred studies were included in this analysis representing 6,407 participants. Statistical improvements were found for the different conditions after the training. We do not find statistical difference between the type of intervention or the delivery mode. No dose-response relationship between the total amount of training and the improvement of cognitive functions was found. CCT is effective in improving cognitive function in patients suffering from neurological conditions and in healthy aging. There is therefore an urgent need for health care systems to recognize its therapeutic potential and to evaluate at a larger scale their integration into the clinical pipeline as preventive and rehabilitation tool.

Keywords: Cognitive Brain Training, mHealth, rehabilitation

Introduction

Cognitive impairments are common symptoms of most neurological diseases¹⁻⁴ and cognitive decline is associated with aging.⁵ Various cognitive training programs have been developed for decades to try to preserve, increase, restore, or slow down the decrease of cognitive functions. Such kind of interventions have been validated to improve cognition of young healthy people,⁶ healthy older adults,⁷ and patients with Mild Cognitive Impairment (MCI).⁸ Cognitive interventions have also been developed and successfully tested in patients with various neurological disorders such as stroke,⁹ Multiple Sclerosis (MS),¹⁰ and Parkinson's disease (PD).¹¹ However classic center-based¹² cognitive training programs may be difficult to implement since they require the presence of healthcare professionals, additional travels to the centre and may seem boring and repetitive for the patients.¹³ The development and generalization of computer and mobile technologies offer new possibilities in the management of cognitive deficits in various populations.¹⁴ In parallel to this rapid growth, the industry of software and mobile apps is exploding. Health-related apps make up an important part of this market, and numerous solutions (including online platforms, and mobile apps) have been developed to 'train' cognition and challenge the brain, what is called computerized cognitive training (CCT). CCT is a digital health application that enables consumers to access interactive cognitive exercises from a personal computer or mobile. These exercises can be designed to enhance general cognition or specific cognitive functions (i.e., learning and memory, attention, processing speed, executive functioning). The advantage over classic center-based cognitive training is that they can be performed as well in center as at home, the exercise level is continuously adapted based on the score obtained in the exercises or based on (self-administered) cognitive tests, and adherence can be monitored remotely.¹⁵

Currently two types of approaches are possible to perform CCT; using commercially available solutions,¹⁶ part of which are not primarily developed for clinical applications, or solutions that

have been developed in-house specifically for rehabilitation of patients with specific disorders. A lot of studies have been performed to validate the use of CCT, these works have been synthesized in different systematic reviews and meta-analyses in various conditions such as aging,^{17,18} MCI,¹⁹ stroke,²⁰ PD,²¹ and MS.²² Despite these analyses, important clinical issues remain unknown such as defining the appropriate duration and frequency of the intervention, the best type of intervention and delivery mode, and the patients most likely to benefit from this type of intervention. Another important question, not been answered yet, is to determine if training performed independently at home is as efficient as centre-based training. Most of the interventions are indeed performed in centres under the supervision of clinicians while one of the potential advantages of such solutions is that they could be used to train patients in home-environment without requiring the continuous presence of healthcare professionals and frequent travels to the center. Therefore this study aims to perform an umbrella meta-analysis to determine if the type of CCT (commercial vs. specific), the delivery mode (at-home or on-site), the amount of training, and participants' age at inclusion influence the improvement of the cognitive function.

Methods

We performed a second analysis of the meta-analyses by synthesizing evidence from multiple meta-analyses performed in this field. We performed an omnibus meta-analysis, which pooled all primary studies across all potential moderators, followed by individual meta-analyses for various potential moderators, to synthesize the effect size across the first order meta-analyses.²⁴

The protocol of the present study was registered in the international prospective register of systematic reviews PROSPERO (registration number CRD42020167321). This systematic review and omnibus meta-analysis were reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) recommendations.²⁵ For the present study, no ethics committee approval was necessary.

Search strategy and selection criteria

We searched the PubMed electronic database, Web of Sciences, Embase, Scopus, and Sciences Direct for relevant meta-analyses published up to the 31st of March 2022. MeSH terms and free words referring to brain training (“cognitive training”, “brain training”, “memory training”, “reasoning training”, “attention training”, “processing speed training”), games (“video games”, “exergames”, “computer training”, “games”, “mobile games”, “cognitive games”) and meta-analysis were used as keywords. The search was limited to meta-analyses published in English in peer-reviewed journals.

The inclusion and exclusion criteria were as follows. No time period threshold was used because CCT is a fairly recent paradigm. A PICOS approach (Population, Intervention, Control, Outcome, and Study design) was used for the inclusion and exclusion criteria, which were predetermined and assessed by the study team from the selected meta-analysis.

- Population: For studies in older adults: cognitively healthy participants (defined in the study as an MMSE $\geq 27.5/30$) aged above 60 years old. For studies in patients with neurological diseases: stroke patients and patients with MS, PD and MCI, denoted later in the text as conditions.
- Intervention: Studies using CCT to perform cognitive training. The duration of the training must be a minimum of 1 month.²⁶ Studies using action-video games, active-video games or a combination of cognitive and physical rehabilitation exercises (i.e., dual-task paradigms) were not included. Indeed, although these types of intervention could also improve cognitive functions,²⁷ we decided to focus in present study on CCT dedicated to train and challenge specifically cognitive functions to reduce heterogeneity across the different types of intervention and training.
- Control: Active brain training or brain-stimulating activities (e.g., reading, board games, etc.)

- Outcome: Outcomes included performance on one or more cognitive tests, administered both before and after training that provide any validated measure of cognitive functions. The outcomes were categorized according to their transfer-level. The transfer effect refers to the capability of using the knowledge and skills learned in one task to improve performance in similar or different task. We defined three types of transfer: direct transfer (performance at tests that are similar or identical to the exercises of the CCT), near transfer (performance at tests assessing cognitive functions trained during the CCT) and far transfer (performance in tasks that are different from the training tasks in nature or in appearance).²⁸ Due to an incomplete description of the content of exercises used during CCT, it was sometimes difficult to determine if tests used to assess cognitive function were similar or not to the task trained during the CCT and, because of that, to define the type of transfer for those studies. Therefore, we end up with lot of studies classified as 'mixed' transfer. The assessment of the type of transfer, according to the tests used, was made independently by the two authors (B.B. and M.K.). In case of disagreement, the final decision was made after discussion between the two authors.
- Study design: Randomized Controlled Trials (RCTs).

A flow diagram of the study selection with the screened meta-analyses and the selection process is presented in Figure 1.

Quality assessment

The Physiotherapy Evidence Database Scale (PEDro) was used to assess methodological quality of included studies. The PEDro scale is an 11-item scale designed for rating methodological quality of RCTs. Each satisfied item contributes one point to the total PEDro score (item 1 omitted for score calculation, range=0 –10 points).²⁹ Total PEDro scores of 0-3 are considered 'poor quality', 4-5 'moderate quality' and 6-10 'good quality'.

Data extraction

The following information was extracted from the included studies: characteristics of the participants (age, severity of the disease and/or MMSE, when available), type of intervention (commercial or specific), duration of the intervention (total duration of the training, number of weeks, duration of one session), delivery mode (at-home or on-site), type of assessment and transfer, and results of the cognitive assessment. Summary of the included studies is presented in Table 1.

Statistical analysis

The measure of treatment effect was the effect size (standardized mean difference (SMD)), defined as the between-group difference in mean values divided by the pooled SD computed using the Hedge's g method (see Eq. 1)

$$g = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{((n_1 - 1) \times s_1^2 + (n_2 - 1) \times s_2^2)}{(n_1 + n_2 - 2)}}} \quad \text{Eq.1}$$

where \bar{X} is the mean, n the sample size and s the variance.

In most of the studies, several tests are used to evaluate the different cognitive functions of the participants. The results of the different tests were combined, using weighted mean, to produce a single SMD according to Cochrane's recommendation.³⁰ A positive SMD implies better therapeutic effects over time in the intervention group compared to the control group.

To assess the risk of publication bias, funnel plots were checked for asymmetry³¹ and Egger's test for the intercept was applied for the different conditions evaluated.³²

To assess the potential effect of the type of CCT and the delivery mode on the results we performed sub-group analysis.

To ease the visualization of the different parameters of the included studies (type of intervention, transfer, delivery mode, and condition), we used network meta-analysis to plot

the results. The distribution of the available evidence was assessed using a network geometry graph in which the size of the nodes was proportional to the number of trials included for each intervention and the width of the continuous line connecting nodes corresponding to the number of trials directly comparing the interventions.³³

Random-effects meta-regression analysis quantified the association of changes in cognitive functions and the amount of training and the age of the participants. Studies were weighted by the inverse of the sum of the within- and between-study variance.³⁴

Statistical analyses were performed at an overall significance level of 0.05, carried out in RStudio (version 1.2.1335), using R version 3.6.1

Results

Eight meta-analysis were included in this umbrella meta-analysis (older adults^{17,18,35}, MS²², MCI¹⁹, stroke^{20,36}, and PD²¹) to finally include 100 studies in this meta-analysis (48 with older adults^{37–84}, 17 with patients with MS^{85–101}, 15 with patients with MCI^{102–116}, 14 with patients after a stroke^{117–132} and 6 with patients with PD^{133–138}), representing in total 6,407 participants (see Supplementary Figure 1). The quality of the studies has been assessed using the PEDro scale¹³⁹, we did not take into consideration the question about the blinding of the therapist, since the intervention are not performed with therapists, the mean score is 6.6 (1.4) out of 9. A one-way ANOVA was performed to compare the effect of the conditions on study quality. The ANOVA revealed that there was not a statistically significant difference in PEDro scores between at least two groups ($F(4,97) = 2.212$, $p = 0.073$).

Characteristics of the participants, type of intervention, delivery mode, training duration and frequency and type of transfer and individual PEDro scores are presented in Table 2. Description of the interventions, tests used to assess cognitive function are further detailed in Supplementary Table 1. To ease the visualization of the amount of studies assessing the

different modalities (type of intervention, type of transfer, delivery mode, initial cognitive function and condition) we plotted network graphs (Figure 2).

First, we checked for potential publications bias, the funnel plot is presented in Supplementary Figure 2 and did not show significant asymmetry (Egger's intercept= 0.67, $p = 0.75$).

By far the most represented participants are older adults ($n = 3,438$; 70.1 ± 4.7 years old; $MMSE = 28.4 \pm 0.6$), followed by patients with stroke ($n = 842$; 60.3 ± 6.8 years old; $MMSE = 21.9 \pm 3.1$), MS ($n = 738$; 45.8 ± 5.2 years old; $MMSE = 26.3 \pm 3.3$), MCI ($n = 590$; 73.1 ± 3.3 years old; $MMSE = 27.1 \pm 0.9$) and PD ($n = 255$; 68.0 ± 4.7 years old; $MMSE = 26.4 \pm 3.7$).

Concerning the type of CCT, there are slightly more studies carried out using commercial solutions ($n = 56$) compared to specific ones ($n = 45$). It is important to note that there are differences depending on the condition investigated with 70% of the studies in stroke done with specific solutions, 67% in PD, 53% in MCI, 41% in older adults, and only 30% in MS. The repartition of the commercial solutions used (tested in at least 3 studies) is presented in Supplementary Figure 3. The three most used apps are RehaCom ($n = 11$), Posit Sciences ($n = 8$) and Lumosity ($n = 6$).

Concerning the delivery mode, most of the studies (60 %) were done on site (clinical centers), 31% at home and 9% using a combination. Again, results show important differences between conditions. For older adults and MS the ratio between on-site and at-home is 50/50 while for MCI (76%), PD (67%) and stroke (73%) most of the studies are performed on-site.

On average the studies includes a median number of 20 [$P25 = 13$; $P75 = 32$] sessions of 45 minutes [$P25 = 30$; $P75 = 60$] over a 6 weeks period [$P25 = 5$; $P75 = 12$]. The total amount of training is 15 hours [$P25 = 10$; $P75 = 20$]. There is no difference in the total amount of training ($p = 0.42$) between conditions (Figure 3) but well for the average duration of one training session ($p = 0.016$). After adjusting for multiple testing (Bonferroni's correction), the

median duration of one session is indeed longer in MCI (60 min [P25 = 52 ; P75 = 67]) compared to stroke (30 min [P25 = 30 ; P75 = 52], $p = 0.047$). There is also a difference in the number of weeks of training ($p = 0.025$), after adjusting for multiple testing, the median duration of the intervention being longer in MCI (12 weeks [P25 = 6 ; P75 = 16]) compared to stroke (5 weeks [P25 = 4 ; P75 = 8], $p = 0.019$).

In all the included studies the overall effect of training is moderate ($g = 0.25$ [95%CI 0.21 – 0.29], $p < 0.001$, fixed effect model due to low heterogeneity [$I^2 = 34\%$]). We then analyse the different conditions individually, the effect of training on cognition was moderate in stroke ($g = 0.48$ [95%CI -0.17 – 0.79], $p = 0.002$), small but still significant in patients with MS ($g = 0.37$ [95%CI 0.23 – 0.52], $p < 0.001$), MCI ($g = 0.31$ [95%CI 0.15 – 0.47], $p < 0.001$), PD ($g = 0.30$ [95%CI 0.03 – 0.57], $p = 0.028$) and older adults ($g = 0.21$ [95%CI 0.13 – 0.28], $p < 0.001$). Forest plot summarizing the effect of training on global cognition for the different conditions is presented in [Figure 4](#).

Concerning the type of intervention, we compared the studies using specific solutions versus commercial solutions and did not find significant differences between the two types of solutions ($g = 0.25$ [0.16 – 0.34] for commercial solutions and $g = 0.22$ [0.13 – 0.32] for specific ones, $p = 0.69$) on the cognitive improvement. We then compared the delivery mode and did not find significant difference either ($g = 0.23$ [0.02 – 0.60] for at-home training and $g = 0.30$ [0.11 – 0.53] for on-site training, $p = 0.33$). Finally, we compared the type of transfer, note that due to the small number of studies assessing direct transfer ($n = 4$) we only compared the near ($n = 47$) and far ($n = 14$) effects and the combination of both ($n = 31$). There is no statistically significant difference for the three types of transfer evaluated ($g = 0.25$ [0.14 – 0.36] for near effect, $g = 0.25$ [0.09 – 0.41] for far effect and $g = 0.27$ [0.18 – 0.35] for the combination, $p = 0.67$).

We then performed meta-regression to determine if the amount of training influences the cognitive outcomes. As we did not find any difference for the type of CCT, delivery mode or transfer type, we performed the meta-regression including all the studies. We did not find any significant association between the total amount of training ($\beta = -0.0043$, $SE = 0.0024$, $p = 0.07$), the number of weeks of training ($\beta = -0.0006$, $SE = 0.0087$, $p = 0.95$), or the average duration of one session and the improvement in cognitive outcomes ($\beta = 0.0001$, $SE = 0.0016$, $p = 0.95$). There is no association with the age of the participant ($\beta = -0.0041$, $SE = 0.0031$, $p = 0.19$). The results for the different conditions are presented in Table 3 and summarized in Figure 5. Individual results for the different cognitive functions are presented in Supplementary Tables 2-6.

Discussion

To the authors' best knowledge this is the first study comparing the efficacy of CCT in different populations with neurological disorders, in healthy older adults or with MCI. The main result of our analysis is that CCT can be used to train cognitive function in various conditions ranging from normal aging to neurological diseases. First, let's discuss the clinical implications of our findings. Interestingly we did not observe statistically significant differences either comparing specific and commercially available CCT or comparing the delivery mode. CCT, commercial or specific, could therefore be interesting tools to integrate in prevention and rehabilitation.¹⁴⁰ It must be noted here that commercial solutions are more easily available than the specific ones, which may explain why there are more used in research and in clinics. Note that for stroke and PD we observed more usage of specific solutions compared to other conditions. That may be due to the fact that those patients may experience some specific limitations (motor or cognitive) which may limit the use of commercial solutions.

From a practical point of view, we did not observe a difference in the average total amount of training between the five conditions investigated. Based on our analysis including all the

conditions, the median duration of an intervention improving significantly cognitive functions is 15 hours over 6 weeks (3 sessions/week). The meta-regression analyses do not show a relationship between the total amount of training, the number of weeks of training, or the duration of one single session and cognitive improvements whether for all participants or when studying conditions individually. It must be pointed out here that the dose-response relationship in cognitive training has not been investigated much and is not obvious.^{63,141–143} This could be explained by the fact that the interventions protocol are heterogeneous¹⁴⁴ and that the difference may only be observed in the long term and for far transfer effect,¹⁴⁵ while most of the research is focusing on the immediate or short term effect.¹⁴⁶

Interestingly we observed statistically significant differences in the median duration of one session with longer sessions found in MCI (60 min) compared to stroke patients (30 min), the duration of the intervention in weeks was also shorter in stroke patients compared to MCI. We could not determine if this difference is related to clinical aspects (e.g., fatigue, lower dexterity level, etc.) or to the specificity of the patient care.

The findings of this study have to be seen in light of some limitations. First, as seen in Supplementary Table 1, there is a huge variety of tests used to assess different cognitive functions and numerous training applications being used (both commercial and specific ones) whose main objectives, type of exercises and/or graphical interface may be somewhat different. This high heterogeneity make comparison between studies more complicated. Therefore, we decided to calculate a global cognitive effect by combining results of the different cognitive tests for each study. In addition, results for the different cognitive functions are presented in Supplementary Materials but are unpowered due to the smaller number of studies per cognitive function. It would then be interesting for further studies to analyse the specific effects and the potential dose-response relationship for the different cognitive functions.¹⁴⁷

Due to an incomplete description of the content of exercises used during CCT, it was sometimes difficult to determine if tests used to assess cognitive functions were similar or not to the task trained during the CCT and, because of that, to define the type of transfer for those studies. Therefore, we end up with a lot of studies classified as ‘mixed’ transfer. The absence of difference in the subgroup analysis according to the type of transfer may thus be biased by the high number of studies classified as ‘mixed effects’. Also, only few studies investigated far-transfer effects and the neural mechanisms involved. Given the fact that far-transfer effects occur when two different tasks share an underlying processing component and neuroanatomical areas or neural circuits,¹⁴⁸ it would be interesting in the future to have more studies focusing on change in brain function after CCT training to investigate this particular aspect.

Another important limitation is the relatively small number of studies, and therefore of patients, included for PD (6 studies representing 255 patients), therefore the results for PD must be interpreted carefully.

We focused our analysis on the effect of the dose of training and later checked if the age of the participants or the conditions could influence the results. But other parameters may influence the ability to train the cognition such as sex,¹⁴⁹ genetic factors,¹⁵⁰ depression,¹⁵¹ and fatigue¹⁵², and other aspects such as the familiarity with the use of new technologies can influence the ease of use of CCTs and adherence.

We could also assume that the amount of rehabilitation exercises and participation in physical activities may influence cognition. It has indeed been demonstrated that other types of activities such as aerobic training, alone or combined with cognitive exercises, can increase cognition in aging,^{153,154} MCI,¹⁵⁵ MS,¹⁵⁶ stroke¹⁵⁷. Note that for PD the results are still unclear.¹⁵⁸ The participation in cultural, social activities and other stimulating activities are also positively associated with cognitive functions.^{159,160} Therefore it would be interesting to evaluate the part

of improvement attributable to the cognitive training and the part attributable to other activities. Finally, most of the participants have a relatively good and preserved cognitive function (as evaluated by a median MMSE score of 28). Further studies are therefore needed to determine if such kinds of training are feasible and effective in patients with lower cognitive function and determine the minimal cognitive level required to achieve significant improvement.

Conclusions

This study highlights the potential of CCT for prevention and rehabilitation of cognitive decline. The development and widespread use of computer and mobile technology now make it possible to integrate different cognitive interventions into CCT. Despite the fact that the effects were small to moderate, such kind of solutions have several advantages: easily accessible thanks to the large availability of mobile technology (smartphones, tablets), more fun and engaging than traditional exercises, allow to record the scores of the patients and follow their improvements, do not require the continuous presence of trained healthcare specialist which offers the possibility to implement this kind of solution in areas with fewer healthcare professionals and/or for patients living far from rehabilitation centers.¹⁶¹ Our results show no difference in cognitive improvements between trainings performed at home and on site and support the possibility to perform the training at home with a remote monitoring. In the absence of pharmaceutical treatment to prevent or slow down cognitive decline and due the low accessibility to classical center-based cognitive interventions, there is an urgent need to assess the therapeutic potential of CCT and to evaluate at a larger scale their integration into the clinical pipeline.

Authors' contribution

BB gathered literature sources. BB wrote the initial manuscript with critical input from MK. BB and MK contributed to writing and approved the final version of the manuscript.

Conflicts of Interest

None declared

Abbreviations

MCI: Mild Cognitive Impairment

MS: Multiple Sclerosis

PD: Parkinson's disease

CCT: Computerized Cognitive Training

RCT: randomized controlled trial

MMSE: Mini-mental state examination

SMD: standardized mean difference

References

1. Baiano C, Barone P, Trojano L, et al. Prevalence and clinical aspects of mild cognitive impairment in Parkinson's disease: A meta-analysis. *Mov Disord* 2020;35(1):45–54; doi: 10.1002/mds.27902.
2. Landmeyer NC, Bürkner P-C, Wiendl H, et al. Disease-modifying treatments and cognition in relapsing-remitting multiple sclerosis: A meta-analysis. *Neurology* 2020;94(22):e2373–e2383; doi: 10.1212/WNL.0000000000009522.
3. Mihaescu AS, Masellis M, Graff-Guerrero A, et al. Brain degeneration in Parkinson's disease patients with cognitive decline: a coordinate-based meta-analysis. *Brain Imaging Behav* 2019;13(4):1021–1034; doi: 10.1007/s11682-018-9922-0.
4. Vega JN, Newhouse PA. Mild cognitive impairment: diagnosis, longitudinal course, and emerging treatments. *Curr Psychiatry Rep* 2014;16(10):490; doi: 10.1007/s11920-014-0490-8.
5. Colangeli S, Boccia M, Verde P, et al. Cognitive Reserve in Healthy Aging and Alzheimer's Disease: A Meta-Analysis of fMRI Studies. *Am J Alzheimers Dis Other Dement* 2016;31(5):443–449; doi: 10.1177/1533317516653826.
6. Au J, Sheehan E, Tsai N, et al. Improving fluid intelligence with training on working memory: a meta-analysis. *Psychon Bull Rev* 2015;22(2):366–377; doi: 10.3758/s13423-014-0699-x.
7. Hudes R, Rich JB, Troyer AK, et al. The impact of memory-strategy training interventions on participant-reported outcomes in healthy older adults: A systematic review and meta-analysis. *Psychol Aging* 2019;34(4):587–597; doi: 10.1037/pag0000340.
8. Sherman DS, Mauser J, Nuno M, et al. The efficacy of cognitive intervention in mild cognitive impairment (MCI): a meta-analysis of outcomes on neuropsychological measures. *Neuropsychol Rev* 2017;27(4):440–484; doi: 10.1007/s11065-017-9363-3.
9. Merriman NA, Sexton E, McCabe G, et al. Addressing cognitive impairment following stroke: systematic review and meta-analysis of non-randomised controlled studies of psychological interventions. *BMJ Open* 2019;9(2):e024429; doi: 10.1136/bmjopen-2018-024429.
10. Gromisch ES, Fiszdon JM, Kurtz MM. The effects of cognitive-focused interventions on cognition and psychological well-being in persons with multiple sclerosis: A meta-analysis. *Neuropsychol Rehabil* 2020;30(4):767–786; doi: 10.1080/09602011.2018.1491408.
11. Leung IHK, Walton CC, Hallock H, et al. Cognitive training in Parkinson disease: A systematic review and meta-analysis. *Neurology* 2015;85(21):1843–1851; doi: 10.1212/WNL.0000000000002145.

12. Kueider AM, Parisi JM, Gross AL, et al. Computerized cognitive training with older adults: a systematic review. *PLoS ONE* 2012;7(7):e40588; doi: 10.1371/journal.pone.0040588.
13. Boendermaker WJ, Sanchez Maceiras S, Boffo M, et al. Attentional Bias Modification With Serious Game Elements: Evaluating the Shots Game. *JMIR Serious Games* 2016;4(2):e20; doi: 10.2196/games.6464.
14. Weicker J, Villringer A, Thöne-Otto A. Can impaired working memory functioning be improved by training? A meta-analysis with a special focus on brain injured patients. *Neuropsychology* 2016;30(2):190–212; doi: 10.1037/neu0000227.
15. Bodner KA, Goldberg TE, Devanand DP, et al. Advancing Computerized Cognitive Training for MCI and Alzheimer’s Disease in a Pandemic and Post-pandemic World. *Front Psychiatry* 2020;11:557571; doi: 10.3389/fpsyt.2020.557571.
16. Bonnechère B, Jansen B, Omelina L, et al. The use of commercial video games in rehabilitation: a systematic review. *Int J Rehabil Res* 2016;39(4):277–290; doi: 10.1097/MRR.000000000000190.
17. Bonnechère B, Langley C, Sahakian BJ. The use of commercial computerised cognitive games in older adults: a meta-analysis. *Scientific Reports* 2020;10(1):15276; doi: 10.1038/s41598-020-72281-3.
18. Lampit A, Hallock H, Valenzuela M. Computerized Cognitive Training in Cognitively Healthy Older Adults: A Systematic Review and Meta-Analysis of Effect Modifiers. Gandy S. ed. *PLoS Med* 2014;11(11):e1001756; doi: 10.1371/journal.pmed.1001756.
19. Zhang H, Huntley J, Bhome R, et al. Effect of computerised cognitive training on cognitive outcomes in mild cognitive impairment: a systematic review and meta-analysis. *BMJ Open* 2019;9(8):e027062; doi: 10.1136/bmjopen-2018-027062.
20. Ye M, Zhao B, Liu Z, et al. Effectiveness of computer-based training on post-stroke cognitive rehabilitation: A systematic review and meta-analysis. *Neuropsychological Rehabilitation* 2020;1–17; doi: 10.1080/09602011.2020.1831555.
21. Orgeta V, McDonald KR, Poliakoff E, et al. Cognitive training interventions for dementia and mild cognitive impairment in Parkinson’s disease. Cochrane Dementia and Cognitive Improvement Group. ed. *Cochrane Database of Systematic Reviews* 2020; doi: 10.1002/14651858.CD011961.pub2.
22. Lampit A, Heine J, Finke C, et al. Computerized Cognitive Training in Multiple Sclerosis: A Systematic Review and Meta-analysis. *Neurorehabil Neural Repair* 2019;33(9):695–706; doi: 10.1177/1545968319860490.
23. Wilms IL. The computerized cognitive training alliance - A proposal for a therapeutic alliance model for home-based computerized cognitive training. *Heliyon* 2020;6(1):e03254; doi: 10.1016/j.heliyon.2020.e03254.

24. Schmidt FL, Oh I-S. Methods for second order meta-analysis and illustrative applications. *Organizational Behavior and Human Decision Processes* 2013;121(2):204–218; doi: 10.1016/j.obhdp.2013.03.002.
25. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *PLOS Medicine* 2009;6(7):e1000100; doi: 10.1371/journal.pmed.1000100.
26. Sahakian BJ, Bruhl AB, Cook J, et al. The impact of neuroscience on society: cognitive enhancement in neuropsychiatric disorders and in healthy people. *Philos Trans R Soc Lond, B, Biol Sci* 2015;370(1677):20140214; doi: 10.1098/rstb.2014.0214.
27. Mura G, Carta MG, Sancassiani F, et al. Active exergames to improve cognitive functioning in neurological disabilities: a systematic review and meta-analysis. *Eur J Phys Rehabil Med* 2018;54(3):450–462; doi: 10.23736/S1973-9087.17.04680-9.
28. Weng W, Liang J, Xue J, et al. The Transfer Effects of Cognitive Training on Working Memory Among Chinese Older Adults With Mild Cognitive Impairment: A Randomized Controlled Trial. *Front Aging Neurosci* 2019;11:212; doi: 10.3389/fnagi.2019.00212.
29. Maher CG, Sherrington C, Herbert RD, et al. Reliability of the PEDro Scale for Rating Quality of Randomized Controlled Trials. *Physical Therapy* 2003;83(8):713–721; doi: 10.1093/ptj/83.8.713.
30. Higgins JPT, Thomas J, Chandler J, et al. *Cochrane Handbook for Systematic Reviews of Interventions*, 2nd Edition. Wiley-Blackwell. Wiley-Blackwell; 2019.
31. Sterne JAC, Sutton AJ, Ioannidis JPA, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011;343:d4002; doi: 10.1136/bmj.d4002.
32. Pustejovsky JE, Rodgers MA. Testing for funnel plot asymmetry of standardized mean differences. *Res Synth Methods* 2019;10(1):57–71; doi: 10.1002/jrsm.1332.
33. Salanti G, Ades AE, Ioannidis JPA. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of Clinical Epidemiology* 2011;64(2):163–171; doi: 10.1016/j.jclinepi.2010.03.016.
34. Borenstein M, Hedges LV, Higgins JPT, et al. *Introduction to Meta-Analysis*. John Wiley & Sons. John Wiley & Sons; 2009.
35. Gates NJ, Rutjes AW, Di Nisio M, et al. Computerised cognitive training for 12 or more weeks for maintaining cognitive function in cognitively healthy people in late life. *Cochrane Database Syst Rev* 2020;CD012277; doi: 10.1002/14651858.CD012277.pub3.

36. Zhou Y, Feng H, Li G, et al. Efficacy of computerized cognitive training on improving cognitive functions of stroke patients: A systematic review and meta-analysis of randomized controlled trials. *Int J Nurs Pract* 2021;e12966; doi: 10.1111/ijn.12966.
37. Ackerman PL, Kanfer R, Calderwood C. Use it or lose it? Wii brain exercise practice and reading for domain knowledge. *Psychol Aging* 2010;25(4):753–766; doi: 10.1037/a0019277.
38. Anguera JA, Boccanfuso J, Rintoul JL, et al. Video game training enhances cognitive control in older adults. *Nature* 2013;501(7465):97–101; doi: 10.1038/nature12486.
39. Ballesteros S, Prieto A, Mayas J, et al. Brain training with non-action video games enhances aspects of cognition in older adults: a randomized controlled trial. *Front Aging Neurosci* 2014;6:277; doi: 10.3389/fnagi.2014.00277.
40. Ballesteros S, Mayas J, Prieto A, et al. Effects of video game training on measures of selective attention and working memory in older adults: results from a randomized controlled trial. *Front Aging Neurosci* 2017;9; doi: 10.3389/fnagi.2017.00354.
41. Barnes DE, Santos-Modesitt W, Poelke G, et al. The Mental Activity and eXercise (MAX) trial: a randomized controlled trial to enhance cognitive function in older adults. *JAMA Intern Med* 2013;173(9):797–804; doi: 10.1001/jamainternmed.2013.189.
42. Basak C, Boot WR, Voss MW, et al. Can training in a real-time strategy video game attenuate cognitive decline in older adults? *Psychol Aging* 2008;23(4):765–777; doi: 10.1037/a0013494.
43. Berry AS, Zanto TP, Clapp WC, et al. The influence of perceptual training on working memory in older adults. *PLoS One* 2010;5(7):e11537; doi: 10.1371/journal.pone.0011537.
44. Boot WR, Champion M, Blakely DP, et al. Video games as a means to reduce age-related cognitive decline: attitudes, compliance, and effectiveness. *Front Psychol* 2013;4:31; doi: 10.3389/fpsyg.2013.00031.
45. Bottiroli S, Cavallini E. Can computer familiarity regulate the benefits of computer-based memory training in normal aging? A study with an Italian sample of older adults. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2009;16(4):401–418; doi: 10.1080/13825580802691763.
46. Bozoki A, Radovanovic M, Winn B, et al. Effects of a computer-based cognitive exercise program on age-related cognitive decline. *Arch Gerontol Geriatr* 2013;57(1):1–7; doi: 10.1016/j.archger.2013.02.009.
47. Brehmer Y, Westerberg H, Bäckman L. Working-memory training in younger and older adults: training gains, transfer, and maintenance. *Front Hum Neurosci* 2012;6:63; doi: 10.3389/fnhum.2012.00063.

48. Bürki CN, Ludwig C, Chicherio C, et al. Individual differences in cognitive plasticity: an investigation of training curves in younger and older adults. *Psychol Res* 2014;78(6):821–835; doi: 10.1007/s00426-014-0559-3.
49. Buschkuehl M, Jaeggi SM, Hutchison S, et al. Impact of working memory training on memory performance in old-old adults. *Psychol Aging* 2008;23(4):743–753; doi: 10.1037/a0014342.
50. Casutt G, Theill N, Martin M, et al. The drive-wise project: driving simulator training increases real driving performance in healthy older drivers. *Front Aging Neurosci* 2014;6:85; doi: 10.3389/fnagi.2014.00085.
51. Colzato LS, van Muijden J, Band GPH, et al. Genetic Modulation of Training and Transfer in Older Adults: BDNF ValMet Polymorphism is Associated with Wider Useful Field of View. *Front Psychol* 2011;2:199; doi: 10.3389/fpsyg.2011.00199.
52. Dahlin E, Nyberg L, Bäckman L, et al. Plasticity of executive functioning in young and older adults: immediate training gains, transfer, and long-term maintenance. *Psychol Aging* 2008;23(4):720–730; doi: 10.1037/a0014296.
53. Edwards JD, Wadley VG, Myers R enee S, et al. Transfer of a speed of processing intervention to near and far cognitive functions. *Gerontology* 2002;48(5):329–340; doi: 10.1159/000065259.
54. Edwards JD, Wadley VG, Vance DE, et al. The impact of speed of processing training on cognitive and everyday performance. *Aging Ment Health* 2005;9(3):262–271; doi: 10.1080/13607860412331336788.
55. Edwards JD, Valdés EG, Peronto C, et al. The Efficacy of InSight Cognitive Training to Improve Useful Field of View Performance: A Brief Report. *J Gerontol B Psychol Sci Soc Sci* 2015;70(3):417–422; doi: 10.1093/geronb/gbt113.
56. Garcia-Campuzano MT, Virues-Ortega J, Smith S, et al. Effect of cognitive training targeting associative memory in the elderly: a small randomized trial and a longitudinal evaluation. *J Am Geriatr Soc* 2013;61(12):2252–2254; doi: 10.1111/jgs.12574.
57. Heinzl S, Schulte S, Onken J, et al. Working memory training improvements and gains in non-trained cognitive tasks in young and older adults. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2014;21(2):146–173; doi: 10.1080/13825585.2013.790338.
58. Lampit A, Hallock H, Moss R, et al. The Timecourse of Global Cognitive Gains from Supervised Computer-Assisted Cognitive Training: A Randomised, Active-Controlled Trial in Elderly with Multiple Dementia Risk Factors. *J Prev Alzheimers Dis* 2014;1(1):33–39; doi: 10.14283/jpad.2014.18.
59. Lee YM, Jang C, Bak IH, et al. Effects of Computer-assisted Cognitive Rehabilitation Training on the Cognition and Static Balance of the Elderly. *J Phys Ther Sci* 2013;25(11):1475–1477; doi: 10.1589/jpts.25.1475.

60. Legault C, Jennings JM, Katula JA, et al. Designing clinical trials for assessing the effects of cognitive training and physical activity interventions on cognitive outcomes: The Seniors Health and Activity Research Program Pilot (SHARP-P) Study, a randomized controlled trial. *BMC Geriatr* 2011;11:27; doi: 10.1186/1471-2318-11-27.
61. Mahncke HW, Connor BB, Appelman J, et al. Memory enhancement in healthy older adults using a brain plasticity-based training program: a randomized, controlled study. *Proc Natl Acad Sci USA* 2006;103(33):12523–12528; doi: 10.1073/pnas.0605194103.
62. Mayas J, Parmentier FBR, Andrés P, et al. Plasticity of attentional functions in older adults after non-action video game training: a randomized controlled trial. *PLoS ONE* 2014;9(3):e92269; doi: 10.1371/journal.pone.0092269.
63. McAvinue LP, Golemme M, Castorina M, et al. An evaluation of a working memory training scheme in older adults. *Front Aging Neurosci* 2013;5:20; doi: 10.3389/fnagi.2013.00020.
64. Miller KJ, Dye RV, Kim J, et al. Effect of a computerized brain exercise program on cognitive performance in older adults. *Am J Geriatr Psychiatry* 2013;21(7):655–663; doi: 10.1016/j.jagp.2013.01.077.
65. Nouchi R, Taki Y, Takeuchi H, et al. Brain training game boosts executive functions, working memory and processing speed in the young adults: a randomized controlled trial. *PLoS ONE* 2013;8(2):e55518; doi: 10.1371/journal.pone.0055518.
66. Nouchi R, Saito T, Nouchi H, et al. Small acute benefits of 4 weeks processing speed training games on processing speed and inhibition performance and depressive mood in the healthy elderly people: evidence from a randomized control trial. *Front Aging Neurosci* 2016;8; doi: 10.3389/fnagi.2016.00302.
67. O'Brien JL, Edwards JD, Maxfield ND, et al. Cognitive training and selective attention in the aging brain: an electrophysiological study. *Clin Neurophysiol* 2013;124(11):2198–2208; doi: 10.1016/j.clinph.2013.05.012.
68. Peng H, Wen J, Wang D, et al. The Impact of Processing Speed Training on Working Memory in Old Adults. 2012; doi: 10.1007/S10804-012-9142-6.
69. Peretz C, Korczyn AD, Shatil E, et al. Computer-based, personalized cognitive training versus classical computer games: a randomized double-blind prospective trial of cognitive stimulation. *Neuroepidemiology* 2011;36(2):91–99; doi: 10.1159/000323950.
70. Richmond LL, Morrison AB, Chein JM, et al. Working memory training and transfer in older adults. *Psychol Aging* 2011;26(4):813–822; doi: 10.1037/a0023631.
71. Sandberg P, Rönnlund M, Nyberg L, et al. Executive process training in young and old adults. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2014;21(5):577–605; doi: 10.1080/13825585.2013.839777.

72. Shatil E. Does combined cognitive training and physical activity training enhance cognitive abilities more than either alone? A four-condition randomized controlled trial among healthy older adults. *Front Aging Neurosci* 2013;5:8; doi: 10.3389/fnagi.2013.00008.
73. Shatil E, Mikulecká J, Bellotti F, et al. Novel television-based cognitive training improves working memory and executive function. *PLoS ONE* 2014;9(7):e101472; doi: 10.1371/journal.pone.0101472.
74. Simpson T, Camfield D, Pipingas A, et al. Improved processing speed: online computer-based cognitive training in older adults. *Educ Gerontol* 2012;38(7):445–458; doi: 10.1080/03601277.2011.559858.
75. Smith GE, Housen P, Yaffe K, et al. A cognitive training program based on principles of brain plasticity: results from the improvement in memory with plasticity-based adaptive cognitive training (IMPACT) study. *J Am Geriatr Soc* 2009;57(4):594–603; doi: 10.1111/j.1532-5415.2008.02167.x.
76. Stern Y, Blumen HM, Rich LW, et al. Space Fortress game training and executive control in older adults: a pilot intervention. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2011;18(6):653–677; doi: 10.1080/13825585.2011.613450.
77. Strenziok M, Parasuraman R, Clarke E, et al. Neurocognitive enhancement in older adults: comparison of three cognitive training tasks to test a hypothesis of training transfer in brain connectivity. *Neuroimage* 2014;85 Pt 3:1027–1039; doi: 10.1016/j.neuroimage.2013.07.069.
78. Ten Brinke LF, Best JR, Chan JLC, et al. The effects of computerized cognitive training with and without physical exercise on cognitive function in older adults: an 8-week randomized controlled trial. *J Gerontol A Biol Sci Med Sci* 2019; doi: 10.1093/gerona/glz115.
79. von Bastian CC, Langer N, Jäncke L, et al. Effects of working memory training in young and old adults. *Mem Cognit* 2013;41(4):611–624; doi: 10.3758/s13421-012-0280-7.
80. van Muijden J, Band GPH, Hommel B. Online games training aging brains: limited transfer to cognitive control functions. *Front Hum Neurosci* 2012;6:221; doi: 10.3389/fnhum.2012.00221.
81. Vance D, Dawson J, Wadley V, et al. The accelerate study: The longitudinal effect of speed of processing training on cognitive performance of older adults. *Rehabilitation Psychology* 2007;52(1):89–96; doi: 10.1037/0090-5550.52.1.89.
82. Walton CC, Kavanagh A, Downey LA, et al. Online cognitive training in healthy older adults: a preliminary study on the effects of single versus multi-domain training. *Transl Neurosci* 2015;6(1):13–19; doi: 10.1515/tnsci-2015-0003.
83. Wang M-Y, Chang C-Y, Su S-Y. What's Cooking? - Cognitive Training of Executive Function in the Elderly. *Front Psychol* 2011;2:228; doi: 10.3389/fpsyg.2011.00228.

84. Wolinsky FD, Weg MWV, Howren MB, et al. Interim analyses from a randomised controlled trial to improve visual processing speed in older adults: the Iowa Healthy and Active Minds Study. *BMJ Open* 2011;1(2):e000225; doi: 10.1136/bmjopen-2011-000225.
85. Amato MP, Goretti B, Viterbo RG, et al. Computer-assisted rehabilitation of attention in patients with multiple sclerosis: results of a randomized, double-blind trial. *Mult Scler* 2014;20(1):91–98; doi: 10.1177/1352458513501571.
86. Campbell J, Langdon D, Cercignani M, et al. A Randomised Controlled Trial of Efficacy of Cognitive Rehabilitation in Multiple Sclerosis: A Cognitive, Behavioural, and MRI Study. *Neural Plast* 2016;2016:4292585; doi: 10.1155/2016/4292585.
87. Cerasa A, Gioia MC, Valentino P, et al. Computer-assisted cognitive rehabilitation of attention deficits for multiple sclerosis: a randomized trial with fMRI correlates. *Neurorehabil Neural Repair* 2013;27(4):284–295; doi: 10.1177/1545968312465194.
88. Charvet LE, Shaw MT, Haider L, et al. Remotely-delivered cognitive remediation in multiple sclerosis (MS): protocol and results from a pilot study. *Mult Scler J Exp Transl Clin* 2015;1:2055217315609629; doi: 10.1177/2055217315609629.
89. Charvet LE, Yang J, Shaw MT, et al. Cognitive function in multiple sclerosis improves with telerehabilitation: Results from a randomized controlled trial. *PLoS One* 2017;12(5):e0177177; doi: 10.1371/journal.pone.0177177.
90. Chiaravalloti ND, Goverover Y, Costa SL, et al. A Pilot Study Examining Speed of Processing Training (SPT) to Improve Processing Speed in Persons With Multiple Sclerosis. *Front Neurol* 2018;9:685; doi: 10.3389/fneur.2018.00685.
91. De Giglio L, De Luca F, Prosperini L, et al. A low-cost cognitive rehabilitation with a commercial video game improves sustained attention and executive functions in multiple sclerosis: a pilot study. *Neurorehabil Neural Repair* 2015;29(5):453–461; doi: 10.1177/1545968314554623.
92. Grasso MG, Broccoli M, Casillo P, et al. Evaluation of the Impact of Cognitive Training on Quality of Life in Patients with Multiple Sclerosis. *Eur Neurol* 2017;78(1–2):111–117; doi: 10.1159/000478726.
93. Hancock LM, Bruce JM, Bruce AS, et al. Processing speed and working memory training in multiple sclerosis: a double-blind randomized controlled pilot study. *J Clin Exp Neuropsychol* 2015;37(2):113–127; doi: 10.1080/13803395.2014.989818.
94. Hubacher M, Kappos L, Weier K, et al. Case-Based fMRI Analysis after Cognitive Rehabilitation in MS: A Novel Approach. *Front Neurol* 2015;6:78; doi: 10.3389/fneur.2015.00078.
95. Mattioli F, Flavia M, Stampatori C, et al. Efficacy and specificity of intensive cognitive rehabilitation of attention and executive functions in multiple sclerosis. *J Neurol Sci* 2010;288(1–2):101–105; doi: 10.1016/j.jns.2009.09.024.

96. Messinis L, Nasios G, Kosmidis MH, et al. Efficacy of a Computer-Assisted Cognitive Rehabilitation Intervention in Relapsing-Remitting Multiple Sclerosis Patients: A Multicenter Randomized Controlled Trial. *Behav Neurol* 2017;2017:5919841; doi: 10.1155/2017/5919841.
97. Pedullà L, Bricchetto G, Tacchino A, et al. Adaptive vs. non-adaptive cognitive training by means of a personalized App: a randomized trial in people with multiple sclerosis. *J Neuroeng Rehabil* 2016;13(1):88; doi: 10.1186/s12984-016-0193-y.
98. Pérez-Martín MY, González-Platas M, Eguía-Del Río P, et al. Efficacy of a short cognitive training program in patients with multiple sclerosis. *Neuropsychiatr Dis Treat* 2017;13:245–252; doi: 10.2147/NDT.S124448.
99. Pusswald G, Mildner C, Zeberholz K, et al. A neuropsychological rehabilitation program for patients with Multiple Sclerosis based on the model of the ICF. *NeuroRehabilitation* 2014;35(3):519–527; doi: 10.3233/NRE-141145.
100. Solari A, Motta A, Mendozzi L, et al. Computer-aided retraining of memory and attention in people with multiple sclerosis: a randomized, double-blind controlled trial. *J Neurol Sci* 2004;222(1–2):99–104; doi: 10.1016/j.jns.2004.04.027.
101. Tesar N, Bandion K, Baumhackl U. Efficacy of a neuropsychological training programme for patients with multiple sclerosis -- a randomised controlled trial. *Wien Klin Wochenschr* 2005;117(21–22):747–754; doi: 10.1007/s00508-005-0470-4.
102. Barban F, Annicchiarico R, Pantelopoulos S, et al. Protecting cognition from aging and Alzheimer's disease: a computerized cognitive training combined with reminiscence therapy. *Int J of Geriatr Psychiatry* 2016;31(4):340–348; doi: 10.1002/gps.4328.
103. Ciarmiello A, Gaeta MC, Benso F, et al. FDG-PET in the Evaluation of Brain Metabolic Changes Induced by Cognitive Stimulation in aMCI Subjects. *Curr Radiopharm* 2015;8(1):69–75; doi: 10.2174/1874471008666150428122924.
104. Djabelkhir L, Wu Y-H, Vidal J-S, et al. Computerized cognitive stimulation and engagement programs in older adults with mild cognitive impairment: comparing feasibility, acceptability, and cognitive and psychosocial effects. *Clin Interv Aging* 2017;12:1967–1975; doi: 10.2147/CIA.S145769.
105. Fiatarone Singh MA, Gates N, Saigal N, et al. The Study of Mental and Resistance Training (SMART) study—resistance training and/or cognitive training in mild cognitive impairment: a randomized, double-blind, double-sham controlled trial. *J Am Med Dir Assoc* 2014;15(12):873–880; doi: 10.1016/j.jamda.2014.09.010.
106. Finn M, McDonald S. Computerised Cognitive Training for Older Persons With Mild Cognitive Impairment: A Pilot Study Using a Randomised Controlled Trial Design. *Brain Impairment* 2011;12(3):187–199; doi: 10.1375/brim.12.3.187.
107. Finn M, McDonald S. Repetition-lag training to improve recollection memory in older people with amnesic mild cognitive impairment. A randomized controlled trial.

- Neuropsychol Dev Cogn B Aging Neuropsychol Cogn 2015;22(2):244–258; doi: 10.1080/13825585.2014.915918.
108. Gagnon LG, Belleville S. Training of attentional control in mild cognitive impairment with executive deficits: results from a double-blind randomised controlled study. *Neuropsychol Rehabil* 2012;22(6):809–835; doi: 10.1080/09602011.2012.691044.
 109. Gooding AL, Choi J, Fiszdon JM, et al. Comparing three methods of computerised cognitive training for older adults with subclinical cognitive decline. *Neuropsychol Rehabil* 2016;26(5–6):810–821; doi: 10.1080/09602011.2015.1118389.
 110. Han JW, Son KL, Byun HJ, et al. Efficacy of the Ubiquitous Spaced Retrieval-based Memory Advancement and Rehabilitation Training (USMART) program among patients with mild cognitive impairment: a randomized controlled crossover trial. *Alzheimers Res Ther* 2017;9(1):39; doi: 10.1186/s13195-017-0264-8.
 111. Herrera C, Chambon C, Michel BF, et al. Positive effects of computer-based cognitive training in adults with mild cognitive impairment. *Neuropsychologia* 2012;50(8):1871–1881; doi: 10.1016/j.neuropsychologia.2012.04.012.
 112. Hyer L, Scott C, Atkinson MM, et al. Cognitive Training Program to Improve Working Memory in Older Adults with MCI. *Clin Gerontol* 2016;39(5):410–427; doi: 10.1080/07317115.2015.1120257.
 113. Lin F, Heffner KL, Ren P, et al. Cognitive and Neural Effects of Vision-Based Speed-of-Processing Training in Older Adults with Amnesic Mild Cognitive Impairment: A Pilot Study. *J Am Geriatr Soc* 2016;64(6):1293–1298; doi: 10.1111/jgs.14132.
 114. Rosen AC, Sugiura L, Kramer JH, et al. Cognitive training changes hippocampal function in mild cognitive impairment: a pilot study. *J Alzheimers Dis* 2011;26 Suppl 3:349–357; doi: 10.3233/JAD-2011-0009.
 115. Rozzini L, Costardi D, Chilovi BV, et al. Efficacy of cognitive rehabilitation in patients with mild cognitive impairment treated with cholinesterase inhibitors. *Int J Geriatr Psychiatry* 2007;22(4):356–360; doi: 10.1002/gps.1681.
 116. Savulich G, Piercy T, Fox C, et al. Cognitive training using a novel Memory Game on an iPad in patients with amnesic mild cognitive impairment (aMCI). *Int J Neuropsychopharmacol* 2017;20(8):624–633; doi: 10.1093/ijnp/pyx040.
 117. Barker-Collo SL, Feigin VL, Lawes CMM, et al. Reducing attention deficits after stroke using attention process training: a randomized controlled trial. *Stroke* 2009;40(10):3293–3298; doi: 10.1161/STROKEAHA.109.558239.
 118. Cho H-Y, Kim K-T, Jung J-H. Effects of computer assisted cognitive rehabilitation on brain wave, memory and attention of stroke patients: a randomized control trial. *J Phys Ther Sci* 2015;27(4):1029–1032; doi: 10.1589/jpts.27.1029.

119. De Luca R, Leonardi S, Spadaro L, et al. Improving Cognitive Function in Patients with Stroke: Can Computerized Training Be the Future? *Journal of Stroke and Cerebrovascular Diseases* 2018;27(4):1055–1060; doi: 10.1016/j.jstrokecerebrovasdis.2017.11.008.
120. Lin Z, Tao J, Gao Y, et al. Analysis of central mechanism of cognitive training on cognitive impairment after stroke: Resting-state functional magnetic resonance imaging study. *J Int Med Res* 2014;42(3):659–668; doi: 10.1177/0300060513505809.
121. Liu X, Huang X, Lin J, et al. Computer Aided Technology-Based Cognitive Rehabilitation Efficacy Against Patients' Cerebral Stroke. *An Interdisciplinary Journal of Neuroscience and Quantum Physics* 2018;Volume 16(No 4):86–92; doi: 10.14704/nq.2018.16.4.1212.
122. Park J-H, Park J-H. The effects of a Korean computer-based cognitive rehabilitation program on cognitive function and visual perception ability of patients with acute stroke. *J Phys Ther Sci* 2015;27(8):2577–2579; doi: 10.1589/jpts.27.2577.
123. Prokopenko SV, Mozheyko EY, Petrova MM, et al. Correction of post-stroke cognitive impairments using computer programs. *Journal of the Neurological Sciences* 2013;325(1):148–153; doi: 10.1016/j.jns.2012.12.024.
124. van de Ven RM, Murre JMJ, Buitenweg JIV, et al. The influence of computer-based cognitive flexibility training on subjective cognitive well-being after stroke: A multi-center randomized controlled trial. *PLoS One* 2017;12(11):e0187582; doi: 10.1371/journal.pone.0187582.
125. Wentink MM, Berger M a. M, de Kloet AJ, et al. The effects of an 8-week computer-based brain training programme on cognitive functioning, QoL and self-efficacy after stroke. *Neuropsychol Rehabil* 2016;26(5–6):847–865; doi: 10.1080/09602011.2016.1162175.
126. Yoo C, Yong M-H, Chung J, et al. Effect of computerized cognitive rehabilitation program on cognitive function and activities of living in stroke patients. *J Phys Ther Sci* 2015;27(8):2487–2489; doi: 10.1589/jpts.27.2487.
127. Zucchella C, Capone A, Codella V, et al. Assessing and restoring cognitive functions early after stroke. *Funct Neurol* 2014;29(4):255–262.
128. Bo W, Lei M, Tao S, et al. Effects of combined intervention of physical exercise and cognitive training on cognitive function in stroke survivors with vascular cognitive impairment: a randomized controlled trial. *Clin Rehabil* 2019;33(1):54–63; doi: 10.1177/0269215518791007.
129. Jiang C, Yang S, Tao J, et al. Clinical Efficacy of Acupuncture Treatment in Combination With RehaCom Cognitive Training for Improving Cognitive Function in Stroke: A 2 × 2 Factorial Design Randomized Controlled Trial. *Journal of the American Medical Directors Association* 2016;17(12):1114–1122; doi: 10.1016/j.jamda.2016.07.021.

130. Si Hyun Kang, Kim D-K, Kyung Mook Seo, et al. A computerized visual perception rehabilitation programme with interactive computer interface using motion tracking technology — a randomized controlled, single-blinded, pilot clinical trial study. *Clin Rehabil* 2009;23(5):434–444; doi: 10.1177/0269215508101732.
131. Prokopenko S, Bezdenezhnykh A, Mozheyko E, et al. A comparative clinical study of the effectiveness of computer cognitive training in patients with post-stroke cognitive impairments without dementia. *Psych Rus* 2018;11(2):55–67; doi: 10.11621/pir.2018.0205.
132. Park I-S, Yoon J-G. The effect of computer-assisted cognitive rehabilitation and repetitive transcranial magnetic stimulation on cognitive function for stroke patients. *J Phys Ther Sci* 2015;27(3):773–776; doi: 10.1589/jpts.27.773.
133. Alloni A, Quaglini S, Panzarasa S, et al. Evaluation of an ontology-based system for computerized cognitive rehabilitation. *Int J Med Inform* 2018;115:64–72; doi: 10.1016/j.ijmedinf.2018.04.005.
134. Cerasa A, Gioia MC, Salsone M, et al. Neurofunctional correlates of attention rehabilitation in Parkinson's disease: an explorative study. *Neurol Sci* 2014;35(8):1173–1180; doi: 10.1007/s10072-014-1666-z.
135. Edwards JD, Hauser RA, O'Connor ML, et al. Randomized trial of cognitive speed of processing training in Parkinson disease. *Neurology* 2013;81(15):1284–1290; doi: 10.1212/WNL.0b013e3182a823ba.
136. Folkerts A-K, Dorn ME, Roheger M, et al. Cognitive Stimulation for Individuals with Parkinson's Disease Dementia Living in Long-Term Care: Preliminary Data from a Randomized Crossover Pilot Study. *Parkinsons Dis* 2018;2018:8104673; doi: 10.1155/2018/8104673.
137. Lawrence BJ, Gasson N, Johnson AR, et al. Cognitive Training and Transcranial Direct Current Stimulation for Mild Cognitive Impairment in Parkinson's Disease: A Randomized Controlled Trial. *Parkinsons Dis* 2018;2018:4318475; doi: 10.1155/2018/4318475.
138. Petrelli A, Kaesberg S, Barbe MT, et al. Effects of cognitive training in Parkinson's disease: a randomized controlled trial. *Parkinsonism Relat Disord* 2014;20(11):1196–1202; doi: 10.1016/j.parkreldis.2014.08.023.
139. Blobaum P. Physiotherapy Evidence Database (PEDro). *J Med Libr Assoc* 2006;94(4):477–478.
140. Sokolov AA, Collignon A, Bieler-Aeschlimann M. Serious video games and virtual reality for prevention and neurorehabilitation of cognitive decline because of aging and neurodegeneration. *Curr Opin Neurol* 2020;33(2):239–248; doi: 10.1097/WCO.0000000000000791.

141. Brum PS, Borella E, Carretti B, et al. Verbal working memory training in older adults: an investigation of dose response. *Aging Ment Health* 2020;24(1):81–91; doi: 10.1080/13607863.2018.1531372.
142. Teixeira-Santos AC, Moreira CS, Magalhães R, et al. Reviewing working memory training gains in healthy older adults: A meta-analytic review of transfer for cognitive outcomes. *Neuroscience & Biobehavioral Reviews* 2019;103:163–177; doi: 10.1016/j.neubiorev.2019.05.009.
143. Weicker J, Hudl N, Hildebrandt H, et al. The effect of high vs. low intensity neuropsychological treatment on working memory in patients with acquired brain injury. *Brain Injury* 2020;34(8):1051–1060; doi: 10.1080/02699052.2020.1773536.
144. Ophey A, Roheger M, Folkerts A-K, et al. A Systematic Review on Predictors of Working Memory Training Responsiveness in Healthy Older Adults: Methodological Challenges and Future Directions. *Front Aging Neurosci* 2020;12:575804; doi: 10.3389/fnagi.2020.575804.
145. Borella E, Carretti B, Sciore R, et al. Training working memory in older adults: Is there an advantage of using strategies? *Psychol Aging* 2017;32(2):178–191; doi: 10.1037/pag0000155.
146. Holmes SC, Johnson CM, Suvak MK, et al. Examining patterns of dose response for clients who do and do not complete cognitive processing therapy. *J Anxiety Disord* 2019;68:102120; doi: 10.1016/j.janxdis.2019.102120.
147. Kamin ST, Lang FR. Cognitive Functions Buffer Age Differences in Technology Ownership. *Gerontology* 2016;62(2):238–246; doi: 10.1159/000437322.
148. Jonides J. How does practice makes perfect? *Nat Neurosci* 2004;7(1):10–11; doi: 10.1038/nn0104-10.
149. Hyde JS. Sex and cognition: gender and cognitive functions. *Curr Opin Neurobiol* 2016;38:53–56; doi: 10.1016/j.conb.2016.02.007.
150. Lee T, Henry JD, Trollor JN, et al. Genetic influences on cognitive functions in the elderly: a selective review of twin studies. *Brain Res Rev* 2010;64(1):1–13; doi: 10.1016/j.brainresrev.2010.02.001.
151. Price RB, Duman R. Neuroplasticity in cognitive and psychological mechanisms of depression: an integrative model. *Mol Psychiatry* 2020;25(3):530–543; doi: 10.1038/s41380-019-0615-x.
152. Wylie GR, Yao B, Genova HM, et al. Using functional connectivity changes associated with cognitive fatigue to delineate a fatigue network. *Sci Rep* 2020;10(1):21927; doi: 10.1038/s41598-020-78768-3.
153. Barha CK, Davis JC, Falck RS, et al. Sex differences in exercise efficacy to improve cognition: A systematic review and meta-analysis of randomized controlled trials in

- older humans. *Front Neuroendocrinol* 2017;46:71–85; doi: 10.1016/j.yfrne.2017.04.002.
154. Northey JM, Cherbuin N, Pumpa KL, et al. Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis. *Br J Sports Med* 2018;52(3):154–160; doi: 10.1136/bjsports-2016-096587.
 155. Karssemeijer EGA, Aaronson JA, Bossers WJ, et al. Positive effects of combined cognitive and physical exercise training on cognitive function in older adults with mild cognitive impairment or dementia: A meta-analysis. *Ageing Res Rev* 2017;40:75–83; doi: 10.1016/j.arr.2017.09.003.
 156. Gharakhanlou R, Wesselmann L, Rademacher A, et al. Exercise training and cognitive performance in persons with multiple sclerosis: A systematic review and multilevel meta-analysis of clinical trials. *Mult Scler* 2020;1352458520917935; doi: 10.1177/1352458520917935.
 157. Veldema J, Jansen P. Ergometer Training in Stroke Rehabilitation: Systematic Review and Meta-analysis. *Arch Phys Med Rehabil* 2020;101(4):674–689; doi: 10.1016/j.apmr.2019.09.017.
 158. Schootemeijer S, van der Kolk NM, Bloem BR, et al. Current Perspectives on Aerobic Exercise in People with Parkinson’s Disease. *Neurotherapeutics* 2020;17(4):1418–1433; doi: 10.1007/s13311-020-00904-8.
 159. Fancourt D, Steptoe A. Cultural engagement predicts changes in cognitive function in older adults over a 10 year period: findings from the English Longitudinal Study of Ageing. *Sci Rep* 2018;8(1):10226; doi: 10.1038/s41598-018-28591-8.
 160. Evans IEM, Martyr A, Collins R, et al. Social Isolation and Cognitive Function in Later Life: A Systematic Review and Meta-Analysis. *J Alzheimers Dis* 2019;70(s1):S119–S144; doi: 10.3233/JAD-180501.
 161. Bonnechère B, Klass M, Langley C, et al. Brain training using cognitive apps can improve cognitive performance and processing speed in older adults. *Sci Rep* 2021;11(1):12313; doi: 10.1038/s41598-021-91867-z.