




Technical Note

# Metazoan Parasites of Mozambique tilapia (*Oreochromis mossambicus*) Native to Lake Urema, Mozambique

Willem J. Smit <sup>1,\*</sup>, Maarten P. M. Vanhove <sup>2</sup>, Ngonidzashe A. G. Moyo <sup>3</sup> and Wilmien J. Luus-Powell <sup>1</sup>

<sup>1</sup> DSI-NRF SARChI Chair (Ecosystem Health), Department of Biodiversity, School of Molecular and Life Sciences, University of Limpopo, Sovenga 0727, South Africa

<sup>2</sup> Research Group Zoology: Biodiversity & Toxicology, Centre for Environmental Sciences, Hasselt University, Agoralaan gebouw D, 3590 Diepenbeek, Belgium

<sup>3</sup> Aquaculture Research Unit, University of Limpopo, Sovenga 0727, South Africa

\* Correspondence: willem.smit@ul.ac.za

**Abstract:** Mozambique tilapia *Oreochromis mossambicus* (Peters, 1852), an important aquaculture species investigated herein, is at risk and listed as vulnerable on the IUCN Red List. During a preliminary survey conducted in 2011, 30 Mozambique tilapia from Lake Urema in Gorongosa National Park in central Mozambique, were examined for metazoan parasites. Two groups of ectoparasites were recovered consisting of two species of copepods and a monogenean. Endoparasites retrieved were in the larval stage and included one trematode and one nematode species. This study provides new parasite records for *O. mossambicus* in Mozambique and forms a baseline for monitoring against the presence of ichthyoparasites associated with the introduction in neighbouring systems of non-native fish such as Nile tilapia *Oreochromis niloticus* (Linnaeus, 1758), a fish of worldwide importance in capture fisheries and especially aquaculture.

**Keywords:** freshwater fish parasites; Gorongosa National Park; invasive; *Oreochromis niloticus*

**Key Contribution:** Tilapias are often kept in aquaculture and represent a major source of protein, especially in Africa. In a time of intense aquaculture and worldwide translocation of fishes, baseline data on the biodiversity and occurrence of fish parasites are of importance for aquaculture and conservation of native fauna. To date, only a few studies have reported fish parasites from Mozambique and this is the first record of parasites from Mozambique tilapia in Lake Urema.



**Citation:** Smit, W.J.; Vanhove, M.P.M.; Moyo, N.A.G.; Luus-Powell, W.J. Metazoan Parasites of Mozambique tilapia (*Oreochromis mossambicus*) Native to Lake Urema, Mozambique. *Fishes* **2023**, *8*, 273. <https://doi.org/10.3390/fishes8050273>

Academic Editors: Pierre Sasal and Craig Stockwell

Received: 10 February 2023

Revised: 11 May 2023

Accepted: 15 May 2023

Published: 20 May 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Understanding parasite communities of aquatic systems can have important implications for conservation [1–3]. Here, we report a preliminary study of the metazoan parasites of the Mozambique tilapia *Oreochromis mossambicus* (Peters, 1852) population occurring in Lake Urema in Gorongosa National Park (GNP). This work is important as this fish species is vulnerable in most of its native range, a troublesome invasive in some countries and an important aquaculture species [4]. Indeed, this cichlid has been widely anthropogenically introduced, leading to the co-introduction of its parasites e.g., in Australia [4] and Madagascar [5].


Since the mandate of GNP is to record, restore and conserve the indigenous fauna and flora therein, this study intended to investigate and compile a record of metazoan parasites associated with *O. mossambicus* from Lake Urema and to establish if the resident population of fish is at risk if alien introductions would occur. As fisheries and aquaculture will play an increasingly important role in providing food and nutrition to a growing world population in the future [6], knowledge regarding their parasites and the effect of parasites on their survival is of utmost importance. Indeed, this baseline information will also be valuable by improving the knowledge concerning parasites that can possibly be transferred

between natural systems and aquaculture facilities and potentially save-guarding both these important entities.

## 2. Methods

### 2.1. Site Description

Lake Urema, a shallow reservoir lake with a mean depth of 1.64 m [7], is located within GNP in Sofala Province, central Mozambique [8,9] at 18°52' S and 34°30' E (Figure 1). The lake forms part of the Pungwe River catchment area of 31,150.5 km<sup>2</sup> with 4.7% of the catchment being situated in Zimbabwe and the remainder in Mozambique [10]. Water enters Lake Urema from rivers meandering across the Bárue Midlands, Cheringoma Plateau, and the Rift Valley. Due to the underlying geology, landscape geomorphology, and climatology, the water surface swells from 10 km<sup>2</sup> during the dry season to 200 km<sup>2</sup> in the rainy season with the overflow draining into the Pungwe River, which flows into the Indian Ocean some 80 km away [7,8].



**Figure 1.** The location of Lake Urema in the Gorongosa National Park (GNP), Sofala Province, Mozambique. Inserts illustrate the locality of Mozambique within Africa and the locality of Sofala Province within Mozambique.

## 2.2. Sampling of Fish and Parasite Collection

Fish were captured using a seine net. This study only intended to collect metazoan parasites. Once removed from the net, specimens were visually examined for the presence of mobile ectoparasites. These were removed using a fine brush or a pair of forceps and placed into glass vials filled with lake water for further processing at a field laboratory. Mucus smears, taken by scraping the length of the body and fins with a microscope slide, were examined using a stereomicroscope (Model: Leica EZ4, Leica Microsystems, Wetzlar, Germany). The total length of fish was measured (mm) and recorded. Fish were euthanised and the gills, eyes, and internal organs removed and placed into separate Petri dishes containing distilled water and examined for parasites using a stereomicroscope. The body cavity, body fat, and muscle tissue were thoroughly scrutinised for encysted parasites.

Monogeneans were mounted onto slides in either glycerine ammonium picrate (GAP) or glycerine jelly. All parasitic crustaceans were preserved in 70% ethanol. Endoparasites were fixed and preserved according to methods outlined in Madanire-Moyo [11]. Monogeneans were identified following standard procedures by Douëllou [12] and Pariselle and Euzet [13], digeneans according to Gibson et al. [14] and Chibwana and Nkwengulila [15], nematodes as prescribed by Anderson [16], and copepods as by Oldewage and van As [17] and Robinson and Avenant-Oldewage [18]. Due to time constraints, the organs of specimens not processed in the field were preserved in ethanol to be examined later.

Parasite prevalence (%) with confidence limits, based on the Clopper and Pearson intervals, and mean abundance (MA) and mean intensity (MI) with bootstrap confidence limits ( $BC_a$ ), for each parasite species, were calculated using Quantitative Parasitology 3.0 (QP 3.0: [19]) following Rózsa et al. [20]. Bootstrapping ( $BC_a$ ) for MA and MI was set at 2000 replications.

## 2.3. Water Parameters

The survey was undertaken prior to the onset of the rainy season. Water temperature, dissolved oxygen, pH, conductivity, and salinity were taken near the surface and bottom of the water column using a handheld YSI multiparameter meter (Model: YSI 556 MPS, Yellow Springs, OH, USA).

## 2.4. Ethical Considerations

The protocol was approved following the GNP mandate and guidelines. Fish were euthanised by severing the spinal cord whilst covering the eyes with a damp cloth. Handling and treatment of animals were in accordance with the guidelines of the South African Council on Animal Care [21].


## 3. Results

Thirty individuals of *O. mossambicus*, having a mean ( $\pm$ standard deviation) total length of 287.6 ( $\pm$ 20.8) mm, were captured, from which five parasite taxa were recorded.

### 3.1. Parasites Retrieved

Ectoparasites recovered included copepods and monogeneans. The monogenean *Cichlidogyrus halli* (Price & Kirk, 1967) (Figure 2), was recovered from the gills with a non-specific distribution. Copepods recorded were *Ergasilus mirabilis* Oldewage & van As, 1987 (Figure 3) recovered from the gills and *Lernaea cyprinacea* Linnaeus, 1758 (Figure 4), from the skin. Specimens of *E. mirabilis* collected were mainly attached close to the gill arch near the base of the filaments, with a small number occurring in close proximity to the distal end of the filaments. For most, tissue damage was noted in the attachment area. *Lernaea cyprinacea* occurred mainly on the ventral and lateral regions with visible tissue damage leading to secondary infections. Two groups of larval endoparasites were recorded, i.e., metacercariae of the trematode *Euclinostomum* Travassos, 1928 (Figure 5A) embedded in muscle tissue and the third stage larvae of the nematode *Contracaecum* Railliet & Henry, 1912 (Figure 5B) from the body cavity in the proximity of the liver.






**Figure 2.** *Cichlidogyrus halli*; (A) = haptor parts; (B) = male copulatory organ (MCO).




**Figure 3.** *Ergasilus mirabilis*; (A) = attached to the gill filament; (B) = second antenna (smooth with single terminal tip).





**Figure 4.** *Lernaean cyprinacea*; (A) = Embedded in the skin (note signs of inflammation); (B) = removed from the fish and still attached to the ventral side of the scale.



**Figure 5.** Larval endoparasites; (A) = *Euclinostomum* sp.; (B) = *Contraecaecum* sp.

The parasite community was dominated by ectoparasites whereby all hosts were infected with *E. mirabilis* and *C. halli* and 90.0% by *L. cyprinacea* (Table 1). The prevalence, MA, and MI of ectoparasites were notably higher than those of endoparasites (Table 1). Although not all the fish examined were infected with *L. cyprinacea*, this parasite had the highest recorded MA and MI, ranging from 2–24 parasites per infected host (Table 1). Fish with no *L. cyprinacea* attached displayed lesions from previous infestations by this parasite, indicating that all fish inspected were infected at some point.

**Table 1.** Prevalence (P), mean abundance (MA), mean intensity (MI), and intensity range (IR) of parasites from 30 *Oreochromis mossambicus* sampled from Lake Urema during a survey undertaken in October 2011. The upper and lower confidence levels (CL) for P, MA, and MI are given in parentheses.

Parasite Group	Parasite Species	Infected Site	P (%)	95% CL for P (%)	MA	95% CL for MA	MI	95% CL for MI	IR
Monogenea	<i>Cichlidogyrus halli</i>	Gills	100.0	(88.4–100.0)	5.9	(4.6–7.1)	5.9	(4.7–7.1)	2–15
Trematoda	<i>Euclinostomum</i> sp. *	Muscle	16.7	(3.8–30.7)	0.2	(0.0–0.3)	1.0	(NC)	1
Nematoda	<i>Contracaecum</i> sp. *	Body cavity	60.0	(40.6–77.3)	0.9	(0.6–1.2)	1.5	(1.2–1.8)	1–3
Copepoda	<i>Ergasilus mirabilis</i>	Gills	100.0	(88.4–100.0)	8.0	(6.0–10.7)	8.0	(6.0–10.4)	2–28
	<i>Lernaea cyprinacea</i>	Skin	90.0	(69.3–96.2)	10.4	(7.3–12.5)	11.6	(8.7–14.0)	2–24

\* larval forms. (NC) = Not calculated due to intensity being constant.

### 3.2. Water Parameters

The mean ( $\pm$ standard deviation) recorded for water temperature, dissolved oxygen, conductivity, and salinity was 27.45 ( $\pm$ 0.16) °C, 3.21 ( $\pm$ 0.04) mg/L, 0.10 ( $\pm$ 0.00) mS·cm<sup>-1</sup> and 0.03 ( $\pm$ 0.00) ppt, respectively. The pH measured was 6.50 throughout the water column of ~1 m.

## 4. Discussion

### 4.1. Lake Condition and Parasite Burden

All water measurements were consistent with pre-flood conditions [7]. Factors from the external environment such as water temperature and water quality may affect the parasite itself or the intermediate hosts that are needed to complete the life cycle. The environmental conditions determine the general character of the parasite fauna as a whole and the absence, presence, and abundance of parasites depends thus on the availability of a suitable environment for them. The water temperature was high which is normally favorable for invertebrate intermediate hosts as well as for most ectoparasite groups such as monogeneans [22,23] and copepods [24]. The effects of seasonal changes in water quality and levels, in combination with fish size, fish gender, and the region within the lake e.g., inlet vs. outlet, on parasite burden of *O. mossambicus* and other fish species require investigation.

### 4.2. Known Parasite Records and Distribution

It is important to gain knowledge of the parasite composition of fish species in their native environment, to attain baseline information for when invasions may occur in the future. As this lake is within a protected area, knowledge gained here will add to the continual need to conserve and protect the biodiversity in Lake Urema. A study by Firmat et al. [25] is the only record of parasites from *O. mossambicus* in Mozambique, however totally isolated from Lake Urema and approximately 700 km apart. During the above-mentioned study they recorded five species of monogenean gill parasites from *O. mossambicus*. *Gyrodactylus* spp. are reported from the skin of this fish from South Africa [26,27] and invasive *O. mossambicus* in the Philippines [28]. Besides gill monogeneans, three species of endoparasitic monogeneans of *Enterogyrus* [11,28] were recorded for *O. mossambicus* in South Africa. More monogenean species were thus expected; however, only *C. halli* was recorded in this study. A potential loss of monogenean species has been proposed to result from the host population going through a genetic bottleneck event [29]. This could have been the case if Lake Urema was originally colonised by only a small number of Mozambique tilapias. Another explanation can be a temporal variation in infection due to an immune response by fish [30]. *Cichlidogyrus halli* has been reported from *O. mossambicus* in Mozambique [25], South Africa [28,31,32], cultured *O. mossambicus* in Japan [33], as well as various cichlid hosts, including introduced populations belonging to various tilapia species, from other African countries [5,13,34,35]. With all fish in this study infected with *C. halli*, the prevalence was notably higher than reported in the aforementioned studies, besides the study by

Firmat et al. [25] where a 100% prevalence was recorded at Linlangalinwe in the Changane River (lower Limpopo Drainage System, Mozambique). The MI levels for *C. halli* reported here were comparable to the findings by Madanire-Moyo et al. [31] and Maneepitaksanti and Nagasawa [33] for the same parasite and host species.

The larval stages of endoparasites collected in this study could not be identified to the species level since the reproductive organs, used to characterise the worms, were not fully developed. The occurrence of larval stages of *Euclinostomum* spp. and *Contracaecum* spp. has been widely reported in cichlids from southern Africa by Khalil and Polling [34], Scholz et al. [35] and Tavakol et al. [36]. The MI for *Contracaecum* sp. recorded here corresponds well with findings by Tavakol et al. [36] for *O. mossambicus* sampled from a number of impoundments in South Africa. The prevalence of *Contracaecum* sp. in this study was however notably higher than that reported by Tavakol et al. [36] but similar to that reported by Sara et al. [30] for the same fish species. *Contracaecum* sp., with a prevalence of 97.5%, was retrieved from *C. gariepinus* in Lake Urema by Dumbo [37]. *Cyprinus carpio* Linnaeus, 1758 was indicated by Boane et al. [38] to have a prevalence of 22.0% and 17.4% of *Contracaecum* sp. infestation from Lagoon Chuáli and the Limpopo River, Mozambique, respectively. The current study is the first to record *Contracaecum* sp. from *O. mossambicus* in Mozambique. The prevalence and MI of *Euclinostomum* sp. recorded reflect well with results by Britz et al. [39] and Olivier et al. [28] for the same species of fish in South Africa. This observation is the first record of this parasite from Mozambique.

Fifteen species of *Ergasilus* have been recorded throughout Africa [35,40–42], however, no published records of *Ergasilus* spp. from *O. mossambicus* exist, making this study a new host record for this parasite genus. Other *Ergasilus* spp. have, however, been recorded from *Oreochromis* spp. as well as other cichlids. *Ergasilus mirabilis* with a prevalence of 7.3% was reported for *C. gariepinus* from Lake Urema by Dumbo [37] and a MI far greater than for *O. mossambicus* recorded here. A range of host fishes throughout southern Africa, including some squeakers [41] and parrotfish [43], is reported to harbour this copepod.

*Lernaea cyprinacea* had an aggregated distribution on the body of the fish. This parasite, commonly known as anchor worm, is an alien species to Africa and an opportunistic host generalist in that it infests various families of fish and amphibians [44–46]. It is a cosmopolitan parasite that is widely distributed due to the translocation of edible and ornamental cyprinids [46–49]. In southern Africa, *L. cyprinacea* has been reported from *O. mossambicus* in Lake Victoria [18] and a number of water bodies in Zimbabwe [50,51] and South Africa [18,52–54]. Although this is the first record of this parasite from *O. mossambicus* in Mozambique, it has been reported for *C. carpio* in this country by Boane et al. [38]. Another lernaeid, *Lernaea barnimiana* (Hartmann, 1865) is also known to occur in the southern African region and has been reported from *O. mossambicus* [35]. High infestation levels of copepods are expected in Lake Urema given that this group is known to thrive in shallow waters at high temperatures [50,55–57]. The prevalence and MI of anchor worms reported from South Africa are lower [18] whereas those in Zimbabwe [50] (associated with shallow and warm water) are higher than in the current study. As with studies by Dezfuli et al. [58], Lester and Hayward [59], and Vinobaba [60], tissue damage that can lead to secondary infections, was observed at the attachment sites for both copepod species reported here. Infection by anchor worms has been associated with poor general health [54,61,62] and thus poses a great threat to native fish in its non-native range as well as to aquaculture practices.

All parasite species recorded here may be transferred to and become established within aquaculture facilities and have a negative impact on aquaculture fish stocks and therefore the economic viability of this sector [6,63]. Parasitic infestation has been associated with a reduction in host weight, growth, fecundity, and in severe cases mortality [5,62,64,65]. Knowledge regarding parasitic threats and potential stressors for fish may reduce economic strain on a volatile sector making it more sustainable in the future.



### 4.3. Zoonotic Importance

The fish-borne zoonotic trematodes and nematodes reported here may pose a health risk as they can be transmitted to humans consuming raw or undercooked fish [66–70]. Dried salted fish is a common product in rural areas and is being offered by local traders [71]. This is of special concern as fishermen from nearby Muaredzi, some 30 km from the lake, have access to catch and supply the village inhabitants with fresh fish. There is also small-scale extensive aquaculture production taking place within Sofala Province where farmers are stocking from natural systems, posing a risk of parasite transfer. Therefore, appropriate measures of control should be established and implemented. Park authorities and educators working at surrounding rural schools should attempt to make the general public and fish farmers aware of the zoonotic nature of some fish parasites and the risks they may pose to human health.

**Author Contributions:** W.J.S.: collected samples; conceptualisation; analysed the data; writing—original draft preparation. W.J.L.-P. and N.A.G.M.: writing—review and editing; funding acquisition; resources. M.P.M.V.: writing—review and editing. All authors have read and agreed to the published version of the manuscript.

**Funding:** This project would not have been possible without the financial assistance from the Gregory C Carr Foundation for fieldwork and in part by the Department of Science and Innovation and the National Research Foundation of South Africa (Grant Number 101054). MPMV is financed by the Special Research Fund of Hasselt University (BOF20TT06).

**Institutional Review Board Statement:** At the time when the study was undertaken neither the University of Limpopo (UL) nor the Gorongosa National Park administration had an ethics committee. The researchers are currently working under UL ethics approval (AREC/01/2020:IR) issued on 14 February 2020. The Animal Research Ethics Committee (AREC) is registered with the National Health Research Ethics Council, Registration Number: AREC-290914-017.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author. The data are not publicly available due to being intellectual property.

**Acknowledgments:** The support from Gorongosa National Park field rangers, LP Olivera, T Castigo; park ecologist A Short and Park Director F Steinbruch for the successful execution of this survey is appreciated. University of Limpopo staff members, J Theron and G Geldenhuys, are also thanked for their technical support and assistance in the field.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Gilbert, B.M.; Avenant-Oldewage, A. Parasites and pollution: The effectiveness of tiny organisms in assessing the quality of aquatic ecosystems, with a focus on Africa. *Environ. Sci. Pollut. Res.* **2017**, *24*, 18742–18769. [[CrossRef](#)] [[PubMed](#)]
2. Gagne, R.B.; Crooks, K.R.; Craft, M.E.; Chiu, E.S.; Fountain-Jones, N.M.; Malmberg, J.L.; Carver, S.; Funk, W.C.; VandeWoude, S. Parasites as conservation tools. *Conserv. Biol.* **2022**, *36*, e13719. [[CrossRef](#)]
3. Marick, J.; Patra, B.K.; Ash, A. Loss of Biodiversity and Ecosystem Services: Told and Untold Stories from Parasite World. *Proc. Zool. Soc.* **2023**. [[CrossRef](#)]
4. Wilson, J.R.; Saunders, R.J.; Hutson, K.S. Parasites of the invasive tilapia *Oreochromis mossambicus*: Evidence for co-introduction. *Aquat. Invasions* **2019**, *14*, 332–349. [[CrossRef](#)]
5. Šimková, A.; Řehulková, E.; Rasoloariniaina, J.R.; Jorissen, M.W.P.; Scholz, T.; Faltýnková, A.; Mašová, Š.; Vanhove, M.P. Transmission of parasites from introduced tilapias: A new threat to endemic Malagasy ichthyofauna. *Biol. Invasions* **2019**, *21*, 803–819. [[CrossRef](#)]
6. Food and Agriculture Organization of the United Nations. *The State of World Fisheries and Aquaculture. Towards Blue Transformation*; FAO: Rome, Italy, 2022.
7. Böhme, B. Geo-ecology of the Lake Urema/Central Moçambique. *Freib. Online Geol.* **2005**, *14*, 1–114.
8. Tinley, K.L. Framework of the Gorongosa Ecosystem. Ph.D. Thesis, University of Pretoria, Pretoria, South Africa, 1977.
9. Arvidsson, K.; Stenberg, L.; Chirindja, F.; Dahlin, T.; Owen, R.; Steinbruch, F. A hydrogeological study of the Nhandugue River, Moçambique—A major groundwater recharge zone. *Phys. Chem. Earth* **2011**, *36*, 789–797. [[CrossRef](#)]



10. SWECO and Associates I. *Development of the Pungwe River Basin. Joint Integrated Water Resources Management Strategy*. Monograph Report. Annex I. Sector Study on: Surface Water Resources of the Pungwe River Basin. Client: Government of the Republic of Mozambique, Government of the Republic of Zimbabwe; Swedish International Development Cooperation Agency (Sida). 2004. Available online: [https://biblioteca.biofund.org.mz/biblioteca\\_virtual/development-of-the-pungwe-river-basin-joint-integrated-water-resources-management-strategy-annex-x-sector-study-on-conservation-areas-wildlife-and-tourism/](https://biblioteca.biofund.org.mz/biblioteca_virtual/development-of-the-pungwe-river-basin-joint-integrated-water-resources-management-strategy-annex-x-sector-study-on-conservation-areas-wildlife-and-tourism/) (accessed on 10 May 2023).
11. Madanire-Moyo, G.N.; Luus-Powell, W.J.; Olivier, P.A. Diversity of metazoan parasites of the Mozambique tilapia; *Oreochromis mossambicus* (Peters, 1852), as indicators of pollution in the Limpopo and Olifants River Systems. *Onderstepoort J. Vet. Res.* **2012**, *79*, 146–152. [CrossRef]
12. Douëllou, L. Monogeneans of the genus *Cichlidogyrus* Paperna, 1960 (Dactylogyridae: Ancyrocephalinae) from cichlid fishes of Lake Kariba (Zimbabwe) with descriptions of five new species. *Syst. Parasitol.* **1993**, *25*, 159–186. [CrossRef]
13. Pariselle, A.; Euzet, L. Systematic revision of dactylogyridean parasites (Monogenea) from cichlid fishes in Africa; the Levant and Madagascar. *Zoosystema* **2009**, *31*, 849–898. [CrossRef]
14. Gibson, D.I.; Bray, R.A.; Jones, A. *Keys to the Trematoda*; Oxford University Press: Oxford, UK, 2008; Volume 3.
15. Chibwana, F.D.; Nkwengulila, G. Variation in the morphometrics of diplostomid metacercariae (Digenea: Trematoda) infecting the catfish, *Clarias gariepinus* in Tanzania. *J. Helminthol.* **2010**, *84*, 61–70. [CrossRef] [PubMed]
16. Anderson, R.C. *Nematode Parasites of Vertebrates: Their Development and Transmission*; CAB International: Wallingford, UK, 1992.
17. Oldewage, W.H.; van As, J.G. A key for the identification of African piscine parasitic Ergasilidae (Copepoda: Poecilostomatoida). *Suid-Afrik. Tydskr. Vir Dierkd.* **1988**, *23*, 42–46.
18. Robinson, J.; Avenant-Oldewage, A. Aspects of the morphology of the parasitic copepod *Lernaea cyprinacea* Linnaeus, 1758 and notes on its distribution in Africa. *Crustaceana* **1996**, *69*, 610–626.
19. Reiczigel, J.; Rózsa, L. *Quantitative Parasitology 3.0*. Budapest, Distributed by the Authors. 2005. Available online: <https://www.zoologia.hu/qp/> (accessed on 25 October 2022).
20. Rózsa, L.; Reiczigel, J.; Majoros, G. Quantifying parasites in samples of hosts. *J. Parasitol.* **2000**, *86*, 228–232. [CrossRef]
21. *South African National Standard 10386; The Care and Use of Animals for Scientific Purposes*. SABS Standards Division: Groenkloof, South Africa, 2008.
22. Koskivaara, M.; Valtonen, E.T.; Vuori, K.M. Microhabitat distribution and coexistence of *Dactylogyrus* species (Monogenea) on the gills of roach. *Parasitology* **1992**, *104*, 273–281. [CrossRef]
23. Villar-Torres, M.; Montero, F.E.; Raga, J.A.; Repullés-Albelda, A. Come rain or come shine: Environmental effects on the infective stages of *Sparicotyle chrysophrii*, a key pathogen in Mediterranean aquaculture. *Parasites Vectors* **2018**, *11*, 558. [CrossRef]
24. Cavaleiro, F.I.; Santos, M.J. Seasonality of metazoan ectoparasites in marine European flounder *Platichthys flesus* (Teleostei: Pleuronectidae). *Parasitology* **2009**, *136*, 855–865. [CrossRef] [PubMed]
25. Firmat, C.; Alibert, P.; Mutin, G.; Losseau, M.; Pariselle, A.; Sasal, P. A case of complete loss of gill parasites in the invasive cichlid *Oreochromis mossambicus*. *Parasitol. Res.* **2016**, *115*, 3657–3661. [CrossRef]
26. García-Vásquez, A.; Hansen, H.; Christison, K.W.; RubioGodoy, M.; Bron, J.E.; Shinn, A.P. Gyrodactylids (Gyrodactylidae; Monogenea) infecting *Oreochromis niloticus niloticus* (L.) and *O. mossambicus* (Peters) (Cichlidae): A pan-global survey. *Acta Parasitol.* **2010**, *55*, 215–229. [CrossRef]
27. Zahradníčková, P.; Barson, M.; Luus-Powell, W.J.; Prikrylová, I. Species of *Gyrodactylus* von Nordmann, 1832 (Platyhelminthes: Monogenea) from cichlids from Zambezi and Limpopo river basins in Zimbabwe and South Africa: Evidence for unexplored species richness. *Syst. Parasitol.* **2016**, *93*, 679–680. [CrossRef]
28. Olivier, P.A.; Luus-Powell, W.J.; Saayman, J.E. Report on some monogenean and clinostomid infestations of freshwater fish and waterbird hosts in Middle Letaba Dam, Limpopo Province, South Africa. *Onderstepoort J. Vet. Res.* **2009**, *76*, 187–199. [CrossRef] [PubMed]
29. Pariselle, A.; Morand, S.; Deveney, M.; Pouyau, L. Parasite species richness of closely related hosts: Historical scenario and “genetic” hypothesis. In *Hommage à Louis Euzet—Taxonomie; Écologie et Évolution des Métazoaires Parasites. Taxonomy; Ecology and Evolution of Metazoan Parasites*; Combes, C., Jourdan, J., Eds.; Presses Universitaires de Perpignan: Perpignan, France, 2003; pp. 147–166.
30. Ilgová, J.; Salát, J.; Kašný, M. Molecular communication between the monogenea and fish immune system. *Fish Shellfish. Immunol.* **2021**, *112*, 179–190. [CrossRef]
31. Madanire-Moyo, G.N.; Matla, M.M.; Olivier, P.A.; Luus-Powell, W.J. Population dynamics and spatial distribution of Monogeneans on the gills of *Oreochromis mossambicus* (Peters, 1852) from two lakes of the Limpopo River System, South Africa. *J. Helminthol.* **2011**, *85*, 146–152. [CrossRef]
32. Sara, J.R.; Smit, W.J.; Erasmus, L.J.; Ramalepe, T.P.; Mogashoa, M.E.; Raphahlelo, M.E.; Theron, J.; Luus-Powell, W.J. Ecological status of Hout River Dam; Limpopo province, South Africa, using fish condition and health assessment index protocols: A preliminary investigation. *Afr. J. Aquat. Sci.* **2014**, *39*, 35–43. [CrossRef]
33. Maneepitaksanti, W.; Nagasawa, K. Monogeneans of *Cichlidogyrus* Paperna, 1960 (Dactylogyridae); gill parasites of tilapias; from Okinawa Prefecture, Japan. *Biogeography* **2012**, *14*, 111–119.
34. Khalil, L.F.; Polling, L. *Checklist of the Helminth Parasites of African Freshwater Fishes*; University of the North: Pietersburg, South Africa, 1997.

35. Scholz, T.; Vanhove, M.P.; Smit, N.; Jayasundera, Z.; Gelnar, M. A Guide to the Parasites of African Freshwater Fishes. *Abc Taxa* **2018**, *18*, 1–425.
36. Tavakol, S.; Smit, W.J.; Sara, J.R.; Halajian, A.; Luus-Powell, W.J. Distribution of *Contracaecum* (Nematoda: Anisakidae) larvae in freshwater fish from the northern regions of South Africa. *Afr. Zool.* **2015**, *50*, 133–139. [[CrossRef](#)]
37. Dumbo, J.C. Estudo de Parasitas Metazoários de *Clarias gariepinus* (Burchell; 1822) que ocorre no Lago Urema, Parque Nacional da Gorongosa. Master's Thesis, Universidade Eduardo Mondlane, Maputo, Mozambique, 2009.
38. Boane, C.; Cruze, C.; Saraiva, A. Metazoan Parasites of *Cyprinus carpio* L. (Cyprinidae) from Mozambique. *Aquaculture* **2008**, *284*, 59–61. [[CrossRef](#)]
39. Britz, J.; van As, J.G.; Saayman, J.E. Occurrence and distribution of *Clinostomum tilapiae* Ukoli, 1966 and *Euclinostomum heterostomum* (Rudolphi; 1809) metacercarial infections of freshwater fish in Venda and Lebowa, southern Africa. *J. Fish Biol.* **1985**, *26*, 21–28. [[CrossRef](#)]
40. Oldewage, W.H.; van As, J.G. A new fish-ectoparasitic ergasilid (Crustacea: Copepoda) from the Pongola River system. *S. Afr. J. Zool.* **1987**, *22*, 62–65. [[CrossRef](#)]
41. Oldewage, W.H.; Avenant-Oldewage, A. Checklist of the piscine parasitic Copepoda (Crustacea) of African fishes. *Zool. Meded.* **1993**, *23*, 2–28.
42. Abdel-Hady, O.K.; Bayoumy, E.M.; Osman, H.A.M. New copepodal ergasilid parasitic on *Tilapia zilli* from Lake Tamsah with special reference to its pathological effect. *Glob. Vet.* **2008**, *2*, 123–129.
43. Douëllou, L.; Erlwanger, K.H. Crustacean parasites of fishes in Lake Kariba, Zimbabwe, preliminary results. *Hidrobiologica* **1994**, *287*, 233–242. [[CrossRef](#)]
44. Yashuv, V.A. On the biology of *Lernaea cyprinacea* in fish ponds. *Bull. Fish Cult.* **1959**, *11*, 80–89.
45. Paperna, I. *Parasites; Infections and Diseases of Fishes in Africa: An Update*; FAO/CIFA Technical Paper No. 31; FAO Publications: Rome, Italy, 1996.
46. Piasecki, W.; Goodwin, A.E.; Eiras, J.C.; Nowak, B.F. Importance of Copepoda in freshwater aquaculture. *Zool. Stud.* **2004**, *43*, 193–205.
47. Hoffman, G.L. Intercontinental and transcontinental dissemination and transfaunation of fish parasites with emphasis on whirling disease (*Myxosoma cerebralis*). In *A Symposium on Diseases of Fish and Shellfish*; Snieszko, S.F., Ed.; American Fisheries Society: Bethesda, MD, USA, 1970; Volume 5, pp. 69–81.
48. Marcogliese, D.J. Seasonal occurrence of *Lernaea cyprinacea* on fishes in Belews Lake, North Carolina. *J. Parasitol.* **1991**, *77*, 326–327. [[CrossRef](#)]
49. Medeiros, E.S.; Maltchik, L. The effects of hydrological disturbance on the intensity of infestation of *Lernaea cyprinacea* in an intermittent stream fish community. *J. Arid. Environ.* **1999**, *43*, 351–356. [[CrossRef](#)]
50. Barson, M.; Mulonga, A.; Nhwatiwa, T. Investigation of a parasitic outbreak of *Lernaea cyprinacea* Linnaeus (Crustacea: Copepoda) in fish from Zimbabwe. *J. Afr. Zool.* **2008**, *43*, 175–183. [[CrossRef](#)]
51. Dalu, T.; Nhwatiwa, T.; Clegg, B.; Barson, M. Impact of *Lernaea cyprinacea* Linnaeus 1758 (Crustacea: Copepoda) almost a decade after an initial parasitic outbreak in fish of Malilangwe Reservoir, Zimbabwe. *Knowl. Manag. Aquat. Ecosyst.* **2012**, *406*, 1–9.
52. Van As, J.G.; Basson, L. Checklist of freshwater parasites from southern Africa. *S. Afr. J. Wildl.* **1984**, *14*, 49–61.
53. Viljoen, B.C.S. A seasonal investigation of the genus *Lernaea* (Crustacea: Copepoda) on cyprinid fish in Boskop Dam; Transvaal; South Africa. *S. Afr. J. Wildl.* **1986**, *16*, 27–31.
54. Welicky, R.L.; de Swart, J.; Gerber, R.; Netherlands, E.; Smit, N.J. Drought associated absence of alien invasive anchorworm; *Lernaea cyprinacea* (Copepoda: Lernaecidae); is related to changes in fish health. *Int. J. Parasitol. Parasites Wildl.* **2017**, *6*, 430–438. [[CrossRef](#)] [[PubMed](#)]
55. Lahav, M.; Sarig, S.; Shilo, M. The eradication of *Lernaea* in storage ponds of carps through destruction of the copepodid stage by *Dipterex*. *Bamidgeh* **1964**, *27*, 341–397.
56. Bulow, F.J.; Winningham, J.R.; Hooper, R.C. Occurrence of the copepod parasite *Lernaea cyprinacea* in a stream fish population. *Trans. Am. Fish. Soc.* **1979**, *108*, 100–102. [[CrossRef](#)]
57. Hoffman, G.L. Copepod parasites of freshwater fish: Ergasilus, Achtheres and Salmincola. *Fish Dis. Leaflet.* **1977**, *48*, 1–10.
58. Dezfuli, B.S.; Glari, L.; Konecny, R.; Jaeger, P.; Manera, M. Immunohistochemistry, ultrastructure and pathology of gills of *Abramis brama* from Lake Mondsee, Austria, in infected with *Ergasilus sieboldi* (Copepoda). *Dis. Aquat. Org.* **2003**, *53*, 257–262. [[CrossRef](#)] [[PubMed](#)]
59. Lester, R.G.; Hayward, C.J. Phylum arthropoda. In *Fish Diseases and Disorders: Protozoan and Metazoan Infections*; Woo, P.T.K., Ed.; CAB International: Wallingford, UK, 2006; Volume 1, pp. 466–565. [[CrossRef](#)]
60. Vinobaba, P. Histopathological changes induced by ergasilid copepod infections on the gills of food fish from Batticaloa lagoon, Sri Lanka. *Sri Lanka J. Aquat. Sci.* **2007**, *12*, 77–87. [[CrossRef](#)]
61. Kabata, Z. Copepoda (Crustacea) parasitic on fishes: Problems and perspectives. *Adv. Parasitol.* **1982**, *19*, 1–71.
62. Khan, M.N.; Aziz, F.; Afzal, M.; Rab, A.; Sahar, L.; Ali, R.; Naqvi, S.M. Parasitic infestation in different fresh water fishes of mini dams of Potohar region, Pakistan. *J. Sci. Technol. (Pak.)* **2003**, *6*, 1092–1095. [[CrossRef](#)]
63. Timi, J.T.; Buchmann, K. A century of parasitology in fisheries and aquaculture. *J. Helminthol.* **2023**, *97*, e4. [[CrossRef](#)]
64. Van De, N.; Waikagul, J.; Dalsgaard, A.; Chai, J.Y.; Sohn, W.M.; Murrell, K.D. Fishborne zoonotic intestinal trematodes, Vietnam. *Emerg. Infect. Dis.* **2007**, *13*, 1828–1833.

65. Vollset, K.W. Parasite induced mortality is context dependent in Atlantic salmon: Insights from an individual based model. *Sci. Rep.* **2019**, *9*, 17377. [[CrossRef](#)] [[PubMed](#)]
66. Cribb, T.H.; Chisholm, L.A.; Bray, R.A. Diversity in the Monogenea and Digenea: Does lifestyle matter? *Int. J. Parasitol.* **2002**, *32*, 321–328.93. [[CrossRef](#)]
67. Kaddumukasa, M.; Kaddu, J.B.; Makanga, B. A Survey of nematode infection in *Oreochromis niloticus* (L.) (Teleostei: Cichlidae) in Lake Kyoga, Uganda. *Afr. J. Trop. Hydrobiol.* **2013**, *13*, 12–13.
68. Bekele, J.; Hussien, D. Prevalence of internal parasites of *Oreochromis niloticus* and *Clarias gariepinus* fish species in Lake Ziway, Ethiopia. *J. Aquac. Res.* **2015**, *6*, 308.
69. Reshid, M.; Adugna, M.; Redda, Y.T.; Awol, N.; Teklu, A. A Study of *Clinostomum* (Trematode) and *Contraecaecum* (Nematode) parasites affecting *Oreochromis niloticus* in Small Abaya Lake, Silite Zone, Ethiopia. *J. Aquac. Res. Dev.* **2015**, *6*, 316.
70. Thu, N.D.; Dalsgaard, A.; Loan, L.T.; Murrell, K.D. Survey for zoonotic liver and intestinal trematode metacercariae in cultured and wild fish in an Giang Province; Vietnam. *Korean J. Parasitol.* **2007**, *45*, 45–54. [[CrossRef](#)]
71. Urban-Econ Development Economists. *Nile & Mozambique Tilapia Feasibility Study*; Department of Agriculture, Forestry and Fisheries Chief Directorate: Vlaeberg, South Africa; Aquaculture and Economic Development: Vlaeberg, South Africa, 2018.

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.