

Pesticide concentration in three selected fish species and human health risk in the Lake Tana sub-basin, Ethiopia

Peer-reviewed author version

Zelalem, Wondie; Anteneh, Wassie; Mingist, Minwyelet; Kibret, Mulugeta; Adgo, Enyew; Erarto, Felegush; De Rop, Jasmine; De Cock, Andree; Spanoghe, Pieter; Goethals, Peter L. M.; Nyssen, Jan; Verleyen, Elie & DE VOCHT, Alain (2023) Pesticide concentration in three selected fish species and human health risk in the Lake Tana sub-basin, Ethiopia. In: ENVIRONMENTAL MONITORING AND ASSESSMENT, 195 (8) (Art N° 988).

DOI: 10.1007/s10661-023-11594-y

Handle: <http://hdl.handle.net/1942/41688>

Pesticide concentration in three selected fish species and human health risk in the Lake Tana sub-basin, Ethiopia

This Accepted Manuscript (AM) is a PDF file of the manuscript accepted for publication after peer review, when applicable, but does not reflect post-acceptance improvements, or any corrections. Use of this AM is subject to the publisher's embargo period and AM terms of use. Under no circumstances may this AM be shared or distributed under a Creative Commons or other form of open access license, nor may it be reformatted or enhanced, whether by the Author or third parties. By using this AM (for example, by accessing or downloading) you agree to abide by Springer Nature's terms of use for AM versions of subscription articles: <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>

The Version of Record (VOR) of this article, as published and maintained by the publisher, is available online at: <https://doi.org/10.1007/s10661-023-11594-y>. The VOR is the version of the article after copy-editing and typesetting, and connected to open research data, open protocols, and open code where available. Any supplementary information can be found on the journal website, connected to the VOR.

For research integrity purposes it is best practice to cite the published Version of Record (VOR), where available (for example, see ICMJE's guidelines on overlapping publications). Where users do not have access to the VOR, any citation must clearly indicate that the reference is to an Accepted Manuscript (AM) version.

1 **Pesticide concentration in three selected fish species and human health risk in the Lake Tana Sub-basin, Ethiopia**

2 Wondie Zelalem^{1,2,*}, Wassie Anteneh³, Minwyelet Mingist², Mulugeta Kibret⁴, Enyew Adgo⁵, Felegush Erarto², Jasmine
3 De Rop⁶, Andrée De Cock⁷, Pieter Spanoghe⁶, Peter L.M Goethals⁷, Jan Nyssen⁸, Elie Verleyen⁹, Alain De Vocht^{1,10}

4 ¹ UHasselt, Centre for Environmental Sciences, Agoralaan, 3590 Diepenbeek, Belgium

5 ² Department of Fisheries and Aquatic Sciences, Bahir Dar University, Bahir Dar, Ethiopia

6 ³ Intergovernmental Authority on Development (IGAD), Djibouti

7 ⁴ Department of Biology, Bahir Dar University, Bahir Dar, Ethiopia

8 ⁵ Department of Natural Resource Management, Bahir Dar University, Bahir Dar, Ethiopia

9 ⁶ Department of Plants and Crops, Ghent University, Ghent, Belgium

10 ⁷ Department of Animal Sciences and Aquatic Ecology, Ghent University, Ghent, Belgium

11 ⁸ Department of Geography, Ghent University, Ghent, Belgium

12 ⁹ Department of Biology, Ghent University, Ghent, Belgium

13 ¹⁰ PXL University College, Hasselt, Belgium

*Correspondence author; Email: wondiehm@gmail.com, ORCID ID: [0000-0001-5322-7100](https://orcid.org/0000-0001-5322-7100)

14 **Abstract**

15 Pesticide use has increased in the Lake Tana sub-basin due to increased agricultural activity, potentially endangering
16 nontargeted organisms. To assess its potential impact on fish health and fish-consuming human populations, pesticide
17 concentrations in the fillet and liver tissue of three fish species, namely *Labeobarbus megastoma*, *L. tsanensis*, and
18 *Oreochromis niloticus*, were investigated in Lake Tana. Fish samples were taken from the lake near the rivers of Ribb and
19 Gumara, which flow through agricultural areas where considerable amounts of pesticides have been applied. A total of 96
20 fish samples were collected. Liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS) revealed the
21 presence of ten pesticides. Pyrimethanil was frequently detected in 96 % of liver and 65 % of fillet samples at a median
22 concentration of 33.9 µg kg⁻¹ and 19.7 µg kg⁻¹, respectively. The highest concentration of pyrimethanil was found in *L.*
23 *megastoma* (1850.0 µg kg⁻¹). *Labeobarbus megastoma* also had the highest concentration of oxamyl (507.0 µg kg⁻¹) and
24 flazasulfuron (60.1 µg kg⁻¹) detected in the liver tissue. The highest concentration of carbaryl (56.5 µg kg⁻¹) was found in
25 the liver tissue of *O. niloticus*. Fish tissue samples from the two study sites contained pyrimethanil, oxamyl, carbaryl, and
26 flazasulfuron. Only pyrimethanil showed a statistically significant difference between the two sites and the species *L.*
27 *megastoma* and *L. tsanensis*. The amounts of pesticides found in the fish species pose no direct risk to the health of fish
28 consumers human population. However, the results show that the lake ecosystem needs immediate attention and regular
29 monitoring of the rising pesticide usage in the lake watershed.

30 **Keywords:** Lake Tana, pesticide, *Labeobarbus*, *Oreochromis*, health risk, toxicity

31 Introduction

32 Pesticides have been used in agriculture for a long time to protect crops from weeds, pests, and diseases while increasing
33 crop yields (Lengai et al., 2020; Sabzevari & Hofman, 2022). They have also been used to reduce crop deterioration during
34 storage and extend shelf life, ensuring food security for the world's ever-growing population (Sharma et al., 2019). About
35 4.2 million tons of pesticides, including herbicides, insecticides, and fungicides, are utilized worldwide annually (FAO,
36 2021). In developing countries such as Ethiopia, the government has encouraged farmers to use pesticides and fertilizers
37 in smallholder farming (WHO & FAO, 2019). However, because farmers and agricultural workers frequently lack proper
38 personal protection equipment and may not understand the labels with safety instructions, human health and environmental
39 hazards associated with pesticide use are commonly noticed in those nations (Sarkar et al., 2021).

40 Although pesticides improve crop yield, their extensive use produces residual effects on food products. Moreover, they
41 could develop problems associated with chemical buildup in aquatic organisms, including fish, and negatively impact
42 aquatic life and human health (Amenyogbe et al., 2021). They potentially affect the growth and reproductive efficiency of
43 non-targeted organisms due to their bio-magnification and persistent nature (Deribe et al., 2013). Some pesticides, such as
44 organochlorine pesticides (OCPs), can last for many years in the environment, are deposited in sediments (MacKay &
45 Fraser, 2000; Deribe et al., 2013), and concentrated in the top predators of food chains (Di et al., 2017). As a result, most
46 pesticide-related ecosystem studies focused on those OCPs (Marchand et al., 2010; Sifakis et al., 2017; Yohannes et al.,
47 2017; Worku et al., 2022). Organophosphate pesticides (OPPs) and other synthetic pesticides, which are less persistent
48 than OCPs, are also widely used by farmers worldwide (Manuelmolina-Ruiz et al., 2014). Most of them are water-soluble
49 and taken up by organisms through skin contact, inhalation, or ingestion as food particles. As a result, these pesticides are
50 hazardous to aquatic species, and their residues might affect the phosphorylation of the acetylcholinesterase enzyme
51 (AChE) at nerve terminals in non-targeted animals, causing malfunctioning (Gultekin et al., 2000; Kushwaha et al., 2016).
52 Pesticides can limit the food sources available to fish through their toxicity effect on invertebrate prey, potentially reducing
53 fish growth and survival. At high concentrations in water bodies, pesticides can affect fish by changing their growth rates,
54 progeny survival, and behavior. This influences fish population stability and structure (Holden, 1972). Toxicity effects on
55 fish can also lead to histological and hormonal changes in females, which may reduce the number of oocytes produced
56 (Forsgren et al., 2013).

57 The fish species of the genus *Labeobarbus* are the most numerous and endemic to Lake Tana. Two of the five *Labeobarbus*
58 species most frequently caught for commercial fishing are *L. tsanensis* and *L. megastoma* (Gebremedhin et al., 2019).
59 *Labeobarbus megastoma* is a piscivores fish that dominantly feeds (up to 67%) on fish, while *L. tsanensis* is feeding
60 dominantly on benthos 43%, 18% mollusks, and 14% detritus/substratum (Sibbing & Nagelkerke, 2001). Similarly,
61 *Oreochromis niloticus*, a highly commercially demanded species in the lake (Nagelkerke et al., 1995; Vijverberg et al.,
62 2013), is an omnivorous fish that feeds on plankton and aquatic plants (Tesfahun & Alebachew, 2023).


63 The Lake Tana sub-basin, one of Ethiopia's most productive places, is suitable for growing a variety of staple food crops,
64 including Teff (*Eragrostis tef*), Finger millet (*Eluesine coracana*), rice (*Oryza sativa*), maize (*Zea mays*), grass pea
65 (*Lathyrus sativus*) and chickpea (*Cicer arietinum*), as well as cash crops like Khat (*Catha edulis*) and other vegetables
66 (tomato, cabbage, onion, garlic, potato, pepper, etc.) (Abera, 2017). Recently, pesticide consumption has increased due to
67 increased agricultural activity in the area, particularly in irrigation-based dry season cash crops and vegetable cultivations
68 (Agmas & Adugna, 2020; Abera et al., 2022). Farmers spray various pesticides on those crops and vegetables in the field

69 and at their home storage, but they are usually not fully informed or aware of the dangers of the pesticides they use (Agmas
70 & Adugna, 2020). Improper spraying techniques, dropping empty containers into water bodies, careless disposal, or
71 unintentional spills of residual solutions have all been noted as contamination routes in the catchment (Agmas & Adugna,
72 2020). Additionally, the lake has a large catchment area (approximately 16,750 km²), and the process of pesticides leaching
73 through soil erosion and surface runoff increases environmental hazards and may have detrimental effects on the lake's
74 biodiversity. For example, due to brain, endocrine, behavioral, and genetic abnormalities, as well as histopathological and
75 hematological changes, fish species may not be able to grow and reproduce normally (Kumari, 2020; Srivastava et al.,
76 2016). Furthermore, pesticide residues may pose a health risk to human communities that consume fish (Agmas & Adugna,
77 2020; Abera et al., 2022). Despite the increasing use of pesticides in the lake catchment, no research has been done yet to
78 determine the concentrations and the possible harmful impacts on non-target organisms except a recent work by Abera et
79 al. (2022), and to the best of our knowledge, there was no work done on the concentration and effect of pesticide residues,
80 particularly on the economically important and endangered fish species in the lake. Therefore, this study aimed to 1)
81 determine the concentration of pesticides routinely used in the fillet and liver tissue of three commercially important fish
82 species in the study area, 2) assess any potential impact of pesticide residues on fish health and 3) assess the effect of the
83 measured pesticides on the health of fish-consuming communities.

84 **Materials and Methods**

85 Study area

86 Lake Tana and its tributaries, such as Ribb and Gumara (Fig. 1), provide important ecosystem services to people living in
87 the catchment. The two river watersheds are the most important agricultural areas for most staple crops through smallholder
88 irrigation farming. These water bodies supply drinking water for humans and animals. They are also important for fisheries.
89 *Labeobarbus* species, *Oreochromis niloticus*, and *Clarias garipinus* are commercially important and frequently consumed
90 fish species in the research area. People living near the lake and along the river use rainwater to grow crops and vegetables
91 during the rainy season. While in the dry season, farmers use the lake and river waters through furrow and pump irrigation.



92

93 **Fig. 1** Map of the study locations in Lake Tana and sampling points (Red circles) at the Ribb River Mouth (RRM) and
 94 Gumara River Mouth (GRM) sites.

95

Fish sampling and preparations

96

97 Fish samples were collected from Lake Tana's shore near the mouths of the Gumara and Ribb Rivers. Sampling was
 98 executed between September and October 2021 using multifilament gillnets (10, 12, and 14 cm mesh sizes) and catches
 99 from fishers at the sampling points. Specimens of three fish species, *Labeobarbus tsanensis*, *L. megastoma*, and
 100 *Oreochromis niloticus*, with a weight of more than 200 grams and with a total length of at least 23 cm were identified and
 101 selected (Table 1). Tissue samples were collected based on the approach stated by Rosseland et al. (2001). Each fish sample
 102 was extracted with stainless steel disposable scalpel blades with a handle, yielding around 50 g (wet weight) of the fillet
 103 (white meat taken from the dorsal muscle) and 2 to 5g of the liver. Samples were wrapped in aluminum foil, frozen (-20°C),
 104 and transported to Belgium in an ice-filled cooling cabinet. The pesticide analysis was done in the laboratory for
 105 phytopharmaceuticals at Ghent University.

105

106 Each fillet and the liver sample were homogenized for 10 minutes with an Ultra-Turrax (T 25 ultra Turrax - IKA). The
 107 homogenizer was cleaned with acetone after each homogenization to prevent cross-contamination. Based on the analyses'
 weight requirements, samples were weighed and stored in falcon tubes.

108 **Table 1** Biometric data and feeding mode of the three fish species sampled from Lake Tana between September and
 109 October 2021.

Fish species	Feeding mode	Gumara river mouth			Ribb river mouth		
		Total Length (cm)*	Weight (g)*	n	Total Length (cm)*	Weight (g)*	n
<i>L. megastoma</i>	Piscivore	29.7 – 42.5	205 - 615	8	31.7 – 43.5	285 - 700	8
<i>L. tsanensis</i>	Insectivore	30.5 – 39.5	270 - 670	8	30.1 – 35.2	250 - 445	8
<i>O. niloticus</i>	Herbivore	23.2 – 26.4	220 - 295	8	23.5 – 26.5	220 - 290	8

110 * minimum-maximum values, n: number of samples analyzed.

111 Pesticide Analysis

112 Based on data from farmers and district (locally called woreda) agricultural experts on local pesticide use in 2019 and
 113 2020, a list of 109 relevant pesticide products was prepared for the multi-residue analysis before the pesticide study. Liquid
 114 chromatography coupled with tandem mass spectrometry (LC-MS/MS) with the setting defined in Table S2 (supplementary
 115 material) was used to detect and quantify all of the screening pesticides (n = 109) (Table S1 (supplementary material)).
 116 The fish tissue samples were extracted using a modified quick, easy, cheap, effective, rugged, and safe (QuEChERS)
 117 approach. This approach has a small number of steps, is highly reliable, and yields excellent recoveries for a wide range of
 118 pesticides from various chemical families (Kim et al., 2019).

119 In a QuEChERS tube, 10 ml of acetonitrile was added to 10 g of **fillet** sample and agitated for one minute. Subsequently,
 120 a salt solution was added (6 g magnesium sulfate, 0.75 g citrate sesquihydrate, 1.5 g citrate dihydrate, 1.5 sodium chloride),
 121 and the tube was vigorously shaken. The sample was then homogenized for 2 minutes in an ultra-turax mixer before being
 122 centrifuged for 5 minutes at 3,000 rpm. In a 10 ml flask, 1 ml of the supernatant was diluted in 9 ml ultrapure/miliQ/ water,
 123 and 1.5 ml of the solution was then transferred to a vial for LC-MS/MS analysis. The liver samples were extracted by
 124 adding 10 ml of acetonitrile to 2 – 5 g of material and then adding water to a total mass of 10 g. The falcon tubes were
 125 shaken for 1 minute before adding the salt mixture. The remaining stages were the same as the **fillet**, except for a clean-up
 126 phase. As a result, 7 ml of the supernatant was transferred to an SPE tube, which was agitated for 5 minutes before being
 127 centrifuged at 3000 rpm for 5 minutes. In a 10 ml flask, 1 ml of the supernatant was diluted in 9 ml ultrapure (miliQ) water.
 128 Finally, 1.5 ml of the sample was subsampled into a vial for pesticide determination.

129 Method validation

130 **The minimal concentration at which the analyte was identified and established by comparing measured signals from**
 131 **samples with known low amounts of the analyte with those of blank samples. The present method was developed and**
 132 **validated using the ICH Q2(R1) guideline.** The pesticide residues in the fish tissue samples were validated and quantified
 133 using a calibration curve from samples spiked with the multi-compound stock standard solution. The linear range of the
 134 calibration curve was developed using five concentration levels between 0.001 and 0.1 mg l⁻¹. The spike-placebo recovery
 135 method was used to determine the validation of the analyte's lowest detectable concentration in a sample (limit of detection
 136 or LOD) and the lowest confirmed level with adequate precision and recovery (limit of quantification or LOQ). A blank

137 sample was spiked and analyzed under the same conditions in four replicates. The LOD and LOQ values for the pesticide
 138 analysis in LC-MS/MS were set as 0.0003 mg l⁻¹ and 0.00086 mg l⁻¹, respectively.

139 Human health risk assessment

140 A human health risk assessment was conducted to evaluate if the pesticides found in the lake threatened those who included
 141 fish in their diet. The Acute Reference Dose (ARfD) and Acceptable Daily Intake (ADI) were utilized as expected, with
 142 no impact levels for acute and chronic pesticide exposure. The Estimated Daily Intake (EDI) for a given pesticide was
 143 calculated using the following formula (Mahmood & Malik, 2014; Hamid et al., 2017)

$$144 \quad EDI = \frac{C_p \times F_c}{bw} \quad (1)$$

145 C_p is the maximum pesticide residue concentration in white fish meat in µg/kg wet weight. F_c is the amount of fish food
 146 consumed by a person per day in kg/day. bw is the fish-consuming adult in kg.

147 The hazard quotient (HQ) for acute and chronic estimations was performed using the following formula (Hamid et al.,
 148 2017)

$$149 \quad HQ_{acute} = \frac{EDI}{ARfD} \quad (2)$$

$$150 \quad HQ_{chronic} = \frac{EDI}{ADI} \quad (3)$$

151 For the Ethiopian community who live around the water bodies and have high access to fish in their diet, the average daily
 152 intake rate was taken as 0.03 kg/day (Yohannes et al., 2014) and assumed the average body weight of 60 kg for an adult
 153 Ethiopian for the calculation (Teklu et al., 2015). As liver tissue is not consumed in the area, the pesticide concentration in
 154 the fillet was only considered for the risk assessment.

155 The Hazard Index (HI) was taken as the sum of the chronic or acute hazard quotients separately of all pesticides detected
 156 in the fish fillet. Pesticide residues could potentially produce acute/chronic health hazards if the percentage of HI is greater
 157 than 100 (Lozowicka, 2015).


158 Data analysis

159 The Kruskal-Wallis ANOVA test was used to examine differences in pesticide residue concentrations in the tissue among
 160 species, and the Mann-Whitney U test between tissue types using the R package version 4.2.0. The Dunn test was used to
 161 evaluate any significant differences between the groups. **As a result of an uneven distribution of the pesticide concentration
 162 data in the fish tissue, the median values were utilized to represent the pesticide concentrations in the different fish species
 163 and tissue types.** For these statistical comparisons, a pesticide concentration below the LOQ was taken to be half of the
 164 corresponding assigned limiting value. Hence the number of > LOQ values recorded among the pesticides detected was
 165 very small (< 60%); the deterministic approach was used to determine the exposure of the fish consumers and the hazard
 166 quotients by taking an average pesticide concentration for values that had two or greater than two > LOQ values. For the
 167 risk assessment, the LOQ cutoff value, 0.00086 mg/kg, was used for fish **fillet** samples with none > LOQ value or those
 168 detected pesticides with < LOQ values as a maximum pesticide concentration, considering the worst-case scenario. The
 169 exposure analysis was performed using Microsoft Excel.

170 **Results**

171 Pesticide concentrations in fish samples

172 Ten pesticides were found in the fish **fillet** and liver tissue samples of the three fish species under investigation. **The**
 173 **pesticide types insecticides and fungicides were found in equal proportion (40% of each) from the detected pesticides,**
 174 **while herbicides constitute 20%.** The level of pesticides quantified ranges from below LOQ to **1850.0 $\mu\text{g kg}^{-1}$ wet weight.**
 175 The pesticide most commonly found and had the highest concentration was pyrimethanil. **The difference between the two**
 176 **research sites was statistically significant ($p < 0.001$) due to the higher median concentration of pyrimethanil (49.7 $\mu\text{g kg}^{-1}$)**
 177 **at the site near the mouth of the Gumara River (GRM) compared to 2.1 $\mu\text{g kg}^{-1}$ at the site near the mouth of the Ribb**
 178 **River (RRM) (Table 2).** Pyrimethanil was detected in 65% of **fillet** samples with a mean concentration of 96.6 $\mu\text{g kg}^{-1}$ and
 179 96% of liver tissue samples with a mean concentration of 104.0 $\mu\text{g kg}^{-1}$ for the three fish species. A statistically significant
 180 difference in pyrimethanil concentration between species ($p = 0.043$) was observed, with the highest concentration in *L.*
 181 *megastoma* and the lowest in *L. tsanensis* (Fig. 2 and Table 3). The mean detected concentration of pyrimethanil was much
 182 higher in the liver (258.8 $\mu\text{g kg}^{-1}$) than in the **fillet** (55.4 $\mu\text{g kg}^{-1}$) of *L. megastoma*. However, pyrimethanil was detected in
 183 higher concentration in the **fillet** of *O. niloticus* (mean = 359.8 $\mu\text{g kg}^{-1}$) and *L. tsanensis* (mean = 99.1 $\mu\text{g kg}^{-1}$) than in the
 184 liver tissue of those species (Table 4). Like pyrimethanil, carbaryl was quantified in the **fillet** and liver tissue of the three
 185 species and found in both studied sites. The highest mean concentration of carbaryl (10.4 $\mu\text{g kg}^{-1}$) was recorded in *O.*
 186 *niloticus*. However, variations among species were not statistically different.



187 **Fig. 2** Boxplot showing pyrimethanil concentration in the tissue of the three studied fish species in the studied sites in
 188 **Lake Tana.**

189 **Table 2** Mean concentrations and range of detected pesticides in Lake Tana fish species sampled between September and
 190 October 2021 at the GRM and RRM sampling site.

Pesticide	Pesticide type	GRM		RRM	
		Median ($\mu\text{g kg}^{-1}$)	Range ($\mu\text{g kg}^{-1}$)	Median ($\mu\text{g kg}^{-1}$)	Range ($\mu\text{g kg}^{-1}$)
Benalaxyl	Fungicide	2.2*	< LOD – 2.2	< LOD	-
Butachlor	Herbicide	< LOD	-	1.1	< LOD – 1.1
Carbaryl	Insecticide	0.4	0.2 – 56.5	0.4	0.2 – 2.2
Difenoconazole	Fungicide	0.2*	< LOD – 2.2	< LOD	-
Flazasulfuron	Herbicide	2.2	2.2 – 60.1	2.2	< LOD – 2.2
Imidacloprid	Insecticide	< LOD	-	0.9	0.4 – 2.2
Oxamyl	Insecticide	216.5	32.0 – 507.0	24.4	2.2 – 136.9
Pirimiphos-methyl	Insecticide	< LOD	-	0.4*	< LOD – 0.4
Pyrimethanil	Fungicide	49.7	2.2 – 1850.0	2.2	0.4 – 171.7
Tebuconazole	Fungicide	0.2*	< LOD – 0.2	< LOD	-

*quantified only in a single sample; LOD = 0.0003 $\mu\text{g kg}^{-1}$; LOQ = 0.00086 $\mu\text{g kg}^{-1}$

192 Oxamyl and flazasulfuron were detected in both study sites (Table 2). However, oxamyl was found in the tissues of the
 193 two *Labeoburbus* species but at a level below LOD in *O. niloticus* (Table 3). Oxamyl was found in *L. megastoma* and *L.*
 194 *tsanensis*, and the highest concentration was detected in the liver tissue of *L. megastoma* (median = 361.7 $\mu\text{g kg}^{-1}$) (Tables
 195 3 & 4). Though its concentration was small compared to liver tissue, oxamyl was also detected in the fillet of *L. megastoma*,
 196 while it was below LOD in *L. tsanensis* and *O. niloticus*. Imidacloprid and flazasulfuron were detected in different
 197 quantities in the tissues of the three fish species under investigation at site RRM (Table 5.4), whereas imidacloprid was
 198 below LOD at site GRM (Table 3). Imidacloprid was only detected in the liver tissue of *L. tsanensis*; however, it was
 199 detected in the fillets of *L. megastoma* and *O. niloticus* at the RRM site (Table 4). However, the concentrations of oxamyl,
 200 imidacloprid, and flazasulfuron in neither species nor the tissue types were statistically different.

201 **Table 3.** The median pesticide concentration detected in the tissues (fillet + liver) of the three fish species at the two
 202 sampling sites in Lake Tana and variations of pesticide types among species is presented in the *p* value.

Sampling site	Pesticide	Pesticide concentrations in fish species ($\mu\text{g kg}^{-1}$)			<i>p</i> value
		<i>L. megastoma</i>	<i>L. tsanensis</i>	<i>O. niloticus</i>	
GRM	Carbaryl	0.2	0.4	4.6	0.041
	Flazasulfuron	2.2	2.2*	2.2	0.446
	Imidacloprid	<LOD	<LOD	<LOD	-
	Oxamyl	361.7	<LOD	< LOD	-
	Pyrimethanil	82.6	30.1	55.1	0.074
RRM	Carbaryl	0.4	-	1.3	0.739
	Flazasulfuron	2.2	2.2	2.2	-
	Imidacloprid	0.4	2.2	0.4	0.223
	Oxamyl	<LOD	24.4	<LOD	-

Pyrimethanil 22.8 2.2 2.2 0.010

* quantified only in a single sample

Benalaxyl, a pesticide that was rarely discovered, was found in the liver tissue of *L. megastoma* at the GRM site (median = 2.2 $\mu\text{g kg}^{-1}$); nevertheless, difenoconazole was found in the fillet tissue (median = 0.22 $\mu\text{g kg}^{-1}$) of *L. megastoma*. Similarly, butachlor and pirimiphos-methyl were detected only in RRM. Butachlor was detected in the liver tissue of *L. tsanensis* (median = 1.1 $\mu\text{g kg}^{-1}$), and pirimiphos-methyl was found in the fillet of *O. niloticus* (median = 0.4 $\mu\text{g kg}^{-1}$). Tebuconazole was also detected in the fillet of *L. megastoma* in the GRM site (Table 4).

Table 4 Detected pesticide concentration in the tissues of the studied fish species sampled between September and October 2021 at the GRM and RRM sampling site in Lake Tana.

Pesticide	Mean pesticide concentration in $\mu\text{g kg}^{-1}$ wet weight					
	<i>L. megastoma</i>		<i>L. tsanensis</i>		<i>O. niloticus</i>	
	liver	fillet	liver	fillet	Liver	fillet
Benalaxyl	2.2*	< LOD	< LOD	< LOD	< LOD	< LOD
Butachlor	< LOD	< LOD	1.1	< LOD	< LOD	< LOD
Carbaryl	2.2*	0.2	2.2*	0.4	29.3	3.1
Difenoconazole	< LOD	0.2*	< LOD	< LOD	< LOD	< LOD
Flazasulfuron	2.2	< LOD	2.2	< LOD	2.2	< LOD
Imidacloprid	< LOD	0.2	2.2*	< LOD	< LOD	0.4*
Oxamyl	361.7	32.0*	24.4	< LOD	< LOD	< LOD
Pirimiphos-methyl	< LOD	< LOD	< LOD	< LOD	< LOD	0.4*
Pyrimethanil	15.0	54.1	7.9	35.6	25.4	94.9
Tebuconazole	< LOD	0.2	< LOD	< LOD	< LOD	< LOD

* quantified only in a single sample; LOD = 0.0003 $\mu\text{g kg}^{-1}$; LOQ = 0.00086 $\mu\text{g kg}^{-1}$

Human health risk assessment

People eating fish from Lake Tana might be exposed to the seven pesticides detected in the fillet of the three fish species (Table 4). The risk assessment estimation considers all values found in the fillet, including < LOQ values. The highest daily exposure of consumers was recorded for oxamyl (0.016 $\mu\text{g/kg bw/d}$) (Table 5). The calculated values for HQ acute and HQ chronic were low, with the highest value of 1.60% found for oxamyl. Similarly, the HI values for both HI acute and HI chronic (Table 5) showed a lower value than the threshold value of 100, indicating that adverse non-cancer health effects are not likely to occur or the health risk for the fish meat-consuming community is very low.

219 **Table 5** Adult human exposure to selected quantified (> LOQ) pesticides in the **fillet** of fish species in the Lake Tana sub-
 220 basin. Maximum detected quantity (Cp), Acceptable daily intake (ADI), Acceptable reference dose (ARfD), Estimated
 221 Daily Intake (EDI), and the percentage of hazard quotient for acute (HQ acute) and chronic (HQ chronic) cases.

Pesticide	ADI ($\mu\text{g}/\text{kg bw/d}$)	ARfD ($\mu\text{g}/\text{kg bw}$)	Max ($\mu\text{g kg}^{-1}$)	EDI ($\mu\text{g}/\text{kg bw/d}$)	HQ acute (%)	HQ chronic (%)
Carbaryl	7.5	10.0	14.0	0.007	0.07	0.10
Oxamyl	1.0	1.0	32.0	0.016	1.60	1.60
Pyrimethanil	170	1000	1206.6	0.603	0.06	0.35
				HI (%)	1.73	2.06

222 Discussion

223 Detected pesticides and the possible effect on fish species

224 The development, survival, and reproduction of aquatic animals are known to be impacted when surface waters are
 225 contaminated by high pesticide concentrations (Vonesh & Kraus, 2009; Akhter, 2019; Kumari, 2020). It has also been
 226 shown that several pesticides in water systems have affected primary and secondary production, including fish species in
 227 the food chain (Seeland et al., 2012; Müller et al., 2019). The effect, however, potentially depends on the physicochemical
 228 environment and may be ecoregion-specific, long-lasting, or delayed, yet most likely exacerbated by indirect effects
 229 resulting from interspecific interactions (Araújo et al., 2012; Müller et al., 2019).

230 The highest concentration and the frequent occurrence of pyrimethanil in the three fish species in Lake Tana indicate that
 231 it is intensively used in the watershed. Pyrimethanil is one of the most widely used fungicides worldwide and has been
 232 found in numerous aquatic environments (Lozowicka, 2015; De Cock et al., 2021). It is used mainly on fruits and vegetables
 233 and was manufactured to treat resistant fungus strains such as *Botrytis* spp, which affects onions and other vegetables and
 234 fruits (Tournas, 2005; Abo-Elyousr et al., 2020). Pyrimethinil is also applied for seed treatment on cereals (EFSA, 2011).
 235 It affects algae, zooplankton, and macroinvertebrates which are potential food sources for fish species. For example, for
 236 the model organism *Scenedemus acutus* in a 48h chronic test and *Daphnia magna* in a 48-h acute test, the average LC50
 237 was 23.2 mg l⁻¹ and 3.6 mg l⁻¹, respectively (Araújo et al., 2012). They also reported that the reproduction of adult *D.*
 238 *magna* was adversely affected when exposed to 0.5 mg l⁻¹ of pyrimethanil over many generations under variable
 239 temperatures between 20 and 27 °C. **Although the pesticide concentration was not directly measured from the water in**
 240 **Lake Tana, the maximum concentration (up to 1.85 mg kg⁻¹) in the fish tissue suggests that the concentration was**
 241 **potentially high. This high concentration of pyrimethanil could affect the normal development of fish organs. According**
 242 **to Bernab et al. (2017), doses up to 50 $\mu\text{g l}^{-1}$ cause a histological alteration in tree frogs (*Hyla intermedia*). The effect of**
 243 **this pesticide also elevated with increasing temperature. For example, Araújo et al. (2012) confirmed that a rise in ambient**
 244 **temperature exacerbated the fungicide's toxicological effects on *D. magna*.** Thus, with the relatively high dry season
 245 temperature, the effect of pyrimethanil on aquatic invertebrates in Lake Tana might be higher, which could indirectly affect
 246 fish species.

247 Metals and pesticides can be absorbed by aquatic macrophytes and algae (Dosnon-Olette et al., 2010, 2009), and organisms
 248 that directly consume those primary sources may have higher concentrations in their tissue. The higher mean concentration

249 of pyrimethanil in the tissue of *O. niloticus* (Table 3) might reflect the assimilation of the fungicide through the food chain.
250 Similarly, the highest concentration in the piscivore species *L. megastoma* (Table 4) could also result from its feeding habit,
251 despite pyrimethanil having less potential in bioaccumulation (Araújo et al., 2015; Lewis et al., 2016).

252 Flazasulfuron, belonging to the sulfonylurea herbicide family, was detected in 29% of tissue samples from the three fish
253 species with a maximum concentration of 0.06 mg kg⁻¹ in *L. megastoma*. This herbicide is known to prevent protein
254 synthesis in plants by blocking branched-chain amino acids protein synthesis (Olette et al., 2008). It has also been shown
255 to impact the chlorophyll pigments and leaf gas exchange (Frankart et al., 2003). While it is not toxic to aquatic animals
256 (US EPA, 2007), it can indirectly have an impact because it might reduce plant and nonvascular aquatic plant productivity
257 (US EPA, 2007). For instance, flazasulfuron reduces the photosynthetic capacity of duckweed (*Lemna minor*) by 16% after
258 seven days of exposure to a 100 µg l⁻¹ concentration (Olette et al., 2008).

259 The other frequently found pesticide in the three sampled fish species and the two study sites in Lake Tana was carbaryl
260 (Tables 2 & 3, Fig. S2 (supplementary material)). This is a carbamate insecticide that has been reported to have an impact
261 on several fish species. For example, Patnaik and Patra (2006) observed its impact on the blood cells in *Clarias batrachus*
262 following a 96-hour acute exposure, with a cell shape modified at 12.6 mg l⁻¹ and an LC50 at 15.6 mg l⁻¹. In addition, under
263 a controlled investigation, Boran et al. (2010) found that rainbow trout had an LC50 at 2.5 mg l⁻¹ and 1.4 mg l⁻¹ over 24
264 and 96 hours of exposure time, respectively. Matos et al. (2007) also report the effect of carbaryl as it resulted in necrosis
265 of the liver tissue of *O. niloticus* when exposed to a concentration of 0.25 mg l⁻¹ for seven days. The pH and temperature
266 of the aquatic environment significantly impact the persistence of carbamate insecticides in natural waters (Aly & El-Dib,
267 1971). Carbaryl generally has a low water persistence (US EPA, 2003). However, the amount recorded in the fish tissue,
268 up to 56.5 µg kg⁻¹ in this study, indicates its excessive pesticide use. The detected amount may not impose risk in the short
269 term. However, prolonged exposure possibly imposes environmental and health impacts, including oxidative stress in the
270 fish species, as observed in several animal groups (Ribera et al., 2001; Matos et al., 2007). In addition, the effect of
271 pesticides is more pronounced on juveniles than on adults of the same species (Lin et al., 2007), so any prolonged and
272 extensive use could affect fish recruitment and population growth.

273 The other carbamate insecticide that was detected in the two *Labeobarbus* species of Lake Tana was oxamyl (Table 3, Fig.
274 S1 (supplementary material)), which is relatively more toxic than carbaryl for aquatic organisms (Lewis et al., 2016;
275 Alvarez et al., 2022). Oxamyl is applied as an insecticide on field crops, vegetables, fruits, and ornamental plants (Alvarez
276 et al., 2022). It is moderately toxic to fish, and its LC50 after 96 h exposure for species such as bluegill sunfish, goldfish,
277 and rainbow trout were found to be 5.6 mg l⁻¹, 27.5 mg l⁻¹, and 42 mg l⁻¹, respectively (Smith, 1982). Less persistent
278 pesticides such as oxamyl are rarely found in fish tissue. For example, Polat et al. (2018) found a 45 µg kg⁻¹ concentration
279 in the fillet of sardine (*Sardine pilchardus*) among nine studied fish species in the Iskenderun Bay of Turkey. However,
280 in this investigation, oxamyl was detected in 7% of the fish tissue sampled from the lake (Table 3), with the highest
281 concentration (507 µg kg⁻¹) in the liver tissue of *L. megastoma*, indicating its excessive use in the catchment. Although
282 additional research on fish sensitivity to pesticides in Lake Tana is needed, the amount of oxamyl in this study showed the
283 necessity for strict regulation and monitoring.

284 The fungicides, benalaxyl, difenoconazole and tebuconazole were found < LOQ in fish sampled from GRM site. Albeit
285 their concentrations were low, the presence of these pesticides in this site indicates that they have been applied in the
286 Gumara River catchment. Benalaxyl and difenoconazole, however, were not listed in Abera et al. (2022)'s list of pesticides
287 used in the lake watershed, indicating that there are still other pesticides available in the local market that may come through

288 a different marketing route and circulate through an informal supply network as reported in Mengistie et al. (2016).
289 Benalaxyl is a widely used fungicide on tomatoes, onions, and potatoes in the field, while difenoconazole and tebuconazole
290 are applied on cereals and other field crops (Lewis et al., 2016). Although these fungicides had a high log K_{ow} value used
291 (Lewis et al., 2016) compared to other pesticides found in Lake Tana, they were only found in one sample of the fillet of
292 *L. megastoma*. This suggests that they are probably applied infrequently in the catchment or, due to their lipophilic nature
293 (Houbraken et al., 2016; Rasool et al., 2022), adsorbed in organic wastes in sediment. Studies showed that tebuconazole
294 could cause a decrease in hemoglobin and the number of red blood cells in cyprinids when applied with a 2.5 mg l⁻¹
295 concentration for a 96-h exposure (Çilingir Yeltekin et al., 2020). The LC50 of tebuconazole for fish is 4.4 mg l⁻¹ (Lewis
296 et al., 2016). Similarly, the broad-spectrum triazole fungicide difenoconazole is extensively used in agriculture (Jiang et
297 al., 2022). It influences immune-related pathways and energy, lipid, and amino acid metabolism in the early stages of
298 zebrafish development (Teng et al., 2018). Although the examined fish species' susceptibility to those pesticides was not
299 assessed, the observed concentration, which was < LOQ, suggests they very likely do not cause any risk to fish at the
300 current application level in the lake catchment.

301 The other commonly used insecticides, imidacloprid and pirimiphos-methyl, are frequently detected in many aquatic
302 environments (Vignet et al., 2019). Both insecticides are moderately toxic to fish (Lewis et al., 2016). Imidacloprid was
303 detected in the three studied fish species sampled from RRM. According to Mhadhbi and Beiras (2012), when imidacloprid
304 concentrations in the water were above 0.04 mg l⁻¹, it posed pericardial edema and spinal deformities, and it also reduced
305 hatching success when its concentration reached above 0.4 mg l⁻¹ in turbot (*Psetta maxima*). In the Ribb and Gumara
306 catchments, crops like rice and other cereals, including maize, are grown and commonly treated with pirimiphos-methyl
307 and imidacloprid. However, the fact that both pirimiphos-methyl and imidacloprid were found in the fish samples taken
308 from the RRM site suggests that farmers in the Ribb River catchment may have used these insecticides in greater quantities
309 than in Gumara.

310 In general, the number of herbicides, fungicides, and insecticides found in the tissue of the three fish species in Lake Tana
311 indicates how much pesticide use has polluted the lake water. Pesticides used by farmers in the Gumara and Ribb rivers to
312 reduce crop loss and increase the quality of their crops, vegetables, and fruits are indispensable. However, reducing
313 agricultural losses by using too many of these agrochemicals should not come at the expense of aquatic life. Increasing
314 production using intensive agrochemicals could not be sustainable because a rise in pesticide use seriously harms an
315 ecosystem's health and natural resource. The best way to recommend sustainable production and an environmentally
316 friendly approach is using integrated pest management, which weighs natural alternatives to chemical pesticides and only
317 uses them as a last resort (Barzman et al., 2015). Using less toxic and less persistent pesticides is also better for creating
318 more resilient, sustainable food systems. Monitoring the type and quantity of pesticides used and assessing their impact on
319 biodiversity should be part of the management plan of the lake and its catchment, even though the concentrations currently
320 seem to have no adverse toxicological effect on fish species at the current application level.

321 Pesticides risk to human health

322 Fish ingest pesticide residues mainly through the epidermis of the gills and ingested food (Braunbeck et al., 1998). When
323 a fish is consumed, the pesticides accumulated in its edible tissue can be transferred to humans. Liver tissue is the
324 detoxification center and stores toxicants in many studied fish species (Braunbeck et al., 1998; Macirella et al., 2022). In
325 this study, however, some pesticides, e.g., pyrimethanil, were found in higher concentrations in the fillet tissue of *L.*

326 *tsanensis* and *O. niloticus* (Table 4). Consequently, humans who consume fish meat from the lake are exposed to these
327 pollutants. Two to three different pesticides were found in some fillet samples, and their impact on the health of fish and
328 its consumers is likely greater than that of just a single pesticide due to their combined effect (Kortenkamp, 2007;
329 Hernández et al., 2013). No clear maximum residual limit (MRL) for the three pesticides was available for the risk
330 assessment (Table 5). However, from the metadata for most of the animal products in Great Britain, a maximum value of
331 50 µg kg⁻¹ for carbaryl and pyrimethanil and 10 µg kg⁻¹ for oxamyl are used (Lewis et al., 2016). Considering those
332 conservative MRL values, the detected concentration of pyrimethanil in 12 samples and one sample for oxamyl was higher
333 than the limit. The risk assessment, however, might reflect the worst-case scenario because we did not consider the impact
334 of washing, soaking with solutions, cooking, and freezing procedures that could lower the concentration of pesticides in
335 the tissue (Witczak, 2009; Alaboudi et al., 2021; Islam et al., 2022). Overall, the HQ and HI values showed that, at the
336 current application practices, pesticides pose no known harm to human health or consumers of fish collected from the lake.
337 **However, the detection of ten pesticides in the fish tissue in this study** shows that the use of pesticides along the river
338 catchments needs urgent attention and close monitoring to safeguard the public health of local communities and the
339 population of both culturally endangered and commercially important fish species.

340 **Conclusion**

341 **The contamination of ten pesticide residues (2 herbicides, 4 fungicides, and 4 insecticides) was detected in the three**
342 **commercially important fish species.** Variations in pesticide concentrations among fish species and between tissue types
343 were not statistically significant, except for pyrimethanil, which revealed a significant difference between *L. megastoma*
344 and *L. tsanensis*. *L. megastoma* was found to have the highest level of pesticide contamination (eight out of ten) among the
345 three studied fish species, which may be related to its' piscivorous feeding behavior. Pyrimethanil > carbaryl > oxamyl >
346 flazasulfuron were the most frequently occurring pesticides in the studied fish species with > LOQ values. Moreover, most
347 pesticides were found in higher concentrations in the fish's liver than in the fillet. For all of the dominantly detected
348 pesticides, the health risk calculations revealed that the health risk index was below the cutoff value, indicating that, given
349 current pesticide use, there is likely no harm to human health from pesticide exposure in fish. **Due to the lack of samples**
350 **from the water and sediment, the study is limited in its ability to evaluate the effects of combined exposure to toxins and**
351 **their influence on the lake's biota. Additionally, the small sample size results in an uneven distribution of data about the**
352 **presence of pesticides in fish tissue.** However, although the study was conducted on a limited sample size, only three
353 species and a single developmental stage, the results could serve as a preliminary step for further research, and the
354 information will be used to generate a management strategy for the fish populations in Lake Tana. **Further investigations**
355 **are recommended, including a wide variety of fish species,** a sizable sample at various stages of development, and
356 measurements of pesticide concentrations in the water and sediment for a more in-depth ecotoxicological assessment.

357 **Ethics statement**

358 The Ethical Committee of Bahir Dar University, College of Agriculture and Environmental Sciences, approved the study.
359 The care and use of animals followed all applicable international, national, and institutional guidelines.

360 Acknowledgment

361 The authors thank the Belgian Development Cooperation (VLIR-UOS) for financially supporting this research through the
362 Inter-University Cooperation of Flemish and Bahir Dar Universities (IUC-BDU). The VLIR-UOS project staff at Bahir
363 Dar University are acknowledged for the support and logistics provided. We thank Mr. Ferede Tsegaw and Mr. Kassahun
364 Erkyehun for their help in fish specimen collections.

365 Author contribution

366 Wondie Zelalem: conceptualization, investigation, methodology, formal analysis, writing–original draft; Wassie Anteneh,
367 Minwelet Mingist: conceptualization, investigation, editing; Alain De Vocht: conceptualization, project administration,
368 methodology, investigation, data curation, editing; Jasmine De Rop, Andrée De Cock, Pieter Spanoghe, Peter L.M
369 Goethals: methodology, data curation, formal analysis, editing; Mulugeta Kibret, Enyew Adgo, Jan Nyssen, Elie Verleyen:
370 investigation, project administration, editing. Felegush Erarto: investigation, editing.

371 Funding

372 No funding was obtained for this study.

373 Data Availability

374 The corresponding author will provide the data upon reasonable request.

375 Competing interests

376 The authors declare no competing interests.

377 References

- 378 Abera, B., Van Echelpoel, W. Van, De Cock, A. De, Tytgat, B., Kibret, M., Spanoghe, P., Mengistu, D., Adgo, E., Nyssen,
379 J., Goethals, P. L. M., & Verleyen, E. (2022). Environmental and Human Health Risks of Pesticide Presence in the
380 Lake Tana Basin (Ethiopia). *Sustainability*, *14*(21), 14008. <https://doi.org/10.3390/su142114008>
- 381 Abera, M. (2017). Agriculture in the Lake Tana Sub-basin of Ethiopia. In A. S. and G. G. Stave K. (Ed.), *Social and*
382 *Ecological System Dynamics* (pp. 375–397). Springer. https://doi.org/10.1007/978-3-319-45755-0_23
- 383 Abo-Elyousr, K. A. M., Alamri, S. A. M., Hussein, M. M. A., Hassan, M. A. H., Abd El-Fatah, B. E. S., & Hashem, M.
384 (2020). Molecular disparities among Botrytis species involved in onion umbel blight disease and its management
385 using *Bacillus subtilis* PHYS7. *Egyptian Journal of Biological Pest Control*, *30*(1), 1–12.
386 <https://doi.org/10.1186/s41938-020-0205-x>

- 387 Agmas, B., & Adugna, M. (2020). Attitudes and practices of farmers with regard to pesticide use in NorthWest Ethiopia.
388 *Cogent Environmental Science*, 6(1), 1791462. <https://doi.org/10.1080/23311843.2020.1791462>
- 389 Akhter, Z. (2019). Pesticide Contamination in Aquatic Ecosystems. *International Journal of Scientific Research and*
390 *Review*, 7(January), 2356–2381. <https://www.researchgate.net/publication/348150568>
- 391 Alaboudi, A. R., Almashhadany, D. A., & Jarrah Bashar Suleiman. (2021). Effect of Cooking and Freezing on Levels of
392 pesticides residues in Local Fresh Fish. *Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj-*
393 *Napoca. Animal Science and Biotechnologies*, 78(2), 28. <https://doi.org/10.15835/buasvmcn-asb:2021.0007>
- 394 Alvarez, F., Arena, M., Auteri, D., Binaglia, M., Federica Castoldi, A., Chiusolo, A., Colagiorgi, A., Colas, M., Crivellente,
395 F., de Lentdecker, C., Egsmose, M., Fait, G., Ferilli, F., Gouliarmou, V., Nogareda, L. H., Ippolito, A., Istace, F.,
396 Jarrah, S., Kardassi, D., ... Villamar□Bouza, L. (2022). Peer review of the pesticide risk assessment of the active
397 substance oxamyl. *EFSA Journal*, 20(5). <https://doi.org/10.2903/j.efsa.2022.7296>
- 398 Aly, O. M., & El-Dib, M. A. (1971). Studies on the persistence of some carbamate insecticides in the Aquatic Environment-
399 -I. Hydrolysis of sevin, baygon, pyrolan, and dimetilan in waters. *Water Research Pergamon Press*, 5, 1191–1205.
- 400 Amenyogbe, E., Huang, J. S., Chen, G., & Wang, Z. (2021). An overview of the pesticides' impacts on fishes and humans.
401 *International Journal of Aquatic Biology*, 9(1), 55–65. <https://doi.org/10.22034/ijab.v9i1.972>
- 402 Araújo, C. V. M., Shinn, C., Müller, R., Moreira-Santos, M., Shinn, C., Espindola, E. L., & Ribeiro, R. (2015). *The*
403 *Ecotoxicity of Pyrimethanil for Aquatic Biota Skin microbiome of Pelophylax perezi frogs View project The*
404 *Ecotoxicity of Pyrimethanil for Aquatic Biota*. <https://doi.org/10.13140/RG.2.1.1568.0485>
- 405 Barzman, M., Bårberi, P., Birch, A. N. E., Boonekamp, P., Dachbrodt-Saaydeh, S., Graf, B., Hommel, B., Jensen, J. E.,
406 Kiss, J., Kudsk, P., Lamichhane, J. R., Messéan, A., Moonen, A. C., Ratnadass, A., Ricci, P., Sarah, J. L., & Sattin,
407 M. (2015). Eight principles of integrated pest management. *Agronomy for Sustainable Development*, 35(4), 1199–
408 1215. <https://doi.org/10.1007/s13593-015-0327-9>
- 409 Bernabò, I., Guardia, A., MacIrella, R., Tripepi, S., & Brunelli, E. (2017). Chronic exposures to fungicide pyrimethanil:
410 Multi-organ effects on Italian tree frog (*Hyla intermedia*). *Scientific Reports*, 7(1). [https://doi.org/10.1038/s41598-](https://doi.org/10.1038/s41598-017-07367-6)
411 [017-07367-6](https://doi.org/10.1038/s41598-017-07367-6)
- 412 Boran, H., Altinok, I., & Capkin, E. (2010). Histopathological changes induced by maneb and carbaryl on some tissues of
413 rainbow trout, *Oncorhynchus mykiss*. *Tissue and Cell*, 42(3), 158–164. <https://doi.org/10.1016/j.tice.2010.03.004>
- 414 Braunbeck, T. ..., Hinton, D. E., & Streit, B. (1998). Bioaccumulation of contaminants in fish. In T. Braunbeck, D. E. Hinton,
415 & B. Streit (Eds.), *Fish Ecotoxicology* (86., Vol. 86). Birkhäuser, Basel. [https://doi.org/10.1007/978-3-0348-8853-](https://doi.org/10.1007/978-3-0348-8853-0_12)
416 [0_12](https://doi.org/10.1007/978-3-0348-8853-0_12)
- 417 Çilingir Yeltekin, A., Oğuz, A. R., Kankaya, E., Özok, N., & Güneş, İ. (2020). Hematological and biochemical response
418 in the blood of *alburnus tarichi* (Actinopterygii: Cypriniformes: Cyprinidae) exposed to tebuconazole. *Acta*
419 *Ichthyologica et Piscatoria*, 50(4), 373–379. <https://doi.org/10.3750/AIEP/02931>
- 420 Andréé, D. C., Marie Anne Eurie, F., Niels, D. T., Isabel, G. A., Arne, D., Wout, V. E., Lenin, R. F., Jasmine, D. R.,
421 Liesbeth, J., Pieter, S., Luis, D. G., & Peter L.M., G. (2021). From field to plate: Agricultural pesticide presence in

- 422 the guayas estuary (Ecuador) and commercial mangrove crabs. *Environmental Pollution*, 289(August).
 423 <https://doi.org/10.1016/j.envpol.2021.117955>
- 424 Deribe, E., Rosseland, B. O., Borgström, R., Salbu, B., Gebremariam, Z., Dadebo, E., Skipperud, L., & Eklo, O. M. (2013).
 425 Biomagnification of DDT and its metabolites in four fish species of a tropical lake. *Ecotoxicology and Environmental*
 426 *Safety*, 95, 10–18. <https://doi.org/10.1016/j.ecoenv.2013.03.020>
- 427 Di, S., Liu, R., Tian, Z., Cheng, C., Chen, L., Zhang, W., Zhou, Z., & Diao, J. (2017). Assessment of tissue-specific
 428 accumulation, elimination and toxic effects of dichlorodiphenyltrichloroethanes (DDTs) in carp through aquatic food
 429 web. *Scientific Reports*, 7(1), 1–15. <https://doi.org/10.1038/s41598-017-02612-4>
- 430 Dosnon-Olette, R., Couderchet, M., & Eullaffroy, P. (2009). Phytoremediation of fungicides by aquatic macrophytes:
 431 Toxicity and removal rate. *Ecotoxicology and Environmental Safety*, 72(8), 2096–2101.
 432 <https://doi.org/10.1016/j.ecoenv.2009.08.010>
- 433 Dosnon-Olette, R., Trotel-Aziz, P., Couderchet, M., & Eullaffroy, P. (2010). Fungicides and herbicide removal in
 434 *Scenedesmus* cell suspensions. *Chemosphere*, 79(2), 117–123. <https://doi.org/10.1016/j.chemosphere.2010.02.005>
- 435 EFSA. (2011). Review of the existing maximum residue levels (MRLs) for pyrimethanil according to Article 12 of
 436 Regulation (EC) No 396/2005. *EFSA Journal*, 9(11). <https://doi.org/10.2903/j.efsa.2011.2454>
- 437 FAO. (2021). *FAOSTAT ANALYTICAL BRIEF 16: Pesticides use Global, regional and country trends*.
 438 <https://www.fao.org/3/cb3411en/cb3411en.pdf>
- 439 Forsgren, K. L., Riar, N., & Schlenk, D. (2013). The effects of the pyrethroid insecticide, bifenthrin, on steroid hormone
 440 levels and gonadal development of steelhead (*Oncorhynchus mykiss*) under hypersaline conditions. *General and*
 441 *Comparative Endocrinology*, 186, 101–107. <https://doi.org/10.1016/j.ygcen.2013.02.047>
- 442 Frankart, C., Eullaffroy, P., & Vernet, G. (2003). Comparative effects of four herbicides on non-photochemical
 443 fluorescence quenching in *Lemna minor*. *Environmental and Experimental Botany*, 49, 159–168.
 444 www.elsevier.com/locate/envexpbot
- 445 Gebremedhin, S., Bruneel, S., Getahun, A., Anteneh, W., & Goethals, P. (2019). The Endemic Species Flock of
 446 *Labeobarbus* spp. in Lake Tana (Ethiopia) Threatened by Extinction: Implications for Conservation Management.
 447 *Water*, 11(12), 2560. <https://doi.org/10.3390/w11122560>
- 448 Gultekin, F., Ozturk, M., & Akdogan, M. (2000). The effect of organophosphate insecticide chlorpyrifos-ethyl on lipid
 449 peroxidation and antioxidant enzymes (in vitro). *Archives of Toxicology*, 74(9), 533–538.
 450 <https://doi.org/10.1007/s002040000167>
- 451 Hamid, A., Yaqub, G., Ahmed, S. R., & Aziz, N. (2017). Assessment of human health risk associated with the presence of
 452 pesticides in chicken eggs. *Food Science and Technology (Brazil)*, 37(3), 378–382. <https://doi.org/10.1590/1678-457x.11616>
- 454 Hernández, A. F., Parrón, T., Tsatsakis, A. M., Requena, M., Alarcón, R., & López-Guarnido, O. (2013). Toxic effects of
 455 pesticide mixtures at a molecular level: Their relevance to human health. *Toxicology*, 307, 136–145.
 456 <https://doi.org/10.1016/j.tox.2012.06.009>

- 457 Holden, A. v. (1972). The Effects of Pesticides on Life in Fresh Waters Author (s): Proceedings of the Royal Society of
458 London . Series B , Biological Sciences , Mar . 21 , 1972 , Vol . 180 , No . 1061 , A Discussion on Freshwater and
459 Estaurine Studie. *Biological Sciences*, 180(1061), 383–394.
- 460 Houbraken, M., Sprangers, T., de Clercq, P., Cooreman-Algoed, M., Couchement, T., de Clercq, G., Verbeke, S., &
461 Spanoghe, P. (2016). Pesticide contamination of *Tenebrio molitor* (Coleoptera: Tenebrionidae) for human
462 consumption. *Food Chemistry*, 201, 264–269. <https://doi.org/10.1016/j.foodchem.2016.01.097>
- 463 Islam, M. A., Nurul Amin, S. M., Brown, C. L., Juraimi, A. S., Uddin, M. K., & Arshad, A. (2022). Determination of the
464 Most Efficient Household Technique for the Reduction of Pesticide Residues from Raw Fish Muscles. *Foods*, 11(9).
465 <https://doi.org/10.3390/foods11091254>
- 466 Jiang, J., Zhang, C., Wang, L., Wang, X., He, H., Wu, S., & Zhao, X. (2022). Insights into the combined effects of
467 environmental concentration of difenoconazole and tebuconazole on zebrafish early life stage. *Science of The Total*
468 *Environment*, 830, 154687. <https://doi.org/10.1016/j.scitotenv.2022.154687>
- 469 Kim, L., Lee, D., Cho, H. K., & Choi, S. D. (2019). Review of the QuEChERS method for the analysis of organic pollutants:
470 Persistent organic pollutants, polycyclic aromatic hydrocarbons, and pharmaceuticals. *Trends in Environmental*
471 *Analytical Chemistry*, 22(e00063). <https://doi.org/10.1016/j.teac.2019.e00063>
- 472 Kortenkamp, A. (2007). Ten years of mixing cocktails: A review of combination effects of endocrine-disrupting chemicals.
473 In *Environmental Health Perspectives* (Vol. 115, Issue SUPPL1, pp. 98–105). <https://doi.org/10.1289/ehp.9357>
- 474 Kumari, K. (2020). Pesticides toxicity in fishes: A review. *Journal of Entomology and Zoology Studies*, 8(5), 1640–1642.
- 475 Kushwaha, M., Verma, S., & Chatterjee, S. (2016). Profenofos, an Acetylcholinesterase-Inhibiting Organophosphorus
476 Pesticide: A Short Review of Its Usage, Toxicity, and Biodegradation. *Journal of Environmental Quality*, 45(5),
477 1478–1489. <https://doi.org/10.2134/jeq2016.03.0100>
- 478 Lengai, G. M. W., Muthomi, J. W., & Mbega, E. R. (2020). Phytochemical activity and role of botanical pesticides in pest
479 management for sustainable agricultural crop production. In *Scientific African* (Vol. 7, Issue e00239). Elsevier B.V.
480 <https://doi.org/10.1016/j.sciaf.2019.e00239>
- 481 Lewis, K. A., Tzilivakis, J., Warner, D. J., & Green, A. (2016). An international database for pesticide risk assessments
482 and management. *Human and Ecological Risk Assessment*, 22(4), 1050–1064.
483 <https://doi.org/10.1080/10807039.2015.1133242>
- 484 Lozowicka, B. (2015). Health risk for children and adults consuming apples with pesticide residue. *Science of the Total*
485 *Environment*, 502, 184–198. <https://doi.org/10.1016/j.scitotenv.2014.09.026>
- 486 Macirella, R., Curcio, V., Ahmed, A. I. M., Pellegrino, D., & Brunelli, E. (2022). Effect of short-term exposure to low
487 concentration of tebuconazole: morphological, histometric and functional modifications in *Danio rerio* liver.
488 *European Zoological Journal*, 89(1), 324–338. <https://doi.org/10.1080/24750263.2022.2043469>
- 489 MacKay, D., & Fraser, A. (2000). Bioaccumulation of persistent organic chemicals: Mechanisms and models.
490 *Environmental Pollution*, 110(3), 375–391. [https://doi.org/10.1016/S0269-7491\(00\)00162-7](https://doi.org/10.1016/S0269-7491(00)00162-7)

- 491 Mahmood, A., & Malik, R. N. (2014). Human health risk assessment of heavy metals via consumption of contaminated
 492 vegetables collected from different irrigation sources in Lahore, Pakistan. *Arabian Journal of Chemistry*, 7(1), 91–
 493 99. <https://doi.org/10.1016/j.arabjc.2013.07.002>
- 494 Manuelmolina-Ruiz, J., Cieslik, E., Cieslik, I., & Walkowska, I. (2014). Determination of pesticide residues in fish tissues
 495 by modified QuEChERS method and dual-d-SPE clean-up coupled to gas chromatography–mass spectrometry.
 496 *Environmental Science and Pollution Research*, 22(1), 369–378. <https://doi.org/10.1007/s11356-014-3361-2>
- 497 Marchand, M. J., Pieterse, G. M., & Barnhoorn, I. E. J. (2010). Sperm motility and testicular histology as reproductive
 498 indicators of fish health of two feral fish species from a currently DDT sprayed area, South Africa. *Journal of Applied*
 499 *Ichthyology*, 26(5), 707–714. <https://doi.org/10.1111/j.1439-0426.2010.01558.x>
- 500 Matos, P., Fontainhas-Fernandes, A., Peixoto, F., Carrola, J., & Rocha, E. (2007). Biochemical and histological hepatic
 501 changes of Nile tilapia *Oreochromis niloticus* exposed to carbaryl. *Pesticide Biochemistry and Physiology*, 89(1),
 502 73–80. <https://doi.org/10.1016/j.pestbp.2007.03.002>
- 503 Mengistie, B. T., Mol, A. P. J., & Oosterveer, P. (2016). Private Environmental Governance in the Ethiopian Pesticide
 504 Supply Chain: Importation, Distribution and Use. *NJAS - Wageningen Journal of Life Sciences*, 76, 65–73.
 505 <https://doi.org/10.1016/j.njas.2015.11.005>
- 506 Mhadhbi, L., & Beiras, R. (2012). Acute toxicity of seven selected pesticides (alachlor, atrazine, dieldrin, diuron,
 507 pirimiphos-methyl, chlorpyrifos, diazinon) to the marine fish (turbot, *Psetta maxima*). *Water, Air, and Soil Pollution*,
 508 223(9), 5917–5930. <https://doi.org/10.1007/s11270-012-1328-9>
- 509 Müller, R., Shinn, C., Waldvogel, A. M., Oehlmann, J., Ribeiro, R., & Moreira-Santos, M. (2019). Long-term effects of
 510 the fungicide pyrimethanil on aquatic primary producers in macrophyte-dominated outdoor mesocosms in two
 511 European ecoregions. *Science of the Total Environment*, 665, 982–994.
 512 <https://doi.org/10.1016/j.scitotenv.2019.02.050>
- 513 Olette, R., Couderchet, M., Biagiante, S., & Eullaffroy, P. (2008). Toxicity and removal of pesticides by selected aquatic
 514 plants. *Chemosphere*, 70(8), 1414–1421. <https://doi.org/10.1016/j.chemosphere.2007.09.016>
- 515 Patnaik, L., & Patra, ; (2006). Haematopoietic Alterations induced by Carbaryl in *Clarias batrachus* (LINN). *J. Appl. Sci.*
 516 *Environ. Mgt.*, 10(3), 5–7. www.bioline.org.br/ja
- 517 Polat, A., Polat, S., Simsek, A., Kurt, T. T., & Ozyurt, G. (2018). Pesticide residues in muscles of some marine fish species
 518 and seaweeds of Iskenderun Bay (Northeastern Mediterranean), Turkey. *Environmental Science and Pollution*
 519 *Research*, 25(4), 3756–3764. <https://doi.org/10.1007/s11356-017-0756-x>
- 520 Rasool, S., Rasool, T., & Gani, K. M. (2022). A review of interactions of pesticides within various interfaces of intrinsic
 521 and organic residue amended soil environment. In *Chemical Engineering Journal Advances* (Vol. 11). Elsevier B.V.
 522 <https://doi.org/10.1016/j.cej.2022.100301>
- 523 Ribera, D., Narbonne, J. F., Arnaud, C., & Saint-Denis, M. (2001). Biochemical responses of the earthworm *Eisenia fetida*
 524 *andrei* exposed to contaminated artificial soil, effects of carbaryl. *Soil Biology & Biochemistry*, 33, 1123–1130.
 525 www.elsevier.com/locate/soilbio

- 526 Rosseland, B. O., Massabuau, J.-C., Grimalt, J., Hofer, R., Lackner, R., Raddum, G., Rognerud, S., & Vives, I. (2001).
 527 *Fish Ecotoxicology, The EMERGE Fish Sampling Manual for Live Fish. The EMERGE project (European mountain*
 528 *lake Ecosystem: regionalisation, diaGnostic and socio-economics valuation)*. 1–7.
- 529 Sabzevari, S., & Hofman, J. (2022). A worldwide review of currently used pesticides' monitoring in agricultural soils. In
 530 *Science of the Total Environment* (Vol. 812, Issue 152344). Elsevier B.V.
 531 <https://doi.org/10.1016/j.scitotenv.2021.152344>
- 532 Sarkar, S., Dias Bernardes Gil, J., Keeley, J., Möhring, N., & Jansen, K. (2021). *The use of pesticides in developing*
 533 *countries and their impact on health and the right to food*: study. (the European union).
 534 <https://doi.org/10.2861/28995>
- 535 Araújo, C. V. M., Shinn, C., Müller, R., Moreira-Santos, M., Shinn, C., Espíndola, E. L., & Ribeiro, R. (2012). Aquatic
 536 ecotoxicity of the fungicide pyrimethanil: Effect profile under optimal and thermal stress conditions. *Environmental*
 537 *Pollution*, 168, 161–169. <https://doi.org/10.1016/j.envpol.2012.04.020>
- 538 Sharma, A., Kumar, V., Shahzad, B., Tanveer, M., Sidhu, G. P. S., Handa, N., Kohli, S. K., Yadav, P., Bali, A. S., Parihar,
 539 R. D., Dar, O. I., Singh, K., Jasrotia, S., Bakshi, P., Ramakrishnan, M., Kumar, S., Bhardwaj, R., & Thukral, A. K.
 540 (2019). Worldwide pesticide usage and its impacts on ecosystem. In *SN Applied Sciences* (Vol. 1, Issue 11). Springer
 541 Nature. <https://doi.org/10.1007/s42452-019-1485-1>
- 542 Sibbing, F. A., & Nagelkerke, L. A. J. (2001). Resource partitioning by Lake Tana barbs predicted from fish morphometrics
 543 and prey characteristics. *Reviews in Fish Biology and Fisheries*, 10(4), 393–437.
 544 <https://doi.org/10.1023/A:1012270422092>
- 545 Sifakis, S., Androutsopoulos, V. P., Tsatsakis, A. M., & Spandidos, D. A. (2017). Human exposure to endocrine disrupting
 546 chemicals: effects on the male and female reproductive systems. *Environmental Toxicology and Pharmacology*, 51,
 547 56–70. <https://doi.org/10.1016/j.etap.2017.02.024>
- 548 Smith, L. W. (1982). Wildlife toxicity studies with Oxamyl. *Environmental Toxicology and Chemistry*, 1, 157–165.
- 549 Srivastava, P., Singh, A., & Pandey, A. K. (2016). Pesticides toxicity in fishes: Biochemical, Physiological and Genotoxic
 550 aspects. *Biochem. Cell. Arch.*, 16(2), 199–218. www.connectjournals.com/bca
- 551 Teklu, B. M., Adriaanse, P. I., ter Horst, M. M. S., Deneer, J. W., & van den Brink, P. J. (2015). Surface water risk
 552 assessment of pesticides in Ethiopia. *Science of the Total Environment*, 508, 566–574.
 553 <https://doi.org/10.1016/j.scitotenv.2014.11.049>
- 554 Teng, M., Zhu, W., Wang, D., Qi, S., Wang, Y., Yan, J., Dong, K., Zheng, M., & Wang, C. (2018). Metabolomics and
 555 transcriptomics reveal the toxicity of difenoconazole to the early life stages of zebrafish (*Danio rerio*). *Aquatic*
 556 *Toxicology*, 194, 112–120. <https://doi.org/10.1016/j.aquatox.2017.11.009>
- 557 Tesfahun, A., & Alebachew, S. (2023). Food and feeding habits of the Nile tilapia *Oreochromis niloticus* (Linnaeus, 1758)
 558 from Ribb reservoir, Lake Tana sub-basin, Ethiopia. *Cogent Food & Agriculture*, 9(1).
 559 <https://doi.org/10.1080/23311932.2023.2212457>
- 560 Tournas, V. H. (2005). Spoilage of vegetable crops by bacteria and fungi and related health hazards. *Critical Reviews in*
 561 *Microbiology*, 31(1), 33–44. <https://doi.org/10.1080/10408410590886024>

- 562 US EPA. (2003). *Revised EFED Risk Assessment of Carbaryl in Support of the Reregistration Eligibility Decision RED*.
563 <https://www3.epa.gov/pesticides/endanger/litstatus/effects/carb-riskass.pdf>
- 564 US EPA. (2007). *Pesticide fact sheet for Flazasulfuron*.
565 https://www3.epa.gov/pesticides/chem_search/reg_actions/registration/fs_PC-119101_17-May-07.pdf
- 566 Vignet, C., Cappello, T., Fu, Q., Lajoie, K., de Marco, G., Clérandeau, C., Mottaz, H., Maisano, M., Hollender, J., Schirmer,
567 K., & Cachot, J. (2019). Imidacloprid induces adverse effects on fish early life stages that are more severe in Japanese
568 medaka (*Oryzias latipes*) than in zebrafish (*Danio rerio*). *Chemosphere*, 225, 470–478.
569 <https://doi.org/10.1016/j.chemosphere.2019.03.002>
- 570 Vonesh, J. R., & Kraus, J. M. (2009). Pesticide alters habitat selection and aquatic community composition. *Oecologia*,
571 160(2), 379–385. <https://doi.org/10.1007/s00442-009-1301-5>
- 572 WHO and FAO. (2019). *Global situation of pesticide management in agriculture and public health*.
573 <http://www.fao.org/3/ca7032en/ca7032en.pdf>
- 574 Witczak, A. (2009). Effect of heat treatment on organochlorine pesticide residues in selected fish species. *Pol. J. Food*
575 *Nutr. Sci*, 59(3), 231–235. <http://journal.pan.olsztyn.pl>
- 576 Worku, S. A., Dubale, Y. M., Woldemariam, E. D., & Eklo, O. M. (2022). Levels of organochlorine pesticides in five
577 species of fish from Lake Ziway, Ethiopia. *Scientific African*, e01252. <https://doi.org/10.1016/j.sciaf.2022.e01252>
- 578 Yohannes, Y. B., Ikenaka, Y., Nakayama, S. M. M., Mizukawa, H., & Ishizuka, M. (2017). DDTs and other organochlorine
579 pesticides in tissues of four bird species from the Rift Valley region, Ethiopia. *Science of the Total Environment*,
580 574, 1389–1395. <https://doi.org/10.1016/j.scitotenv.2016.08.056>
- 581 Yohannes, Y. B., Ikenaka, Y., Saengtienchai, A., Watanabe, K. P., Nakayama, S. M. M., & Ishizuka, M. (2014).
582 Concentrations and human health risk assessment of organochlorine pesticides in edible fish species from a Rift
583 Valley lake-Lake Ziway, Ethiopia. *Ecotoxicology and Environmental Safety*, 106, 95–101.
584 <https://doi.org/10.1016/j.ecoenv.2014.04.014>