

REVIEW

Open Access



# Effect of virtual reality-based upper limb training on activity of daily living and quality of life among stroke survivors: a systematic review and meta-analysis

Diriba Dereje Olana<sup>1,2\*</sup>, Teklu Gemechu Abessa<sup>2,3</sup>, Dheeraj Lamba<sup>4</sup>, Lisa Tedesco Triccas<sup>2,5</sup> and Bruno Bonnechere<sup>2,6,7</sup>

## Abstract

**Background** Stroke is a leading cause of disability worldwide, significantly impairing upper limb (UL) function and reducing patients' ability to perform activities of daily living (ADL) and quality of life (QoL). Virtual reality (VR) has emerged as a promising tool for UL rehabilitation, offering immersive and engaging environments for motor recovery. However, the effectiveness of VR, its integration with conventional therapy, and their efficacy across different stroke recovery stages remain unclear. Therefore, this systematic review and meta-analysis aimed to evaluate the effectiveness of VR-based UL interventions in improving ADL and QoL among stroke survivors.

**Method** This study adhered to PRISMA guidelines and was registered on PROSPERO (CRD42023426256). A systematic search of PubMed, Scopus, and Web of Science identified randomized controlled trials (RCTs) published in English. Inclusion criteria focused on studies using immersive VR (IVR) and non-immersive VR (NIVR) interventions to assess ADL and QoL in stroke survivors. Data extraction and quality assessment were performed independently by two reviewers using the PEDro scale to assess quality. Meta-analyses were conducted to determine the efficacy. Subgroup analyses were performed to compare IVR and NIVR, VR combined with conventional therapy versus standalone VR, and potential differences between stroke recovery stages.

**Result** Thirty RCTs, representing 1,661 participants, were included. Overall, VR interventions significantly improved ADL (SMD = 0.27, 95% CI [0.11; 0.43],  $p < 0.001$ ) and QoL (SMD = 0.94 [0.09; 1.79],  $p = 0.035$ ) compared to conventional therapy. IVR demonstrated superior outcomes for ADL compared to NIVR (SMD = 0.54 [0.13; 0.95] Vs. 0.17 [0.02; 0.36],  $p = 0.03$ ). Subacute stroke survivors exhibited the most significant gains in ADL (SMD = 0.52 [0.16; 0.88],  $p = 0.004$ ), compared to chronic (SMD = 0.05 [-0.36; 0.46]) or acute patients (SMD = 0.08 [-0.11; 0.27]).

**Conclusion** VR interventions, particularly IVR and VR combined with conventional therapy, significantly enhance ADL and QoL in stroke survivors with moderate certainty of evidence. These findings underscore the value of VR in rehabilitation, especially during the subacute phase, but highlight the need for further research into long-term effects and implementation in low-resource settings.

\*Correspondence:  
Diriba Dereje Olana  
[diriba.dereje@ju.edu.et](mailto:diriba.dereje@ju.edu.et)

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

**Keywords** Stroke, Upper extremity, Exercise, Independent living, psychological wellbeing, Rehabilitation, Immersive technology

## Introduction

As the prevalence of chronic diseases is rising globally, there is a great need for acute and long-term rehabilitation services [1, 2]. Stroke, the second most common cause of morbidity worldwide, significantly impairs individuals' lives and poses a rising prevalence. According to the latest Global Burden of Disease report, there were 12.2 million new stroke cases and 101 million existing cases globally, underscoring the urgent need for immediate action to address this multifaceted public health challenge [3, 4]. Stroke survivors commonly experience upper limb (UL) impairment, with 50–80% affected in the acute, 40–60% in the subacute, and 30–50% in the chronic phase [5, 6]. Decreased UL function is a common post-stroke impairment, restricting activities of daily living (ADL), with around 30% requiring assistance, and also negatively impacting quality of life (QoL) for up to two-thirds of stroke patients [7, 8].

Targeting improvement in patients' ability to carry out ADL independently are main objectives of UL rehabilitation programs and a main driver of independence and QoL. One key factor for a successful rehabilitation is the dosage. The total amount of rehabilitation, refers to the cumulative duration of rehabilitation interventions administered to a patient, typically measured in minutes and calculated from the frequency and duration of therapy sessions was the strongest predictors of UL motor function recovery after stroke, with high-intensity having better outcome [9–11]. However it is to note that, currently, the need for rehabilitation is most of the time unmet and the patients do not benefit from enough rehabilitation, especially in Low and Middle-Income Countries (LMICs), leading to suboptimal and poor recovery [12]. Therefore, there is a huge need to develop and implement additional complementary or alternative solutions to the current rehabilitation services provided to people with stroke [4, 5]. Virtual Reality (VR) based interventions have the potential to positively improve ADL and QOL which could be the most promising substitute or addition to the current global stroke rehabilitation programs [13, 14]. VR is a computer-human interface that allows users to interact with computers, generating virtual environments where users can perform different tasks in real time [15].

VR offers numerous benefits, such as simulating real environments for training, enabling 3D visualization and modification of the body (embodiment), and enhancing motivation through fun [16, 17]. Immersive virtual reality (IVR) fully immerses users in a simulated environment using head-mounted displays, minimizing awareness of

their physical surroundings. However, non-immersive VR (NIVR) relies on traditional displays like smartphone screens or monitors. Semi-Immersive VR allows interaction with virtual content while informing users of the real world, often utilizing large screens or projection systems [18]. VR therapy has improved compliance by enhancing patient engagement in rehabilitation [19], leading to greater exercise adherence while reducing physical and mental fatigue [20], even though it may cause cybersickness in some patients [21]. Few systematic reviews focused on the effect of game-based VR on upper limb impairment and function without considering ADL or QoL. They did not comprehensively examine VR's effectiveness across stroke recovery stages [22–26]. VR, as an adjunct to conventional therapy, is reported to be safe for stroke since it significantly enhances the quality of life compared to standard rehabilitation, specifically with immersive VR [27]. Current evidence demonstrates the superiority of VR, both with or without conventional therapy, over conventional therapy alone. However, there is currently no evidence regarding the comparative effectiveness of combining conventional treatment compared to solely VR [28–30]. Furthermore, the optimal timing for initiating VR interventions after a stroke, as well as identifying the most effective types of VR interventions for upper limb rehabilitation, remain unclear. Therefore, the primary objective was to examine the effect of upper-limb VR training on ADL and QoL in stroke survivors. Secondary objectives included assessing the impact of total rehabilitation time on ADL and QoL; comparing the effectiveness of IVR and NIVR, and evaluating the added benefit of combining conventional rehabilitation with VR.

## Methods

This systematic review reporting was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [31], and the protocol was registered on PROSPERO (CRD42023426256).

## Search strategy

The literature search was conducted in PubMed, Web of Science, and Scopus by two independent researchers. Key search terms included “stroke”, “virtual reality”, “upper limb”, “quality of life” and “activity of daily living” using Boolean operators (AND/OR). The search term was (“virtual reality” OR “video game” OR “immersive” OR “non-immersive”) AND (“upper limb” OR “upper extremity”) AND (exercise OR training OR rehabilitation

OR “quality of life\*” OR “activity of daily living”) AND (stroke OR “cerebrovascular disorder”) for the three databases (the complete search strategy is presented in Supplementary Material 1). The search was completed on February 4, 2025, and was not limited based on the year of publication given the recent development of this technology. Additionally, remaining references were manually retrieved using the snowball method from previous relevant reviews and literatures.

### Study selection

The identified articles were imported into Rayyan [32], where duplicates were eliminated. Two reviewers independently reviewed titles and abstracts to select eligible articles for the review. Both reviewers then independently examined the full-text articles. Final decisions on each article were reached through consensus, with a third reviewer consulted in cases of disagreement. The study selection process done by using a predefined inclusion criteria based on the Population Intervention Comparison Outcome Study Design (PICO) method [33].

- Population: Participants aged 18 years or older with any type of upper limb stroke. Studies involving healthy individuals or non-stroke patients were excluded.
- Intervention: Studies utilized VR for upper limb intervention, either of IVR and NIVR, as well as VR combined with neuromuscular stimulation and robotic assistance.
- Comparison: Conventional rehabilitation without VR intervention. Conventional rehabilitation that utilized any feature of VR as a comparison group were excluded.
- Outcome: Studies measured either activity of daily living (by Barthel Index, BI; Modified Barthel Index, MBI; or Functional Independence Measure, FIM) or quality of life (by EQ VAS, Euro QoL Visual Analogue Scale; SS QoL, Stroke Specific Quality of Life; SIS, Stroke Impact Scale; or SF-36, Short-Form 36 Health Survey). Studies that did not measure at least one of the above outcomes were excluded.
- Study design: Randomized controlled trials (RCTs).

The analysis focused exclusively on peer-reviewed articles published in English. This language restriction was implemented to maintain consistency in data extraction, analysis, and interpretation, given the researchers' language proficiency. Limiting the scope to English-language publications also mitigated potential bias introduced by language barriers. Protocols, reviews, conference proceedings, theses, letters, unpublished studies, and studies involving nonhuman subjects were excluded.

### Methodological quality assessment

The studies' methodological quality was evaluated using the PEDro scale. The strength of the PEDro scale lies in its comprehensive criteria for quality assessment, specifically designed for RCTs only, making it highly suitable to use in this systematic review study. Furthermore, this scale provides a clear structured format that is not complex to apply while it is comprehensive enough. The PEDro scale has 11 items scale, and each satisfied item contributes one point to the total PEDro score. Item one is omitted for score calculation so that the score ranges from zero to ten points. Total PEDro scale scores of zero to three are considered 'poor quality', four to five 'moderate quality', and six to ten 'good quality' for each RCT study, and this category was used in this study [34–36]. Additionally, the Cochrane Risk of Bias tool (RoB 2) was also used to assess the risk of bias that may have been under evaluated using the PEDro scale [37]. Two researchers conducted the methodological quality assessment. In cases of disagreement, a third researcher was consulted, and a decision was reached upon agreement. PEDro scale was conducted to assess the methodological quality [36], to further determine the level of evidence associated with these new interventions using the GRADE system. GRADE summary of evidence and recommendation was conducted by using GRADEPro software [38, 39].

### Data extraction and synthesis

Two reviewers independently extracted data from the included studies. A third independent reviewer verified the accuracy of the extracted data. Discussions and consultations with the primary author resolved discrepancies and inconsistencies as needed. The extracted information from the included studies encompassed age, gender, authors' names, countries, year of publication, type of intervention, type of stroke, VR type, study design, sample size, type and name of the devices, outcomes on ADL and QoL, and main characteristics of training and interventions. Regarding outcome data, if the single outcome measured by two different tools, we included both outcome in meta-analysis.

### Statistical analysis

The potential effect of the VR for UL stroke rehabilitation was examined with meta-analysis. The measure of treatment effect was the standardized mean difference effect size (standardized mean difference (SMD)), defined as the between-group difference in mean values divided by the pooled SD computed using the Hedge's *g* method. The SMD is categorized as small (0.2), moderate (0.5), or large (0.8), considering the effect size between the two group means in terms of standard deviation units [40]. Change from baseline data was computed for both

control and intervention from pre and post-sample size, mean, and standard deviation by using the Wan et al. formula [41]. If several tests were used to evaluate ADL or QoL in the same study, the results of the different tests were combined, using weighted mean, to produce a single SMD according to Cochrane's recommendation [42]. If the median and range were reported instead of the mean and standard deviation, these values were estimated using the methods developed by Luo et al. and Wan et al. [41, 43]. We assessed the heterogeneity in stratified analyses by type of VR and stroke classification using  $I^2$ , computed using a restricted maximum likelihood method (REML). We categorized  $I^2$  using 25%, 50%, and 75%, representing low, moderate, and high heterogeneity [44]. To deal with high or moderate heterogeneity, we used random-effect models and presented forest plots [45]. We checked for publication bias using a funnel plot [50] and Egger's test for the intercept was applied to check the asymmetry [51]. Random-effects meta-regression analysis quantified the association of changes in ADL and QoL and the total amount of training (number of sessions multiplied by the duration of one session). Studies were weighted by the inverse of the sum of the within- and between-study variance. Also, we reported stroke classification as acute, subacute, and chronic. Sensitivity analysis was performed using the leave-one-out method to assess the robustness of the meta-analysis results. This approach involved systematically removing one study at a time from the analysis and recalculating the pooled effect estimate. This process helped us to identify whether any single study had a disproportionate influence on the overall results and evaluate the stability of the findings. Finally, a GRADE assessment of the level of evidence and recommendations was performed by systematically identifying the clinical question, population, interventions, and outcomes. The level of evidence for each outcome was based on factors such as risk of bias, imprecision, inconsistency, indirectness, and publication bias. Then, GRADE recommendations were subsequently categorized as high, moderate, or low, considering the balance of benefits and harms, the level of evidence, and other factors [46]. Statistical analyses were performed at an overall significance level of 0.05 in the R program (version 4.4.1).

## Results

### Search results

A total of 3030 research papers were identified from the three databases with 1294 records were filtered using the titles and abstracts. After, 673 full texts were checked and 30 RCTs were finally included in this analysis. The complete flow chart of study selection is presented in Fig. 1.

### Characteristics of the participants

1661 patients were included in this review, with a predominance of male (60%) compared to women. The mean age was  $59.8 \pm 5.4$  years (complete socio-demographic characteristics of the patients are presented in Table 1).

Concerning stroke's stage, the most frequently assessed population was subacute stroke survivor with 10 studies [47–56]. Four studies involved people in the acute stage [57–60], and in the chronic phase [61–64]. In 9 studies, a combination of acute, subacute or chronic stages were involved [58, 65–72], the stage was not clearly specified in 3 studies [73–75]. Seventeen studies reported the ratios of ischemic to hemorrhagic strokes, with ischemic strokes being the majority in all cases [48–50, 54–57, 61, 64, 68–73, 76, 77]. The breakdown of ischemic versus hemorrhagic stroke types was not reported in the other studies.

Twenty studies were undertaken in various Asian countries [47–51, 53, 54, 56, 58, 59, 61, 64–67, 72–75], accounting for the majority of the studies conducted. Europe accounted for the second-highest number of studies with 9 studies [55, 57, 60, 68–71, 76]. Both North [52] and South America [62] each contributed one study. In addition, there were no studies conducted in Africa.

### Quality of the study


The quality of the research included in the current review exhibited a range of scores on the PEDro scale, spanning from 5 to 8 out of a maximum of 10 points with mean score of  $6.27 \pm 0.83$ , indicating moderate to high quality (individual results are presented in Table 1 and Supplementary Material 2). Figure 2 presents the RoB 2 score of the different individuals studies. Subject blinding, therapist blinding, and intention to treat were commonly unmet in the PEDro scale. Based on Rob2 tool for risk of bias assessment, the main cause of bias is deviation from intended interventions.

### Type of intervention

Most of the included studies, 77% ( $n = 23$ ) [50–54, 57–60, 62–68, 70–72, 74–77] used NIVR, while the 7 others used IVR [47–49, 56, 59, 61, 69]. More recent studies showed a higher prevalence of IVR use (Table 1). Another important aspect is how VR was integrated in the care: VR was either used alone or in combination with conventional therapy. The majority of the studies 66% ( $n = 20$ ), combined VR with conventional rehabilitation (Table 1).

### Immersive vs. non-immersive

The review outlined comprehensive VR exercises and rehabilitation programs for the ULs. Among these were VR video games that allowed patients to reach, grab, and manipulate things in interactive scenarios, such as “Traffic Control [53, 54],” “Mouse Mayhem [54],” and “Balloon



**Fig. 1** PRISMA flow diagram of study screening and selection procedures

Buster [53, 54]” Playing games like table tennis [53], bowling [47, 53, 78], bingo [52], reaching [60, 73, 76], gripping [48], moving [47, 78], and releasing balls [56, 65], and other tabletop activities [56, 74, 78]. Patients also practiced specific movement patterns like flexion, extension, and abduction through targeted exercises [64, 69, 72]. IVR games such as “Underwater Fire” [72, 75] and “Bug Hunter” [72, 75] offered engaging landscapes that test one’s ability to lift, reach, and coordinate their hands and fingers. Other exercises concentrated on movement speed, duration, and difficulty. Some systems incorporated advanced technology for rehabilitation, such as motion tracking and exoskeletons, to facilitate accurate

and personalized instruction in reaching, gripping, and manipulating objects. In general, the data showed various and creative ways of VR being used to provide efficient and engaging UL therapy through interactive tasks, games, and immersive experiences (Table 1).

#### Virtual reality devices

The studies included a wide range of VR hardware and software platforms in programs for the rehabilitation of ULs. These include VR systems connected to well-known gaming consoles like Xbox Kinect (30%,  $n = 9$ ) [53–55, 58, 61, 64, 67, 71, 76] and the Nintendo Wii (10%,  $n = 3$ ) [49, 52, 62], enabling patients to participate in interactive,



**Table 1** Characteristics of the included studies

Authors	Country of the study	Stroke stages	Stroke types in I/H ratio	Sex in M/F ratio	Mean age in years	Sample Size in C/E ratio	Exercise or training given	Types of VR	VR device used	Total PEDro score out of ten	Cochrane risk of Bias Over all (RoB2)
Laffonte et al.,2020 [55]	France	SA	C = 16/10 E = 22/3	C = 15/11 E = 16/9	C = 55.8 ± 14.0 E = 60.8 ± 14.1	C = 26 E = 25	Passive and active mobilization of limbs	NIVR	VR based video game device	8	Low
Afsar et al., 2018 [54]	Turkey	SA	C = 12/4 E = 15/4	C = 8/8 E = 12/7	C = 63.44 ± 15.73 E = 69.42 ± 8.55	C = 16 E = 19	Mouse Mayhem, traffic control, Balloon Buster and Mather sizing games	NIVR	VR based Xbox Kinect Game system	6	High
Gueye et al.,2021 [60]	Czech Republic	A	NS	C = 15/10 E = 14/11	C = 68.12 ± 11.97 E = 66.56 ± 12.26	C = 25 E = 25	Reach and grasping exercise	NIVR	Arneo spring upper limb exoskeleton	5	High
Leng et al.,2022 [53]	China	SA	NS	C = 8/18 E = 6/25	C = 59.12 ± 11.62 E = 59.25 ± 10.70	C = 26 E = 31	Balloon buster, table tennis, bowling and traffic control games	NIVR	Microsoft Xbox 360 Kinect	6	Low
Keskin et al.,2020 [67]	Turkey	M	NS	C = 6/8 E = 5/7	C = 63.6 ± 7.1 E = 63.6 ± 9.2	C = 12 E = 12	Difficulty, duration and speed exercise	NIVR	VR based with leap motion camera device	7	High
Rong et al.,2021 [74]	China	NS	NS	C = 14/6 E = 17/3	C = 62.30 ± 13.10 E = 56.25 ± 12.29	C = 20 E = 20	Object reaching, grasping and placing	NIVR	Arm rehabilitation robot and a customized camera based MVR	6	High
Saposnik et al.,2016 [80]	Canada	SA	NS	NS	NS	C = 70 E = 71	Playing cards, bingo, Jenga or ball games	NIVR	VR based Nintendo Wii gaming	7	Some concerns
Choi et al.,2014 [49]	South Korea	SA	C = 8/2 E = 6/4	C = 5/5 E = 5/5	C = 64.70 ± 11.3 E = 64.30 ± 10.3	C = 10 E = 10	Stretching and strengthening exercise	NIVR	Nintendo Wii based commercial game with VR	7	High
Shin et al.,2014 [75]	South Korea	M	NS	C = 3/4 E = 5/4	C = 46.6 ± 5.8 E = 52.0 ± 11.9	C = 7 E = 9	Underwater fire, bug hunter, and roller coaster game	NIVR	A task specific interactive game based VR	5	High
Kiper et al.,2013 [71]	Italy	M	C = 11/10 E = 13/10	C = 15/6 E = 14/9	C = 21 E = 23	C = 21 E = 23	Grasping ball, disk or glass	NIVR	Virtual reality rehabilitation system	5	High
Lee et al.,2016 [64]	South Korea	Ch	C = 5/8 E = 5/8	C = 10/3 E = 8/5	C = 69.92 ± 7.18 E = 66.46 ± 7.26	C = 13 E = 13	Flexion, diagonal extension and diagonal flexion	NIVR	VR with motion capture sensor (KINECT)	5	Some concerns
Long et al.,2020 [66]	China	M	NS	C = 16/11 E = 18/7	C = 54.11 ± 14.81 E = 53.28 ± 15.30	C = 27 E = 25	Arm flexion, abduction, gold coin picking, and cross and mixed training	NIVR	VR based game system with KINECT	7	Some concerns
Yin et al.,2014 [58]	Singapore	A	NS	C = 10/2 E = 6/5	C = 62(only mean) E = 56(only mean)	C = 12 E = 11	NS	NIVR	Non immersive VR based game	6	High
Park et al.,2019 [73]	South Korea	NS	C = 5/7 E = 8/5	C = 7/5 E = 8/5	C = 51.5 ± 16.7 E = 53.5 ± 13.0	C = 12 E = 13	Point to point reaching, circle drawing, and free exploration	NIVR	VR based planar motion	7	High
Turolla et al.,2013 [76]	Italy	M	C = 82/31 E = 188/75	C = 72/41 E = 157/106	C = 65.4 ± 12.5 E = 60.2 ± 14.3	C = 113 E = 263	Reaching movement, putting glass on shelf, and holding objects	NIVR	VR rehabilitation system with 3D motion tracking	5	High
Piron et al., 2014 [63]	Italy	Ch	NS	C = 12/11 E = 17/10	C = 62.2(9.75) E = 58.8 (8.3)	C = 23 E = 23	Touching, recognizing different arm position, and follow trajectories	NIVR	VR based reinforced feedback (RFVE)	8	High

**Table 1** (continued)

Authors	Country of the study	Stroke stages	Stroke types in I/H ratio	Sex in M/F ratio	Mean age in years	Sample Size in C/E ratio	Exercise or training given	Types of VR	VR device used	Total PEDro score out of ten	Cochrane risk of Bias Over all (RoB2)
Cameirão et al., 2011 [57]	Spain	A	C=6/2 E=6/2	C=4/4 E=4/4	C=65.22±11.46 E=58.77±11.42	C=8 E=8	Hitting, grasping, and placing	NIVR	VR based rehabilitation gaming	6	Some concerns
Ballester et al., 2016 [70]	Spain	M	C=6/3 E=8/1	C=8/1 E=7/2	C=54.80 (12.00) E=63.40 (9.40)	C=9 E=9	NS	NIVR	Rehabilitation gaming system	7	High
Zheng et al., 2015 [50]	China	SA	C=35/19 E=834/24	C=33/21 E=35/23	C=66.2±13.1 E=65.4±13.5	C=54 E=58	NS	NIVR	Transcranial magnetic stimulation with VR	6	High
Shin et al., 2016 [72]	South Korea	NS	C=14/8 E=15/7	C=17/5 E=19/5	C=54.67±13.4 E=53.37±11.8	C=22 E=24	Forearm pronation, wrist flexion, and finger flexion	NIVR	VR based RAPEL SMART glove	6	High
Rodríguez-Hernández et al., 2021 [68]	Spain	M	C=21/4 E=18/2	C=18/5 E=17/3	C=63.6±12.2 E=62.6±13.5	C=23 E=20	Resistant with ball, moving objects on table, and elevation	NIVR	VR with 3D Tutor	5	High
Ribeiro et al., 2015 [62]	Brazil	Ch	NS	C=10/5 E=9/6	C=52.8 (8.6) E=53.7 (6.1)	C=15 E=15	Transforming objects	NIVR	VR based Nintendo Wii	7	Low
Ali et al., 2024 [65]	India	M	C=47/9 E=53/11	C=41/15 E=50/14	C=57.7±10.9 E=54.4±11.7	C=56 E=64	Delicious dosa, stride the mosquito, champion cleaner, bouncing ball and fresh fruit catcher	NIVR	VR-based Arm Able TM device	7	Low
Mekbib et al., 2021 [56]	China	SA	C=8/3 E=9/3	C=8/3 E=9/3	C=61.00±7.69 E=52.17±13.26	C=11 E=12	top tabletop ball grasping, transporting, and releasing	IVR	VR based mounted head-mounted display with motion tracking	7	High
Huang et al., 2023 [48]	China	SA	C=18/2 E=19/1	C=11/9 E=13/7	C=65.05±6.14 E=63.3±14.32	C=20 E=20	Gripping and selective finger movement, gross movement, strength, and stretching	IVR	Immersive VR device not specified	7	Some concerns
Ogun et al., 2019 [61]	Turkey	Ch	C=32/0 E=33/0	C=23/9 E=28/5	C=59.75±8.07 E=61.48±10.92	C=32 E=33	A cube handling, decorating the tree with leaves, drumming, and picking up vegetables games	IVR	based VR-based game with leap motion tracking	6	High
Choi et al., 2021 [51]	South Korea	SA	NS	C=6/6 E=5/7	C=61.58±9.99 E=63.00±10.02	C=12 E=12	Visual tracking, writing, drawing, and puzzling	IVR	Ocular rift Dk2 and leap motion	7	Low
Sip et al., 2023 [69]	Poland	M	C=10/0 E=10/0	NS	C=59.2±4.34 E=54.9±9.38	C=10 E=10	Flexion, opening and closing fist, spreading and bringing together	IVR	Oculus Quest VR device	7	High
Amin et al., 2024 [47]	Pakistan	SA	NS	C=18/8 E=16/10	C=49.8±9.9 E=51.8±12.9	C=26 E=26	Hitting rolling balls, grasping balls, and swapping hand position	IVR	Oculus Quest 2 VR device	7	Low
Kwon et al. 2012 [59]	South Korea	A	NS	C=5/8 E=8/5	C=57.92±12.32 E=57.15±15.42	C=13 E=13	Reaching and lifting at different angles	IVR	Immersive VR game with video capture	5	High

A = Acute, SA = Sub-acute, Ch = Chronic, M = Mixed, NS = Not Specified, I = Ischemic, H = Hemorrhagic, C = Control, E = Experimental, M = Male, F = Female, and NIVR = Non-Immersive Virtual Reality

	Risk of bias domains					
	D1	D2	D3	D4	D5	Overall
Laffonte et al.,2020	+	+	+	+	+	+
Afsar et al., 2018	+	X	X	+	+	X
Gueye et al.,2021	X	X	+	+	+	X
Mekbib et al.,2021	+	X	+	+	+	X
Leng et al.,2022	+	+	+	+	+	+
Keskin et al.,2020	+	X	+	+	+	X
Rong et al.,2021	X	X	+	-	+	X
Saposnik et al.,2016	+	-	+	+	+	-
Choi et al.,2014	-	X	+	+	+	X
Shin et al.,2014	X	X	+	+	+	X
J.S. Kwon et al. 2012	X	X	+	+	+	X
Kiper et al.,2013	X	X	+	+	+	X
Lee et al.,2016	+	-	+	+	+	-
Long et al.,2020	+	-	+	+	+	-
Yin et al.,2014	-	X	+	+	+	X
Huang et al.,2023	+	-	+	-	+	-
Ogun et al.,2019	X	X	X	+	+	X
Choi et al.,2021	+	+	+	+	+	+
Park et al.,2019	X	-	X	-	+	X
Turolla et al.,2013	X	X	+	+	+	X
Piron et al., 2010	+	-	+	-	+	-
Cameirao et al.,2011	-	+	+	+	+	-
Ballester et al., 2016	X	-	+	+	+	X
Zheng et al., 2015	-	X	+	+	+	X
Amin et al., 2024	+	+	+	+	+	+
Shin et al., 2016	X	X	X	+	+	X
Sip et al., 2023	X	+	X	+	+	X
Rodríguez-Hernández et al.,2021	-	X	X	+	+	X
Ribeiro et al., 2015	+	+	+	+	+	+
Ali et al., 2024	+	+	+	+	+	+

Study

Domains:  
D1: Bias arising from the randomization process.  
D2: Bias due to deviations from intended intervention.  
D3: Bias due to missing outcome data.  
D4: Bias in measurement of the outcome.  
D5: Bias in selection of the reported result.

Judgement  
X High  
- Some concerns  
+ Low

**Fig. 2** : Risk of bias assessment (RoB 2) for included study



motion-controlled activities. Other systems combined VR settings with specialist rehabilitation robots, like the Armeo Spring exoskeleton [60], to facilitate passive and active limb motion. Advanced motion tracking-equipped head-mounted VR displays [47, 49, 56, 59, 61, 69], were also used to generate engaging, interactive training experiences. Interactive rehabilitation including task-specific games [75], reinforced feedback systems [63], and unique camera-based motion capture were also used [56, 58, 61, 67]. (Table 1).

### Intervention dose

The trials' median duration was 4 weeks, ranging from 2 to 12 weeks. The control groups' average daily exercise session length ranged from 20 to 150 min. Session lengths of 30 to 60 min of conventional therapy per week were included in 70% of the studies [47–52, 54–56, 58, 59, 61, 63, 64, 66, 67, 73]. Although they were less prevalent, longer sessions ranging from 90 to 150 min per day were also used [68, 71, 76, 77], mainly when the overall duration of the intervention was shorter (i.e., 2 to 3 weeks). The intervention lasted a median of 4 weeks (IQR: 3 to 6), with median sessions of 52.5 min (IQR: 30 to 60). (Table 2)

### Clinical efficacy

To assess the clinical efficacy of VR for UL stroke rehabilitation on ADL and QoL, different analyses were performed.

### The activity of daily living

Twenty-five studies were included in this meta-analysis, in addition to 2 studies using two different measures (FIM, MBI or BI) to assess ADL [52, 66], resulting in 25 outcomes included in this section. ADL was assessed using FIM in 9 studies [52, 54, 58, 60, 61, 63, 71, 74, 76], MBI in 10 studies [47, 50, 51, 59, 64, 66, 73–75], and BI in 8 studies [48, 52, 53, 55–57, 67, 70].

The overall SMD indicates a statistically significant larger effect of VR in comparison with conventional therapy (SMD=0.27 [95% CI 0.11; 0.43],  $p<0.001$ ). We then perform subgroup analysis to compare IVR and NIVR. A statistically significant difference was found between the two types of interventions ( $p=0.03$ ), with larger effect reported for the IVR group (SMD=0.54 [95% CI 0.13; 0.95],  $p<0.001$ ) in comparison with NIVR (SMD=0.19 [95% CI 0.02; 0.36],  $p=0.004$ ). (Fig. 3).

We then compared the efficacy of VR alone and in combination with conventional therapy. No statistically significant difference was found between the two groups ( $p=0.39$ ), but a tendency was found for a larger effect in the combined group (SMD: 0.31, [95% CI 0.12; 0.49]) compared to VR only intervention (SMD: 0.15, [95% CI

-0.26; 0.56]), see Supplementary Material 3 for complete results.

Lastly, we compared the efficacy according to the stroke stage. Statistically significant differences were found between the different strokes' stages ( $p=0.048$ ) with larger effect obtained in the subacute phase (SMD=0.52 [95% CI 0.16; 0.88]), in comparison with acute (SMD=0.08 [95% CI -0.11; 0.27]) or chronic (SMD=0.05 [95% CI -0.36; 0.46]). The forest plot is presented in Supplementary Material 4.

### Quality of life

Eight studies were included in this meta-analysis using 4 different tools to measure QoL; SF-36 in 3 studies [62, 69, 73], SIS in 3 studies [50, 65, 72], EQ-VAS in one study [78], and SSQoL in one study [47].

When compared to conventional treatment, an overall statistically significant effect was found in favor of VR (SMD=0.94 (95% CI: [ 0.09; 1.79],  $p=0.035$ ). When comparing IVR and NIVR, no statistically significant difference was found ( $p=0.98$ ), as presented in Fig. 4.

We then compared the efficacy of VR alone or in combination with conventional therapy. A statistically significant difference was found between the two groups ( $p=0.007$ ) with larger effect observed for the combination group (SMD=1.39 [95% CI 0.13; 2.64]) compared to VR alone (SMD=0.08 [95% CI -0.63; 0.80], see complete results in Supplementary Material 5. No significant difference was observed based different stages of stroke see Supplementary Material 6.

According to the GRADE summary of the evidence, we are moderately confident in the effect estimate for both ADL and QoL outcomes, and we can only conditionally recommend the intervention. (Supplementary Materials 7 and 8).

### Dose response

We performed meta-regression to determine if the total amount of rehabilitation influences the outcome. We did not find a statistically significant association between the total duration of rehabilitation and the clinical outcome for neither ADL ( $\beta = -0.0000$ , Standard Error [SE]=0.0001,  $p=0.88$ ), nor for QoL ( $\beta = 0.001$ , SE=0.0006,  $p=0.17$ ) (Fig. 5). Note that for QoL this analysis may be underpowered due to the low number of included studies ( $n=8$ ).

### Risk of publication bias and sensitivity analysis

Lastly, to test the robustness of our results, we performed sensitivity analysis to detect any potential study with extreme large effect and assessed the risk of publication bias. The analysis of the funnel plot did not reveal significant asymmetry (Supplementary Materials 9 and 10 for ADL and QoL respectively). Furthermore, the statistical

**Table 2** Intervention characteristics for the comparative effectiveness of virtual reality-based upper limb rehabilitation versus conventional therapy on ADL and QoL in stroke patients

Authors	Charac- teristics of exercise in control	Characteristics of exercise in experimental	Primary outcome	Is the VR intervention combined with conventional?	Duration of inter- ventions (weeks)	Duration and frequency Control	Intervention
Laffonte et al.,2020 [55]	CT	CT plus VR	BI	Yes	6	45 Min/session, 5 days /week	45 Min/session, 5 days /week
Afsar et al., 2018 [54]	CT	CT Plus Xbox Kinect game system	FIM	Yes	4	60 Min /session, 6 days/week	60 Min /session, 6 days/week
Gueye et al.,2021 [60]	CT	CT plus VR Arneo Spring	FIM	Yes	3	40 Min/session, 4 days/week	40 Min/session, 4 days/week
Mekbib et al.,2021 [56]	CT	CT for plus IVR	BI	Yes	2	60 Min/session, 4 days/week	60 Min/session, 4 days/week
Leng et al.,2022 [53]	CT	CT plus VR-based game	BI	Yes	3	30 Min/session, 5 days a week	30 Min /session, 5 days a week
Shin et al.,2014 [75]	CT	CT plus VR-based training	MBI	Yes	6	60 Min/session, 3days/week	60 Min/session, five 3days/week
Rong et al.,2021 [74]	CT with Mirror visual feedback	CT plus Robot-assisted training combination with base camera- based Mirror-Visual feedback	FIM and MBI	Yes	4	90 Min/session, 5 days/week	90 Min/session, 5 days/week
Kwon et al. 2012 [59]	CT	CT combined with an intensive VR program	MBI	Yes	4	70 Min/session, 5 days/week	30 Min/session, 5 days/week
Long et al.,2020 [66]	CT	CT plus VR program	MBI	Yes	3	45 Min/session, 5 days/week	45 Min/session, 5 days/ week
Yin et al.,2014 (58)	Ct	CT plus VR therapy	FIM	Yes	2	30Min/session, 5days/week	30Min/session, 5days/week
Huang et al.,2023 [48]	CT	CT plus VR	BI	Yes	3	60Min/session, 5days/week	30Min/session,5days/week
Park et al.,2019 [73]	CT	CT plus based VR-based therapy	MBI and SF-36	Yes	4	30 min/session, 5 days/week	30 min/session, 5 days/week
Turola et al.,2013 [76]	CT	CT plus VR	FIM	Yes	4	120Min/session, 5 days /week	120Min/session, 5 days /week
Cameir'ao et al., 2011 [57]	CT	CT plus VR-based games	BI	Yes	12	20 Min/session, 3days/ week	20 Min/session, 3days/ week
Zheng et al., 2015 [50]	CT	CT plus VR	MBI and SIS	Yes	4	30 Min/ session, 6 days/week	30 Min/ session, 6 days/week
Amin et al., 2024 [47]	CT	CT plus VR	MBI and SS QOL	Yes	6	48 Min/session for 2 weeks then 80 Min/ session for 4 weeks, 4 days/week	24 Min/per session for 2, then 40 Min/per session for 4 weeks, 4 days/per week
Shin et al., 2016 [75]	CT	CT plus VR	SIS	Yes	4	30 Min /session, 5 days/week	30 Min /session, 5 days/week
Sip et al., 2023 [69]	CT	CT plus VR	SF-36	Yes	3	30 Min/session, 6days/week, 18days only	30 Min/session, 6days/week, 18days only
Rodríguez-Hernández et al., 2021 [68]	CT	CT plus VR	EQ-VAS	Yes	3	150 Min/session, 5days week	150 Min/session, 5days week
Ribeiro et al., 2015 [62]	CT	CT plus VR based therapy	SF-36	Yes	8	60 Min/session, 2 days/week	60 Min/session, 2 days/week
Ali et al., 2024 [65]	CT	CT plus VR-based gamified therapy	SIS	Yes	6	120 Min/session, 6 days /week	120 Min/session, 6 days /week
Keskin et al.,2020 [67]	CT	VR-based therapy	BI	No	6	60 Min/session, 5 days/weekdays/week	60 Min/session, 5 days/week
Saposnik et al.,2016 [80]	CT	VR-based Nintendo Wii	BI and FIM	No	2	60 Min/session, 5 days/week	60 Min/session, 5 days/week

**Table 2** (continued)

Authors	Charac- teristics of exercise in control	Characteristics of exercise in experimental	Primary outcome	Is the VR intervention combined with conventional?	Duration of inter- ventions (weeks)	Duration and frequency Control	Intervention
Choi et al., 2014 [49]	CT	Commercial gaming-based VR therapy using Wii (Nintendo, Tokyo, Japan)	MBI	No	4	30 Min/session, 5 days/4 week	30 Min/session, 5 days/4 week
Kiper et al., 2013 [71]	CT	VR-based therapy	FIM	No	5	120/session, 5 days/week	120/session, 5 days/week
Lee et al., 2016 [64]	CT	Individual-based VR-based therapy	MBI	No	8	30 Min/session, 3 days/week	30 Min/session, 3 days/week
Ogun et al., 2019 [61]	CT	VR based therapy	FIM	No	6	60 Min/session, 3 days/week	60 Min/session, 3 days/week
Choi et al., 2021 [51]	CT	VR based therapy	MBI	No	4	30 Min/session, 3 days/ week	30 Min/session, 3 days/ week
Piron et al., 2010 [63]	CT	VR based therapy	FIM	No	4	60 Min/session, 5 days /week	60 Min/session, 5 days /week
Ballester et al., 2016 [70]	CT	VR-based therapy	BI	No	6	30 Min/session, 5 days /week	30 Min/session, 5 days /week


CT = Conventional Therapy, BI = Barthel Index, MBI = Modified Barthel Index, FIM = Functional Independence Measure, Min = Minute/s, EQ VAS = Euro QoL Visual Analogue Scale; SS QoL = Stroke Specific Quality of Life, SIS = Stroke Impact Scale, and SF-36 = Short-Form 36 Health Survey

assessment using Egger's intercept yielded a value of 0.21 (SE = 0.68), with a corresponding  $p$ -value of 0.76 for ADL and 2.63 (3.55),  $p = 0.48$  for QoL. Furthermore, the sensitivity analysis (Supplementary Materials 11 and 12 for ADL and QoL respectively) did not identify any study that had an extreme influence on the overall results.

## Discussion

The main goal of this research was to evaluate the effectiveness of VR therapies designed for UL rehabilitation in comparison to conventional rehabilitation. The presented systematic review and meta-analysis thoroughly analyzed data from 30 RCTs, which included a cohort of 1661 stroke survivors. The review identified a diverse, versatile range of immersion ideal for upper limb rehabilitation. In general, VR-based technology illustrates an innovative approach to providing UL rehabilitation. In current review, 77% of the studies used NIVR systems, the average duration of session length was 30 to 60 min, and the median duration of the intervention programs was 4 weeks.

Our meta-analysis revealed that VR therapy demonstrated statistically significant superiority over conventional therapy. VR therapy showed a moderate improvement in ADL (SMD = 0.27 [0.11; 0.43] and a large effect on QoL (SMD = 0.94 [0.09; 1.79]. Notably, the QoL results were derived from aggregating data from only eight individual studies. The VR therapy has the ability to improve patients' motivation and enjoyment [19], improve their compliance with rehabilitation, and reduce fatigue [79]. In addition, numerous advantages of VR-based programs have been suggested, such as their affordability, ability to improve treatment outcomes, and ability to immerse stroke survivors in a world that closely resembles real objects and events - by integrating multiple sensory stimuli, such as tactile, visual, auditory, and somatosensory systems [62, 80–83]. From this review, it was also shown that VR was effective in improving both ADL and QoL when combined with conventional therapy. This could be explained by the fact that VR enhances movement quality through motor learning and repetitive practice, which can be effectively transferred to ADL [84–86]. The immersive environment that IVR creates contributes to realistic and engaging rehabilitation tasks, which could facilitate greater neuroplasticity and recovery [87, 88]. Immersive VR improves motor learning principles by providing realistic sensory feedback, enhancing task specificity through engaging environments, and facilitating error augmentation that allows users to identify and correct mistakes in a controlled setting, leading to improved skill acquisition compared to non-immersive VR [89]. In addition, IVR may enhance motor learning outcomes through VR-induced sensory integration, which encourages a more cohesive




**Fig. 3** Forest plot result for the comparative effectiveness of virtual reality-based upper limb rehabilitation versus conventional therapy on ADL in stroke survivors -based VR type


perception of body movements and environment; activation of the mirror neuron system, which improves the simulation of observed actions, and enhanced embodiment creates a sense of presence that strengthens the connection between mental imagery; and physical execution, collectively leading to improved neural pathways for skill acquisition and retention [90]. This approach focuses on improving movement efficiency, particularly within rehabilitation settings. Additionally, VR based rehabilitation promotes movement efficiency by allowing for tailored and repetitive practice, which is especially beneficial for individuals with motor impairments, ultimately improving balance and motor function and enhancing overall quality of life [84, 91].

Similarly, prior research indicates that conventional rehabilitation combined with a particular VR technology may be more beneficial than conventional programs alone in enhancing motor recovery and activity among stroke survivors [78, 92, 93]. Particularly in people with subacute stroke, who showed the most significant improvements with high effect in ADL based on our finding, VR offers a promising tool to accelerate recovery during this critical rehabilitation window [94, 95]. Given the relative difficulty in engaging chronic-phase patients in conventional therapies, VR's gamified and immersive nature offers an innovative alternative that sustains motivation and participation, which may otherwise decline over time. Clinicians can leverage VR to provide a more dynamic, individualized rehabilitation experience. For





**Fig. 4** Forest plot result for the comparative effectiveness of virtual reality-based upper limb rehabilitation versus conventional therapy on QoL in stroke-based VR type subgroup



**Fig. 5** Bubble plot showing the relationship between the total number of rehabilitation (minutes) and the effect on ADL and QoL (meta-regression). The size is proportional to the study weight

example, VR systems that simulate real-life tasks, such as grasping and manipulating objects, can offer functional and relevant training that directly transfers to improved ADL performance in patients' daily lives [96, 97]. Moreover, the accessibility of non-immersive VR systems (e.g., those using devices such as Microsoft Kinect or Nintendo Wii) makes them suitable for home-based rehabilitation programs [98, 99]. This opens new avenues for post-discharge rehabilitation, which can help maintain gains achieved during inpatient therapy and potentially reduce hospital readmissions. This is particularly critical in regions where access to rehabilitation services is limited or where healthcare resources are strained [100, 101].

The broad spectrum of VR-based options in this review highlights the creative ways that are being investigated to provide efficient, engaging upper limb rehabilitation through immersive, motion-driven experiences as supported by another review [102]. Numerous VR-based hardware and software technologies that have been included into UL rehabilitation programs as highlighted in the review. Training situations that are immersive and engaging are frequently created using head-mounted virtual reality displays that have motion tracking capabilities. IVR demonstrated superior outcomes compared to NIVR for ADL improvements, suggesting that IVR should be prioritized in clinical settings where possible, to maximize the rehabilitation outcomes for UL function



so that it will add sensitive measures to current clinically available ones [103]. IVR provides considerable benefits for therapy targeting upper limb recovery after a stroke by increasing patient involvement, mimicking everyday tasks, fostering improved neuroplasticity, enabling real-time treatment monitoring, and enhancing gross motor skills [104, 105].

There was no significant association between the total amount of rehabilitation, ADL, and QoL. This may be due to differences in individuals responsive to VR interventions, the possibility that the amount of rehabilitation is not solely adequate, and the influence of other psychosocial factors that affect recovery beyond the rehabilitation process. Additionally, it may be because the duration of rehabilitation was not long enough [106, 107]. The actual effect is likely to be close to the effect estimate as per our GRADE summary of the evidence, and we are moderately confident in the effect estimate for both ADL and QoL outcomes; however, there is still a possibility that it is substantially different. Further research could have a substantial impact, which may change the effect estimates. Moreover, we can only conditionally recommend the intervention, and the desirable effects probably outweigh the undesirable effects, but still, we need more confidence.

### Strengths and limitations

The results of this study have to be analyzed in light of some limitations. First, even though most of the studies that were included had high methodological quality, most of them lacked blinded allocation and an intention-to-treat analysis, this may have contributed to bias in the included trials. Lack of blinding is one of the most important potential source of bias in rehabilitation research [108–111]. Other limitations including that the results of the study may be difficult to apply to stroke survivors due to high heterogeneity between the studies with important variations in the age and gender of the participants, both ischemic and hemorrhagic strokes survivors were included while their recovery path and rehabilitation process may differ. Additionally, the training parameters, and the treatment durations allocated in the different individual studies, still concerning the intervention, as we also included a few research that combined NIVR with robotic exoskeletons [50, 74, 112, 113], which makes it challenging to conclude the observed difference was solely due to NIVR. Furthermore, beside this high heterogeneity in term of patients and intervention, most of the studies included a relatively low number of participants. There is also a lack of long-term follow-up, many studies had relatively short follow-up periods, limiting the understanding of the long-term effects of VR-based rehabilitation. Thirdly, due to the low number of included studies assessing QoL, these results

should be interpreted cautiously, especially for the risk of bias assessment and the meta-regression since less than 10 studies were included [114]. Fourthly, we only included studies assessed activities of daily living using specific tools like BI, MBI, and FIM and did not consider other more general activity measuring tools such as the Chedoke Arm and Hand Activity Inventory. Finally, geographical limitations with LMICs account for about 90% of all stroke-related deaths and disabilities, with sub-Saharan Africa bearing a disproportionately large burden [3]. Thus, research on the feasibility and efficacy of VR-based therapy must be carried out in LMICs, particularly in Africa, to fill the gap.

Despite these limitations, this study demonstrates several strengths. First, it provides a comprehensive analysis of the current literature on the use of VR in upper limb rehabilitation for stroke survivors, clearly articulating the differential impacts of IVR and NIVR. Furthermore, the detailed subgroup analyses offer valuable insights into how the integration of VR and conventional therapy can enhance rehabilitation outcomes. Lastly, the inclusion of both ADL and QoL as outcome measures provides a holistic assessment of the impact of VR interventions.

### Conclusion

This systematic review and meta-analysis found that VR upper limb interventions combined with conventional therapy significantly improve ADL and QoL in stroke survivors compared to conventional rehabilitation methods with moderate certainty of evidence. The findings highlight the potential of VR as a beneficial tool in stroke rehabilitation, particularly in enhancing patient motivation and engagement through immersive and interactive environments even though the total amount of VR rehabilitation does not seem to have a statistically significant impact on the clinical outcomes. However, the review also identified several limitations, including high heterogeneity among studies, short follow-up periods, and a lack of research from LMICs. To address these gaps, future research should focus on conducting large-scale studies with diverse populations and extended follow-up durations. Despite the promising results, further research is necessary to establish the long-term benefits of VR-based rehabilitation, particularly in the chronic phase of stroke recovery. More large-scale, high-quality RCTs are needed to assess the durability of VR's effects on ADL and QoL over time, and to determine the optimal dose and intensity of VR interventions. Additionally, studies should explore the integration of more advanced technologies, such as VR combined with robotic devices or neurostimulation, to further enhance motor recovery outcomes.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12984-025-01603-1>.

Supplementary Material 1

## Acknowledgements

The authors would like to express appreciation to all the NASCERE project coordinators and their teams, at Ghent University, Hasselt University and Jimma University, for the funding and other facilities.

## Author contributions

Conceptualization: DDO, BB, DL, TG. Data curation: DDO, BB. Formal analysis: DDO, BB. Funding acquisition: BB. Investigation: DDO, BB, DL, TG, LTT. Supervision: BB, TG, DL. Validation: BB, TG, DL, LTT. Visualization: BB, DDO. Writing – original draft: DD, BB. Writing – review & editing: DDO, BB, DL, TG, LTT. All authors have read and agreed to the published version of the manuscript.

## Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article. This study was supported by the NASCERE program.

## Data availability

No datasets were generated or analysed during the current study.

## Declarations

## Ethics approval and consent to participate

Not applicable.

## Consent for publication

All authors have approved this manuscript for publication. This manuscript has not previously been published and is not pending publication elsewhere.

## Competing interests

The authors declare no competing interests.

## Author details

<sup>1</sup>Department of Biomedical Sciences, Faculty of Medical Sciences, Institute of Health, Jimma University, Jimma, Ethiopia

<sup>2</sup>REVAL Rehabilitation Research Center, Faculty of Rehabilitation Sciences, Hasselt University, Diepenbeek, Belgium

<sup>3</sup>Department of Special Needs and Inclusive Education, Jimma University, Jimma, Ethiopia

<sup>4</sup>Department of Physiotherapy, Faculty of Medical Sciences, Institute of Health, Jimma University, Jimma, Ethiopia

<sup>5</sup>Department of Clinical and Movement Neurosciences, University College London, London, UK

<sup>6</sup>Data Sciences Institute, Technology-Supported and Data-Driven Rehabilitation, Hasselt University, Diepenbeek, Belgium

<sup>7</sup>Department of PXL—Healthcare, PXL University of Applied Sciences and Arts, Hasselt 3500, Belgium

Received: 20 November 2024 / Accepted: 4 March 2025

Published online: 24 April 2025

## References

- Cieza A. Rehabilitation the health strategy of the 21st century. really? Arch Phys Med Rehabil. 2019;100(11):2212–4.
- Stucki G, Bickenbach J, Gutenbrunner C, Melvin JL. Rehabilitation: The health strategy of the 21st century. 2018.
- eClinicalMedicine. The rising global burden of stroke. eClinicalMedicine. 2023 May 1 [cited 2023 Sep 25];59. Available from: [https://www.thelancet.com/journals/echinm/article/PIIS2589-5370\(23\)00205-5/fulltext](https://www.thelancet.com/journals/echinm/article/PIIS2589-5370(23)00205-5/fulltext)
- Feigin VL, Stark BA, Johnson CO, Roth GA, Bisignano C, Abady GG, et al. Global, regional, and National burden of stroke and its risk factors, 1990–2019: a systematic analysis for the global burden of disease study 2019. Lancet Neurol. 2021;20(10):795–820.
- Rafsten L, Meirrelles C, Danielsson A, Sunnerhagen KS. Impaired Motor Function in the Affected Arm Predicts Impaired Postural Balance After Stroke: A Cross Sectional Study. Front Neurol. 2019 Aug 21 [cited 2024 Aug 13];10. Available from: <https://www.frontiersin.org/journals/neurology/articles/https://doi.org/10.3389/fneur.2019.00912/full>
- Hussain N, Alt Murphy M, Sunnerhagen KS. Upper limb kinematics in stroke and healthy controls using Target-to-Target task in virtual reality. Front Neurol. 2018;9:300.
- Niama Natta DD, Lejeune T, Detrembleur C, Yarou B, Sogbossi ES, Alagnidé E, et al. Effectiveness of a self-rehabilitation program to improve upper-extremity function after stroke in developing countries: A randomized controlled trial. Ann Phys Rehabil Med. 2021;64(1):101413.
- Ingram LA, Butler AA, Brodie MA, Lord SR, Gandevia SC. Quantifying upper limb motor impairment in chronic stroke: a physiological profiling approach. J Appl Physiol. 2021;131(3):949–65.
- Doumen S, Sorba L, Feys P, Tedesco Triccas L. Efficacy and dose of rehabilitation approaches for severe upper limb impairments and disability during early acute and subacute stroke: a systematic review. Phys Ther. 2023;103(4):pzad002.
- Salvalaggio S, Cacciantè L, Maistrello L, Turolla A. Clinical predictors for upper limb recovery after stroke rehabilitation: retrospective cohort study. Health-care. 2023;11(3):335.
- Hiraga Y, Hayashi T. Mediating effect of upper limb use on the relationship between upper limb performance and activities of daily living: a longitudinal mediation analysis. Cureus. 2022;14(10).
- Rehabilitation. [cited 2025 Feb 12]. Available from: <https://www.who.int/new-s-room/fact-sheets/detail/rehabilitation>
- Bindawas SM, Vennu VS. Stroke rehabilitation. Neurosciences. 2016;21(4):297–305.
- Wurzinger E, Abzhandadze H, Rafsten T, Sunnerhagen L. KS. Dependency in Activities of Daily Living During the First Year After Stroke. Front Neurol. 2021 [cited 2023 Sep 25];12. Available from: <https://www.frontiersin.org/articles/https://doi.org/10.3389/fneur.2021.736684>
- Tuominen PP, Saarni LA. The use of virtual technologies with music in rehabilitation: a scoping systematic review. Front Virtual Real. 2024;5:1290396.
- Kern F, Winter C, Gall D, Käthner I, Pauli P, Latoschik ME. Immersive virtual reality and gamification within procedurally generated environments to increase motivation during gait rehabilitation. In: 2019 IEEE Conference on Virtual Reality and 3D User Interfaces (VR). IEEE. [cited 2023 Nov 13]. 2019;500–9. Available from: <https://ieeexplore.ieee.org/abstract/document/8797828/>
- Demeco A, Zola L, Frizziero A, Martini C, Palumbo A, Foresti R, et al. Immersive virtual reality in Post-Stroke rehabilitation: A systematic review. Sensors. 2023;23(3):1712.
- Diriba Kenea C, Gemechu Abessa T, Lamba D, Bonnechère B. Technological features of immersive virtual reality systems for upper limb stroke rehabilitation: A systematic review. Sensors. 2024;24(11):3546.
- Zhang C, Yu S. The technology to enhance patient motivation in virtual reality rehabilitation: A review. Games Health J. 2024;13(4):215–33.
- Micheluzzi V, Casu G, Sanna GD, Canu A, Iovino P, Caggianelli G, et al. Improving adherence to rehabilitation for heart failure patients through immersive virtual reality (VIRTUAL-HF): A protocol for a randomized controlled trial. Contemp Clin Trials. 2024;138:107463.
- Lundin RM, Yeap Y, Menkes DB. Adverse effects of virtual and augmented reality interventions in psychiatry: systematic review. JMIR Ment Health. 2023;10:e43240.
- Domínguez-Téllez P, Moral-Muñoz JA, Salazar A, Casado-Fernández E, Lucena-Antón D. Game-Based virtual reality interventions to improve upper limb motor function and quality of life after stroke: systematic review and Meta-analysis. Games Health J. 2020;9(1):1–10.
- Chen X, Liu F, Lin S, Yu L, Lin R. Effects of virtual reality rehabilitation training on cognitive function and activities of daily living of patients with poststroke cognitive impairment: A systematic review and Meta-Analysis. Arch Phys Med Rehabil. 2022;103(7):1422–35.
- Leong SC, Tang YM, Toh FM, Fong KNK. Examining the effectiveness of virtual, augmented, and mixed reality (VAMR) therapy for upper limb recovery and activities of daily living in stroke patients: a systematic review and meta-analysis. J Neuroeng Rehabil. 2022;19(1):93.
- Gao Y, Ma L, Lin C, Zhu S, Yao L, Fan H, et al. Effects of virtual Reality-Based intervention on cognition, motor function, mood, and activities of daily living

- in patients with chronic stroke: A systematic review and Meta-Analysis of randomized controlled trials. *Front Aging Neurosci.* 2021;13:766525.
26. Zhang B, Wong KP, Qin J. Effects of virtual reality on the limb motor function, balance, gait, and daily function of patients with stroke: systematic review. *Med Kaunas Lith.* 2023;59(4):813.
27. Jin M, Pei J, Bai Z, Zhang J, He T, Xu X, et al. Effects of virtual reality in improving upper extremity function after stroke: A systematic review and meta-analysis of randomized controlled trials. *Clin Rehabil.* 2022;36(5):573–96.
28. Khan A, Imam YZ, Muneer M, Al Jerdi S, Gill SK. Virtual reality in stroke recovery: a meta-review of systematic reviews. *Bioelectron Med.* 2024;10(1):23.
29. Barger S, Scalea S, Agosta F, Banfi G, Corbetta D, Filippi M et al. Effectiveness and safety of virtual reality rehabilitation after stroke: an overview of systematic reviews. *eClinicalMedicine.* 2023 Oct 1 [cited 2025 Jan 22];64. Available from: [https://www.thelancet.com/journals/eclinm/article/PIIS2589-5370\(23\)00397-8/fulltext](https://www.thelancet.com/journals/eclinm/article/PIIS2589-5370(23)00397-8/fulltext)
30. Wang S, Meng H, Zhang Y, Mao J, Zhang C, Qian C et al. Effect of Virtual Reality-Based Rehabilitation on Mental Health and Quality of Life of Stroke Patients: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Arch Phys Med Rehabil.* 2024 Nov 2 [cited 2025 Jan 22]; Available from: <https://www.sciencedirect.com/science/article/pii/S000399932401311X>
31. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Syst Rev.* 2021;10(1):89.
32. Johnson N, Phillips M. Rayyan for systematic reviews. *J Electron Resour Librariansh.* 2018;30(1):46–8.
33. Amir-Behghadami M, Janati A, Population. Intervention, Comparison, Outcomes and Study (PICOS) design as a framework to formulate eligibility criteria in systematic reviews. *Emerg Med J.* 2020 [cited 2024 Jun 24]; Available from: <https://emj.bmj.com/content/early/2020/04/05/emjmed-2020-09567?versioned=true>
34. Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther.* 2003;83(8):713–21.
35. PEDro scale - PEDro. 2016 [cited 2023 Oct 24]. Available from: <https://pedro.org.au/english/resources/pedro-scale/>
36. de Morton NA. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Aust J Physiother.* 2009;55(2):129–33.
37. Sterne JA, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *bmj.* 2019 [cited 2025 Feb 4];366. Available from: <https://www.bmj.com/content/366/bmj.1489.8.short>
38. Evidence Prime I, GRADEpro GDT. GRADEpro guideline development [software] [Software]. McMaster Univ. 2015;140.
39. Schünemann H, Brożek J, Guyatt G, Oxman A. The GRADE handbook. Cochrane Collaboration London, UK; 2013 [cited 2024 Aug 14]. Available from: [https://hero.epa.gov/hero/index.cfm/reference/details/reference\\_id/10284249](https://hero.epa.gov/hero/index.cfm/reference/details/reference_id/10284249)
40. Gallardo-Gómez D, Richardson R, Dwan K. Standardized mean differences in meta-analysis: A tutorial. *Cochrane Evid Synth Methods.* 2024;2(3):e12047.
41. Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol.* 2014;14(1):135.
42. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. Cochrane handbook for systematic reviews of interventions version 6.5 (updated August 2024). Cochrane; 2024. Available from: <https://www.training.cochrane.org/handbook>.
43. Luo D, Wan X, Liu J, Tong T. Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range. *Stat Methods Med Res.* 2018;27(6):1785–805.
44. Melsen WG, Bootsma MCJ, Rovers MM, Bonten MJM. The effects of clinical and statistical heterogeneity on the predictive values of results from meta-analyses. *Clin Microbiol Infect.* 2014;20(2):123–9.
45. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327(7414):557–60.
46. Bezerra CT, Grande AJ, Galvão VK, dos Santos DHM, Atallah ÁN, Silva V. Assessment of the strength of recommendation and quality of evidence: GRADE checklist. A descriptive study. *São Paulo Med J.* 2022;140(6):829–36.
47. Amin F, Waris A, Syed S, Amjad I, Umar M, Iqbal J et al. Effectiveness of Immersive Virtual Reality Based Hand Rehabilitation Games for Improving Hand Motor Functions in Subacute Stroke Patients. *IEEE Trans Neural Syst Rehabil Eng.* 2024 [cited 2024 Jun 9]; Available from: <https://ieeexplore.ieee.org/abstract/document/10539093/>
48. Huang Q, Jiang X, Jin Y, Wu B, Vigotsky AD, Fan L, et al. Immersive virtual reality-based rehabilitation for subacute stroke: a randomized controlled trial. *J Neurol.* 2024;271(3):1256–66. <https://doi.org/10.1007/s00415-023-12060-y>. Epub 2023 Nov 10. PMID: 37947856; PMCID: PMC10896795.
49. Choi JH, Han EY, Kim BR, Kim SM, Im SH, Lee SY, et al. Effectiveness of commercial gaming-based virtual reality movement therapy on functional recovery of upper extremity in subacute stroke patients. *Ann Rehabil Med.* 2014;38(4):485–93.
50. Zheng CJuan, Liao W, jing, Xia Wguang. Effect of combined low-frequency repetitive transcranial magnetic stimulation and virtual reality training on upper limb function in subacute stroke: a double-blind randomized controlled trial. *J Huazhong Univ Sci Technolog Med Sci.* 2015;35(2):248–54.
51. Choi HS, Shin WS, Bang DH. Application of digital practice to improve head movement, visual perception and activities of daily living for subacute stroke patients with unilateral Spatial neglect: preliminary results of a single-blinded, randomized controlled trial. *Med (Baltim).* 2021;100(6):e24637.
52. Saposnik G, Cohen LG, Mamdani M, Pooyania S, Ploughman M, Cheung D, et al. Efficacy and safety of non-immersive virtual reality exercising in stroke rehabilitation (EVREST): a randomised, multicentre, single-blind, controlled trial. *Lancet Neurol.* 2016;15(10):1019–27.
53. Leng Y, Lo WLA, Mao YR, Bian R, Zhao JL, Xu Z, et al. The impact of cognitive function on virtual reality intervention for upper extremity rehabilitation of patients with subacute stroke: prospective randomized controlled trial with 6-month follow-up. *JMIR Serious Games.* 2022;10(3):e33755.
54. Afsar SI, Mirzayev I, Yemisci OU, Saracgil SNC. Virtual reality in upper extremity rehabilitation of stroke patients: a randomized controlled trial. *J Stroke Cerebrovasc Dis.* 2018;27(12):3473–8.
55. Laffont I, Froger J, Jourdan C, Bakhti K, van Dokkum LE, Gouaich A, et al. Rehabilitation of the upper arm early after stroke: video games versus conventional rehabilitation. A randomized controlled trial. *Ann Phys Rehabil Med.* 2020;63(3):173–80.
56. Mekbib DB, Debeli DK, Zhang L, Fang S, Shao Y, Yang W, et al. A novel fully immersive virtual reality environment for upper extremity rehabilitation in patients with stroke. *Ann N Y Acad Sci.* 2021;1493(1):75–89.
57. da Silva Cameirão M, Bermudez I, Badia S, Duarte E, Verschure PF. Virtual reality based rehabilitation speeds up functional recovery of the upper extremities after stroke: a randomized controlled pilot study in the acute phase of stroke using the rehabilitation gaming system. *Restor Neurol Neurosci.* 2011;29(5):287–98.
58. Yin CW, Sien NY, Ying LA, Chung SFCM, Tan May Leng D. Virtual reality for upper extremity rehabilitation in early stroke: a pilot randomized controlled trial. *Clin Rehabil.* 2014;28(11):1107–14.
59. Kwon JS, Park MJ, Yoon JJ, Park SH. Effects of virtual reality on upper extremity function and activities of daily living performance in acute stroke: a double-blind randomized clinical trial. *NeuroRehabilitation.* 2012;31(4):379–85.
60. Gueye T, Dedkova M, Rogalewicz V, Grunerova-Lippertova M, Angerova Y. Early post-stroke rehabilitation for upper limb motor function using virtual reality and exoskeleton: equally efficient in older patients. *Neurol Neurochir Pol.* 2021;55(1):91–6.
61. Ögün MN, Kurul R, Yaşar MF, Turkoglu SA, Avci Ş, Yıldız N. Effect of leap motion-based 3D immersive virtual reality usage on upper extremity function in ischemic stroke patients. *Arq Neuropsiquiatr.* 2019;77:681–8.
62. Da Silva Ribeiro NM, Ferraz DD, Pedreira É, Pinheiro Í, Da Silva Pinto AC, Neto MG, et al. Virtual rehabilitation via Nintendo Wii® and conventional physical therapy effectively treat post-stroke hemiparetic patients. *Top Stroke Rehabil.* 2015;22(4):299–305.
63. Piron L, Turolla A, Agostini M, Zucconi CS, Ventura L, Tonin P, et al. Motor learning principles for rehabilitation: A pilot randomized controlled study in poststroke patients. *Neurorehabil Neural Repair.* 2010;24(6):501–8.
64. Lee M, Son J, Kim J, Pyun SB, Eun SD, Yoon B. Comparison of individualized virtual reality-and group-based rehabilitation in older adults with chronic stroke in community settings: a pilot randomized controlled trial. *Eur J Integr Med.* 2016;8(5):738–46.
65. Ali F, Suleman R, Noor A, Ahmad I, Shakeel M, Aqeel M. Effectiveness of a virtual Reality-Based rehabilitation program versus conventional physical therapy in improving motor function and balance in stroke survivors: A randomized controlled trial. *J Health Rehabil Res.* 2023;3(2):817–21.
66. Long Y, Ouyang R, ge, Zhang J. qi. Effects of virtual reality training on occupational performance and self-efficacy of patients with stroke: a randomized controlled trial. *J NeuroEngineering Rehabil.* 2020;17(1):150.

67. KESKİN Y, Atci A, Urkmez B, Akgül Y, Ozaras N, Aydin T. Efficacy of a video-based physical therapy and rehabilitation system in patients with post-stroke hemiplegia: A randomized, controlled, pilot study. *Türk J Geriatr-Türk Geriatri Derg.* [cited 2023 Oct 28] 2020;;23(1). Available from: <https://avesis.bezmialem.edu.tr/yayin/5267f536-86b6-45e4-a762-a0323300ed48/efficacy-of-a-video-based-physical-therapy-and-rehabilitation-system-in-patients-with-post-stroke-hemiplegia-a-randomized-controlled-pilot-study>
68. Rodríguez-Hernández M, Criado-Álvarez JJ, Corregidor-Sánchez AI, Martín-Conty JL, Mohedano-Moriano A, Polonio-López B. Effects of virtual reality-based therapy on quality of life of patients with subacute stroke: a three-month follow-up randomized controlled trial. *Int J Environ Res Public Health.* 2021;18(6):2810.
69. Sip P, Kozłowska M, Czysz D, Daroszewski P, Lisiński P. Perspectives of motor functional upper extremity recovery with the use of immersive virtual reality in stroke patients. *Sensors.* 2023;23(2):712.
70. Ballester BR, Maier M, San Segundo Mozo RM, Castañeda V, Duff A, Verschure MJ. Counteracting learned non-use in chronic stroke patients with reinforcement-induced movement therapy. *J Neuroeng Rehabil.* 2016;13(1):74.
71. Kiper P, Agostini M, Luque-Moreno C, Tonin P, Turola A. Reinforced feedback in virtual environment for rehabilitation of upper extremity dysfunction after stroke: preliminary data from a randomized controlled trial. *BioMed Res Int.* 2014 [cited 2023 Oct 28];2014. Available from: <https://www.hindawi.com/journals/bmri/2014/752128/abs/>
72. Shin JH, Park SB, Jang SH. Effects of game-based virtual reality on health-related quality of life in chronic stroke patients: a randomized, controlled study. *Comput Biol Med.* 2015;63:92–8.
73. Park M, Ko MH, Oh SW, Lee JY, Ham Y, Yi H, et al. Effects of virtual reality-based planar motion exercises on upper extremity function, range of motion, and health-related quality of life: a multicenter, single-blinded, randomized, controlled pilot study. *J Neuroeng Rehabil.* 2019;16(1):122.
74. Rong J, Ding L, Xiong L, Zhang W, Wang W, Deng M, et al. Mirror visual feedback prior to robot-assisted training facilitates rehabilitation after stroke: a randomized controlled study. *Front Neurol.* 2021;12:683703.
75. Shin JH, Ryu H, Jang SH. A task-specific interactive game-based virtual reality rehabilitation system for patients with stroke: a usability test and two clinical experiments. *J Neuroeng Rehabil.* 2014;11(1):32.
76. Turola A, Dam M, Ventura L, Tonin P, Agostini M, Zucconi C, et al. Virtual reality for the rehabilitation of the upper limb motor function after stroke: a prospective controlled trial. *J Neuroeng Rehabil.* 2013;10(1):85.
77. Ali M, Dekker L, Daems JD, Ali M, Van Zwet EW, Steyerberg EW, et al. Sex differences in prehospital identification of large vessel occlusion in patients with suspected stroke. *Stroke.* 2024;55(3):548–54.
78. Rodríguez-Hernández M, Polonio-López B, Corregidor-Sánchez AI, Martín-Conty JL, Mohedano-Moriano A, Criado-Álvarez JJ. Can specific virtual reality combined with conventional rehabilitation improve poststroke hand motor function? A randomized clinical trial. *J Neuroeng Rehabil.* 2023;20(1):38.
79. Soleimani M, Ghazisaedi M, Heydari S. The efficacy of virtual reality for upper limb rehabilitation in stroke patients: a systematic review and meta-analysis. *BMC Med Inf Decis Mak.* 2024;24:135.
80. Gibbons EM, Thomson AN, De Noronha M, Joseph S. Are virtual reality technologies effective in improving lower limb outcomes for patients following stroke – a systematic review with meta-analysis. *Top Stroke Rehabil.* 2016;23(6):440–57.
81. Iruthayarajah J, McIntyre A, Cotoi A, Macaluso S, Teasell R. The use of virtual reality for balance among individuals with chronic stroke: a systematic review and meta-analysis. *Top Stroke Rehabil.* 2017;24(1):68–79.
82. Bonnechère B. Serious Games in Rehabilitation. In: Bonnechère B, editor. *Serious Games in Physical Rehabilitation: From Theory to Practice*. Cham: Springer International Publishing. [cited 2024 Jun 11]. 2018;41–109. Available from: [https://doi.org/10.1007/978-3-319-66122-3\\_4](https://doi.org/10.1007/978-3-319-66122-3_4)
83. Tinga AM, Visser-Meily JMA, Van Der Smagt MJ, Van Der Stigchel S, Van Ee R, Nijboer TCW. Multisensory stimulation to improve Low- and Higher-Level sensory deficits after stroke: A systematic review. *Neuropsychol Rev.* 2016;26(1):73–91.
84. Levac DE, Huber ME, Sternad D. Learning and transfer of complex motor skills in virtual reality: a perspective review. *J Neuroeng Rehabil.* 2019;16(1):121.
85. Grewal J, Eng JJ, Sakakibara BM, Schmidt J. The use of virtual reality for activities of daily living rehabilitation after brain injury: A scoping review. *Aust Occup Ther J.* 2024;71(5):868–93.
86. Peng Qcheng, Yin L, Cao Y. Effectiveness of virtual reality in the rehabilitation of motor function of patients with subacute stroke: A Meta-Analysis. *Front Neurol.* 2021;12:639535.
87. Hao J, Xie H, Harp K, Chen Z, Siu KC. Effects of virtual reality intervention on neural plasticity in stroke rehabilitation: A systematic review. *Arch Phys Med Rehabil.* 2022;103(3):523–41.
88. Huang CY, Chiang WC, Yeh YC, Fan SC, Yang WH, Kuo HC, et al. Effects of virtual reality-based motor control training on inflammation, oxidative stress, neuroplasticity and upper limb motor function in patients with chronic stroke: a randomized controlled trial. *BMC Neurol.* 2022;22(1):21.
89. Demers M, Fung K, Subramanian SK, Lemay M, Robert MT. Integration of motor learning principles into virtual reality interventions for individuals with cerebral palsy: systematic review. *JMIR Serious Games.* 2021;9(2):e23822.
90. Hadjipanayi C, Banakou D, Michael-Grigoriou D. Virtual reality exergames for enhancing engagement in stroke rehabilitation: A narrative review. *Heliyon.* 2024;10(18):e37581.
91. Komariah M, Amirah S, Abdurrahman MF, Handimulya MFS, Platini H, Maulana S, et al. Effectivity of virtual reality to improve balance, motor function, activities of daily living, and upper limb function in children with cerebral palsy: A systematic review and Meta-Analysis. *Ther Clin Risk Manag.* 2024;20:95–109.
92. Virtual Reality in Stroke Recovery. A meta-review of Systematic Reviews. [cited 2024 Aug 17]. Available from: <http://ouci.dntb.gov.ua/en/works/4arpjz-al/>
93. Dixit P, Phalswal U, Kalal N, Srivastava SP. Effectiveness of virtual reality-supported exercise therapy in improving upper extremity function and activities of daily living among patients after stroke: a systematic review of randomized control trials. *Osong Public Health Res Perspect.* 2024;15(3):189–200.
94. Hussain MA, Waris A, Gilani SO, Mushtaq S, Pujari AN, Khan NB, et al. Virtual reality as a non-conventional rehabilitation for stroke: A comprehensive review. *J Neurorestorology.* 2024;12(3):100135.
95. Rodríguez-Hernández M, Polonio-López B, Corregidor-Sánchez AI, Martín-Conty JL, Mohedano-Moriano A, Criado-Álvarez JJ. Can specific virtual reality combined with conventional rehabilitation improve poststroke hand motor function? A randomized clinical trial. *J Neuroeng Rehabil.* 2023;20:38.
96. Kouijzer MMTE, Kip H, Bouman YHA, Kelders SM. Implementation of virtual reality in healthcare: a scoping review on the implementation process of virtual reality in various healthcare settings. *Implement Sci Commun.* 2023;4:67.
97. Aderinto N, Olatunji G, Abdulbasit MO, Edun M, Aboderin G, Egbunu E. Exploring the efficacy of virtual reality-based rehabilitation in stroke: a narrative review of current evidence. *Ann Med.* 2023;55(2):2285907.
98. Maier M, Ballester BR, Duff A, Oller ED, Verschure PFMJ. Effect of specific over nonspecific VR-Based rehabilitation on poststroke motor recovery: A systematic Meta-analysis. *Neurorehabil Neural Repair.* 2019;33(2):112.
99. Massetti T, da Silva TD, Crocetta TB, Guarnieri R, de Freitas BL, Bianchi Lopes P, et al. The clinical utility of virtual reality in neurorehabilitation: A systematic review. *J Cent Nerv Syst Dis.* 2018;10:1179573518813541.
100. Sun Y, Hunt CL, Lamounier EA, Soares AB. Neurorehabilitation with Virtual and Augmented Reality Tools. In: Thakor NV, editor. *Handbook of Neuroengineering*. Singapore: Springer Nature. [cited 2024 Nov 7]. 2020;1–41. Available from: [https://doi.org/10.1007/978-981-15-2848-4\\_49-1](https://doi.org/10.1007/978-981-15-2848-4_49-1)
101. Naro A, Calabrò RS. What do we know about the use of virtual reality in the rehabilitation field? A brief overview. *Electronics.* 2021;10(9):1042.
102. Kim WS, Cho S, Ku J, Kim Y, Lee K, Hwang HJ, et al. Clinical application of virtual reality for upper limb motor rehabilitation in stroke: review of technologies and clinical evidence. *J Clin Med.* 2020;9(10):3369.
103. Kwakkel G, Van Wegen EEH, Burridge JH, Winstein CJ, Van Dokkum LEH, Alt Murphy M, et al. Standardized measurement of quality of upper limb movement after stroke: Consensus-Based core recommendations from the second stroke recovery and rehabilitation roundtable. *Neurorehabil Neural Repair.* 2019;33(11):951–8.
104. Kiper P, Godart N, Cavalier M, Berard C, Cieślak B, Federico S, et al. Effects of immersive virtual reality on Upper-Extremity stroke rehabilitation: A systematic review with Meta-Analysis. *J Clin Med.* 2023;13(1):146.
105. Soleimani M, Ghazisaedi M, Heydari S. The efficacy of virtual reality for upper limb rehabilitation in stroke patients: a systematic review and meta-analysis. *BMC Med Inf Decis Mak.* 2024;24(1):135.
106. Cheong MJ, Kang Y, Kang HW. Psychosocial factors related to stroke patients' rehabilitation motivation: A scoping review and Meta-Analysis focused on South Korea. *Healthcare.* 2021;9(9):1211.
107. Clark B, Whittall J, Kwakkel G, Mehrholz J, Ewings S, Burridge J. Time spent in rehabilitation and effect on measures of activity after stroke. *Cochrane Database Syst Rev.* 2017;2017(3):CD012612.

108. Zhang B, Li D, Liu Y, Wang J, Xiao Q. Virtual reality for limb motor function, balance, gait, cognition and daily function of stroke patients: A systematic review and meta-analysis. *J Adv Nurs*. 2021;77(8):3255–73.
109. Khan A, Podlasek A, Somaa F. Virtual reality in post-stroke neurorehabilitation – a systematic review and meta-analysis. *Top Stroke Rehabil*. 2023;30(1):53–72.
110. Alashram AR, Padua E, Hammash AK, Lombardo M, Annino G. Effectiveness of virtual reality on balance ability in individuals with incomplete spinal cord injury: A systematic review. *J Clin Neurosci*. 2020;72:322–7.
111. Garay-Sánchez A, Suarez-Serrano C, Ferrando-Margelí M, Jimenez-Rejano JJ, Marcén-Román Y. Effects of immersive and Non-Immersive virtual reality on the static and dynamic balance of stroke patients: A systematic review and Meta-Analysis. *J Clin Med*. 2021;10(19):4473.
112. Morone G, Capone F, Iosa M, Cruciani A, Paolucci M, Martino Cinnera A, et al. May dual transcranial direct current stimulation enhance the efficacy of Robot-Assisted therapy for promoting upper limb recovery in chronic stroke?? *Neurorehabil Neural Repair*. 2022;36(12):800–9.
113. Alashram AR. Combined robot-assisted therapy virtual reality for upper limb rehabilitation in stroke survivors: a systematic review of randomized controlled trials. *Neurol Sci*. 2024;45(11):5141–55.
114. Borgonovo E, Plischke E. Sensitivity analysis: A review of recent advances. *Eur J Oper Res*. 2016;248(3):869–87.

### Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.