

TASK-RELATED MODULATION OF CEREBELLAR BRAIN INHIBITION IS MAINTAINED IN OLDER ADULTS

Shanti Van Malderen 1,2,3, Melina Hehl 1,2,3, Marten Nuyts 1, Stephan P. Swinnen 2,3, and Koen Cuypers 1,2,3

(1) Neuroplasticity and Movement Control Research Group, Rehabilitation Research Institute (REVAL), Hasselt University, Diepenbeek, Belgium

(2) KU Leuven, Leuven Brain Institute (LBI), Leuven, Belgium

(3) Movement Control and Neuroplasticity Research Group, Department of Movement Sciences, Group Biomedical Sciences, KU Leuven, Heverlee, Belgium





INTRODUCTION

With the global increase in life expectancy but the rise in healthy life expectancy lagging behind (WHO), it has become crucial to understand the neural processes associated with age-related changes in motor function. While age-related impairments in motor functioning and learning involve various structural (1), functional (2), and biochemical (3) changes in the brain, the integrity of the cerebellum emerges as a key factor influencing motor function in older adults. Consequently, investigating cerebellar neural interactions presents a unique opportunity to comprehend the underlying mechanisms that contribute to age-related deficits in motor control.


OBJECTIVE

The primary aim of this study was to shed light on age-related alterations in cerebellar-M1 interactions in the context of bimanual coordination. To this end, a dual-site transcranial magnetic stimulation (ds-TMS) paradigm was employed to measure cerebellar brain inhibition (CBI), both in a resting state and during the preparation phase of a bimanual motor task.


RESULTS



METHODOLOGY



- Participants: 40 right-handed adults aged 20-40 (younger) or 60-80 (older)
- Protocol: Dual-site transcranial magnetic stimulation (ds-TMS)
- Stimulation Parameters:
 - Conditioning stimulus (CS; double cone coil): 65% MSO
 - Test stimulus (TS; figure-of-eight coil): MEPs of 1 mV
 - ISI: 5 ms
- Experimental Design: One block of 40 trials (20 TS and 20 CS+TS) at rest, and 6 task blocks of 44 trials (20 TS, 20 CS+TS and 4 no TMS) performed in a pseudo-randomized order
- Tolerability Assessment: Visual analogue scale (VAS) used after completion of all experimental blocks



CONCLUSION

- Younger adults perform better relative to older adults on all motor tasks
- No age-related difference in the amount of CBI at rest
- Task-related modulation of CBI (from rest towards task preparation) was maintained in older adults
- During the BTT, low scoring participants showed increased preparatory inhibition in block 2 compared to block 1, whereas high scoring participants exhibited stronger disinhibition/facilitation from baseline to the preparatory period in block 2 relative to block 1
- Perceived discomfort was mild
- Preservation of CBI modulation in older adults might suggest the presence of neural mechanisms (i.e., cerebellar reserve) that counteract the effects of aging on the cerebellum to maintain dexterous movement control (4)

REFERENCES

- Inano, S., H. Takao, N. Hayashi, O. Abe and K. Ohtomo (2011). "Effects of age and gender on white matter integrity." *AJNR Am J Neuroradiol* 32(11): 2103-2109.
- King, B. R., P. van Ruitenbeek, I. Leunissen, K. Cuypers, K. F. Heise, T. Santos Monteiro, L. Hermans, O. Levin, G. Albouy, D. Martini and S. P. Swinnen (2018). "Age-Related Declines in Motor Performance are Associated With Decreased Segregation of Large-Scale Resting State Brain Networks." *Cereb Cortex* 28(12): 4390-4402.
- Gao, F., R. A. Edden, M. Li, N. A. Puts, G. Wang, C. Liu, B. Zhao, H. Wang, X. Bai, C. Zhao, X. Wang and P. B. Barker (2013). "Edited magnetic resonance spectroscopy detects an age-related decline in brain GABA levels." *Neuroimage* 78: 75-82.
- Kawato, M., K. Furukawa and R. Suzuki (1987). "A hierarchical neural-network model for control and learning of voluntary movement." *Biol Cybern* 57(3): 169-185.

TALK TO ME

