



## Research Article

# Specifying time courses of subtypes of spatial neglect after stroke: Necessary or not?

Elissa Embrechts<sup>1,2,3</sup> , Charlotte van der Waal<sup>1</sup> , Tamaya Van Crieckinge<sup>4</sup>, Jonas Schröder<sup>1,5</sup>, Christophe Lafosse<sup>2</sup>, Steven Truijen<sup>1</sup>, Wim Saeys<sup>1,2</sup> and Tanja C.W. Nijboer<sup>3</sup>

<sup>1</sup>Research Group MOVANT, Department of Rehabilitation Sciences and Physical Therapy, University of Antwerp, Wilrijk, Belgium, <sup>2</sup>Department of Neurorehabilitation, Revarte Rehabilitation Hospital, Edegem, Belgium, <sup>3</sup>Helmholtz Institute, Department of Experimental Psychology, Utrecht University, Utrecht, The Netherlands, <sup>4</sup>Department Physical Medicine and Rehabilitation, AZ Delta Roeselare, Roeselare, Belgium and <sup>5</sup>REVAL Research Group, Faculty of Rehabilitation Sciences, University of Hasselt, Diepenbeek, Belgium

## Abstract

**Objective:** Spatial neglect is a heterogeneous post-stroke disorder with subtypes differing in reference frames, processing stages, and spatial domains. While egocentric peri-personal neglect recovery has been studied, recovery trajectories of allocentric peri-personal visuospatial and personal neglect remain unclear. This study investigated recovery time courses of egocentric and allocentric peri-personal visuospatial and personal neglect during the first 12 weeks post-stroke; whether initial severity predicts recovery and defines distinct patient clusters; and how subtypes interrelate over time. **Method:** Forty-one first-ever stroke patients were evaluated at weeks 3, 5, 8, and 12 post-stroke using the Broken Hearts Test, Line Bisection Test, Visuospatial Search Time Test, and Fluff Test. Recovery was analyzed using linear mixed models, clustering with Gaussian finite mixture models, and interrelationships using Spearman correlations. **Results:** Significant improvements occurred in egocentric and allocentric peri-personal visuospatial and personal neglect, primarily between weeks 3 and 5, followed by a plateau. The Line Bisection Test detected no changes. Higher initial severity predicted greater residual impairment. Cluster analysis identified near-normal, mild, and moderate-to-severe baseline subgroups with distinct recovery trajectories. Moderate-to-good correlations ( $\rho = 0.33 - 0.55$ ) emerged between egocentric and allocentric neglect at week 3 and when timepoints were pooled. **Conclusion:** Neglect improved mainly between weeks 3 and 5 after which recovery plateaued, mirroring motor and language recovery and suggesting a shared time-limited window. Initial severity was a determinant of recovery, highlighting the value of early severity stratification to monitor and support recovery potential after stroke. As subtypes are distinctive, assessment should include multiple neglect tests.

**Keywords:** Visuospatial neglect; personal neglect; recovery; time course; stroke; spatial neglect

(Received 10 April 2025; final revision 14 November 2025; accepted 18 November 2025)

## Statement of Research Significance

**Research Question(s) or Topic(s):** This study examined recovery time courses of egocentric and allocentric peri-personal visuospatial and personal neglect; whether initial severity predicts recovery and defines patient clusters; and how subtypes interrelate throughout the first 12 weeks post-stroke. **Main Findings:** Significant improvements occurred across all neglect subtypes, mainly between weeks 3 and 5 post-stroke, followed by a plateau. Higher initial severity predicted greater impairment at later timepoints. Cluster analysis identified near-normal, mild, and moderate-to-severe baseline subgroups with distinct recovery trajectories. Moderate-to-strong correlations emerged only between egocentric and allocentric neglect at week 3 and with pooled data. **Study Contributions:** Unlike prior research, this study examined recovery across multiple neglect subtypes, revealing that recovery time courses in visuospatial and personal neglect parallel those of motor and language recovery, supporting a shared, time-limited recovery window. Findings emphasize the value of early severity stratification and comprehensive assessment using multiple neglect tests.

## Introduction

Spatial neglect is a post-stroke cognitive disorder involving asymmetric attention to space, most often manifesting as reduced

awareness of stimuli opposite the lesion and, less frequently, reduced awareness on the same side (Fellrath et al., 2012; Heilman & Valenstein, 1979; Van der Stigchel & Nijboer, 2010). Rather than

**Corresponding author:** Elissa Embrechts; Email: [elissa.embrechts@uantwerpen.be](mailto:elissa.embrechts@uantwerpen.be)

**Cite this article:** Embrechts E., van der Waal C., Van Crieckinge T., Schröder J., Lafosse C., Truijen S., Saeys W., & Nijboer T.C.W. Specifying time courses of subtypes of spatial neglect after stroke: Necessary or not?. *Journal of the International Neuropsychological Society*, 1–12, <https://doi.org/10.1017/S1355617725101689>

© The Author(s), 2025. Published by Cambridge University Press on behalf of International Neuropsychological Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

being a uniform condition, neglect encompasses various subtypes that differ in their affected *reference frames* (egocentric/viewer-centered, allocentric/object-centered), *processing stages* (sensory (visual/auditory/tactile), representational, or motor), and *spatial domains* (personal, peri-personal, or extra-personal space) (Demeyere & Gillebert, 2019; Williams et al., 2021). Prevalence estimates range from 18% to 80%, reflecting differences in assessment methods, stroke location and severity, and the timing of post-stroke evaluation (Esposito et al., 2021; Williams et al., 2021).

The disorder's heterogeneity poses significant challenges for establishing standardized diagnostic and therapeutic frameworks that address the full spectrum of neglect. Moreover, the limited evidence for the effectiveness of current cognitive interventions (Bowen et al., 2013) may, at least in part, stem from variable treatment responses across subtypes (Carter & Barrett, 2023). A better understanding of how distinct subtypes recover could help refine interventions by identifying whether some follow predictable recovery trajectories while others rather remain resistant to improvement over time. Such insights may also guide the timing of treatment, aligning rehabilitation with subtype-specific periods of heightened recovery potential (Bernhardt et al., 2017).

Prior studies that examined neglect recovery within the first six months post-stroke (i.e., the period of greatest expected recovery (Bernhardt et al., 2017)) have largely focused on peri-personal visuospatial neglect assessed with conventional tools such as cancellation or line bisection tasks (Cassidy et al., 1998; Jehkonen et al., 2000; Jehkonen et al., 2007; Levine et al., 1986; Nijboer et al., 2013; Overman et al., 2024; Samuelsson et al., 1997; Stone et al., 1992). Most of these demonstrate that the greatest improvements occur within the first 12 – 14 weeks post-stroke, followed by a gradual plateau (Cassidy et al., 1998; Jehkonen et al., 2000; Jehkonen et al., 2007; Levine et al., 1986; Nijboer et al., 2013; Overman et al., 2024; Samuelsson et al., 1997; Stone et al., 1992). However, while informative, their narrow focus on (mostly egocentric) peri-personal visuospatial neglect overlooks potential differences in recovery time courses across other neglect subtypes (e.g., allocentric or personal neglect), or other dimensions of the disorder, such as its temporal aspects (e.g., spatial reaction times). Moreover, prior research indicates that initial egocentric visuospatial neglect severity is a significant predictor of neglect recovery (Marchi et al., 2017; Moore et al., 2021; Stone et al., 1992). Yet, it is not known whether this relationship generalizes to other neglect subtypes or dimensions or how its predictive value evolves over the subacute recovery phase. Furthermore, recovery may not be uniform across individuals; some may exhibit distinct patterns of improvement or persistent deficits depending on their initial severity profile.

Another limitation of current literature lies in the limited understanding of how neglect subtypes relate to one another during recovery. While behavioral and neural dissociations between subtypes have been established (Chechlacz et al., 2010; Demeyere & Gillebert, 2019), in the early post-stroke period, deficits are often diffuse due to widespread network disruption or diaschisis (Feeney & Baron, 1986). This raises the possibility that different subtypes may initially co-occur before diverging into the dissociable patterns. Therefore, examining how correlations between subtypes change during recovery can provide a complementary perspective to dissociation studies, capturing transient overlaps that static dissociation analyses alone cannot explain. However, these temporal dynamics during recovery remain unexplored.

Thus, key fundamental questions remain unanswered: Do different neglect subtypes follow similar recovery time courses? Is

initial neglect severity a universal predictor of recovery? And do neglect subtypes covary over time? To address these, this exploratory, non-hypothesis-driven study prospectively investigated the time course of recovery of neglect during the first 12 weeks post-stroke (i.e., early subacute post-stroke phase (Bernhardt et al., 2017)). A comprehensive battery of assessments was employed to capture variations in recovery courses across neglect subtypes. The study had three objectives:

1. To examine the recovery time courses of egocentric and allocentric peri-personal visuospatial neglect (both spatial and temporal aspects), as well as personal neglect, during the first 12 weeks post-stroke,
2. To evaluate whether initial neglect severity is a predictor of their time courses of recovery, to identify distinct clusters based on this initial severity, and to examine transitions between these clusters over time;
3. To investigate how the subtypes interact over time.

## Material and methods

### Study design and setting

This longitudinal cohort study is part of a larger research project, entitled TARGET (Temporal Analyses of hemiplegic Gait and standing balance Early post sTroke; see (Schröder et al., 2022)). This study was approved by the Medical Ethics Committee of the University Hospital Antwerp (No. 18/25/305; Belgium Trial Registration No. B300201837010 and BUN B3002021000098). Additional approval was obtained from the medical ethics committees of the other clinical sites. All procedures were conducted in accordance with the principles of the Declaration of Helsinki. The study protocol was designed in accordance with the STROBE guidelines (von Elm et al., 2008) and was registered online (ClinicalTrials.gov ID NCT05060458).

### Participants

Individuals admitted to one of the four cooperating acute hospitals (Universitair Ziekenhuis Antwerpen, GZA Sint-Vincentius, GZA Sint-Augustinus, Algemeen Ziekenhuis Geel) and 2 rehabilitation facilities (RevArte, AZ Monica), all situated in the larger Antwerp region, Belgium, after an acute stroke were screened for participation between August 2020 and May 2024. For eligibility, they had to meet the following criteria: (1) CT and/or MRI confirmed first-ever unilateral ischemic or hemorrhagic supratentorial stroke, (2) aged between 18 and 90 years, (3) (corrected to) normal visual acuity, (4) premorbid independence in daily life activities (i.e., modified Rankin Scale score of 0 – 1), (5) no prior diagnosis of pre-stroke neurological disease, (6) no severe cognitive or communication deficits that interfere with understanding instructions and procedures. This was determined during recruitment interviews by consulting with the eligible person and/or their caregiver(s), assessing the person's capacity to comprehend instructions and participate in the study, and (7) ability to provide written informed consent. All participants received usual care including physical, occupational, speech, and neuropsychological therapy, depending upon individual needs.

### Protocol, data collection, and outcome measures

Recruitment and screening were performed by EE, CvdW, and JS together with the (para)medical staff employed at the stroke units and rehabilitation facilities. During intake at 3 weeks post-stroke,

the participants' sex, age, and stroke information (type: ischemic/hemorrhagic; affected side: left/right) as well as clinical severity of lower limb motor impairment (Lower Limb Motricity Index) (Demeurisse et al., 1980) and functional mobility (Rivermead Mobility Index) (Collen et al., 1991) were noted. Serial measurements of neglect were employed at weeks 3, 5, 8, and 12 post-stroke onset. A trained assessor (EE or CvdW) administered all follow-up assessments of the same participant.

#### Peri-personal visuospatial neglect tests

**Broken hearts test (BHT).** We used the BHT, part of the Oxford Cognitive Screen (Demeyere et al., 2015), or its variation (Apples test) (Bickerton et al., 2011). These three parallel versions were varied across time points to reduce learning effects. Participants had to cancel complete hearts/apples ( $n = 50$ ) among distractors with left- or right-sided gaps ( $n = 50$  each). The test was presented on standardized A4 landscape sheets and had to be completed within three minutes (Demeyere et al., 2015; Demeyere et al., 2016). For more details and outcome measures, see Table 1.

**Computerized Schenkenberg line bisection test (LBT).** Participants bisected 20 horizontal lines using their less-affected hand. Lines were presented centrally or shifted left/right, with equal peripheral starting points (Vaes et al., 2015). The test was administered using the Metrisquare DiaDiag software ([www.metrisquare.com](http://www.metrisquare.com)) on a Wacom® tablet ( $40 \times 65$  cm) (Vaes et al., 2015). For more details and outcome measures, see Table 1.

**Computerized visuospatial search time test (VSTT).** The VSTT consists of 16 consecutive grids containing 20 different stimuli centered around one stimulus in the middle, presented in a green square (Vaes et al., 2015). The participant had to cross out a stimulus identical to the central stimulus as quickly as possible. Directly afterwards, the next grid was shown with a different central stimulus, and in this way, the task continued. The task recorded ipsilesional and contralesional search times, using the same DiaDiag software and Wacom® tablet ( $40 \times 65$  cm) (Vaes et al., 2015). For more details and outcome measures, see Table 1.

#### Personal neglect test

**Fluff test.** Fifteen targets were placed on the contralesional side of the body (six on the arm, six on the leg, and three on the trunk) and nine on the ipsilesional side (six on the leg and three on the trunk). Participants were blindfolded and unaware of the targets being attached, as placement occurred during a separate sensory (tactile) test not included in this study. They were then instructed to remove all targets using their less-affected hand (Cochinini & Beschin, 2020). If motor deficits (e.g., limited trunk control) interfered with target removal, the examiner assisted with positioning (e.g., supporting a seated posture) while minimizing sensory feedback on the limbs and trunk. For details on outcome measures, see Table 1.

#### Statistical analyses

Analyses included participants with at least two consecutive follow-up assessments. Individuals with missing data between intermediate timepoints (e.g., weeks 3–8) were excluded, though dropouts before week 12 were permitted.

Demographic, clinical, and neglect outcome variables (BHT egocentric and allocentric asymmetry, VSTT index, LBT deviation,

Fluff test asymmetry) were summarized as means  $\pm$  SD at weeks 3, 5, 8, and 12 (Table 2).

#### Objective 1: Recovery time courses

We first analyzed data from the entire cohort, regardless of whether participants met predefined cutoffs for neglect at baseline (week 3, see Table 1) to capture the natural symptom variability in a clinically representative population and avoid a strict baseline dichotomy that could obscure meaningful fluctuations in severity across the spectrum of impairment. Because longitudinal effects in the full sample are likely driven by those with initial neglect, also test-specific subgroup analyses restricted to individuals with clinically significant impairment (i.e., outside normative ranges) at baseline were performed.

For each neglect outcome, linear mixed models (LMMs) were fitted with TIME (categorical: weeks 3, 5, 8, 12) as a fixed effect and a subject-specific random intercept for repeated measures. To further examine the effect of lesion laterality, given that neglect is typically more frequent and severe after right-hemisphere lesions (Cazzoli et al., 2025; Esposito et al., 2021), additional LMMs were conducted in those with clinically significant neglect, including lesion side and its interaction with TIME as fixed factors.

To prevent misinterpreting directional shifts over time (e.g., from ipsilesional to contralesional and vice versa) as improvement or deterioration of neglect, absolute/side-neutral values were used for each dependent variable. Model assumptions were checked via histograms, Q-Q plots, and residuals-versus-predicted plots. Due to violations, BHT allocentric, BHT egocentric, VSTT index, and LBT deviation were log-transformed; a constant of 1 was added to BHT asymmetry scores prior to transformation to handle zeros. Post-hoc Tukey's HSD tests estimated changes (regression coefficients  $\beta$ ) across the full period (weeks 3–12) and individual epochs (weeks 3–5, 5–8, 8–12). Log-transformed  $\beta$  estimates were back-transformed using  $Exp(\beta)$ . These analyses were performed using JMP Pro® version 16.

#### Objective 2: Baseline severity and clustering

The same LMM structure was applied to the entire cohort, with baseline neglect severity (absolute score at 3 weeks) and its interaction with TIME as additional covariates.

Cluster analysis was conducted in R (version 2025.09.1) using the mclust package. For clustering, only baseline measurements of the VSTT index and BHT egocentric asymmetry were included (absolute/side-neutral values) and standardized. These were chosen to capture core dimensions of neglect represented in the largest subsamples of our cohort (both spatial and temporal bias of egocentric neglect). Gaussian finite mixture models with different covariance structures (VVE, VEV, VVV, EEE, EVV) were fitted, with optimal model selection based on Bayesian Information Criterion (BIC) and Integrated Completed Likelihood (ICL). Longitudinal trajectories were examined by linking cluster assignments from week 3 to subsequent timepoints, and individual progress was visualized in plots with cluster-colored trajectories, where arrows indicated change between sessions. Cluster characteristics were summarized with means  $\pm$  SD for continuous variables and proportions for categorical variable (See Supplementary material).

#### Objective 3: Associations between neglect outcome measures

Spearman correlations assessed associations between neglect measures across pooled timepoints and at each timepoint. Non-parametric methods were used due to non-normal data

**Table 1.** Neglect tests and their corresponding characteristics, outcome measures and cut-off scores

Neglect test	Subtype assessed	Processing stage	Reference frames	Spatial domain	Test type	Outcome	Outcome explanation	Neglect considered present when
BHT	Egocentric peri-personal visuo-spatial neglect	Sensory: visual (Williams et al., 2021)	Egocentric (Williams et al., 2021)	Peri-personal (Williams et al., 2021)	P&P	cancellation	Egocentric asymmetry (spatial)	Difference between the cancelled full outlines of the CL and IL sides of the paper
	Asymmetry $> 2$ (IL neglect) or $< -2$ (CL neglect) (Demeyere et al., 2015)							
BHT	Allocentric peri-personal visuo-spatial neglect	Sensory: visual (Williams et al., 2021)	Allocentric (Williams et al., 2021)	Peri-personal (Williams et al., 2021)	P&P	cancellation	Allocentric asymmetry (spatial)	Calculated by subtracting the number of IL with CL gap false positives
	Asymmetry $> 1$ (IL neglect) or $< -1$ (CL neglect) (Demeyere et al., 2015)							
LBT	Peri-personal visuo-spatial neglect (depends on both an egocentric and allocentric space representation (Rorden et al., 2006))	Sensory: visual (Williams et al., 2021)	Ego - and allocentric (Williams et al., 2021)	Peri-personal (Williams et al., 2021)	Bisection	computerized	Deviation from midline (spatial)	Mean percentage of total deviation from centers of IL, CL, and centrally placed lines
Outside	normative range (Larger deviation than $0.4 \pm 3.89$ ) (Vaes et al., 2015)							
VSTT	Egocentric peri-personal visuo-spatial neglect	Sensory: visual (Williams et al., 2021)	Egocentric (Williams et al., 2021)	Peri-personal (Williams et al., 2021)	Search time test	computerized	VSTT index (temporal)	Ratio between CL and IL visuospatial search times
Outside	normative range (Larger than $1.1 \pm 0.39$ ) (Vaes et al., 2015)							
Fluff Test	Personal neglect	Represen-tational: body (Williams et al., 2021)	NA	Personal (Williams et al., 2021)	Bodily	Fluff asymmetry (spatial)	Performance on ipsilesional side (number of targets found compared to total possible ipsilesional targets, in % ( $n = 9$ )) compared with performance on contralateral side (number of targets found compared to total possible contralateral targets, in % ( $n = 15$ )))	Difference of $> 13.3\%$ in asymmetry (positive sign = CL, negative sign = IL). Higher asymmetry indicates more severe neglect (Cocchini & Beschin, 2020)

Abbreviations: BHT = broken hearts test, CL = contralesional, IL = ipsilesional, LBT = line bisection test, NA = not applicable, P&P = paper-and-pencil test, VSTT = visuospatial search time test

**Table 2.** Demographic and clinical characteristics of the participants at each timepoint

	Week 3	Week 5	Week 8	Week 12
Number of participants (N)	41	41	40	27
Time post-stroke (days)	25.17 (1.79)	38.41 (2.61)	58.61 (2.53)	85.42 (2.80)
<b>Neglect information</b>				
Egocentric asymmetry on the BHT (0-20, absolute values) in complete group, left-sided strokes and right-sided strokes	3.17 (4.67), L: 2.25 (4.04), R: 3.76 (5.03)	1.54 (2.31), L: 0.44 (0.63), R: 2.24 (2.71)	1.58 (2.08), L: 0.94 (1.18), R: 2.05 (2.46)	1.41 (2.49), L: 1.09 (0.94), R: 1.63 (3.16)
Number of participants with/without egocentric visuospatial neglect based on egocentric asymmetry (BHT)	15/26 [of which 11CL and 4IL]	6/35 [of which 5CL and 2IL]	9/31 [of which 7L and 2R sided]	3/24 [of which 2L and 1R sided]
Allocentric asymmetry on the BHT (0-20, absolute values)	1.70 (4.79), L: 0.69 (0.95), R: 2.38 (6.10)	0.78 (2.51), L: 1.00 (3.74), R: 0.64 (1.29)	0.61 (1.50), L: 0.81 (1.87), R: 0.46 (1.18)	0.52 (1.05), L: 0.36 (0.92), R: 0.63 (1.15)
Number of participants with/without allocentric visuospatial neglect based on allocentric asymmetry (BHT)	7/34 [of which 6CL and 1IL]	6/35 [of which 5CL and 1IL]	6/34 [all CL]	4/23 [of which 3CL and 1IL]
VSTT index (side neutral values)	2.18 (1.46), L: 1.68 (0.79), R: 2.53 (1.71)	1.60 (0.68), L: 1.42 (0.41), R: 1.72 (0.80)	1.58 (0.65), L: 1.42 (0.33), R: 1.69 (0.79)	1.46 (0.44), L: 1.26 (0.24), R: 1.61 (0.50)
Number of participants with/without egocentric visuospatial neglect based on VSTT index	24/17 [of which 19CL and 5IL]	16/25 [of which 12CL and 4IL]	18/22 [of which 14CL and 4IL]	10/17 [of which 8CL and 2IL]
LBT deviation (%), absolute values)	5.34 (4.48), L: 6.27 (3.26), R: 4.75 (5.10)	4.81 (6.03), L: 4.41 (3.60), R: 5.04 (7.14)	4.54 (3.17), L: 4.28 (2.82), R: 4.71 (3.43)	5.24 (3.60), L: 4.80 (2.28), R: 5.56 (4.36)
Number of participants with/without visuospatial neglect based on LBT deviation	17/24 [of which 12CL and 5IL]	15/26 [of which 7CL and 5IL]	20/20 [of which 1CL and 10IL]	13/14 [of which 2CL and 11IL]
Fluff test asymmetry (%), absolute values)	10.22 (15.70), L: 11.11 (15.90), R: 9.63 (15.89)	5.33 (12.90), L: 6.39 (13.67), R: 4.63 (12.60)	5.07 (13.40), L: 4.86 (13.29), R: 5.22 (13.77)	1.71 (4.76), L: 3.03 (6.91), R: 0.74 (2.00)
Number of participants with/without personal neglect based on Fluff test	13/28 [all CL]	6/35 [of which 5CL and 1IL]	5/35 [all CL]	2/25 [all CL]
<b>Clinical severity</b>				
Lower limb motricity index score (on 99 + 1)	62.26 (25.54)	70.75 (23.76)	74.70 (23.41)	72.96 (21.57)
Rivermead Mobility Index (on 24)	7.49 (4.39)	9.56 (4.63)	10.49 (4.30)	11.00 (3.83)

Abbreviations. BHT = Broken Hearts Test; CL = contralesional; IL = ipsilesional; L = left; R = right. Unsigned/absolute values were used for the parameters presented. Values are mean (standard deviation).

distributions. Correlation matrix heatmaps were generated, and coefficients were interpreted as:  $\leq .30$  (no meaningful relationship),  $.30 - .50$  (moderate),  $.50 - .70$  (good),  $> .70$  (very good) (Portney, 2020). Bonferroni correction adjusted the significance threshold to  $\alpha = .005$  ( $\alpha = .05/10$  correlations). Analyses used JMP Pro® version 16.

## Results

### Participants

A total of 210 potentially eligible individuals were screened, of which 42 were enrolled. Of these, 41 successfully participated in at least two serial measurements and were included in the analysis (Figure 1).

### Descriptive data

Participants had a mean age of  $59.2 \pm 15.6$  years; 17 (41.5%) were female, 11 (26.8%) had left-sided strokes, and 33 (80.5%) ischemic strokes. Mean time post-stroke was  $25.3 \pm 2.0$  days at week 3,  $38.4 \pm 2.6$  at week 5,  $58.6 \pm 2.5$  at week 8, and  $85.4 \pm 2.8$  at week 12.

Table 2 presents mean absolute/side-neutral scores for all neglect measures, the number of participants outside unimpaired ranges (per subtype) and with ipsilesional or contralesional neglect, and clinical severity scores (Lower Limb Motricity Index, Rivermead Mobility Index).

A dropout rate of 32.5% occurred from 8 weeks post-stroke onward (Figure 1). To evaluate potential bias, neglect severity was compared between those who completed all sessions and those who dropped out after week 8 with the Wilcoxon signed rank test.

This showed no significant differences in neglect severity between completers and dropouts across timepoints for any neglect measure. Full statistical details are provided in Supplementary Table 1.

### Objective 1: Time course of visuospatial peri-personal neglect and personal neglect recovery

Figure 2 illustrates individual recovery time courses across different tests. As indicated in Figure 2 and Table 2, the mean values for all tests fall outside unimpaired ranges at week 3. By 5 weeks, most scores returned to within normal limits, except for LBT deviation and Fluff test asymmetry. (See also Figure 3 in the section 'Objective 3: Association between Neglect Outcomes over Time', for mean  $\pm$  SD with corresponding correlation heatmaps over time)

We first examined neglect severity across the full cohort. Egocentric visuospatial neglect severity (BHT egocentric asymmetry, VSTT index) significantly decreased between weeks 3 and 12 post-stroke (BHT:  $\text{Exp}(\beta)-1 = -0.41$ , 95%CI [-0.99, -0.01],  $P = .049$ ; VSTT index:  $\text{Exp}(\beta) = -1.35$ , 95%CI [-1.65, -1.10],  $P = .001$ ). Post-hoc analysis revealed a significant decrease in severity between weeks 3 and 5 only (BHT:  $\text{Exp}(\beta)-1 = -0.41$ , 95%CI [-0.90, -0.05],  $P = .017$ ; VSTT index:  $\text{Exp}(\beta) = -1.24$ , 95%CI [-1.65, -1.05],  $P = .006$ ). No significant changes were observed beyond this period. Additionally, visuospatial neglect severity did not decrease over time when assessed using BHT allocentric asymmetry ( $P = .453$ ) or LBT deviation ( $P = .315$ ) (Table 3).

Personal neglect severity (Fluff Test asymmetry) decreased significantly from weeks 3 – 12 ( $\beta = -6.51$ , 95%CI [-12.01, -1.00],




Figure 1. Screening, recruitment and follow-up flowchart.




Figure 2. Individual time courses of recovery for each neglect test, presented by lesion side.


Note. Individual trajectories are shown as lines, with solid blue lines representing participants with right-sided lesions and dotted red lines representing participants with left-sided lesions. Grey bands indicate the normative (non-impaired) performance range.

$P = .014$ ), driven by improvement between weeks 3 – 5 ( $\beta = -4.89$ , 95% CI  $[-9.63, -0.15]$ ,  $P = .014$ ), with no further changes thereafter (Table 3).

#### Subgroup analyses: Individuals outside normative ranges at week 3

Additional analyses focused on test-specific subgroups (i.e., participants with baseline scores outside normative ranges) to evaluate recovery in individuals with clinically significant neglect at 3 weeks post-stroke. These show significant decrease in neglect

severity (weeks 3 – 12) for BHT egocentric asymmetry ( $\text{Exp}(\beta) - 1 = -4.80$ , 95%CI  $[-8.09, -1.52]$ ,  $P = .002$ ), BHT allocentric asymmetry ( $\text{Exp}(\beta) - 1 = -8.15$ , 95%CI  $[-15.15, -1.14]$ ,  $P = .019$ ), VSTT index ( $\text{Exp}(\beta) = -1.04$ , 95%CI  $[-1.79, -0.29]$ ,  $P = .003$ ), and Fluff Test asymmetry ( $\beta = -2.58$ , 95%CI  $[-4.06, -1.10]$ ,  $P = .001$ ). Post-hoc analyses revealed significant decreases between weeks 3 and 5 for BHT egocentric asymmetry ( $(\text{Exp}(\beta) - 1 = -4.27$ , 95%CI  $[-7.13, -1.41]$ ,  $P = .002$ ), BHT allocentric asymmetry ( $(\text{Exp}(\beta) - 1 = -8.24$ , 95%CI  $[-14.36, -2.12]$ ,  $P = .006$ ), VSTT index ( $\text{Exp}(\beta) = -0.73$ , 95%CI  $[-1.40, -0.05]$ ,  $P = .029$ ), and Fluff test



**Figure 3.** Clusters that emerged from the Gaussian finite mixture model, with longitudinal trajectory from week 3 to 12.  
Note. Three initial severity clusters are shown, with color-coded cluster membership at week 3. Arrows indicate individual changes between week 3 and 12.

**Table 3.** Linear mixed model results for neglect outcomes across the full period (weeks 3–12) and individual epochs

		Week 3 – 12	Week 3 – 5	Week 5 – 8	Week 8 – 12
Visuospatial neglect variables (back-transformed)					
Δ Egocentric asymmetry on the BHT (0–20)	((exp) $\beta$ )-1	−0.41	−0.41	+ 0.08	−0.08
	((exp)SE)-1	0.13	0.12	0.12	0.13
	(exp)95% CI	[−2.00;−1.00]	[−1.90;−1.05]	[1.26;1.47]	[−1.54;0.76]
	P-value	.027*	.005*	.827	.929
Δ VSTT index	((exp) $\beta$ )	−1.37	−1.27	−1.00	−1.08
	((exp)SE)	0.08	1.07	1.07	1.08
	(exp)95% CI	[−1.67;−1.12]	[−1.50;−1.07]	[−1.18;0.84]	[−1.32;0.89]
	P-value	< .001*	.006*	1.000	1.000
Δ Allocentric asymmetry on the BHT (0–20)	((exp) $\beta$ )-1	−0.45	−0.47	+ 0.09	−0.08
	((exp)SE)-1	0.14	0.12	0.12	0.14
	(exp)95% CI	[−0.19;0.84]	[−0.12;0.85]	[−0.30;0.52]	[−0.39;0.41]
	P-value	.598	.330	.996	.963
Δ Midline deviation on the LBT	((exp) $\beta$ )	−1.22	−1.59	+ 0.72	−1.06
	((exp)SE)	1.33	1.28	1.28	1.32
	(exp)95% CI	[−0.58; 2.58]	[−0.83;3.06]	[−0.38;1.37]	[−0.51;2.21]
	P-value	.754	.525	.802	.938
Personal neglect variables (non-transformed)					
Δ Asymmetry on the Fluff test (%)	$\beta$	−6.51	−4.89	−0.27	−1.89
	SE	2.11	1.81	1.84	2.12
	95% CI	[−12.01; −1.01]	[−9.63; −0.15]	[−7.12; 3.88]	[−7.41;3.64]
	P-value	.015*	.040*	.999	.925

BHT = Broken Hearts Test, CI = confidence interval, LBT = Line Bisection Test, SE = standard error, VSTT = Visuospatial Search Time Test,  $\Delta$  = difference,  $\beta$  = estimate,  $(\exp)\beta$  = back-transformed value, \* $P < .05$ .

asymmetry ( $\beta = -6.05$ , 95%CI  $[-28.91, -1.27]$ ,  $P = .019$ ), with a plateau afterwards. No significant decrease was observed for LBT deviation ( $P = .208$ ).

Regarding lesion-side effects within this subset, none were observed for BHT egocentric asymmetry, LBT deviation, or Fluff test asymmetry. However, for BHT allocentric asymmetry, right-hemisphere lesions were associated with larger asymmetry ( $F = 18.29$ ,  $p = .008$ ;  $\text{Exp}(\beta) = 0.66$ ,  $p = .008$ ), without a time interaction. For the VSTT index, a significant time  $\times$  lesion side interaction was found ( $F(3, 60.8) = 2.9$ ,  $p = .048$ ), with participants with left-sided lesions showing lower (i.e., better) VSTT index scores at 3 weeks ( $\text{Exp}(\beta) = -0.81$ ,  $p = .012$ ) than right-sided lesions, whereas no significant differences were observed at 5 or 8 weeks.

### **Objective 2: Influence of initial neglect severity on neglect recovery time courses and cluster analysis**

Initial neglect severity significantly interacted with TIME for BHT egocentric asymmetry ( $F = 9.49$ ,  $P < .0001$ ), BHT allocentric asymmetry ( $F = 25.14$ ,  $P < .0001$ ), VSTT index ( $F = 8.97$ ,  $P < .0001$ ), and Fluff Test asymmetry ( $F = 13.00$ ,  $P < .0001$ ). Higher initial severity predicted higher BHT egocentric and allocentric asymmetry and VSTT index scores at 12 weeks ( $\text{Exp}(\beta-1) = 0.07$ , 95%CI  $[0.03, 0.12]$ ,  $P = .002$ ;  $\text{Exp}(\beta-1) = 0.05$ , 95%CI  $[0.01, 0.08]$ ,  $P = .008$ , and  $\text{Exp}(\beta) = 1.29$ , 95%CI  $[1.20, 1.39]$ ,  $P < .0001$ , respectively), and higher Fluff test asymmetry at 5 weeks ( $\text{Exp}(\beta-1) = 0.05$ , 95%CI  $[0.02, 0.08]$ ,  $P = .002$ ) and 8 weeks ( $\text{Exp}(\beta-1) = 0.04$ , 95%CI  $[0.01, 0.07]$ ,  $P = .014$ ). Adding initial neglect severity to the model did not influence LBT deviation ( $P = .770$ ).

Cluster analysis indicated that a VEV model, allowing variable cluster volume and orientation but equal shape, provided the best fit ( $\text{BIC} = -147.09$ ;  $\text{ICL} = -148.76$ ). Three clusters were identified: Cluster 1 ( $n = 20$ ) showed mild neglect (VSTT index  $M = 1.73$ ,  $SD = 0.60$ ; BHT egocentric asymmetry  $M = 2.20$ ,  $SD = 1.15$ ), Cluster 2 ( $n = 9$ ) showed moderate-to-severe deficits (VSTT  $M = 4.17$ ,  $SD = 1.79$ ; BHT egocentric asymmetry  $M = 9.33$ ,  $SD = 6.87$ ), and Cluster 3 ( $n = 10$ ) demonstrated near-normal performance (VSTT  $M = 1.29$ ,  $SD = 0.17$ ; BHT egocentric asymmetry  $M = 0$ ). As shown in Figure 3, those in Clusters 1 and 3 generally showed either modest improvement (Cluster 1  $\rightarrow$  3) or stable near-normal performance (Cluster 3). In contrast, Cluster 2 showed recovery, with nearly all individuals moving toward the performance range of Cluster 1 by week 12 (See Supplementary Material for more details on cluster characteristics and epoch-by-epoch trajectories).

### **Objective 3: Association between neglect outcomes over time**

Figure 4 shows time courses of neglect for each outcome measure (mean  $\pm$  SD), including correlation heatmaps per timepoint. At 3 weeks post-stroke, a strong correlation was observed between the VSTT index and BHT allocentric asymmetry ( $\rho = 0.54$ ,  $P < .001$ ), and a moderate correlation between the VSTT index and LBT deviation ( $\rho = 0.33$ ,  $P = .002$ ). At 5, 8, and 12 weeks, no significant or meaningful correlations were found.

Across all timepoints, statistically significant ( $\alpha = 0.005$ ) moderate correlations were observed between the VSTT index and BHT egocentric ( $\rho = 0.38$ ,  $P < .001$ ) and allocentric asymmetry ( $\rho = 0.45$ ,  $P < .001$ ). Remaining correlations were not significant ( $P > .005$ ) (Figures in Supplementary Files).

## **Discussion**


This prospective study examined recovery time courses of neglect subtypes during the first 12 weeks post-stroke, the role of initial neglect severity, and associations between neglect subtypes over time. Results show that egocentric peri-personal neglect (BHT asymmetry, VSTT index), allocentric peri-personal neglect (BHT asymmetry), and personal neglect (Fluff Test asymmetry) improved between weeks 3 and 5, after which recovery plateaued. In contrast, line bisection deviation showed no recovery overall. Higher baseline severity predicted poorer outcomes across multiple neglect subtypes, though effects were modest. Cluster analysis based on this initial severity identified three groups (near-normal, mild, moderate-to-severe) that followed distinct recovery trajectories, with initially more impaired individuals showing gradual improvement, yet rarely reaching near-normal performance by week 12. Correlations between some neglect measures were present early but disappeared after 5 weeks.

### **Time courses of recovery across neglect subtypes**

Analyses demonstrated recovery of egocentric peri-personal neglect within the first 12 weeks post-stroke, with the most pronounced improvements in the first five weeks, after which recovery plateaued. This aligns with Stone *et al.* (1992), who also examined visuospatial neglect recovery over the first 12 weeks post-stroke, though they reported stabilization later, around week 8. The parallel improvements in spatial (BHT) and temporal (VSTT) dimensions of egocentric visuospatial neglect suggest that reductions in spatial attentional bias are accompanied by faster visuospatial information processing in the neglected hemispace. Moreover, it shows that both a paper-and-pencil cancellation test (BHT) and digitized task (VSTT) are equally informative for monitoring recovery. Personal neglect followed a similar time course, whereas allocentric neglect showed significant recovery only in the subgroup of participants with clinically relevant impairments at baseline. In both cases, improvements were confined to weeks 3 to 5, followed by a plateau. The observed recovery time courses mirror those reported for motor and language deficits (Duncan *et al.*, 1994; Kwakkel *et al.*, 2006; Lazar *et al.*, 2010), underscoring a critical early phase in which recovery is most pronounced and supporting the existence of a time-limited window of recovery that extends across neurological domains.

Additional analyses considering lesion side revealed that right-hemisphere lesions were associated with greater allocentric asymmetry over time and more pronounced early search time deficits, consistent with prior research showing more severe visuospatial deficits in right-hemisphere lesions (Chechlacz *et al.*, 2012), although lesion effects were limited to specific tasks.

In contrast, LBT deviation did not show change over time. This differs from Nijboer *et al.* (2013), who reported gradual recovery up to 14 weeks post-stroke. The discrepancy likely reflects methodological differences: our four assessments versus their 11, a digitized large-screen LBT (40  $\times$  65 cm, 20 lines) versus their paper version (A4, 10 lines), and our smaller sample with milder baseline deviation (13 – 20 participants, mean 5° vs. 52 participants, mean 7°) (Nijboer *et al.*, 2013). Moreover, despite selecting the subgroup of individuals with neglect symptoms at baseline and thereby reducing baseline variability in our study, LBT scores remained highly variable across timepoints, which may have contributed as well by masking potential recovery at the group level.



**Figure 4.** Time courses of recovery per neglect test with correlation heatmaps per timepoint. Note. Mean (standard deviation) recovery time courses are shown for each neglect test, separately for participants with left-sided and right-sided lesions. At each timepoint, a correlation heatmap is included, showing correlation strength and significance between the neglect tests.

### Influence of initial neglect severity on recovery and cluster analysis

Greater initial neglect severity predicted more severe egocentric, allocentric, and personal neglect at later timepoints, whereas LBT deviation remained unaffected by this. These findings align partly with previous work showing baseline severity as a prognostic factor for egocentric neglect recovery (Marchi et al., 2017; Margaret J. Moore et al., 2021; Stone et al., 1992). We also observed a similar effect in allocentric neglect, contrasting with Moore et al. (2021),

who reported no such relationship despite using the same test. Baseline severity further predicted outcomes in personal neglect, suggesting that its prognostic value may extend across neglect subtypes. Nevertheless, effects were modest: initial severity explained only a small proportion of variance in later test scores.

Cluster analysis provides a complementary perspective, suggesting that initial neglect severity may meaningfully differentiate patient subgroups. Takamura et al. (2021) similarly used multivariate clustering to identify behavioral subgroups within the neglect population, though their cross-sectional approach focused

on lateralized versus non-lateralized attention deficits. Our longitudinal approach extends this work by examining how neglect subgroups evolve over time based on lateralized spatial and temporal characteristics, providing additional insight into individual recovery. Individuals with initially severe deficits tended to shift toward the mild cluster by week 12 but remained distinct from the near-normal cluster. Conversely, those with milder initial impairments progressed toward near-normal performance. These findings demonstrate that early severity may capture clinically relevant heterogeneity in recovery and suggest that normalization rarely occurs in those with severe initial deficits.

### **Correlations between neglect subtypes over time**

Despite mostly similar recovery time courses of the neglect subtypes, significant correlations between their measures were generally absent, particularly beyond five weeks post-stroke. Moderate-to-strong correlations were observed only between egocentric and allocentric peri-personal neglect measures, and only when all timepoints were pooled or at 3 weeks post-stroke, when neglect severity and between-subject variability were greatest. Beyond 5 weeks, when many participants had recovered to unimpaired ranges, correlations were no longer detectable. This indicates that neglect subtypes may overlap shortly after stroke, likely reflecting shared vulnerability to diffuse disruption (Feeney & Baron, 1986), but become increasingly distinct as recovery progresses, in line with evidence for their dissociable nature (Guilbert, 2023; Williams *et al.*, 2021).

### **Strengths and limitations**

This study's primary strength is its longitudinal design with fixed timepoints relative to stroke onset, aligning with Stroke Recovery and Rehabilitation Roundtable recommendations (Bernhardt *et al.*, 2017). This approach controls variability in post-stroke timing. The incorporation of both traditional and digitized assessment tools for neglect further strengthened our approach, though several considerations merit discussion regarding our findings.

The modest sample size ( $n = 42$ ) and 32.5% dropout rate from week 8 onward reflect common challenges in longitudinal stroke research. Dropouts were primarily attributed to difficulties rescheduling follow-up sessions after discharge and COVID-19 restrictions limiting outpatient access. Yet, most dropouts occurred after recovery had plateaued, and post-hoc analyses confirmed no differences in early neglect severity between completers and dropouts, suggesting minimal impact on our core findings.

Our assessment timeline, beginning 3 weeks post-stroke, may have missed very early improvements, and the lack of detailed neuroimaging data (e.g., lesion location and size) prevented examination of their impact on recovery. These limitations stemmed from our recruitment setting, as most participants were enrolled in rehabilitation facilities after acute hospital discharge, where access to acute neuroimaging data was rarely accessible to the research team. This highlights the need for future work to integrate acute-phase imaging or lesion-symptom mapping approaches. Moreover, the lack of systematic monitoring of rehabilitation content and dosage (occupational, physical, and neuropsychological therapy) across hospitals prevented evaluation of how therapy variations might impact outcomes. Moreover, our stroke sample was relatively young (mean age 59.2 years) in comparison to the average age for stroke reported by prevalence

studies (Béjot, 2023; Li *et al.*, 2018; Retho *et al.*, 2023). This limits the generalizability of our findings to older stroke populations, as younger individuals could exhibit faster or more complete recovery trajectories, greater neuroplasticity, or different responses to rehabilitation (Yoo *et al.*, 2020).

We used asymmetry scores for BHT metrics, as they are clinically interpretable and widely applied. Yet, they lack the granularity of alternative measures such as Centre of Cancellation (Rorden & Karnath, 2010) and proportional allocentric scores (M. J. Moore *et al.*, 2021), which more sensitively capture spatial distribution and can distinguish qualitatively different cancellation patterns that yield similar asymmetry values. In addition, neglect subtypes were assessed with a targeted set of validated tests, reflecting the broader design of the TARGET project (Schröder *et al.*, 2022), which primarily focused on motor recovery. While this approach does not capture the full complexity of neglect, it enabled us to identify meaningful recovery patterns and subtype-specific trajectories within a protocol that remains feasible and implementable in clinical contexts.

### **Implications for clinical practice and future research**

This study shows that most participants appeared to have recovered from neglect by 5 weeks post-stroke. However, because conventional assessments may overlook more subtle, persistent symptoms (Menon-Nair *et al.*, 2006), this should not be taken as evidence of complete resolution. Instead, it may highlight limitations in the sensitivity of these assessments. Future research on neglect recovery should therefore employ more fine-grained and/or ecologically valid assessments to capture potential residual symptoms. Although not evaluated in our study, digital platforms can enhance sensitivity by automatically computing advanced metrics (e.g., Centre of Cancellation, proportional allocentric indices) and by capturing continuous performance. Eye-tracking may provide a further complement in detecting residual impairments and compensatory scanning strategies that endpoint scores alone cannot (Embrechts *et al.*, 2025). Moreover, our test battery primarily assessed neglect at the body function level of the International Classification of Functioning, Disability and Health (ICF) (WHO, 2001; Williams *et al.*, 2021), rather than at the activity level where real-life performance occurs (Williams *et al.*, 2021). Future research should therefore incorporate assessments with higher ecological validity, such as the Catherine Bergego Scale (Azouvi, 2017), which evaluates neglect during activities of daily living. Innovative approaches using virtual or augmented reality offer promising avenues for simulating complex, dynamic real-world scenarios, enabling assessment across multiple reference frames and spatial dimensions under more naturalistic task demands (Cavedoni *et al.*, 2022).

Finally, to better understand factors influencing recovery, future studies should recruit larger, more diverse samples that include balanced representation across biological sex, age, neglect severity, and left- versus right-lateralized presentations. This will allow systematic investigation of the patient- and task-specific variables that differentiate individuals who, for example, achieve full recovery from those who exhibit persistent deficits.

### **Conclusion**

This study demonstrated significant recovery in egocentric and allocentric peri-personal visuospatial neglect and personal neglect during the first 12 weeks post-stroke. Most improvement occurred within the initial 3 – 5 weeks, after which it plateaued, mirroring

trajectories reported for motor and language impairments (Duncan et al., 1994; Kwakkel et al., 2006; Lazar et al., 2010). This supports a shared, time-limited recovery window across neurological domains.

Initial neglect severity modestly predicted later outcomes, with greater initial impairment linked to more severe deficits later on. Cluster analysis identified near-normal, mild, and moderate-to-severe baseline severity groups, each following distinct recovery trajectories. Participants in the higher-severity cluster showed often transitioned to the milder cluster, while those within the mild cluster tended to migrate toward the near-normal cluster.

The absence of strong correlations between neglect subtypes underscores their distinctiveness and highlights the need for assessment using multiple neglect measures to capture the disorder's heterogeneity.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S1355617725101689>.

**Data availability.** The data supporting the findings of this study are available from the corresponding author, EE, upon reasonable request.

**Acknowledgements.** The authors thank Prof. Dr Erik Fransen (StatUA, UAntwerpen) for the statistical guidance.

**Funding statement.** This work was supported by the BOF University Research Fund under a DOC PRO Grant [40,180], a post-doc FWO grant [1232425N] and a doctoral FWO grant for strategic basic research [1S64819N]. Open access funding provided by Utrecht University.

**Competing interests.** The authors declare that they have no conflict of interest.

## References

Azouvi, P. (2017). The ecological assessment of unilateral neglect. *Annals of Physical and Rehabilitation Medicine*, 60(3), 186–190.

Béjot, Y. (2023). Age gap between stroke patients included in randomized clinical trials of acute revascularization therapy and those in population-based studies: A review. *Neuroepidemiology*, 57(2), 65–77.

Bernhardt, J., Hayward, K. S., Kwakkel, G., Ward, N. S., Wolf, S. L., Borschmann, K., Krakauer, J. W., Boyd, L. A., Carmichael, S. T., Corbett, D., & Cramer, S. C. (2017). Agreed definitions and a shared vision for new standards in stroke recovery research: The stroke recovery and rehabilitation roundtable taskforce. *International Journal of Stroke*, 12(5), 444–450.

Bickerton, W. L., Samson, D., Williamson, J., & Humphreys, G. W. (2011). Separating forms of neglect using the Apples test: Validation and functional prediction in chronic and acute stroke. *Neuropsychology*, 25(5), 567–580.

Bowen, A., Hazelton, C., Pollock, A., & Lincoln, N. B. (2013). Cognitive rehabilitation for spatial neglect following stroke. *Cochrane Database of Systematic Reviews*, 7, CD003586. <https://doi.org/10.1002/14651858.CD003586.pub3>

Carter, A. R., & Barrett, A. M. (2023). Recent advances in treatment of spatial neglect: Networks and neuropsychology. *Expert Review of Neurotherapeutics*, 23(7), 587–601.

Cassidy, T. P., Lewis, S., & Gray, C. S. (1998). Recovery from visuospatial neglect in stroke patients. *Journal of Neurology, Neurosurgery & Psychiatry*, 64(4), 555–557.

Cavedoni, S., Cipresso, P., Mancuso, V., Bruni, F., & Pedroli, E. (2022). Virtual reality for the assessment and rehabilitation of neglect: where are we now? A 6-year review update. *Virtual Reality*, 26(4), 1663–1704.

Cazzoli, D., Kaufmann, B. C., Rühe, H., Geiser, N., & Nyffeler, T. (2025). Incidence of visuospatial neglect in acute stroke: Assessment and stroke characteristics in an unselected 1-year cohort. *Stroke*, 56(7), 1693–1703.

Chechlacz, M., Rotshtein, P., Bickerton, W. L., Hansen, P. C., Deb, S., & Humphreys, G. W. (2010). Separating neural correlates of allocentric and egocentric neglect: Distinct cortical sites and common white matter disconnections. *Cognitive Neuropsychology*, 27(3), 277–303.

Chechlacz, M., Rotshtein, P., Roberts, K. L., Bickerton, W.-L., Lau, J. K. L., Humphreys, G. W., & Baron, J.-C. (2012). The prognosis of allocentric and egocentric neglect: Evidence from clinical scans. *PLOS ONE*, 7(11), e47821.

Cocchini, G., & Beschin, N. (2020). The fluff test: Improved scoring system to account for different degrees of contralesional and ipsilesional personal neglect in brain damaged patients. *Neuropsychological Rehabilitation*, 32(1), 1–15.

Collen, F. M., Wade, D. T., Robb, G. F., & Bradshaw, C. M. (1991). The rivermead mobility index: A further development of the rivermead motor assessment. *International Disability Studies*, 13(2), 50–54.

Demeurisse, G., Demol, O., & Robaye, E. (1980). Motor evaluation in vascular hemiplegia. *European Neurology*, 19(6), 382–389.

Demeyere, N., & Gillebert, C. R. (2019). Ego- and allocentric visuospatial neglect: Dissociations, prevalence, and laterality in acute stroke. *Neuropsychology*, 33(4), 490–498.

Demeyere, N., Riddoch, M. J., Slavkova, E. D., Bickerton, W. L., & Humphreys, G. W. (2015). The Oxford Cognitive Screen (OCS): Validation of a stroke-specific short cognitive screening tool. *Psychological Assessment*, 27(3), 883–894.

Demeyere, N., Riddoch, M. J., Slavkova, E. D., Jones, K., Reckless, I., Mathieson, P., & Humphreys, G. W. (2016). Domain-specific versus generalized cognitive screening in acute stroke. *Journal of Neurology*, 263(2), 306–315.

Duncan, P. W., Goldstein, L. B., Horner, R. D., Landsman, P. B., Samsa, G. P., & Matchar, D. B. (1994). Similar motor recovery of upper and lower extremities after stroke. *Stroke*, 25(6), 1181–1188.

Embrechts, E., De Boi, I., Schatteman, Q., Nijboer, T. C. W., Truijen, S., & Saeys, W. (2025). Use of immersive virtual reality to explore visual search behaviour in individuals with visuospatial neglect after stroke. *Neuropsychological Rehabilitation*, 1–28. <https://doi.org/10.1080/09602011.2025.2511193>

Esposito, E., Shekhtman, G., & Chen, P. (2021). Prevalence of spatial neglect post-stroke: A systematic review. *Annals of Physical and Rehabilitation Medicine*, 64(5), 101459.

Feehey, D. M., & Baron, J. C. (1986). Diaschisis. *Stroke*, 17(5), 817–830.

Fellrath, J., Blanche-Durbec, V., Schnider, A., Jacquemoud, A. S., & Ptak, R. (2012). Visual search in spatial neglect studied with a preview paradigm. *Frontiers in Human Neuroscience*, 6, 93.

Guilbert, A. (2023). Clinical assessment of unilateral spatial neglect dissociations and heterogeneities: A narrative synthesis. *Neuropsychology*, 37(4), 450–462.

Heilman, K. M., & Valenstein, E. (1979). Mechanisms underlying hemispatial neglect. *Annals of Neurology*, 5(2), 166–170.

Jehkonen, M., Ahonen, J. P., Dastidar, P., Koivisto, A. M., Laippala, P., Vilkki, J., & Molnár, G. (2000). Visual neglect as a predictor of functional outcome one year after stroke. *Acta Neurologica Scandinavica*, 101(3), 195–201.

Jehkonen, M., Laihosalo, M., Koivisto, A. M., Dastidar, P., & Ahonen, J. P. (2007). Fluctuation in spontaneous recovery of left visual neglect: A 1-year follow-up. *European Neurology*, 58(4), 210–214.

Kwakkel, G., Kollen, B., & Twisk, J. (2006). Impact of time on improvement of outcome after stroke. *Stroke*, 37(9), 2348–2353.

Lazar, R. M., Minzer, B., Antoniello, D., Festa, J. R., Krakauer, J. W., & Marshall, R. S. (2010). Improvement in aphasia scores after stroke is well predicted by initial severity. *Stroke*, 41(7), 1485–1488.

Levine, D. N., Warach, J. D., Benowitz, L., & Calvano, R. (1986). Left spatial neglect: Effects of lesion size and premorbid brain atrophy on severity and recovery following right cerebral infarction. *Neurology*, 36(3), 362–366.

Li, C., Baek, J., Sanchez, B. N., Morgenstern, L. B., & Lisabeth, L. D. (2018). Temporal trends in age at ischemic stroke onset by ethnicity. *Annals of Epidemiology*, 28(10), 686–690.e682.

Marchi, N. A., Ptak, R., Di Pietro, M., Schnider, A., & Guggisberg, A. G. (2017). Principles of proportional recovery after stroke generalize to neglect and aphasia. *European Journal of Neurology*, 24(8), 1084–1087.

Menon-Nair, A., Korner-Bitensky, N., Wood-Dauphinee, S., & Robertson, E. (2006). Assessment of unilateral spatial neglect post stroke in Canadian acute care hospitals: Are we neglecting neglect? *Clinical Rehabilitation*, 20(7), 623–634.

Moore, M. J., Gillebert, C. R., & Demeyere, N. (2021). Right and left neglect are not anatomically homologous: A voxel-lesion symptom mapping study. *Neuropsychologia*, 162, 108024.

Moore, M. J., Vancleef, K., Riddoch, M. J., Gillebert, C. R., & Demeyere, N. (2021). Recovery of visuospatial neglect subtypes and relationship to functional outcome six months after stroke. *Neurorehabilitation and Neural Repair*, 35(9), 823–835.

Nijboer, T. C., Kollen, B. J., & Kwakkel, G. (2013). Time course of visuospatial neglect early after stroke: A longitudinal cohort study. *Cortex*, 49(8), 2021–2027.

Overman, M. J., Binns, E., Milosevich, E. T., & Demeyere, N. (2024). Recovery of visuospatial neglect with standard treatment: A systematic review and meta-analysis. *Stroke*, 55(9), 2325–2339.

Portney, L. G. (2020). Foundations of clinical research: Applications to evidence-based practice, 4e. F. A. Davis Company. [fadavispt.mhmedical.com/content.aspx?aid=1171753091](http://fadavispt.mhmedical.com/content.aspx?aid=1171753091)

Retho, E., Tasseng, Y., Consigny, M., Le Bourhis, L., Leblanc, A., Jourdain, A., Merrien, F. M., Rouhart, F., Viakhireva-Dovganyuk, I., Goas, P., Lavenant, C., Bruguet, M., & Timsit, S. (2023). Increased incidence of ischemic stroke in young: A population-based stroke registry study from 2008 to 2018. *Revue Neurologique*, 180(3), 182–194.

Rorden, C., Berger, Fruhmann M., Karnath, H. O. (2006). Disturbed line bisection is associated with posterior brain lesions. *Brain Research*, 1080(1), 17–25. <https://doi.org/10.1016/j.brainres.2004.10.071>. Epub 2006 March 7. PMID: 16519881.

Rorden, C., & Karnath, H. O. (2010). A simple measure of neglect severity. *Neuropsychologia*, 48(9), 2758–2763.

Samuelsson, H., Jensen, C., Ekholm, S., Naver, H., & Blomstrand, C. (1997). Anatomical and neurological correlates of acute and chronic visuospatial neglect following right hemisphere stroke. *Cortex*, 33(2), 271–285.

Schröder, J., Saeys, W., Yperzeele, L., Kwakkel, G., & Truijen, S. (2022). Time course and mechanisms underlying standing balance recovery early after stroke: Design of a prospective cohort study with repeated measurements. *Frontiers in Neurology*, 13, 781416.

Stone, S. P., Patel, P., Greenwood, R. J., & Halligan, P. W. (1992). Measuring visual neglect in acute stroke and predicting its recovery: The visual neglect recovery index. *Journal of Neurology, Neurosurgery & Psychiatry*, 55(6), 431–436.

Takamura, Y., Fujii, S., Ohmatsu, S., Morioka, S., & Kawashima, N. (2021). Pathological structure of visuospatial neglect: A comprehensive multivariate analysis of spatial and non-spatial aspects. *iScience*, 24(4), 102316.

Vaes, N., Lafosse, C., Nys, G., Schevernels, H., Dereymaeker, L., Oostra, K., Hemelsoet, D., & Vingerhoets, G. (2015). Capturing peripersonal spatial neglect: An electronic method to quantify visuospatial processes. *Behavior Research Methods*, 47(1), 27–44.

Van der Stigchel, S., & Nijboer, T. C. (2010). The imbalance of oculomotor capture in unilateral visual neglect. *Consciousness and Cognition*, 19(1), 186–197.

von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Götzsche, P. C., & Vandenbroucke, J. P. (2008). The strengthening the reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Journal of Clinical Epidemiology*, 61(4), 344–349.

Williams, L. J., Kernot, J., Hillier, S. L., & Loetscher, T. (2021). Spatial Neglect Subtypes, Definitions and Assessment Tools: A Scoping Review. *Frontiers in Neurology*, 12, 742365.

World Health Organisation (WHO). (2001, May 21). International Classification of Functioning, Disability and Health: ICF. <https://www.who.int/classifications/international-classification-of-functioning-disability-and-health>

Yoo, J. W., Hong, B. Y., Jo, L., Kim, J. S., Park, J. G., Shin, B. K., & Lim, S. H. (2020). Effects of age on long-term functional recovery in patients with stroke. *Medicina (Kaunas)*, 56(9), 451.