



Letter

Finite-resolution limits on quantum uniqueness

Seramika Ariwahjoedi^a, Sameer Ahmad Mir^{b,c,*}, Mir Faizal^{b,d,e,f}^a Research Center for Quantum Physics, National Research and Innovation Agency (BRIN), South Tangerang, Banten, 15314, Indonesia^b Canadian Quantum Research Center, 204-3002 32 Ave, Vernon, BC V1T 2L7, Canada^c Department of Computer Sciences, Asian School of Business, Noida, Uttar Pradesh, 201303, India^d Irving K. Barber School of Arts and Sciences, University of British Columbia Okanagan, Kelowna, British Columbia, V1V 1V7, Canada^e Department of Mathematical Sciences, Durham University, Upper Mountjoy, Stockton Road, Durham, DH1 3LE, UK^f Faculty of Sciences, Hasselt University Agoralaan Gebouw D, Diepenbeek, 3590, Belgium

ARTICLE INFO

Editor: Robert Mann

Keywords:

Weyl algebra (Weyl relations)
 Stone-von Neumann theorem
 Strong continuity / regular representations
 Polymer representation (polymer quantization)
 Canonical quantization and uniqueness
 Finite-resolution operational protocols

ABSTRACT

The Stone-von Neumann theorem makes canonical quantization unique only after one imposes strong continuity of the Weyl translation groups. We show that this continuity assumption cannot be certified by any finite-resolution interrogation of the Weyl relations. A finite protocol probes only finitely many phase-space displacements and bounded expectation values, thereby determining a protocol algebra C_S generated by a finitely generated subgroup of Weyl translations, but not the regularity of an extension to the full Weyl algebra. Using the polymer representation as a canonical non-regular representation, we prove that for every regular Weyl representation, every normal state, and every finite family of bounded protocol observables, a polymer state reproduces the same expectation values to arbitrary prescribed accuracy. Thus finite agreement with Schrödinger quantum mechanics does not operationally certify the regularity hypothesis entering uniqueness. The result does not assert physical equivalence between regular and polymer quantizations, it identifies where additional input, such as Hamiltonian dynamics, energy regularity, semiclassicality, or a continuum limit, is required to distinguish or select representations. In this precise sense, non-regular polymer kinematics are not excluded by finite Weyl data alone, a point of direct relevance to quantum-gravity motivated quantizations.

1. Introduction

Canonical quantization of a single degree of freedom is most naturally formulated in terms of Weyl operators implementing phase-space translations and satisfying the Weyl form of the canonical commutation relations [1]. The standard uniqueness result in this setting is the Stone-von Neumann theorem, which asserts that at all irreducible representations of the Weyl relations are unitarily equivalent provided the translation groups are strongly continuous [2,3]. Strong continuity is the sole analytic hypothesis of the theorem; it permits the use of Stone's theorem to obtain self-adjoint generators and hence the familiar position and momentum operators [4,5]. While mathematically natural, strong continuity is not an operational requirement implied by finite experiments. Laboratory protocols implement translations through finitely many calibrated control settings with finite resolution and finite statistics, so only a finite family of Weyl operators and their finite compositions can be prepared and distinguished, up to bounded tolerances in their parameters. The Stone-von Neumann theorem does not address whether strong

continuity can be certified, or excluded, from such finite operational data.

In this Letter we formulate finite experimental resolution directly at the level of accessible Weyl operators, without modifying the Weyl relations and without privileging any representation a priori. The point is to separate two logically distinct ingredients of the usual uniqueness statement. The Weyl relations fix the algebraic composition law of phase-space translations, whereas regularity is an additional strong-continuity condition imposed on a representation of that algebra. This distinction belongs naturally to the C^* -algebraic formulation of canonical commutation relations and unitary group representations [6–9]. A finite experiment probes only finitely many displacement settings and finitely many expectation values. It therefore determines only a finite operational algebra C_S , not the strong-continuity behavior of the full one-parameter Weyl translation groups.

Our main result shows that this limitation is structural, not merely practical. For any finite protocol algebra C_S , any regular Weyl representation, any normal state, and any finite family of bounded observables

* Corresponding author at: Canadian Quantum Research Center, 204-3002 32 Ave, Vernon, BC V1T 2L7, Canada .

E-mail addresses: ceramya@gmail.com (S. Ariwahjoedi), sameerphst@gmail.com, sameerphst@candqrc.ca (S.A. Mir), mirfaizalmir@gmail.com (M. Faizal).

<https://doi.org/10.1016/j.physletb.2026.140853>

Received 4 March 2026; Received in revised form 16 May 2026; Accepted 11 August 2026

Available online 17 August 2026

0370-2693/© 2026 The Author(s). Published by Elsevier B.V. Funded by SCOAP³. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

in C_S , the same expectation values can be reproduced to arbitrary prescribed accuracy. This reproduction is achieved by a suitable state in the polymer representation. The polymer representation is non-regular. The approximation mechanism uses only standard C^* -algebraic tools. The protocol algebra is a twisted group C^* -algebra of the finitely generated abelian group Γ_S . Amenability identifies the relevant full and reduced twisted group C^* -algebras. Fell approximation then allows states to be approximated on finite sets by vector states in a faithful representation [9–11]. The polymer representation is used here only as a canonical non-regular realization of the Weyl relations. The representation-theoretic approximation argument relies on the standard algebraic facts described above [7,9,11]. This does not contradict the Stone-von Neumann theorem. That theorem remains exactly valid within the class of irreducible regular representations. Our result identifies instead the operational status of its continuity hypothesis. Regularity is an additional analytic input. It is not something certified by finite access to Weyl translations alone.

2. Experimental relevance and finite operational access

Let $\xi = (\lambda, \mu) \in \mathbb{R}^2$ denote a phase-space displacement label, and let $W(\xi)$ denote the corresponding Weyl displacement unitary. In experiments, such phase-space displacements are implemented by calibrated control pulses with finite resolution and are inferred from finitely many measurement shots. The accessible information is therefore not the full continuum of Weyl labels $\xi \in \mathbb{R}^2$, but a finite data table consisting of discrete control settings, uncertain displacement parameters, and finitely sampled expectation values. Thus a finite protocol can constrain only finitely many bounded operational quantities, each within finite statistical and calibration tolerances.

In continuous-variable systems this restriction is explicit. Displacements, realized as Gaussian primitives, are characterized indirectly through rotated-quadrature measurements, homodyne sampling, and tomographic reconstructions employing finitely many local-oscillator phases and finitely many samples. These procedures constrain only a finite family of accessible displacement operators, or coarse-grained approximants to them, together with finite moment or correlation data [12–15]. More generally, validation of implemented dynamics relies on process tomography and self-consistent variants, yielding only finite input-output probabilities and a finite reconstructed model [16–18], while randomized benchmarking provides finite-data estimates of average error rates without pointwise continuity information [19,20].

This operational finiteness motivates our formulation. A protocol specifies a finite set of calibrated displacement labels

$$S = \{\xi_1, \dots, \xi_m\} \subset \mathbb{R}^2, \quad (1)$$

and finitely many expectation values estimated within finite tolerance. The subgroup algebraically generated by the accessible labels is $\Gamma_S = \text{span}_{\mathbb{Z}}(S) \subset \mathbb{R}^2$. The physically available control operations generate the unital $*$ -algebra spanned by finite products of $\{W(\xi_j)\}_{j=1}^m$. Equivalently, the relevant Weyl labels lie in Γ_S . The corresponding norm-closed protocol algebra is

$$C_S = C^*(\{W(\gamma) : \gamma \in \Gamma_S\}) \subset \mathcal{W}, \quad (2)$$

where $\mathcal{W}$ denotes the full Weyl C^* -algebra. Thus C_S is the part of the Weyl algebra generated by the control operations available to the protocol. The distinction is important: Γ_S , and hence C_S , may be infinite even though S is finite. However, any actual finite data set tests only finitely many bounded observables in C_S , and only up to finite tolerance. The notation $\xi_n \rightarrow 0$ refers to convergence in the Euclidean topology of the full parameter space $\mathbb{R}^2$. It should not be confused with the experimental act of enlarging the finite set S . Strong continuity is a representation-theoretic condition: for a representation π , it requires

$$\begin{aligned} \lim_{\lambda \rightarrow 0} \|\pi(W(\lambda, 0))\psi - \psi\| &= 0, \\ \lim_{\mu \rightarrow 0} \|\pi(W(0, \mu))\psi - \psi\| &= 0 \end{aligned} \quad (3)$$

for every vector ψ in the representation Hilbert space. Equivalently, since the Weyl parameter space is metrizable, regularity requires the same convergence along every sequence $\xi_n \rightarrow 0$ in the Euclidean topology of $\mathbb{R}^2$. This is an infinite limiting condition at the identity of the displacement group. A finite protocol supplies no such limiting test. It samples only the displacement labels generated by finitely many chosen settings and compares only finitely many bounded expectation values.

Allowing adaptive refinement does not alter the logical point. If a later experiment uses additional settings, S is replaced by a larger finite set and C_S by a larger finitely generated protocol algebra. Each finite stage may provide a better operational approximation to a continuum model, but no finite stage certifies strong continuity of the full one-parameter Weyl translation groups

$$\lambda \mapsto \pi(W(\lambda, 0)), \quad \mu \mapsto \pi(W(0, \mu)). \quad (4)$$

Accordingly, C_S encodes the discrete access structure of the protocol, whereas regularity concerns an extension of a representation from C_S to the full Weyl algebra together with the Euclidean topology of its label space. Regularity may therefore be a natural and physically useful modeling assumption, but it is not forced by finite operational access to Weyl translations. This does not mean that regular and non-regular representations are physically identical in all respects. It means only that finite Weyl-displacement data, by themselves, do not decide the strong-continuity hypothesis required by the Stone-von Neumann uniqueness theorem. Additional physical structure, such as Hamiltonian dynamics, energy regularity, semiclassicality, or a continuum limit, may distinguish or select representations. Such structure, however, goes beyond the finite protocol algebra C_S . This finite-resolution limitation is summarized schematically in Fig. 1. Finite control precision selects only sampled displacement labels with error bars, and finite statistics determine only finitely many noisy expectation values. A regular representation and a non-regular representation may agree on all such sampled data within the prescribed tolerance, even though they differ in their behavior under the infinite limiting test required for strong continuity.

3. Weyl algebra and regularity

Let $\xi = (\lambda, \mu)$ and $\eta = (\lambda', \mu')$ be elements of $\mathbb{R}^2$. We use the standard symplectic form

$$\sigma(\xi, \eta) = \lambda\mu' - \mu\lambda'. \quad (5)$$

The Weyl C^* -algebra $\mathcal{W}$ is the universal C^* -algebra generated by unitaries $\{W(\xi)\}_{\xi \in \mathbb{R}^2}$ satisfying

$$W(\xi)W(\eta) = e^{-\frac{i}{2}\sigma(\xi, \eta)}W(\xi + \eta), \quad W(\xi)^* = W(-\xi). \quad (6)$$

These relations are the Weyl form of the canonical commutation relations [1]. A representation of $\mathcal{W}$ is a nondegenerate $*$ -homomorphism $\pi : \mathcal{W} \rightarrow B(\mathcal{H})$, where $B(\mathcal{H})$ denotes the bounded operators on the representation Hilbert space $\mathcal{H}$. Such a representation is called regular if the one-parameter unitary groups as in Eq. (4), are strongly continuous. Equivalently, regularity is strong continuity at the identity in the sense stated in Eq. (3): for every $\psi \in \mathcal{H}$, the vectors $\pi(W(\lambda, 0))\psi$ and $\pi(W(0, \mu))\psi$ converge in Hilbert-space norm to ψ as $\lambda \rightarrow 0$ and $\mu \rightarrow 0$, respectively. Thus regularity is not an algebraic consequence of the Weyl relations alone. It is an additional continuity requirement on the representation. Under this requirement, Stone's theorem gives self-adjoint generators Q and P for the two one-parameter unitary groups. The Stone-von Neumann theorem then states that every irreducible regular representation of the Weyl relations for one canonical degree of freedom is unitarily equivalent to the Schrödinger representation [2–5].

Therefore the theorem proves uniqueness inside the regular sector. It does not prove that finite operational data, restricted to a protocol algebra C_S , force a representation of the full Weyl algebra $\mathcal{W}$ to be regular.




Fig. 1. Finite-resolution access to Weyl translations (schematic). (a) Finite control precision restricts experimental access to a discrete family of displacement parameters $\{\xi_k\} \subset \mathbb{R}^2$ with $\xi = (q, p)$, each realized only within a tolerance $\delta\xi$ (disks). (b) Finite statistics determine only noisy estimates of an operational quantity, e.g. $f(\xi) = \langle \psi | W(\xi) | \psi \rangle$, at sampled settings; a regular (strongly continuous) and a non-regular realization can agree on all sampled points within error bars, so strong continuity is not operationally decidable from finite data.

4. Operational resolution

With the notation above, a finite protocol specified by $S = \{\xi_1, \dots, \xi_m\}$ determines the algebraically generated subgroup $\Gamma_S = \text{span}_{\mathbb{Z}}(S)$, and hence, through Eq. (2), the associated protocol algebra C_S . Although C_S may contain infinitely many algebraic and norm-limit elements, any actual finite experimental data set tests only finitely many bounded observables in C_S , and only within finite tolerance. Thus the operational comparison between two representations is necessarily a finite-set comparison of expectation values.

Let

$$\pi_1 : \mathcal{W} \rightarrow B(H_1), \quad \pi_2 : \mathcal{W} \rightarrow B(H_2) \quad (7)$$

be two representations. We say that π_2 is (S, η) -operationally compatible with π_1 if, for every density operator ρ_1 on H_1 and every finite collection of test observables $A_1, \dots, A_n \in C_S$ with $\|A_j\| \leq 1$, there exists a density

operator ρ_2 on H_2 such that

$$\left| \text{Tr}(\rho_1 \pi_1(A_j)) - \text{Tr}(\rho_2 \pi_2(A_j)) \right| \leq \eta, \quad j = 1, \dots, n. \quad (8)$$

This is intentionally a one-sided notion. It formalizes the statement that π_2 cannot be ruled out by any finite (C_S, η) data set compatible with π_1 . In the application below, π_1 will be a regular Weyl representation and π_2 will be the polymer representation.

The notation $\xi_n \rightarrow 0$ always refers to convergence in the Euclidean topology of the full parameter space $\mathbb{R}^2$. This is precisely the topology used in the definition of regularity. A finite protocol does not implement an arbitrary sequence approaching the identity in this topology. It accesses only the subgroup Γ_S generated by finitely many chosen settings, and its finite data constrain only finitely many expectation values on C_S . Hence C_S records the algebraic information available to the protocol, whereas regularity concerns an extension of a representation from C_S to the full Weyl algebra $\mathcal{W}$ together with the Euclidean topology of its label space. Therefore the Stone-von Neumann theorem yields uniqueness only after the additional continuity assumption of regularity has been imposed. Finite operational data on C_S may be consistent with a regular representation, but they do not by themselves certify that every compatible extension to the full Weyl algebra is regular.

5. Operational continuity and the scope of uniqueness

For a fixed protocol S , the accessible parameter set is Γ_S : only finitely many generators and their finite products are queried, and no operation internal to the protocol implements a limiting procedure $\xi_n \rightarrow 0$ in $\mathbb{R}^2$. Accordingly, the protocol induces no topology on Γ_S beyond the discrete one, and in particular does not operationally test strong continuity of $\lambda \mapsto \pi(W(\lambda, 0))$ or $\mu \mapsto \pi(W(0, \mu))$ at the identity. Regularity is therefore not a property of the abstract operational algebra C_S , but of an extension of a representation of C_S to a representation of $\mathcal{W}$ together with the Euclidean topology on $\mathbb{R}^2$. The Stone-von Neumann theorem thus yields uniqueness only within the class of irreducible regular representations, while leaving regularity itself undetermined by operational data restricted to C_S .

6. Polymer representation

Let $\mathbb{R}_d$ denote the real line equipped with the discrete topology, and define

$$\mathcal{H}_{\text{pol}} = \ell^2(\mathbb{R}_d). \quad (9)$$

Here $\mathbb{R}_d$ has the same underlying set as $\mathbb{R}$, but it is regarded as a discrete measure space with counting measure. Thus $\mathcal{H}_{\text{pol}}$ is not the Schrödinger Hilbert space $L^2(\mathbb{R})$. Its vectors are square-summable functions on the set $\mathbb{R}$ with respect to counting measure. Consequently the vectors $\{|x\rangle\}_{x \in \mathbb{R}}$ are genuine normalizable basis vectors of $\mathcal{H}_{\text{pol}}$, satisfying $\langle x | x' \rangle = \delta_{x, x'}$, where $\delta_{x, x'}$ is the Kronecker delta. This is different from the usual generalized position eigenvectors of $L^2(\mathbb{R})$, which are distributions and only formally satisfy Dirac-delta normalization. This distinction is essential: in $\mathcal{H}_{\text{pol}}$, two different basis vectors $|x\rangle$ and $|x'\rangle$ are orthogonal whenever $x \neq x'$.

The polymer action of the Weyl generators is given by

$$\pi_{\text{pol}}(W(\lambda, \mu))|x\rangle = e^{i\mu(x+\lambda/2)}|x+\lambda\rangle. \quad (10)$$

Each $\pi_{\text{pol}}(W(\lambda, \mu))$ is unitary, since it shifts the orthonormal basis by $x \mapsto x + \lambda$ and multiplies each basis vector by a phase. A direct evaluation on basis vectors gives

$$\pi_{\text{pol}}(W(\xi))\pi_{\text{pol}}(W(\eta)) = e^{-\frac{i}{2}\sigma(\xi, \eta)}\pi_{\text{pol}}(W(\xi + \eta)). \quad (11)$$

Thus the polymer construction is a genuine representation of the Weyl relations. It is nevertheless non-regular [9,11]. Indeed, for any $x \in \mathbb{R}$ and any nonzero λ , the vectors $|x + \lambda\rangle$ and $|x\rangle$ are orthogonal in $\mathcal{H}_{\text{pol}}$. Hence

$$\left\| \pi_{\text{pol}}(W(\lambda, 0))|x\rangle - |x\rangle \right\|^2 = \left\| |x + \lambda\rangle - |x\rangle \right\|^2 = 2. \quad (12)$$

Therefore $\lambda \mapsto \pi_{\text{pol}}(W(\lambda, 0))$ is discontinuous at $\lambda = 0$ in the strong operator topology. The failure of continuity is not due to a failure of the Weyl relations; rather, it follows from the discrete Hilbert-space topology of the polymer representation.

For the fixed protocol S , the shift subgroup is defined as

$$\Lambda_S = \{\lambda \in \mathbb{R} : (\lambda, \mu) \in \Gamma_S \text{ for some } \mu\}. \quad (13)$$

For each $x_0 \in \mathbb{R}$, we define

$$\mathcal{H}_{x_0, S} = \overline{\text{span}}\{|x_0 + \lambda\rangle : \lambda \in \Lambda_S\} \subset \mathcal{H}_{\text{pol}}. \quad (14)$$

Each $\mathcal{H}_{x_0, S}$ is invariant under $\pi_{\text{pol}}(C_S)$, because every element of C_S shifts a basis vector only by an element of Λ_S . More precisely, the Weyl generators with labels in Γ_S shift basis vectors only by elements of Λ_S , and the invariance then extends from finite algebraic combinations to the norm-closed algebra C_S because $\mathcal{H}_{x_0, S}$ is closed. Explicitly, for $(\lambda, \mu) \in \Gamma_S$,

$$\begin{aligned} \pi_{\text{pol}}(W(\lambda, \mu))|x_0 + \lambda'\rangle &= e^{i\mu(x_0 + \lambda' + \lambda/2)}|x_0 + \lambda' + \lambda\rangle, \\ \lambda' &\in \Lambda_S. \end{aligned} \quad (15)$$

Since $\lambda \in \Lambda_S$, the vector on the right again belongs to $\mathcal{H}_{x_0, S}$. Thus $\mathcal{H}_{\text{pol}}$ decomposes as an orthogonal direct sum of invariant protocol sectors indexed by the cosets of Λ_S in $\mathbb{R}$. The proof below uses only this sector decomposition and the standard finite-set form of Fell approximation for faithful representations of C^* -algebras.

7. Main theorem

Fix a finite protocol S , let $\Gamma_S = \text{span}_{\mathbb{Z}}(S)$, $C_S = C^*(\{W(\gamma) : \gamma \in \Gamma_S\})$, and let $\eta > 0$. Let $\pi : \mathcal{W} \rightarrow B(\mathcal{H})$ be a regular Weyl representation, and let ρ be a density operator on $\mathcal{H}$. For every finite family $A_1, \dots, A_n \in C_S$ with $\|A_j\| \leq 1$, there exists a density operator ρ_{pol} on $\mathcal{H}_{\text{pol}}$, supported on a finite direct sum of invariant polymer sectors, such that

$$\left| \text{Tr}(\rho \pi(A_j)) - \text{Tr}(\rho_{\text{pol}} \pi_{\text{pol}}(A_j)) \right| \leq \eta, \quad j = 1, \dots, n. \quad (16)$$

Thus no finite collection of bounded observables in the protocol algebra C_S can operationally exclude the polymer representation on the basis of expectation values alone. The regularity of π is not used as an approximation assumption; it specifies the usual Stone-von Neumann sector against which the non-regular polymer representation is being compared.

8. Proof

The group Γ_S is a finitely generated abelian group, considered as a discrete group. The restriction of the Weyl multiplier to Γ_S is $c(\gamma, \gamma') = e^{-\frac{i}{2}\sigma(\gamma, \gamma')}$. By the universal property of twisted group C^* -algebras, the protocol algebra C_S is canonically isomorphic to the twisted group C^* -algebra $C_S \simeq C^*(\Gamma_S, c)$, with product $u_\gamma u_{\gamma'} = c(\gamma, \gamma')u_{\gamma+\gamma'}$. Since Γ_S is abelian, it is amenable. Hence the full and reduced twisted group C^* -algebras coincide [10,11].

We use the following standard finite-set form of Fell's approximation theorem. If σ is a faithful representation of a C^* -algebra A , then every state ω on A can be approximated, on any prescribed finite set of elements of A and to any prescribed accuracy, by finite convex combinations of vector states belonging to σ [7,8]. Equivalently, for every finite set $F \subset A$ and every $\varepsilon > 0$, there exist unit vectors $\psi_1, \dots, \psi_r$ in the representation space of σ and weights $t_k \geq 0$, with $\sum_{k=1}^r t_k = 1$, such that

$$\left| \omega(A) - \sum_{k=1}^r t_k \langle \psi_k, \sigma(A) \psi_k \rangle \right| < \varepsilon, \quad A \in F. \quad (17)$$

In physical terms, this means that a faithful representation contains enough vector states to reproduce arbitrary states on any finite list

of tested observables. The Weyl C^* -algebra associated with a non-degenerate finite-dimensional symplectic vector space is simple [5,6,11]. Therefore the nonzero polymer representation $\pi_{\text{pol}} : \mathcal{W} \rightarrow B(\mathcal{H}_{\text{pol}})$ is faithful, and its restriction to the subalgebra C_S is faithful. Now define the state on C_S $\omega(A) = \text{Tr}(\rho \pi(A))$, $A \in C_S$. Applying Fell's approximation theorem to the faithful representation $\pi_{\text{pol}}|_{C_S}$, with $F = \{A_1, \dots, A_n\}$, gives unit vectors $\psi_1, \dots, \psi_r \in \mathcal{H}_{\text{pol}}$ and weights $t_k \geq 0$, $\sum_{k=1}^r t_k = 1$, such that

$$\left| \omega(A_j) - \sum_{k=1}^r t_k \langle \psi_k, \pi_{\text{pol}}(A_j) \psi_k \rangle \right| < \frac{\eta}{2}, \quad j = 1, \dots, n. \quad (18)$$

It remains to ensure that the approximating polymer state is supported on finitely many invariant protocol sectors. Finitely supported vectors are dense in $\ell^2(\mathbb{R}_d)$. Therefore each ψ_k may be replaced by a sufficiently close finitely supported unit vector. This replacement changes all vector-state expectation values of the contractions A_j by a controlled amount: if $\|\psi\| = \|\phi\| = 1$ and $\|A_j\| \leq 1$, then

$$|\langle \psi, A_j \psi \rangle - \langle \phi, A_j \phi \rangle| \leq 2\|\psi - \phi\|. \quad (19)$$

Choosing the replacements close enough, the total additional error can be made smaller than $\eta/2$ for all $j = 1, \dots, n$.

The union of the finite supports of these approximating vectors is contained in finitely many cosets of the shift subgroup Λ_S defined in Eq. (13). Let $x_1, \dots, x_N$ be representatives of these cosets. Using the invariant sectors $\mathcal{H}_{x_a, S}$ defined in Eq. (14), the approximating vectors therefore lie in the finite direct sum

$$\mathcal{H}_S^{\text{pol}} = \bigoplus_{a=1}^N \mathcal{H}_{x_a, S}. \quad (20)$$

Each $\mathcal{H}_{x_a, S}$ is invariant under $\pi_{\text{pol}}(C_S)$. Indeed, the Weyl generators with labels in Γ_S shift basis vectors only by elements of Λ_S , and this invariance extends from finite algebraic combinations to the norm-closed algebra C_S because $\mathcal{H}_{x_a, S}$ is closed.

Consequently, $\mathcal{H}_S^{\text{pol}}$ is an invariant finite direct sum of protocol sectors. Finally set

$$\rho_{\text{pol}} = \sum_{k=1}^r t_k |\psi_k\rangle\langle\psi_k|. \quad (21)$$

This is a density operator supported on $\mathcal{H}_S^{\text{pol}}$. Combining the Fell approximation error with the finite-support replacement error gives Eq. (16). This proves the theorem.

9. Finite tomography and operational consequences

The theorem has a direct interpretation for continuous-variable tomography. A finite homodyne or displacement-tomography protocol uses finitely many local-oscillator phases, finitely many displacement settings, finitely many coarse-graining choices, and finitely many samples. The resulting data form a finite list of experimentally estimated bounded statistics. When these statistics are expressed, or approximated within the experimental tolerance, by a finite family $A_1, \dots, A_n \in C_S$, the theorem implies that their expectation values in a regular representation can be reproduced by a suitable polymer state within the same prescribed tolerance. Such a protocol may test many features of a state, a gate, or measurement model, but it cannot certify the strong-continuity condition required by the Stone-von Neumann theorem.

This also addresses the apparent operational concern raised by the existence of the polymer approximation. The result does not say that regular and polymer quantizations are physically identical in every respect. It proves the more precise statement that finite Weyl-displacement data alone cannot decide regularity. If additional physical structure is imposed, such as a strongly continuous time-evolution group, a Hamiltonian with appropriate domain properties, energy regularity, semiclassicality conditions, or a continuum limit, then regular and polymer representations may become physically distinguishable. Such distinctions

arise from extra dynamical or analytic input, not from finite access to the Weyl relations alone.

The relation with polymer quantization and loop quantum gravity is structural rather than merely analogical. In the standard background-independent loop representation, polymer-type kinematics arise because the elementary variables are not the infinitesimal canonical fields themselves, but their exponentiated and smeared counterparts: holonomies of the Ashtekar-Barbero connection and fluxes of the densitized triad. The Ashtekar-Lewandowski representation therefore does not realize the connection as a weakly continuous operator-valued field in the same sense in which the Schrödinger representation realizes canonical position and momentum through strongly continuous Weyl translations. Instead, it realizes finite holonomy transformations and flux operators on a non-regular kinematical Hilbert space, and the usual Stone-von Neumann uniqueness mechanism is replaced by uniqueness theorems adapted to diffeomorphism-covariant holonomy-flux algebras [21–23]. The polymer representation studied here is the finite-dimensional prototype of this same mechanism: finite exponentiated translations are represented exactly, while the corresponding infinitesimal generators need not exist because weak or strong continuity fails [24–26].

This observation motivates the precise question addressed in the present work. If polymer kinematics are not an arbitrary deformation but the natural kinematical language of loop quantized geometry, then one may ask whether analogous non-regular representations are operationally excluded for matter degrees of freedom. Standard quantum mechanics assumes the regular Schrödinger representation, in which Weyl translations are strongly continuous and Stone's theorem reconstructs the usual self-adjoint canonical operators. Polymer quantization keeps the Weyl relations but drops this continuity requirement. The issue is therefore not whether the algebraic commutation relations are modified, but whether finite experimental access to Weyl observables can force the regularity assumption that selects the Schrödinger sector. Our theorem answers this question in a restricted but sharp sense: for every regular Weyl representation, every normal state, and every finite family of bounded observables in a finitely generated protocol algebra, there exists a polymer state reproducing the same expectation values to arbitrary prescribed accuracy. Thus finite kinematical Weyl data alone do not operationally exclude polymer quantization for matter fields.

The result should not be read as claiming that matter polymer quantization is already physically established, or that polymer and Schrödinger dynamics are phenomenologically equivalent. It says something more precise. The distinction between regular and polymer quantization cannot be decided by finite kinematical tests of the Weyl relations alone. If matter fields were fundamentally polymer quantized, ordinary Schrödinger quantum mechanics would have to appear as an effective description selected by additional structure: Hamiltonian dynamics, semiclassical states, energy regularity, coarse graining, continuum reconstruction, or a low-energy limit. Conversely, possible deviations from standard quantum mechanics would not be expected merely from the algebraic Weyl relations, which both representations satisfy, but from the non-regular realization of dynamics and limits. In this sense, the present work clarifies the operational relation between standard quantum mechanics and the polymer kinematics underlying loop quantum gravity: regularity is not an experimentally forced consequence of finite Weyl data, but an additional analytic and physical assumption whose justification must come from dynamics and semiclassical recovery.

10. Conclusion

We have shown that the strong-continuity hypothesis entering the Stone-von Neumann uniqueness theorem cannot be certified by any fixed finite-resolution protocol probing Weyl translations. A finite protocol accesses only the norm-closed algebra C_S generated by a finitely generated subgroup of displacement labels. On this protocol algebra,

expectation values produced by a regular Weyl representation can be reproduced, to arbitrary prescribed accuracy and for any finite family of bounded observables, by suitable polymer states. The polymer representation therefore supplies a concrete non-regular extension satisfying the Weyl relations exactly while failing the continuity assumption required for Stone-von Neumann uniqueness. This conclusion leaves the Stone-von Neumann theorem intact. The theorem remains the correct uniqueness statement for irreducible regular representations. What our analysis clarifies is the operational status of regularity, it is an additional analytic and physical assumption, not something enforced by finite Weyl-displacement data. Finite agreement with Schrödinger quantum mechanics therefore does not, by itself, exclude non-regular representations. To exclude or select among such representations one must impose further structure, for example Hamiltonian dynamics, energy regularity, semiclassicality, or a continuum limit. This is precisely the point at which physics beyond the finite Weyl protocol enters. In this restricted but important sense, polymer kinematics in quantum-gravity motivated quantizations are not ruled out by finite Weyl data alone, their physical viability is instead a question of additional dynamical and semiclassical input.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

References

- [1] H. Weyl, *The Theory of Groups and Quantum Mechanics*, Dover, 1950.
- [2] M.H. Stone, *Linear Transformations in Hilbert Space*, Vol. 15 AMS Colloquium Publications, American Mathematical Society, 1932. <https://doi.org/10.1090/coll/015>.
- [3] J. von Neumann, Die Eindeutigkeit der Schrödingerschen Operatoren, *Math. Ann.* 104 (1931) 570–578. <https://doi.org/10.1007/BF01457956>.
- [4] M. Reed, B. Simon, *Methods of Modern Mathematical Physics I: Functional Analysis*, Academic Press, 1980. <https://doi.org/10.1016/B978-0-12-585001-8.X5001-6>.
- [5] J. Dixmier, *C*-Algebras*, North-Holland, 1977. https://archive.org/details/_penalty0calgebras0000dixm.
- [6] O. Bratteli, D.W. Robinson, *Operator Algebras and Quantum Statistical Mechanics I*, Springer, 2nd ed., 1987. <https://doi.org/10.1007/978-3-662-02520-8>.
- [7] J.M.G. Fell, R.S. Doran, *Representations of *-Algebras, Locally Compact Groups, and Banach *-Algebraic Bundles*, Vol. 1–2, Academic Press, 1988.
- [8] R.V. Kadison, J.R. Ringrose, *Fundamentals of the Theory of Operator Algebras*, Vol. I-II, American Mathematical Society, 1997.
- [9] I.E. Segal, Irreducible representations of operator algebras, *Bull. Amer. Math. Soc.* 53 (1947) 73–88. <https://doi.org/10.1090/S0002-9904-1947-08742-5>.
- [10] A. Hulanicki, Means and Følner condition on locally compact groups, *Stud. Math.* 27 (2) (1966) 87–104. <http://eudml.org/doc/217185>.
- [11] M.A. Rieffel, Induced representations of C*-algebras, *Adv. Math.* 13 (1974) 176–257. [https://doi.org/10.1016/0001-8708\(74\)90068-1](https://doi.org/10.1016/0001-8708(74)90068-1).
- [12] K. Vogel, H. Risken, Determination of quasiprobability distributions in terms of probability distributions for the rotated quadrature phase, *Phys. Rev. A* 40 (1989) 2847. <https://doi.org/10.1103/PhysRevA.40.2847>.
- [13] D.T. Smithey, M. Beck, M.G. Raymer, A. Faridani, Measurement of the Wigner distribution and the density matrix of a light mode using optical homodyne tomography, *Phys. Rev. Lett.* 70 (1993) 1244. <https://doi.org/10.1103/PhysRevLett.70.1244>.
- [14] A.I. Lvovsky, M.G. Raymer, Continuous-variable optical quantum-state tomography, *Rev. Mod. Phys.* 81 (2009) 299. <https://doi.org/10.1103/RevModPhys.81.299>.
- [15] C. Weedbrook, S. Pirandola, R. García-Patrón, N.J. Cerf, T.C. Ralph, J.H. Shapiro, S. Lloyd, Gaussian quantum information, *Rev. Mod. Phys.* 84 (2012) 621. <https://doi.org/10.1103/RevModPhys.84.621>.
- [16] I.L. Chuang, M.A. Nielsen, Prescription for experimental determination of the dynamics of a quantum black box, *J. Mod. Opt.* 44 (1997) 2455. <https://doi.org/10.1080/09500349708231894>.
- [17] S.T. Merkel, J.M. Gambetta, J.A. Smolin, S. Poletto, A.D. Córcoles, B.R. Johnson, C.A. Ryan, M. Steffen, Self-consistent quantum process tomography, *Phys. Rev. A* 87 (6) (2013) 062119. [arXiv:1211.0322](https://arxiv.org/abs/1211.0322) [quant-ph], <https://doi.org/10.1103/PhysRevA.87.062119>.
- [18] R. Blume-Kohout, J.K. Gamble, E. Nielsen, K. Rudinger, J. Mizrahi, K. Fortier, P. Maunz, Demonstration of qubit operations below a rigorous fault tolerance threshold with gate set tomography, *Nat. Commun.* 8 (1) (2017) 14485. [arXiv:1605.07674](https://arxiv.org/abs/1605.07674) [quant-ph], <https://doi.org/10.1038/ncomms14485>.

- [19] E. Knill, D. Leibfried, R. Reichle, J. Britton, R.B. Blakestad, J.D. Jost, C. Langer, R. Ozeri, S. Seidelin, D.J. Wineland, Randomized benchmarking of quantum gates, *Phys. Rev. A* 77 (1) (2008) 012307. [arXiv:0707.0963](#) [quant-ph], <https://doi.org/10.1103/PhysRevA.77.012307>.
- [20] E. Magesan, J.M. Gambetta, J. Emerson, Scalable and robust randomized benchmarking of quantum processes, *Phys. Rev. Lett.* 106 (18) (2011) 180504. [arXiv:1009.3639](#) [quant-ph], <https://doi.org/10.1103/PhysRevLett.106.180504>.
- [21] C. Rovelli, L. Smolin, Discreteness of area and volume in quantum gravity, *Nucl. Phys. B* 442 (1995) 593–619. [arXiv:gr-qc/9411005](#), [https://doi.org/10.1016/0550-3213\(95\)00150-Q](https://doi.org/10.1016/0550-3213(95)00150-Q).
- [22] A. Ashtekar, J. Lewandowski, Background independent quantum gravity: a status report, *Class. Quant. Grav.* 21 (2004) R53–R152. [arXiv:gr-qc/0404018](#), <https://doi.org/10.1088/0264-9381/21/15/R01>.
- [23] J. Lewandowski, A. Okolow, H. Sahlmann, T. Thiemann, Uniqueness of diffeomorphism invariant states on holonomy-flux algebras, *Commun. Math. Phys.* 267 (2006) 703–733. [arXiv:gr-qc/0504147](#), <https://doi.org/10.1007/s00220-006-0100-7>.
- [24] A. Ashtekar, S. Fairhurst, J.L. Willis, Quantum gravity, shadow states, and quantum mechanics, *Class. Quant. Grav.* 20 (2003) 1031–1062. [arXiv:gr-qc/0207106](#), <https://doi.org/10.1088/0264-9381/20/6/302>.
- [25] H. Halvorson, Complementarity of representations in quantum mechanics, *Stud. Hist. Philos. Mod. Phys.* 35 (2004) 45–56. [arXiv:quant-ph/0110102](#), <https://doi.org/10.1016/j.shpsb.2003.08.002>.
- [26] A. Corichi, T. Vukasinac, J.A. Zapata, Polymer quantum mechanics and its continuum limit, *Phys. Rev. D* 76 (2007) 044016. [arXiv:0704.0007](#) [gr-qc], <https://doi.org/10.1103/PhysRevD.76.044016>.